



POSTER PRESENTATION

Open Access

Investigation of acute febrile illness outbreak- Asyaita and Dupti districts, Afar Region, Ethiopia, February 2011

Abyot Bekele Woyessa*, Worknesh Ayele, Abdi Ahmed, A Nega

From 17th International Symposium on HIV and Emerging Infectious Diseases (ISHEID)
Marseille, France. 23-25 May 2012

Introduction

Acute Febrile Illnesses (AFIs) due to different etiologic agents are the most common causes of morbidity and mortality in developing tropical and subtropical countries. Afar region reported unidentified AFI outbreak on 10-Aug-2011. We investigated to identify etiologic agent, risk factors and to recommend prevention and control measures.

Methods

Unmatched case control study was employed. Study subjects (57 cases and 57 controls) were obtained and interviewed. Cases were defined as any person with fever $\geq 38.3^{\circ}\text{C}$, headache, pains in joints, muscles and back, anorexia and weakness. Medical records were reviewed and suspected AFI cases were identified from 07-Aug-2011 to 11-Sep-2011 in Asyaita and Dupti districts. Active cases were searched house to house. Blood-samples, blood-serums and throat-swabs were collected and analyzed for heamoparasites, bacterial-pathogens, hemorrhagic fevers and respiratory viruses at national and CDC Kenya laboratories. Environmental scanning was performed. Odd-Ratio (OR) in 95% Confidence-Interval (CI) was calculated using Epi-Info version-3.5.1.

Results

A total of 12816 suspected AFI cases with no death were identified. Of the cases 9107(71%) were male and 3709 (29%) were female. Attack-Rate (AR) was 8.7% (11.5% in male and 5.4% in female) and 13.8% among 15-44 age-groups. On bivariate analysis factors associated with illness were living with sick family member (OR: 2.8; 95%CI: 1.3-

6.2), contact with patient (OR: 3.8; 95%CI: 1.5-9.6) and drinking deep-well water (OR: 2.7; 95% CI: 1.2-5.8). However, on multivariate analysis only having contact with patient (OR: 4.1; 95%CI: 1.5-11.3) was associated with illness. Twenty-five specimens were tested negative for malaria parasites, *Salmonella* species, *brucella* species, dengue fever, yellow fever and rift valley fever.

Conclusions

An outbreak of suspected AFI occurred in 2 districts affecting primarily males and older age. Having contact history with patient was risk factor to contract the illness. Etiologic agent and source of the outbreak was not identified yet. Large-scale investigation is recommended.

Published: 25 May 2012

doi:10.1186/1742-4690-9-S1-P46

Cite this article as: Woyessa et al.: Investigation of acute febrile illness outbreak- Asyaita and Dupti districts, Afar Region, Ethiopia, February 2011. *Retrovirology* 2012 **9**(Suppl 1):P46.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: sifanbashu@yahoo.com
Ethiopian Health and Nutrition Research Institute, Ethiopia, Addis Ababa, Ethiopia