



# Respiratory healthcare professionals' views on long-term recommendations of interventions to prevent acute respiratory illnesses after the COVID-19 pandemic

Copyright ©The authors 2022

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact [permissions@ersnet.org](mailto:permissions@ersnet.org)

Received: 28 Feb 2022  
Accepted: 20 July 2022

## To the Editor:

During the COVID-19 pandemic, healthcare authorities adopted measures to reduce the spread of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). These were lockdowns, social distancing, using facemasks and increased hand sanitising. People with chronic respiratory conditions were encouraged to comply with these measures [1]. Furthermore, it was suggested that nebulisation may facilitate the transmission of COVID-19 [2, 3]; therefore, respiratory societies recommended using inhalers where appropriate [2, 4, 5]. During 2020, significant reductions in the incidences of acute respiratory illnesses including influenza [6–8] and acute exacerbations of chronic airway diseases [9–11] were reported. These reductions may be attributed to the effects of the measures implemented for COVID-19 prevention. It is undetermined whether these measures should be extended to prevent acute respiratory illnesses in the future, when the threat of COVID-19 has lifted.

To obtain respiratory professionals' views, we conducted an online anonymous survey (<https://www.surveymonkey.com/r/T6B3HS2>) regarding recommending continued measures to reduce respiratory illnesses after the end of the COVID-19 pandemic. The survey asked respondents under which conditions they would recommend wearing a facemask (indoors, on flights and at airports, in crowded places, in places of concentrations of patients, and/or during physiotherapy). It also specified three groups of people to adopt continued hygiene measures: 1) patients with a chronic lung disease; 2) patients with conditions associated with COVID-19 adverse outcomes (cardiovascular disease, diabetes and overweight) [12]; and 3) immune-suppressed individuals. We also asked about circumstances under which using inhalers should be preferred over nebulisers to prevent aerosolisation. Respondents were asked to choose their answers from a Likert-type scale of strongly agree/agree/neither agree nor disagree/disagree/not relevant to my occupation, with given statements.

The survey was promoted in the European Respiratory Society (ERS) Newsletter and by e-mail to members of ERS assemblies (Airway Diseases, Interstitial Lung Disease, Paediatric Respiratory Diseases, Pulmonary Vascular Diseases, Respiratory Infections and Thoracic Oncology assemblies). It was available between 19 July and 31 October 2021.

Results were analysed using IBM statistics SPSS version 24. Recommendations for infection control measures were compared between the physician's seniority and physician's specialisation using the Chi-squared test or Fisher exact test, as appropriate. Variables with more than two categories were evaluated with pairwise comparisons followed by Bonferonni correction. The McNemar test was used to determine whether there were differences between the recommendations to use facemasks or hand sanitising and the recommendation for social isolation. A p-value <0.05 was considered statistically significant.

320 respiratory professionals answered the survey. 73% of them self-identified as pulmonology specialists, 24% as paediatric pulmonologists and the rest as nurse practitioners in respiratory medicine. 70% treated mostly adults, 24% mostly children and 6% both adults and children. Among physicians, 59% had



## Shareable abstract (@ERSpublications)

**Respiratory professionals support the continuing use of protective measures for respiratory patients following the #COVID19 pandemic. The optimal use of these measures should be considered in clinical guidelines and public health recommendations.** <https://bit.ly/3IVL2pQ>

**Cite this article as:** Yaacoby-Bianu K, Livnat G, Chalmers JD, *et al.* Respiratory healthcare professionals' views on long-term recommendations of interventions to prevent acute respiratory illnesses after the COVID-19 pandemic. *ERJ Open Res* 2022; 8: 00114-2022 [DOI: 10.1183/23120541.00114-2022].



experience of >15 years, 26% 5–15 years, 12% ≤5 years and 3% were in training. 62% worked in a hospital, 18% in a clinic and 20% in integrated (inpatient and outpatient) care. Professionals from 65 countries answered the survey, mostly from the UK (11%), Israel (10%), Australia (7%), Italy (6%), and Germany (5%).


Recommendations to wear facemasks was agreed/strongly agreed (combined) to by 307 (96%) in places where ill people are concentrating, such as clinic waiting rooms; 298 (95%) on flights and in airports; 292 (91%) in other crowded places (like buses, concert halls and prayer houses); 288 (90%) at concentrations of respiratory patients (physiotherapy or pulmonary rehabilitation centres); and 215 (68%) indoors. Replies from adult and paediatric pulmonologists were similar except for the recommendation to wear a mask on flights (agreed/strongly agreed: adult pulmonologists 195 (95%) out of 201 (97%); paediatric pulmonologists 59 (88%) out of 67 (p=0.009)).

We further asked which group of people to whom respondents would recommend continuing sanitation measures. This item considered a) wearing facemasks, b) disinfection of hands and c) continuing partial social isolation. Examples given were avoiding the use of public transportation and attending movie theatres or concert halls. The majority of respondents strongly agreed or agreed that they would recommend these measures to the following: 1) patients with conditions or medications that cause immunosuppression (a) 300 (94%), b) 285 (90%) and c) 238(74%); 2) patients with a chronic lung disease (a) 295 (93%), b) 275 (87%) and c) 189 (60%), respectively); 3) patients at high risk for COVID-19 such as those with a cardiovascular disease (a) 268 (85%), b) 264 (83%) and c) 184 (58%).

When we compared the recommendations for different infection control measures (defining a recommendation as a “strongly agreed” response), recommendations for wearing facemasks and hand disinfection were significantly more likely to be endorsed in comparison to continuing partial social isolation (p<0.0001 in all disease groups) (figure 1).

Regarding circumstances in which using inhalers are preferred over nebulisers, most respondents strongly agreed or agreed to recommend use of inhalers in inpatient wards (n=213, 68%), in emergency departments (n=209, 66%) and in patients’ homes with shared rooms (n=182,58%) but not in the patients’ homes with separate rooms and open windows (n=128, 41%).

The overall results did not change when comparing with different years of experience, by stratifying physician’s seniority with >15 years’ experience *versus* all others (residents, specialists with ≤5 years’ experience and specialists with 5–15 years’ experience). The only statistical difference was by looking at the “strongly agree” group *versus* all others (agree, neither agree nor disagree and disagree) where 126 (72%) of the more experienced respondents recommended wearing a facemask for physiotherapy, as opposed to 68 (57%) of the less experienced respondents (p=0.006).



**FIGURE 1** Comparison between different infection control measures in various health conditions based on respiratory healthcare professionals’ survey recommendations. The vertical axis represents the percentage of respondents who strongly agree to continuing the measure on horizontal axis after the end of COVID-19. #: p<0.0001 for comparison.

We have demonstrated that most respiratory physicians recommend adopting some of the measures that were originally taken to reduce the spread of COVID-19 after the end of the pandemic. Continued social isolation was significantly less likely to be recommended in comparison to wearing facemasks and hand disinfection ( $p < 0.0001$ ), probably due to consideration of the emotional consequences of social isolation. The COVID-19 pandemic is entering its third year at the time of writing and it is therefore possible that, while the survey asked about recommendations to prevent non-COVID-19 illness, some of the respondents' reasoning may be the expectation that COVID-19 may become endemic and the restrictions may be required to prevent COVID-19 as well as other illnesses.

There is growing literature on the impact of personal protective measures on the transmissibility of SARS-CoV-2 and other respiratory infections [13]. Transmission of viruses is reduced with physical distancing [14] and using facemasks was associated with reductions in infections [14, 15]. A meta-analysis that included eight studies found a reduction in the incidence of COVID-19 with handwashing, mask wearing and physical distancing [16]. Hand sanitising was found to be effective in reducing transmission of non-COVID-19 acute respiratory infections [13].

A recent online survey among people living with respiratory diseases by the Asthma UK–British Lung Foundation Partnership has shown good acceptability of increased handwashing (79.5%) and social distancing (68.6%), and less acceptability for wearing facemasks in indoor public places such as public transport (45.7%) [17]. These findings in patients and our findings in professionals are in contrast with pre-COVID-19 conventions. A summary of recommendations for travelling with bronchiectasis developed through a collaboration between patients and respiratory specialists did not support the use of a facemask during air travel due to inconvenience and concern of the associated stigma [18]. Our results and those of others suggest that physicians and patients have changed their views concerning this precaution, not only for the COVID-19 pandemic but thereafter.

Our survey was conducted online for a few months in the second half of 2021, so there was representation of recommendations that varied within and between countries, and over time, but we acknowledge these findings do not take into account any change in attitudes that may accompany the emergence of the more transmissible but less severe omicron variant. Another limitation is that some respondents could interpret the questions in different ways, which could influence the results. Moreover, we did not address seasonal risks and how attitudes may vary with regards to seasons, as the seasonality in acute infections changed considerably during the COVID-19 pandemic.

In summary, respiratory professionals support the continuing use of protective measures for respiratory patients following the COVID-19 pandemic. The optimal use of these measures should be considered in clinical guidelines and public health recommendations.

**Karin Yaacoby-Bianu<sup>1</sup>, Galit Livnat<sup>1,2</sup>, James D. Chalmers<sup>3</sup> and Michal Shteinberg<sup>2,4</sup>**

<sup>1</sup>Pediatric Pulmonology Unit and CF Center, Carmel Medical Center, Haifa, Israel. <sup>2</sup>B. Rappaport Faculty of Medicine, Technion – Israel Institute of Technology, Haifa, Israel. <sup>3</sup>Scottish Centre for Respiratory Research, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK. <sup>4</sup>Pulmonology Institute and CF Center, Carmel Medical Center, Haifa, Israel.

Corresponding author: Michal Shteinberg ([michalsh@technion.ac.il](mailto:michalsh@technion.ac.il))

Provenance: Submitted article, peer reviewed.

Acknowledgements: We thank the many respondents of the survey for their time and consideration. The statistical analysis of Nili Stein, MPH is gratefully acknowledged.

Conflict of interest: K. Yaacoby-Bianu reports grants from Vertex Pharmaceuticals and Gilead outside the submitted work. G. Livnat reports grants and nonfinancial support from Vertex Pharmaceuticals, and grants from Enanta Pharmaceuticals, MedImmune and DoseNTRX, outside the submitted work. J.D. Chalmers reports grants and personal fees from AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline, Novartis and Insmmed, personal fees from Chiesi, Zambon, Janssen and Grifols, and grants from Gilead Science, outside the submitted work; and is an associate editor of this journal. M. Shteinberg reports grants, personal fees and nonfinancial support from GSK, grants and personal fees from Novartis, personal fees from Boehringer Ingelheim, Zambon, AstraZeneca, Kamada,

Vertex Pharmaceuticals, Teva and Bonus Therapeutics, and nonfinancial support from Actelion and Rafa, outside the submitted work.

## References

- 1 American College of Chest Physicians. Joint Statement on Importance of Patients with Chronic Lung Disease Wearing Facial Coverings During COVID-19 Pandemic. <https://www.chestnet.org/newsroom/press-releases/2020/07/joint-statement-on-importance-of-facial-coverings> Date last accessed: 25 August 2022. Date last updated: 17 July 2020.
- 2 Centers for Disease Control and Prevention. Infection Control: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). <https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html> Date last accessed: 25 August 2022. Date last updated: 2 February 2022.
- 3 Tran K, Cimon K, Severn M, *et al*. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS ONE* 2012; 7: e35797.
- 4 Cazzola M, Ora J, Bianco A, *et al*. Guidance on nebulization during the current COVID-19 pandemic. *Respir Med* 2020; 176: 106236.
- 5 Sethi S, Barjaktarevic IZ, Tashkin DP. The use of nebulized pharmacotherapies during the COVID-19 pandemic. *Ther Adv Respir Dis* 2020; 14: 1753466620954366.
- 6 Redlberger-Fritz M, Kundi M, Aberle SW, *et al*. Significant impact of nationwide SARS-CoV-2 lockdown measures on the circulation of other respiratory virus infections in Austria. *J Clin Virol* 2021; 137: 104795.
- 7 Haapanen M, Renko M, Artama M, *et al*. The impact of the lockdown and the re-opening of schools and day cares on the epidemiology of SARS-CoV-2 and other respiratory infections in children – a nationwide register study in Finland. *EClinicalMedicine* 2021; 34: 100807.
- 8 Kuitunen I. Influenza season 2020–2021 did not begin in Finland despite the looser social restrictions during the second wave of COVID-19: a nationwide register study. *J Med Virol* 2021; 93: 5626–5629.
- 9 Sigala I, Giannakas T, Giannakoulis VG, *et al*. Effect of COVID-19-related lockdown on hospital admissions for asthma and COPD exacerbations: associations with air pollution and patient characteristics. *J Pers Med* 2021; 11: 867.
- 10 Alsallakh MA, Sivakumaran S, Kennedy S, *et al*. Impact of COVID-19 lockdown on the incidence and mortality of acute exacerbations of chronic obstructive pulmonary disease: national interrupted time series analyses for Scotland and Wales. *BMC Med* 2021; 19: 124.
- 11 Shah SA, Quint JK, Nwaru BI, *et al*. Impact of COVID-19 national lockdown on asthma exacerbations: interrupted time-series analysis of English primary care data. *Thorax* 2021; 76: 860–866.
- 12 Dorjee K, Kim H, Bonomo E, *et al*. Prevalence and predictors of death and severe disease in patients hospitalized due to COVID-19: A comprehensive systematic review and meta-analysis of 77 studies and 38,000 patients. *PLoS One* 2020; 15: e0243191.
- 13 Jefferson T, Del Mar CB, Dooley L, *et al*. Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database Syst Rev* 2020; 11: CD006207.
- 14 Chu DK, Akl EA, Duda S, *et al*. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* 2020; 395: 1973–1987.
- 15 Li Y, Liang M, Gao L, *et al*. Face masks to prevent transmission of COVID-19: a systematic review and meta-analysis. *Am J Infect Control* 2021; 49: 900–906.
- 16 Talic S, Shah S, Wild H, *et al*. Effectiveness of public health measures in reducing the incidence of covid-19, SARS-CoV-2 transmission, and covid-19 mortality: systematic review and meta-analysis. *BMJ* 2021; 375: e068302.
- 17 Hurst JR, Cumella A, Niklewicz CN, *et al*. Acceptability of hygiene, face covering and social distancing interventions to prevent exacerbations in people living with airways diseases. *Thorax* 2021; 77: 505–507.
- 18 Shteinberg M, Crossley B, Lavie T, *et al*. Recommendations for travelling with bronchiectasis: a joint ELF/EMBARC/ERN-Lung collaboration. *ERJ Open Res* 2019; 5: 00113-2019.