



Aging changes the expression of adenosine receptors, insulin-like growth factor 1 (IGF1), and hypoxia-inducible factor 1 α (HIF1 α) in hypothalamic astrocyte cultures

Camila Leite Santos^a, Larissa Daniele Bobermin^b, André Quincozes-Santos^{a,b,c,*}

^a Programa de Pós-Graduação em Ciências Biológicas: Bioquímica, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

^b Programa de Pós-Graduação em Neurociências, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

^c Departamento de Bioquímica, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

ARTICLE INFO

Keywords:

Hypothalamic astrocytes
Aging
Adenosine receptors
Hypoxia-inducible factor 1 α
Insulin-like growth factor 1

ABSTRACT

The aging process induces neurochemical alterations in different brain regions, including hypothalamus. This pivotal area of the central nervous system (CNS) is crucial for detection and integration of nutritional and hormonal signals from the periphery of the body to maintain metabolic homeostasis. Astrocytes support the CNS homeostasis, energy metabolism, and inflammatory response, as well as increasing evidence has highlighted a critical role of astrocytes in orchestrating hypothalamic functions and in gliocrine system. In this study, we aimed to investigate the age-dependent mRNA expression of adenosine receptors, the insulin-like growth factor 1 receptor (IGF1R), and the hypoxia-inducible factor 1 α (HIF1 α), in addition to the levels of IGF1 and HIF1 α in hypothalamic astrocyte cultures derived from newborn, adult, and aged rats. Our results revealed age-dependent changes in adenosine receptors, as well as a decrease in IGF1R/IGF1 and HIF1 α . Of note, adenosine receptors, IGF1, and HIF1 α are affected by inflammatory, redox, and metabolic processes, which can remodel hypothalamic properties, as observed in aging brain, reinforcing the role of hypothalamic astrocytes as targets for understanding the onset and/or progression of age-related diseases.

1. Introduction

The aging process is characterized by a progressive loss of physiological integrity, leading to increased risk for several diseases [22]. Aging can also markedly affect both the periphery of the body and the hypothalamus, whose neural networks have emerged as key orchestrators of systemic metabolism, food intake, and body weight [13,17]. Hypothalamic astrocytes are versatile cells that can regulate energy metabolism due to its capacity to detect peripheral signals that inform about energy status. They are also part of the gliocrine system, because they are secretory cells and targets of hormone action, such as leptin and insulin [13,24,30,40].

In line with this, the brain, in particular hypothalamus, plays a role in regulating glucose homeostasis. Insulin-like growth factor 1

* Corresponding author at: Laboratório de Neurotoxicidade e Glioproteção (LABGLIO), Departamento de Bioquímica, Programa de Pós-Graduação em Ciências Biológicas: Bioquímica, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Rua Ramiro Barcelos, 2600 – Anexo, Bairro Santa Cecília, 90035-003 Porto Alegre, RS, Brazil.

E-mail address: andrequincozes@ufrgs.br (A. Quincozes-Santos).

<https://doi.org/10.1016/j.nbas.2023.100104>

Received 29 May 2023; Received in revised form 30 November 2023; Accepted 18 December 2023

2589-9589/© 2023 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

(IGF1) and its receptor (IGF1R) are also crucial to maintain neural functions, including development, neurogenesis, and plasticity [1,28]. IGF1 might also induce hypoxia-inducible factor 1 (HIF1), an essential transcription factor for the regulation of oxygen homeostasis, which more recently was associated with inflammation [3,10]. Accumulated evidence indicates that inflammatory process might have a key role in the obesity, as well as in the onset and progression of age-related diseases [7,26]. In this regard, the adenosinergic system can modulate astroglial responses, including those related to inflammation, and changes in the expression of adenosine receptors are associated with aging [4,15].

Our previous publications showed that aging significantly affect the hypothalamic functionality, due to changes in metabolic, inflammatory response, and leptin signaling [18,32,31]; therefore, here, we aimed to expand the knowledge of how aging modulates the expression of adenosine receptors, IGF1, and HIF1 α . Using a protocol of hypothalamic astrocyte cultures derived from newborn (2 days old), adult (90 days old), and aged (180 days old) Wistar rats, we concluded that aging changes these parameters, which are related to etiological factors of diabetes, obesity, and neurodegeneration.

2. Materials and methods

2.1. Reagents

Dulbecco's modified Eagle's medium/F12 (DMEM/F12) and other materials for cell cultures were purchased from Gibco/Invitrogen (USA). TRIzol, High Capacity complementary DNA (cDNA) Reverse Transcription Kit and TaqMan real-time RT-PCR system were purchased from Applied Biosystems (USA). ELISA kit for IGF1 was purchased from Invitrogen (USA) and for HIF1 α from Elabscience (China). Poly-L-lysine and TRI Reagent were purchased from Sigma-Aldrich (USA). All other chemicals were from common commercial suppliers.

2.2. Animals

Male Wistar rats were obtained from the breeding colony of Department of Biochemistry (UFRGS, Porto Alegre, Brazil) and maintained under a controlled environment (12-h light/12-h dark cycle; 22 ± 1 °C; *ad libitum* access to water and diet – regular laboratory chow). All animal experiments were performed in accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals. The Federal University of Rio Grande do Sul Animal Care and Use Committee approved the use of Laboratory animals for this project (process number 29420).

2.3. Hypothalamic primary astrocyte cultures

The protocol was previously described by Santos et al [32]. Wistar rats at 2, 90, and 180 days old were euthanized by decapitation. They subsequently had their hypothalamus dissected, and the meninges removed. A pool of 3 hypothalamus was used to perform a single astrocyte culture (6 independent cultures were used, each culture was derived from 3 hypothalamus). The sample size was 18 Wistar rats per age/group. The tissue was enzymatically digested in Hank's balanced salt solution (HBSS) containing 0.05 % trypsin at 37 °C for 7 min. The tissue was then mechanically dissociated for 7 min and centrifuged at 100 g for 5 min. The pellet was resuspended in HBSS and again mechanically dissociated until complete homogenization, and then centrifuged at 100 g for 5 min. Then, cells were resuspended in DMEM/F12, supplemented with 10 % fetal bovine serum (FBS), 15 mM HEPES, 14.3 mM NaHCO₃, 1 % Fungizone®, and 0.04 % gentamicin. Cells were seeded (approximately $2 - 4 \times 10^5$ cells/cm²) into 6-well plates pre-coated with poly-L-lysine and cultured at 37 °C in a 5 % CO₂ incubator. After 24 h, the culture medium was exchanged; during the first week, the medium was replaced once every 2 days, and from the second week on, once every 4 days. From the second week on, the astrocytes received medium supplemented with 20 % FBS until they reached confluence (at approximately the fourth week). No dibutyryl-cAMP was added to the culture medium. To determine whether the culture contained microglia or neurons after reaching confluence, we used anti- β -tubulin III, anti-NeuN, and anti-CD11.

2.4. RNA extraction and Quantitative RT-PCR

Total RNA was isolated from newborn, adult, and aged astrocyte cultures using TRIzol reagent. The concentration and purity of the RNA were determined spectrophotometrically at a ratio of 260:280. Then, 1 μ g of total RNA was reverse transcribed using Applied Biosystems High-Capacity complementary DNA (cDNA) Reverse Transcription Kit in a 20 μ L reaction according to the manufacturer's instructions. The messenger RNA (mRNA) encoding adenosine receptors A₁ (#Rn00567668_m1), A_{2A} (#Rn00583935_m1), A_{2B} (#Rn00567697_m1), and A₃ (#Rn00563680_m1), IGF1R (#Rn00583837_m1), HIF1 α (#Rn01472831_m1), and β -actin (#Rn00667869_m1) were quantified using the TaqMan real-time RT-PCR system using inventory primers and probes purchased from Applied Biosystems. Quantitative RT-PCR was performed using the Applied Biosystems 7500 Fast system. The mRNA levels were expressed relative to the levels of newborn (1 day old) cultures using the $2^{-\Delta\Delta Ct}$ method [20].

2.5. IGF and HIF1 α measurement

IGF1 and HIF1 α levels were measured in the extracellular medium and intracellular content, respectively, using commercial ELISA kits following manufacturer's instructions with some modifications. IGF1: Invitrogen (#ERIGF1), analytical sensitivity: 30.0 pg/ml,

detection range: 30.72 – 7500 pg/ml; HIF1 α : Elabscience (#E-EL-R0513), analytical sensitivity: 4.7 pg/ml, detection range: 7.81 – 500 pg/ml. The results are expressed in pg/mL for IGF1 and in pg/mg of protein for HIF1 α .

2.6. Statistical analysis

The results are presented as mean \pm standard error of the mean (SEM). To ensure adequate power for statistical analysis at least six independent experiments were performed in triplicate. All attempts at replication were successful. For each experiment, replicates are noted in the figure legends. The normal distribution was confirmed by Shapiro-Wilk test and variance homogeneity was tested using Bartlett's test. Considering the normality and homogeneity of variance, differences among groups were statistically analyzed using one-way analysis of variance (ANOVA), followed by Tukey's test. All analyses were performed using the GraphPad Prism software version 9 (GraphPad Software). Adjusted P values of post hoc analysis are described in the Results section, only for significant results. P values < 0.05 were considered significant (*a* refers to statistically significant differences from newborn rats – 2 days old; *b* refers to statistically significant differences from 90 days old rats). All graphs were also performed by using GraphPad Prism 9.

3. Results

First, we examined the expression of adenosine receptors. A significant age-dependent decrease in mRNA levels of A₁ was observed ($P = 0.0013$ (newborn vs. 90 days old rats); $P < 0.0001$ (newborn vs. 180 days old rats and 90 days old rats vs. 180 days old rats), Fig. 1A). However, the mRNA levels of A_{2A} presented an age-dependent increase ($P = 0.0003$ (newborn vs. 90 days old rats); $P < 0.0001$ (newborn vs. 180 days old rats and 90 days old rats vs. 180 days old rats), Fig. 1B). The mRNA of A_{2B} did not change in cultured astrocytes and A₃ decreased only in astrocytes from 180 days old ($P < 0.0001$ (newborn vs. 180 days old rats and 90 days old rats vs. 180 days old rats); Fig. 1C and D, respectively).

We also demonstrated an age-dependent decrease in the mRNA expression of IGF1R ($P = 0.0001$ (newborn vs. 90 days old rats); $P < 0.0001$ (newborn vs. 180 days old rats); $P = 0.0009$ (90 days old rats vs. 180 days old rats) and in the extracellular levels of IGF1 ($P < 0.0001$ (newborn vs. 90 days old rats, newborn vs. 180 days old rats and 90 days old rats vs. 180 days old rats), Fig. 2A and B, respectively). The same response was observed for the mRNA expression of HIF1 α ($P < 0.0001$ (newborn vs. 90 days old rats and newborn vs. 180 days old rats); $P = 0.0005$ (90 days old rats vs. 180 days old rats), Fig. 2C), as well as for its intracellular levels ($P < 0.0001$ (newborn vs. 90 days old rats, newborn vs. 180 days old rats and 90 days old rats vs. 180 days old rats), Fig. 2D).




Fig. 1. Age-related changes of adenosine receptors in hypothalamic astrocyte cultures. RT-PCR quantification of mRNA expression of A₁ (A), A_{2A} (B), A_{2B} (C), and A₃ (D). The data represent the mean \pm SEM of triplicate samples from 6 independent cultures (each culture derived from 3 hypothalamus) and statically analyzed by one-way ANOVA followed by Tukey's test. *a* refers to statistically significant differences from newborn rats (2 days old); *b* refers to statistically significant differences from 90 days old rats.




Fig. 2. Aging decreases expression and levels of IGF1 and HIF1 α in hypothalamic astrocyte cultures. RT-PCR quantification of mRNA expression of IGF1R (A) and HIF1 α (C), extracellular levels of IGF1 (B) and intracellular content of HIF1 α (D). For RT-PCR and ELISA measurements, the data represent the mean \pm SEM of triplicate samples from 6 independent cultures (each culture derived from 3 hypothalamus). The data were statically analyzed by one-way ANOVA followed by Tukey's test. *a* refers to statistically significant differences from newborn rats (2 days old); *b* refers to statistically significant differences from 90 days old rats.

4. Discussion

Astrocytes are dynamic cells that maintain brain homeostasis and regulate hypothalamic-feeding circuits [2,38]. These cells respond to metabolic signals interacting with neurons, and this communication can alter neural survival and development, reinforcing the role of hypothalamic astrocytes as target for understanding the pathomechanisms of the communication between periphery and brain, including during the aging process [43]. Our previous studies showed that aging promotes changes in neurochemical properties of hypothalamic astrocytes and, here, we described important age-dependent alterations in the expression of adenosine receptors, IGF1, and HIF1 α , which could be associated with neurodegeneration/neurodegenerative diseases.

Adenosinergic system can control neuroendocrine signals and participates in the mechanisms of blood pressure variability, sleep disorders, obesity, inflammation, and aging [5,6,8,11,33,35,41]. Hypothalamic astrocytes express adenosine receptors (A₁, A_{2A}, A_{2B}, and A₃), which can orchestrate different cellular responses. Our findings showed that A₁ and A₃ receptors decreased with aging, potentially indicating a dysfunction in inflammatory and metabolic states, since their downregulation promote the production of pro-inflammatory mediators and changes in energy balance [15,16,42]. Moreover, A₁ and A₃ adenosine receptors may be also related to A_{2A} signaling, which presented an age-dependent increase in astrocytes. This receptor is strongly associated with neuroinflammation, obesity, sleep disorder, and cognitive decline [9,19]. Therefore, our results may suggest alterations of the adenosine receptor-mediated signaling and functions in hypothalamic astrocytes during aging. In addition, we recently demonstrated that chronic resveratrol treatment mediates glioprotection in aged astrocytes regulating adenosine receptors [37].

The hormone IGF1 and its receptor, IGF1R, are critical for hypothalamic function, because they are involved in the regulation of synaptic activity and attenuation of neuroinflammation [45]. During aging, the concentration of IGF1 and the expression of IGF1R decrease in different brain areas, and their deficiencies, as well as the defective insulin signaling, have been implicated in the cognitive dysfunction and Alzheimer's disease (Lyra E [23,36]. Although in the Alzheimer's disease the role of IGF1/IGF1R is associated with decline in hippocampus-dependent learning and memory [21], our findings point out hypothalamic astrocytes as key players in the pathophysiology of brain metabolism, with potential impact in neurometabolic disorders. Furthermore, defective IGF1 gliosignaling can induce mitochondrial dysfunction and production of reactive oxygen species [21], which actively participates in the pathomechanisms of neurodegenerative diseases. Our findings are also in accordance with a recent publication that indicated an age-dependent reduction in IGF1R that may represent a link between the progression of synaptic dysfunction and inflammation in paraventricular hypothalamic nucleus in an experimental model of hypertension, a common condition observed during aging [29]. The IGF signaling can also modulate the activity of HIF1 α [3,10]. In this sense, astrocytes from mice with insulin signaling deficiency show higher levels of reactive oxidant species, leading to stimulation of HIF1 α , with consequent impact in neurochemical functions [12].

HIF1 α is a transcriptional factor that acts as a master regulator of systemic and cellular homeostatic response to hypoxia, consequently impacting on oxygen delivery and metabolic adaptation [34]. In the CNS, HIF 1 and 2 are critically involved in the regulation

of neurogenesis, nerve cell differentiation, and neuronal apoptosis [27]. Particularly in the hypothalamus, HIF1 α also controls the expression of pro-opiomelanocortin (POMC) in neurons, with significant impact on the energy homeostasis of the body [39,44]. Here, we demonstrated the age-dependent decrease in the expression and intracellular levels of HIF1 α , which can be associated with metabolic, oxidative, and inflammatory functions, which, in turn, are linked to pathomechanisms of neurometabolic diseases [14]. HIF1 α Moreover, HIF1 α is interconnected with adenosinergic system [13,25] and other signaling pathways, including nuclear factor kappa B (NF κ B) and erythroid-derived 2 like 2 (Nrf2), which we recently demonstrated that changed with age in hypothalamic astrocytes [18,32]. It is noteworthy that regulation of the expression and activity of HIF1 may represent a promising neuroprotective tool during ischemic injury [27]. Considering that hypoxia and ischemia are also common diseases in elderly population, the reduction of HIF1 α during aging corroborates this hypothesis, particularly due the relevance of astrocytes in the signaling mechanisms related to adaptive functions.

5. Conclusions

In summary, we demonstrated that adenosine receptors, IGF1, and HIF1 α changed during aging, impacting the hypothalamic astrocyte functionality, and consequently, the development and progression of age-dependent metabolic brain dysfunctions. Our findings reinforce/complement the knowledge from our previous report, in which we found that cultured hypothalamic astrocytes undergo neurochemical remodeling with age [32]. Therefore, hypothalamic astrocytes may emerge as targets for understanding the brain aging, particularly about neurometabolic conditions.

Funding

This study was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS), Universidade Federal do Rio Grande do Sul and Instituto Nacional de Ciência e Tecnologia para Excitotoxicidade e Neuroproteção (INCTEN/CNPq).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This study was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS), Universidade Federal do Rio Grande do Sul and Instituto Nacional de Ciência e Tecnologia para Excitotoxicidade e Neuroproteção (INCTEN/CNPq).

References

- [1] Åberg ND, Bryve KG, Isgaard J. Aspects of Growth Hormone and Insulin-Like Growth Factor-I Related to Neuroprotection, Regeneration, and Functional Plasticity in the Adult Brain. *Scientific World J* 2006;6:53–80. <https://doi.org/10.1100/tsw.2006.22>.
- [2] Argente-Arízón P, Guerra-Cantera S, Garcia-Segura LM, Argente J, Chowen JA. Glial cells and energy balance. *J Mol Endocrinol* 2017;58:R59–71. <https://doi.org/10.1530/JME-16-0182>.
- [3] Balamurugan K. HIF-1 at the crossroads of hypoxia, inflammation, and cancer: Hypoxia, Inflammation, and Cancer. *Int J Cancer* 2016;138:1058–66. <https://doi.org/10.1002/ijc.29519>.
- [4] Bobermin, L.D., Roppa, R.H.A., Quincozes-Santos, A., 2019. Adenosine receptors as a new target for resveratrol-mediated glioprotection. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* 1865, 634–647. doi: 10.1016/j.bbadis.2019.01.004.
- [5] Borea PA, Gessi S, Merighi S, Varani K. Adenosine as a Multi-Signalling Guardian Angel in Human Diseases: When, Where and How Does it Exert its Protective Effects? *Trends Pharmacol Sci* 2016;37:419–34. <https://doi.org/10.1016/j.tips.2016.02.006>.
- [6] Burnstock G, Gentile D. The involvement of purinergic signalling in obesity. *Purinergic Signalling* 2018;14:97–108. <https://doi.org/10.1007/s11302-018-9605-8>.
- [7] Cai D, Khor S. “Hypothalamic Microinflammation” Paradigm in Aging and Metabolic Diseases. *Cell Metab* 2019;30:19–35. <https://doi.org/10.1016/j.cmet.2019.05.021>.
- [8] Castillo CA, Albasanz JL, León D, Jordán J, Pallàs M, Camins A, et al. Age-related expression of adenosine receptors in brain from the senescence-accelerated mouse. *Exp Gerontol* 2009;44:453–61. <https://doi.org/10.1016/j.exger.2009.04.006>.
- [9] Chen W, Ostrowski RP, Obenaus A, Zhang JH. Prodeath or prosurvival: Two facets of hypoxia inducible factor-1 in perinatal brain injury. *Exp Neurol* 2009;216:7–15. <https://doi.org/10.1016/j.expneurol.2008.10.016>.
- [10] Corcoran SE, O'Neill LAJ. HIF1 α and metabolic reprogramming in inflammation. *J Clin Investig* 2016;126:3699–707. <https://doi.org/10.1172/JCI84431>.
- [11] Cunha RA. How does adenosine control neuronal dysfunction and neurodegeneration? *J Neurochem* 2016;139:1019–55. <https://doi.org/10.1111/jnc.13724>.
- [12] Fernandez, A.M., Martínez-Rachadell, L., Navarrete, M., Pose-Utrilla, J., Davila, J.C., Pignatelli, J., Diaz-Pacheco, S., Guerra-Cantera, S., Viedma-Moreno, E., Palenzuela, R., Ruiz De Martin Esteban, S., Mostany, R., Garcia-Caceres, C., Tschöp, M., Iglesias, T., De Ceballos, M.L., Gutierrez, A., Torres Aleman, I., 2022. Insulin regulates neurovascular coupling through astrocytes. *Proc Natl Acad Sci USA* 119, e2204527119. doi: 10.1073/pnas.2204527119.
- [13] García-Cáceres C, Balland E, Prevot V, Luquet S, Woods SC, Koch M, et al. Role of astrocytes, microglia, and tanyocytes in brain control of systemic metabolism. *Nat Neurosci* 2019;22:7–14. <https://doi.org/10.1038/s41593-018-0286-y>.
- [14] Gaspar JM, Velloso LA. Hypoxia Inducible Factor as a Central Regulator of Metabolism – Implications for the Development of Obesity. *Front Neurosci* 2018;12:813. <https://doi.org/10.3389/fnins.2018.00813>.

- [15] Gessi S, Merighi S, Stefanelli A, Fazzi D, Varani K, Borea PA. A1 and A3 adenosine receptors inhibit LPS-induced hypoxia-inducible factor-1 accumulation in murine astrocytes. *Pharmacol Res* 2013;76:157–70. <https://doi.org/10.1016/j.phrs.2013.08.002>.
- [16] Haselkorn ML, Shellington DK, Jackson EK, Vagni VA, Janesko-Feldman K, Dubey RK, et al. Adenosine A₁ Receptor Activation as a Brake on the Microglial Response after Experimental Traumatic Brain Injury in Mice. *J Neurotrauma* 2010;27:901–10. <https://doi.org/10.1089/neu.2009.1075>.
- [17] Kim K-S, Seeley RJ, Sandoval DA. Signalling from the periphery to the brain that regulates energy homeostasis. *Nat Rev Neurosci* 2018;19:185–96. <https://doi.org/10.1038/nrn.2018.8>.
- [18] Leite Santos C, Vizuete AFK, Becker FW, Thomaz NK, Bobermin LD, Goncalves CA, et al. Age-dependent effects of resveratrol in hypothalamic astrocyte cultures. *Neuroreport* 2023;34:419–25. <https://doi.org/10.1097/WNR.0000000000001906>.
- [19] Li R, Wang Y-Q, Liu W-Y, Zhang M-Q, Li L, Cherasse Y, et al. Activation of adenosine A_{2A} receptors in the olfactory tubercle promotes sleep in rodents. *Neuropharmacology* 2020;168:107923. <https://doi.org/10.1016/j.neuropharm.2019.107923>.
- [20] Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2^{-ΔΔCT} Method. *Methods* 2001;25:402–8. <https://doi.org/10.1006/meth.2001.1262>.
- [21] Logan S, Pharaoh GA, Marlin MC, Masser DR, Matsuzaki S, Wronowski B, et al. Insulin-like growth factor receptor signaling regulates working memory, mitochondrial metabolism, and amyloid-β uptake in astrocytes. *Molecular Metabolism* 2018;9:141–55. <https://doi.org/10.1016/j.molmet.2018.01.013>.
- [22] López-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The Hallmarks of Aging. *Cell* 2013;153:1194–217. <https://doi.org/10.1016/j.cell.2013.05.039>.
- [23] Lyra E Silva, N.D.M., Gonçalves, R.A., Boehnke, S.E., Forny-Germano, L., Munoz, D.P., De Felice, F.G., 2019. Understanding the link between insulin resistance and Alzheimer's disease: Insights from animal models. *Experimental Neurology* 316, 1–11. doi: 10.1016/j.expneurol.2019.03.016.
- [24] MacDonald AJ, Robb JL, Morrissey NA, Beall C, Ellacott KJ. Astrocytes in neuroendocrine systems: An overview. *J Neuroendocrinol* 2019;31:e12726.
- [25] Merighi S, Borea PA, Stefanelli A, Bencivenni S, Castillo CA, Varani K, et al. A_{2a} and a_{2b} adenosine receptors affect HIF-1α signaling in activated primary microglial cells: Adenosine Receptors in Microglial Cells. *Glia* 2015;63:1933–52. <https://doi.org/10.1002/glia.22861>.
- [26] Miller AA, Spencer SJ. Obesity and neuroinflammation: A pathway to cognitive impairment. *Brain Behav Immun* 2014;42:10–21. <https://doi.org/10.1016/j.bbi.2014.04.001>.
- [27] Mitroshina EV, Savyuk MO, Ponimaskin E, Vedunova MV. Hypoxia-Inducible Factor (HIF) in Ischemic Stroke and Neurodegenerative Disease. *Front Cell Dev Biol* 2021;9:703084. <https://doi.org/10.3389/fcell.2021.703084>.
- [28] Nieto-Estévez V, Ouelati-Morales CO, Li L, Pickel J, Morales AV, Vicario-Abejón C. Brain Insulin-Like Growth Factor-I Directs the Transition from Stem Cells to Mature Neurons During Postnatal/Adult Hippocampal Neurogenesis. *Stem Cells* 2016;34:2194–209. <https://doi.org/10.1002/stem.2397>.
- [29] Ogundele OM, Lee CC, Francis J. Age-dependent alterations to paraventricular nucleus insulin-like growth factor 1 receptor as a possible link between sympathoexcitation and inflammation. *J Neurochem* 2016;139:706–21. <https://doi.org/10.1111/jnc.13842>.
- [30] Ortiz-Rodríguez A, Arevalo M-A. The Contribution of Astrocyte Autophagy to Systemic Metabolism. *IJMS* 2020;21:2479. <https://doi.org/10.3390/ijms21072479>.
- [31] Santos CL, Bobermin LD, Souza DO, Quincozes-Santos A. Leptin stimulates the release of pro-inflammatory cytokines in hypothalamic astrocyte cultures from adult and aged rats. *Metab Brain Dis* 2018;33:2059–63. <https://doi.org/10.1007/s11011-018-0311-6>.
- [32] Santos CL, Roppa PHA, Truccolo P, Fontella FU, Souza DO, Bobermin LD, et al. Age-Dependent Neurochemical Remodeling of Hypothalamic Astrocytes. *Mol Neurobiol* 2018;55:5565–79. <https://doi.org/10.1007/s12035-017-0786-x>.
- [33] Schindler CW, Karcz-Kubicha M, Thorndike EB, Müller CE, Tella SR, Ferré S, et al. Role of central and peripheral adenosine receptors in the cardiovascular responses to intraperitoneal injections of adenosine A₁ and A_{2A} subtype receptor agonists: Cardiovascular effects of adenosine agonists. *Br J Pharmacol* 2005;144:642–50. <https://doi.org/10.1038/sj.bjp.0706043>.
- [34] Semenza GL. Oxygen Sensing, Hypoxia-Inducible Factors, and Disease Pathophysiology. *Annu Rev Pathol Mech Dis* 2014;9:47–71. <https://doi.org/10.1146/annurev-pathol-012513-104720>.
- [35] Shen F-M, Su D-F. The Effect of Adenosine on Blood Pressure Variability in Sinoaortic Denervated Rats is Mediated by Adenosine A_{2A}-Receptor. *J Cardiovasc Pharmacol* 2000;36:681–6. <https://doi.org/10.1097/00005344-200011000-00019>.
- [36] Sonntag WE, Ramsey M, Carter CS. Growth hormone and insulin-like growth factor-1 (IGF-1) and their influence on cognitive aging. *Ageing Res Rev* 2005;4:195–212. <https://doi.org/10.1016/j.arr.2005.02.001>.
- [37] Sovrani V, Bobermin LD, Santos CL, Brondani M, Gonçalves C-A, Leipnitz G, et al. Effects of long-term resveratrol treatment in hypothalamic astrocyte cultures from aged rats. *Mol Cell Biochem* 2022. <https://doi.org/10.1007/s11010-022-04585-z>.
- [38] Teschemacher AG, Gourine AV, Kasparov S. A Role for Astrocytes in Sensing the Brain Microenvironment and Neuro-Metabolic Integration. *Neurochem Res* 2015;40:2386–93. <https://doi.org/10.1007/s11064-015-1562-9>.
- [39] Varela L, Suyama S, Huang Y, Shanabrough M, Tschöp MH, Gao X-B, et al. Endothelial HIF-1α Enables Hypothalamic Glucose Uptake to Drive POMC Neurons. *Diabetes* 2017;66:1511–20. <https://doi.org/10.2337/db16-1106>.
- [40] Verkhatsky, A., Ho, M.S., Zorec, R., Parpura, V. (Eds.), 2019. Neuroglia in Neurodegenerative Diseases, Advances in Experimental Medicine and Biology. Springer Singapore, Singapore. doi: 10.1007/978-981-13-9913-8.
- [41] Willems L, Ashton K, Headrick J. Adenosine-mediated cardioprotection in the aging myocardium. *Cardiovasc Res* 2005;66:245–55. <https://doi.org/10.1016/j.cardiores.2004.11.008>.
- [42] Wu L, Meng J, Shen Q, Zhang Y, Pan S, Chen Z, et al. Caffeine inhibits hypothalamic A1R to excite oxytocin neuron and ameliorate dietary obesity in mice. *Nat Commun* 2017;8:15904. <https://doi.org/10.1038/ncomms15904>.
- [43] Yan J, Zhang H, Yin Y, Li J, Tang Y, Purkayastha S, et al. Obesity- and aging-induced excess of central transforming growth factor-β potentiates diabetic development via an RNA stress response. *Nat Med* 2014;20:1001–8. <https://doi.org/10.1038/nm.3616>.
- [44] Zhang Z, Yan J, Chang Y, ShiDu Yan S, Shi H. Hypoxia Inducible Factor-1 as a Target for Neurodegenerative Diseases. *CMC* 2011;18:4335–43. <https://doi.org/10.2174/092986711797200426>.
- [45] Zhao H, Zhao X, Cao X, Wu G. Age-Dependent Neuroimmune Modulation of IGF-1R in the Traumatic Mice. *Immun Ageing* 2012;9:12. <https://doi.org/10.1186/1742-4933-9-12>.