

# Adipose Tissue Transfer in Dynamic Definition Liposculpture Part II. The Lower Limb: Gastrocnemius, Vastus Medialis, Vastus Lateralis, and Rectus Femoris Muscles

Alfredo E. Hoyos, MD\*†‡  
 Matt Stefanelli, MD§¶  
 Mauricio E. Perez, MD||  
 Mauricio Padilla, MD\*\*††  
 Rodrigo Dominguez-Millan,  
 MD‡‡§§

**Background:** One big challenge of body contouring surgery is the liposculpture of the lower limbs, probably because of the imperative symmetry and the risk of contour irregularities. We are reporting our experience in fat grafting of the thighs and calves for men and women undergoing dynamic definition liposculpture.

**Methods:** We did cadaveric dissections of the vastus lateralis, vastus medialis, rectus femoris, and gastrocnemius muscles and identified each of their primary pedicles. We also performed fat grafting of these muscles in the contralateral virgin cadaveric specimen. We searched our records for patients who underwent fat grafting of the lower extremity in addition to dynamic definition liposculpture, from January 2016 to May 2022 at a single center in Bogotá, Colombia.

**Results:** Seventy-three consecutive patients met the inclusion criteria (26 men and 47 women). We grafted 102 gastrocnemius muscles, 86 vastus medialis muscles, 98 vastus lateralis muscles, and 22 rectus femoris muscles. Mean age was 34 and 41 years for men and women, respectively. Range of the fat graft volume was 50–200 mL. No complications were recorded related to fat grafting. Almost all patients were satisfied with the procedure (89%). Follow-up period ranged from 2 to 36 months.

**Conclusions:** Fat grafting of the lower limb muscles should be considered an alternative operative technique to enhance volume and athletic appearance of this body segment. Based on cadaveric dissections and clinical evidence, our technique is considered reliable and reproducible with remarkable outcomes and a very low complication rate. (*Plast Reconstr Surg Glob Open* 2023; 11:e4765; doi: 10.1097/GOX.0000000000004765; Published online 27 January 2023.)

From the \*Department of Plastic Surgery, Total Definer Research Group, Bogota, Colombia; †Member of the Colombian Society of Plastic, Aesthetic, Maxillofacial, and Hand Surgery; ‡Member of the American Society of Plastic Surgeons (ASPS), Bogota, Colombia; §Member of the French College of Plastic Surgery (CFCPRE), Paris, France; ¶Department of Plastic Surgery, The European Board of Plastic Reconstructive and Aesthetic Surgery (EBOPRAS), Paris, France; ||Department of Plastic Surgery, Total Definer Research Group, Rochester, Minn.; \*\*Member of the Mexican Association of Aesthetic and Reconstructive Plastic Surgery (AMCPER), Ciudad de México, Mexico; ††Total Definer Research Group, Ciudad de México, Mexico; ‡‡Department of Plastic Surgery, Corporea “Health & Aesthetics”; and §§Department of Plastic Surgery, Plenus Medical Group, Mexico City, Mexico.

Received for publication July 6, 2022; accepted September 20, 2022.

Presented at the 53rd Annual Baker & Gordon Educational Symposium, February 7–9, 2019, Miami, Florida, and at the Aesthetics Meeting, ASAPS, April 20–24, 2022, San Diego, California.

Copyright © 2023 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/GOX.0000000000004765

## INTRODUCTION

Concepts of beauty have been linked to humanity since dawn of civilizations; in particular, well-developed limbs exemplify strength, athleticism, youthfulness, and beauty in both men and women. As goals for symmetry are set, liposculpture of the lower limbs is a challenging body contouring surgery as it carries a high risk of contour irregularities, since the shape of the thighs and legs depends on the balance and proper development of muscles, bones, and subcutaneous fat.<sup>1,2</sup>

**Disclosure:** Dr. Hoyos was an unpaid consultant and speaker for the product development team of Sound Surgical Technologies (SST) system and Cannulas (now: VASER 2018 Solta Medical–Bausch Health Companies Inc.) up to May 2013. He receives royalties for the liposuction kits named after him. The other authors have no financial interest to declare.

Related Digital Media are available in the full-text version of the article on [www.PRSGlobalOpen.com](http://www.PRSGlobalOpen.com).

Thigh lift and liposuction are the most widely used techniques to enhance the aesthetic appearance of the leg and to correct deformities caused by congenital defects, trauma, disease, or infections,<sup>1,2</sup> although aging and/or massive weight loss patients do get the greatest benefit.<sup>3,4</sup> The use of silicone implants for calf and thigh volumizing has also been described, although case series are limited and have reported a considerable number of complications.<sup>5-7</sup> Although they may improve the overall appearance of the leg, their use has several potential complications, including hyperpigmentation, seromas, extrusion, infection, capsular contracture, secondary surgery, and compartment syndrome.<sup>1,8-10</sup> Comparatively, fat grafting is likely to volumize the leg with a reduced likelihood of complications due to the autologous nature of the tissue transfer. We did an anatomical review and cadaveric dissections of the quadriceps and the gastrocnemius muscles before and after our fat grafting technique to provide safety recommendations. We are also reporting our surgical experience with intramuscular (IM) fat grafting of such muscles.

#### Artistic Anatomy

Thighs have three muscle compartments (anterior, posterior, and medial), similar to the legs (anterior, posterior, and lateral).<sup>11</sup> All of the muscles from these compartments get their blood supply from two main arteries: the femoral artery and the tibial artery.

The vastus lateralis is one of the four muscles of the quadriceps and occupies two-thirds of the lateral thigh surface. Its size is near 10 cm wide and 26 cm long, extending from the proximal femur to the patella, running between the vastus intermedius and biceps femoris muscles, and beneath the tensor fascia latae muscle.<sup>12,13</sup> The descending branch of the lateral circumflex femoral artery corresponds to its dominant pedicle<sup>14,15</sup> (Fig. 1), which also provides the blood supply for the rectus femoris muscle (Table 1). The vastus medialis muscle is located at the medial aspect of the thigh, deep to the sartorius and medial to the rectus femoris.<sup>16</sup> Its length and width are 15 cm and 8 cm in average, respectively. Surface anatomy of the posterior thigh is mostly configured by the biceps femoris on the lateral region and the hamstrings on the medial one. Both these muscles receive their innervation from muscular branches of the sciatic nerve and blood supply from perforating branches from the deep femoral artery (Fig. 2).

The gastrocnemius is the most superficial muscle of the posterior upper third of the leg; its size is about 20 cm long and 8 cm width,<sup>15</sup> composed by two muscular bellies, from which the medial one is a bit longer and extends more inferiorly (Fig. 3).

The venous system of the lower limb has a special configuration and comprises three different plexuses that are connected between each other: (1) the subcutaneous venous network (between the deep muscular fascia and the skin); (2) the deep venous network (main veins); and (3) the perforating veins, which are short vessels that penetrate the deep fascia and connect the superficial and deep systems.<sup>13-17</sup>

#### Takeaways

**Question:** Is there any type of surgery to improve both the leg contour and its definition for men and women?


**Findings:** We did cadaveric dissections to identify the relevant anatomic structures to achieve a safe fat grafting technique of the lower limb muscles. Furthermore, we completed over 140 lipoinjections of different muscles with great results and a very low rate of complications.

**Meaning:** Intramuscular and subcutaneous fat grafting of the lower limb can be considered a great alternative for both muscular enhancement and definition.

#### MATERIALS AND METHODS

We performed cadaveric dissections of the lower limb in an attempt to support the safe and reliable approach of IM fat grafting of the muscles from the thigh and the leg. The vastus medialis, the vastus lateralis, the rectus femoris, the gastrocnemius, and the soleus muscles were all dissected following the anatomic models to locate the main pedicles and their distribution within the muscle. A scalpel with 15/20 blades, tissue scissors, and tissue forceps were used to separate the anatomic layers, vascular retractors, and markers that were used to denote the main pedicles and vessels. First, we located the pedicle, and then did a lipoinjection test on the contralateral virgin muscle with methylene blue to identify its proper placement at the IM layer for the thigh muscles and the gastrocnemius muscle. (See Video 1 [online], which displays cadaveric dissection of the thigh. Beware of the “danger triangle” where most vital structures of the thigh are located. Fat grafting of the vastus medialis and vastus lateralis muscles shows the superficial location of the graft, which is an actual premise for a safe approach.) (See Video 2 [online], which displays cadaveric dissection and fat grafting of the gastrocnemius muscle. The lateral head and the medial head have different access points to avoid the popliteal fossa, where critical neurovascular structures are present. The distal access [Achilles region] may be too close to neurovascular structures; hence, it should be avoided.)


The senior author has incorporated the fat grafting technique for the rectus femoris, vastus medialis, vastus lateralis, and gastrocnemius muscles into dynamic definition liposculpture (HD2) since mid 2015; hence, we conducted a retrospective review of the medical records from January 2016 to May 2022 at a single center (Dhara Clinic) in Bogotá, Colombia. Inclusion criteria were any patient who underwent high definition liposculpture (HDL) or HD2 in addition to fat grafting of any muscle of the lower limb. Exclusion criteria included active smokers, patients with body mass index above 30 kg/m<sup>2</sup>, patients with PMH with blood clotting disorders or any thrombotic event (ie, DVT and PE), and patients with American Society of Anesthesiologists (ASA) risk classification of III or higher. Cardiology assessment, including EKG and chest X-ray, was required for patients older than 40 years old. All patients were subject to protocols for safe large-volume liposuction including those for thromboembolic



**Fig. 1.** Muscles and neurovascular structures related to the anterior thigh: sartorius muscle (cut) (1), femoral nerve (2), rectus femoris muscle (3), lateral circumflex femoral artery (4), deep femoral artery (5), vastus lateralis muscle (6), femoral vein (7), femoral artery (8), vastus medialis muscle (9).

**Table 1. Power Muscles of the Leg**

Muscle	Anatomy	Description
Vastus lateralis m.	Main arterial pedicle	Descending branch of the lateral circumflex femoral artery is approximately 3 cm long and 1.5 mm in diameter and is located in the superior third of the muscle, extending along its medial border.
	Secondary arterial pedicle	1. Transverse branch of the lateral circumflex femoral artery: 3 cm long, 1 mm in diameter, and enters the muscle at the superior-posterior surface. 2. Posterior branch of the deep femoris artery: 1–2 cm long, 0.4 cm in diameter, and located in the inferior half of the muscle at the arterial intermuscular septum. 3. Superficial branch of the lateral superior genicular artery: 3 cm long, 0.5 mm in diameter, and located at the lateral condyle of the knee.
	Innervation	Femoral nerve
Vastus medialis m.	Main arterial pedicle	Descending branch of the Superficial femoral artery: 1–3 cm long and 1–1.5 mm in diameter, and enters the muscle at the junction of the mid third with the upper third.
	Secondary arterial pedicle	Branches coming from the superficial femoral artery (about 1–3 cm long and 1 mm in diameter) located deep and medial to the muscle. Some musculoarticular branches of the descending genicular artery (about 2 cm long and 1 mm in diameter) are located deep to the distal portion of the muscle.
Rectus femoris m.	Innervation	Femoral nerve
	Main arterial pedicle	Profunda femoris and descending branch of the lateral circumflex femoral artery.
	Secondary arterial pedicle	Branches coming from the superficial femoral artery.
Gastrocnemius m.	Innervation	Femoral nerve
	Main arterial pedicle	Medial and lateral sural arteries, branches from the posterior tibial artery and the popliteal artery, both 6 cm long and 2 mm in diameter
	Secondary arterial pedicle	Posterior and superficial branches of the peroneal artery.
Innervation	Tibial nerve	



**Fig. 2.** Neurovascular structures related to the posterior thigh: sciatic nerve (1) and its muscular branches (2); perforating branches of the deep femoral artery (3). Semitendinosus (4) and semimembranosus muscles (5), biceps femoris (6), popliteal vein (7), popliteal artery (8), common fibular nerve (9). *Note: The incision for liposuction and fat grafting of the posterior lower limb should be away from both the center (popliteal fossa) and the bony prominences (common fibular nerve).*



**Fig. 3.** Neurovascular structures related to the leg: medial head (1) and lateral head (2) of the gastrocnemius muscle. Popliteal artery (3), tibial nerve (4), anterior tibial artery (5), soleus muscle (6), posterior tibial artery (7), fibular artery (8).

event prevention, blood conservation, and hypothermia prevention.<sup>18</sup>

IV medications used were antibiotic prophylaxis with cefazolin (2 gr IV, 60 minutes before incision); dexamethasone, 8 mg; metoclopramide, 10 mg; diclofenac, 50 mg; and ranitidine, 50 mg. Photographic records were taken before and during follow-up at 2 days and 1, 3, 6, and 12 months after surgery.

### Surgical Technique

#### Markings

Markings are done in the standing position.

1. Gastrocnemius muscle: Ask the patient to stand on tip toes, then palpate the limits of the muscle mass and draw a line for demarcation, particularly the medial head in women and the lateral head in men. Address any lack of projection, the muscle volume, and tone as well.
2. Quadriceps: Ask the patient to do an isometric contraction and feel/see the muscular bellies. Delineate the distal portion of the vastus lateralis, the vastus medialis, and the rectus femoris muscles. Any muscle requiring volume augmentation is marked with a crossed pattern, so as to recognize which one has to be grafted once at the operating table (OT).
3. The “danger triangle” between the sartorius muscle, the adductor group, and inguinal ligament is drawn to prevent any injury of the main neurovascular pedicles.

#### Dynamic Definition Liposculpture

Liposculpture is achieved by a three-step process.

1. Infiltration: Tumescence solution (1000 mL of saline and 1 mL of 1:1000 epinephrine) + lidocaine (10 mg/kg), for both lower limb and the arm liposuction, which will help to ameliorate the exquisite postoperative pain of the extremities.
2. Fat emulsification: Third-generation ultrasound (VASER system) in pulsed (60%–70% power) and continuous modes (60–80% power) for the superficial and deep adipose layers, respectively. Pulsed mode at 60% power for all four extremities.
3. Microaire-assisted liposuction: Thorough debulking of the adipose tissue with 4.0-mm and/or 5.0-mm Mercedes cannulas over the deep layer. We use 3.0-mm Mercedes cannulas for detailed muscular definition (superficial layer), which is achieved based on the Basic, Moderate, and Xtreme definition algorithm.<sup>19</sup>

#### Fat Grafting

The adipose graft is harvested from the lipoaspirate, mainly from the abdomen and inner thighs. We add hypochlorous acid into the harvesting canister, which serves as both a bactericide and a cleansing solution. The Vibratory Tissue Separator (VTS Wells Johnson, Tucson, Ariz.) accelerates the process of decantation; then we transfer the supernatant to 60-mL syringes in preparation for adipose grafting.

#### Vastus Medialis and Lateralis

Patient must be in the supine position over the OT.

1. The kneecap incision gives access to both vastus muscles.
2. The tip of the cannula must be perpendicular to and far from the pedicle: about 45 degrees and 30 degrees away from the thigh’s midline for the vastus medialis and vastus lateralis, respectively.
3. Pinch the muscle bulk with almost a complete hand grip.
4. Place the tip at the center of the muscle bulk and drive the cannula into the muscle (by means of a gentle stab). One should feel a click or a decrease in resistance once inside.

Keep the tip stable and place the graft superficially through fanning and retrograde movements. (See **Video 3 [online]**, which displays fat grafting of the male lower extremity. A 35-year-old male patient undergoes fat grafting of his thighs in addition to HDL. Rectus femoris muscle and both vastus muscles were all grafted with ≈120 mL of adipose tissue. Then, a 27-year-old male patient with a congenital asymmetric body undergoes fat grafting of his right calf in addition to HDL. Both heads of the right gastrocnemius muscle were grafted with 80 mL of adipose tissue. The subcutaneous layer was also grafted to compensate the overall lack of tissue. He actually underwent fat grafting of most of the ipsilateral muscles of his body.) (See **Video 4 [online]**, which displays fat grafting of the female lower extremity. A 42-year-old female patient undergoes fat grafting of her thighs in addition to HDL. Both vastus muscles were grafted with ≈80 mL of adipose tissue. Then, a 29-year-old female patient undergoes fat grafting of her thighs and calves in addition to HDL. The vastus medialis and the gastrocnemius muscles were all grafted with 50–60 mL of adipose tissue. Subcutaneous fat grafting was also performed below the calves to avoid a masculine appearance of the leg.)

#### Rectus Femoris

1. Ipsilateral pubic incision allows its access (patient in supine position).
2. Direct the tip of the cannula perpendicular to the pedicle’s course: about 30 degrees away from the thigh’s midline. (See **Video 3 [online]**).
3. Avoid the “danger triangle”: Always point the tip up, from the incision until surpassing the sartorius muscle.
4. Pinch the muscle bulk.
5. Go as distal as possible from the pedicle (distal third of the thigh) before the maneuver. Then, point the tip horizontally and stab the muscle gently.
6. Place the graft.

#### Gastrocnemius

The patient must be in the prone position over the OT.

1. The medial and lateral popliteal-crease incisions allow access to the medial and lateral heads of the

gastrocnemius muscle, respectively (away from the main neurovascular structures of the leg to avoid unexpected injuries).

2. Point the tip of the cannula about 45 degrees away from the midline of the leg for both heads (perpendicular to the pedicle).
3. Pinch the muscle bulk.
4. Point the tip horizontally and drive the cannula into the muscle.
5. Then, point the tip of the cannula up, keep it stable, and place the adipose graft in small superficial strips inside the muscle through retrograde movements. (See Video 3 [online]). (See Video 4 [online]).
6. Most graft volume should be placed within the medial head for women and the lateral head for men.

**Outcomes Evaluation**

A nonstandardized survey aimed to evaluate the overall satisfaction with the procedure by asking the patients to rate their results on a scale from 1 to 5 (1 = poor results, 2 = below expectations, 3 = average results, 4 = good results, and 5 = above expectations) during the postoperative follow-up appointment.

**Ethical Considerations**

Each patient was informed of the purpose, methods, experimental technique, potential risks and benefits specific to fat grafting and HDL/HD2, sources of funding, any possible conflicts of interest, institutional affiliations of the authors, anticipated benefits and potential risks of

the study and the discomfort it may entail, poststudy provisions, and outcomes according to the Helsinki declaration. They were also informed of the right to refuse to participate in the study or to withdraw consent to participate at any time without reprisal. A freely given informed consent was signed for each patient participating in our report. Dissections were performed at a center for research and surgical training, which holds the ethical rights/standards and legal credentials to provide corpses for medical academic/private institutions for either training or research purposes.

**RESULTS**

We found 73 consecutive patients (26 men and 47 women) who underwent either HDL or HD2 and fat grafting of either the gastrocnemius (102 muscles), the vastus medialis (86 muscles), the vastus lateralis (98 muscles), the rectus femoris (22 muscles), or a combination of them (Table 2). Mean age was 34 and 41 years for men and women, respectively. The range of the fat graft volume was 50–200 mL. Combination of vastus lateralis + vastus medialis was more common among men than women. Almost all patients underwent fat grafting of muscles other than those from the lower limb. There were no major complications; however, two patients did have temporary numbness over the lateral thigh skin surface after fat grafting of the vastus lateralis muscle, which resolved spontaneously after 1 week without sequelae. Follow-up period ranged from 2 to 36 months (average = 18 months). The non-standardized survey showed that most patients were satisfied with their postoperative appearance (Table 3).

**Table 2. Patient Demographics**

n = 73	Men			Women		
	n = 26 (35%)	Average*	Range	n = 47 (65%)	Average	Range
Age		41	28–64		34	25–58
Weight (kg)		73	64–95		62	53–78
Height (m)		174	160–194		161	157–180
BMI (kg/m <sup>2</sup> )		24.2	23.5–28.2		23.7	22.4–28.6
Smoking history (2 wk before)	4 (15)			2 (4)		
		Volume (mL)			Volume (mL)	
<b>Surgery</b>						
HD2 + vastus medialis FG	21 (81)	120	50–180	22 (46)	100	50–120
HD2 + vastus lateralis FG	14 (54)	120	50–200	35 (75)	100	50–150
HD2 + rectus femoris FG	9 (35)	80	50–100	2 (4)	60	50–80
HD2 + VM + VL	15 (58)			19 (40)		
HD2 + VM + VL + RF	9 (35)			2 (4)		
HD2 + gastrocnemius	17 (65)	120	50–180	34 (72)	90	50–130
HD2 + gastrocnemius + thigh	16 (62)			31 (66)		
HD2 + leg FG and other zones	25 (96)			47 (100)		
Infiltration (mL)		5800	3600–10,400		5200	3200–9000
Lipoaspirate (mL)		4200	2900–8600		3950	2100–7600
<b>Complications</b>						
Infection	0 (0)			0 (0)		
Dysesthesia	2 (8)			0 (0)		
Ischemia	0 (0)			0 (0)		
Hematoma	0 (0)			0 (0)		
Necrosis	0 (0)			0 (0)		


\*Average volume and ranges were calculated per muscle unit. The contralateral side was grafted with the same amount of adipose tissue, only with specific exceptions.

BMI, body mass index; Hb, hemoglobin; HCT, hematocrit; FG, fat grafting.

**Table 3. Survey for Outcomes Evaluation**

Grading	Data, n = 65 (89%)	Timing of the Survey		
		3 months, n = 22 (34%)	6 months, n = 35 (54%)	9 months, n = 8 (12%)
1—Poor results	0 (0)	—	—	—
2—Below expectations	0 (0)	—	—	—
3—Average results	2 (3)	1 (1.5)	1 (1.5)	—
4—Good results*	34 (52)	10 (15)	18 (28)	6 (9)
5—Above expectations*	29 (45)	11 (17)	16 (25)	2 (3)

\*89% (n = 65) of patients rated their results as favorable and above the average. There were no complaints or results below “average” score.



**Fig. 4.** A 45-year-old male patient presenting with moderate lipodystrophy of the torso, the arms, and the lower limbs, who underwent dynamic definition liposculpture in addition to fat grafting of the thighs. 150 cc, 120 cc, and 60 cc were grafted into the vastus medialis, vastus lateralis and rectus femoris muscles, respectively. There is evident improvement of the muscle volume and definition in the 3-month postoperative pictures (C-D) compared to the preoperative ones (A-B).

## DISCUSSION


HDL involves a 3D multilayer concept of body contouring, while its evolution, HD2, incorporated concepts about muscle movement and all-or-nothing lipoplasty; hence, both are considered an effective and reproducible method of lipoplasty.<sup>19-21</sup> Fat grafting has been a cornerstone for these techniques as it allows both the volume enhancement and definition of body areas in which shape

is established by muscular structures like the lower limb (Figs. 4–6). (See figure, Supplemental Digital Content 1, A 31-year-old female patient with severe lipodystrophy of the arms, the torso and the lower limbs who specifically complained of her thin calves. She underwent dynamic definition liposculpture in addition to fat grafting of the gastrocnemius muscles. Its medial head and the subcutaneous tissue below it were grafted with 60 cc and 40 cc of



**Fig. 5.** A 27-year-old male patient with congenital asymmetric body and mild lipodystrophy of the torso, who underwent dynamic definition liposculpture in addition to fat grafting of the calves. Both heads of the gastrocnemius muscle were grafted with 80 cc of adipose tissue, while the subcutaneous layer was with 60 cc. The preoperative photographs (A-B) display a severe asymmetry of the gastrocnemius muscles and overall leg contour; fat grafting helped to achieve a more symmetric appearance of the entire leg (C-D) 4 months after surgery.





**Fig. 6.** A 42-year-old female patient with moderate lipodystrophy of the lower extremity in addition to moderate dermatochalasis and also mild lipodystrophy of the abdomen. She underwent dynamic definition miniabdominoplasty, liposculpture and fat grafting of the thighs. Both vastus muscles (medialis and lateralis) were grafted with 80 cc of adipose tissue. Comparatively to the male patient, the female thigh needs contour improvement with fat grafting, rather than just volume and definition, which can be appreciated in the 3-month postoperative photographs (C-D) in comparison to the preoperative ones (A-B).

fat, respectively. The leg shows a new muscular but also soft contour in the 6-week postoperative pictures (C-D) compared to the preoperative ones (A-B). <http://links.lww.com/PRSGO/C347>.

The most representative compartments in terms of contour and aesthetic appeal of the lower limbs are the anterior for the thigh and the posterior for the leg. Our

results show that men preferred augmentation of the vastus medialis while women preferred the vastus lateralis. Likewise, women would rather have the gastrocnemius muscle grafted compared with men, and the overall leg contour was greatly improved by grafting more fat at the medial head (compared with the lateral head) for women, but vice versa for the male patient.

In recent years, both women and men have shown an increasing interest in the appearance of their thighs and calves, which has run along with the increasing demand of aesthetic procedures of this area.<sup>2-22</sup> In the US in 2018, there were 10,100 thigh lifts performed, 9614 in women (95.1%) and 486 in men (4.8%), and also 655 calf augmentations, 377 in men (58%) and 278 in women (42%).<sup>23</sup> Surprisingly, our population showed a different distribution, as men required less calf augmentations compared with women, and women actually sought the appearance of stronger legs and more voluminous thighs compared with men.

Although it might be difficult to set a unique aesthetic ideal of the lower extremity, we may well state that the male leg is characterized by strong muscular indentations that highlight the definition of the leg, compared with the interaction between convexities and concavities of the female leg.<sup>20</sup> In any case, the surgeon must be aware of any unexpected trauma to the leg (including to varicose veins<sup>12-17</sup>) that might end up in acute compartment syndrome (ACS), so we should not only prevent possible injuries to the muscle's vascular pedicle but also avoid the very superficial planes to prevent severe and devastating complications.<sup>24,25</sup> In addition, the newest recommendations for gluteal adipose grafting (after the 2018 practice advisory from the Multi-Society Task Force for Safety in Gluteal Fat Grafting) include the use of ultrasound-guided documentation of cannula placement before and during fat injection and the limitation of three BBL cases as a maximum amount of total operative cases per day.<sup>26</sup> To our knowledge, there has not been any fatality or pulmonary fat embolism reported in the medical literature after IM fat grafting in a muscle other than the gluteus major,<sup>27-31</sup> or any case report of ACS following aesthetic procedures of the lower limb.<sup>32-34</sup> Animal studies have shown that IM fat grafts have no association with long-/short-term events of fat embolism after autologous fat transplantation, which further supports the special case of the gluteus major muscle. In that sense, the cadaveric dissections allowed us to formulate five strategic tips to perform a safe technique for autologous IM fat grafting (Table 4), as the muscle bulk has healthy vessels and few vascular structures. Moreover, the authors' experience with IM fat grafting techniques goes over 1200 cases for the volume improvement of the pectoral and deltoid muscles. In brief, we believe that HD2 liposculpture + a proper technique for IM fat grafting allowed the improvement of overall aesthetics of the lower limb, by improving the volume of the gastrocnemius, the vastus medialis, the

vastus lateralis, the rectus femoris, and at some point, the hamstrings/biceps femoris muscles.

### Limitations

The lack of inferential statistics and the retrospective nature of the study decrease its power and also preclude a strong association between the procedure and the outcomes, although these were clinically evident. The sample size might look small, since the population undergoing/seeking such procedures is limited; nonetheless, more than 300 muscles were grafted. The assessment of the muscular compartment's pressure is a critical aspect that should be studied in future clinical studies to support our findings and the reliability of the technique.

### CONCLUSIONS

Fat grafting could be considered an alternative surgical procedure that improves the athletic and aesthetic appearance of the lower limb. Both the proper identification of neurovascular structures and a structured technique make this approach reliable and reproducible with outstanding outcomes and a very low complication rate.

Alfredo E. Hoyos, MD

Department of Plastic Surgery, Dhara Clinic  
Carrera 15, no. 83-33, Suite 304  
Bogotá, Colombia  
E-mail: alhoyos@gmail.com

### REFERENCES

- Mundinger GS, Vogel JE. Calf augmentation and reshaping with autologous fat grafting. *Aesthet Surg J.* 2016;36:211-220.
- Erol O, Gürlek A, Agaoglu G. Calf augmentation with autologous tissue injection. *Plast Reconstr Surg.* 2008;121:2127-2133.
- Mathes DW, Kenkel JM. Current concepts in medial thighplasty. *Clin Plast Surg.* 2008;35:151-163.
- Armijo BS, Campbell CF, Rohrich RJ. Four-step medial thighplasty: refined and reproducible. *Plast Reconstr Surg.* 2014;134:717e-725e.
- Hoppmann R, Meruane M, Gonza D, et al. Calf lipo-reshaping. *J Plast Reconstr Aesthet Surg.* 2013;66:956-961.
- Kon M, Baert C, Lange M. Thigh augmentation. *Ann Plast Surg.* 1995;35:519-521.
- Anger J. Thigh augmentation: submuscular placement of silicone gel-filled prosthesis. *Aesthet Surg J.* 2005;25:44-48.
- Andjelkov K, Sforza M. Safety and efficacy of subfascial calf augmentation. *Plast Reconstr Surg.* 2017;139:657e-669e.
- Seo BF, Choi JY, Kim J, et al. Capsular contracture after calf augmentation with silicone implant insertion. *Arch Plast Surg.* 2015;42:642-645.
- Ahuai A, Texeira DC. Implantes de panturrilha: complicações, prevenção e tratamento. *Rev Bras Cir Plást.* 2010;25:547-550.
- Ginard I, Kleiber G, Kleiber H. Comprehensive lower extremity anatomy. In: Neligan PC, ed. *Plastic Surgery.* 3rd ed. Vol IV. New York: Elsevier; 2013:1-62.
- Toia F, D'Arpa S, Brenner E, et al. Segmental anatomy of the vastus lateralis: guidelines for muscle-sparing flap harvest. *Plast Reconstr Surg.* 2015;135:185e-198e.
- Bergaj J, Bunke-Paquette N. *The Vein Book.* 2nd ed. Oxford: Oxford University Press; 2014:37-42.
- Caggiati A. Clinical anatomy of the venous system of the lower extremity. In Lanzer P (ed.), *Pan Vascular Medicine.* 2nd ed. Berlin, Germany: Springer; 2015:4279-88.

**Table 4. The "Fav Five": Tips for a Safe Approach to IM Fat Grafting in HD2**

- The access should be as far from the pedicle as possible.
- The tip of the cannula should be directed in a perpendicular fashion in relation to the anatomical axis of the pedicle.
- Place the graft at the superficial layer of the muscle.
- Fat grafting is done in a retrograde fashion.
- Use 3- to 4-mm diameter cannulas for fat grafting (bigger than the diameter of the pedicle).

15. Zenn MR, Jones GE. Part II. Regional flaps. Lower extremity. In: *Reconstructive Surgery: Anatomy, Technique, and Clinical Applications*. Vol 2. 1st ed. St. Louis, MO: Quality Medical Pub.; 2012:1682–1705.
16. Mathes S, Nahai F. *Reconstructive Surgery Principles, Anatomy & Technique*. London: Churchill Livingstone; 1997:1307–1316.
17. Kim R, Lee W, Park E. Anatomic variations of lower extremity venous system in varicose vein patients: demonstration by three-dimensional CT venography. *Acta Radiol*. 2017;58:542–549.
18. Hoyos AE, Stefanelli M, Perez ME, et al. Adipose tissue transfer in dynamic definition liposculpture. PART I. The back: latissimus dorsi and trapezius muscles. *Plast Reconstr Surg Global Open*. 2022:e4587.
19. Hoyos AE, Perez ME, Domínguez-Millán R. Variable sculpting in dynamic definition body contouring: procedure selection and management algorithm. *Aesthet Surg J*. 2021;41:318–332.
20. Hoyos AE, Millard JA. VASER-assisted high-definition liposculpture. *Aesthet Surg J*. 2007;27:594–604.
21. Hoyos A, Prendergast P. *High Definition Body Sculpting.pdf*. Berlin: Springer; 2014: 193–204.
22. Karacaoglu EZ. Calf contouring with endoscopic fascial release, calf implant, and structural fat grafting. *Plast Reconstr Surg Glob Open*. 2013;1:e35.
23. American Society of Plastic Surgeons. 2018 plastic surgery statistics report. Published April 2019. Available at <https://www.plasticsurgery.org/documents/News/Statistics/2018/plastic-surgery-statistics-full-report-2018.pdf>. Accessed March 10, 2022.
24. Bomberg BC, Hurley PE, Clark CA, et al. Complications associated with the use of an infusion pump during knee arthroscopy. *Arthroscopy*. 1992;8:224–228.
25. Seiler JG, III, Valadie AL, III, Drvaric DM, et al. Perioperative compartment syndrome. A report of four cases. *J Bone Joint Surg Am*. 1996;78:600–602.
26. Del Vecchio D, Kenkel J. Practice advisory on gluteal fat grafting. *Aesthet Surg J*. 2022;42:1019–1029.
27. Chopan M, White J. Autogenous fat grafting to the breast and gluteal regions: safety profile including risks and complications. *Plast Reconstr Surg*. 2019;143:1625–1632.
28. Cárdenas-camarena L, Bayter JE, Aguirre-Serrano H, et al. Deaths caused by gluteal lipoinjection: what are we doing wrong? *Plast Reconstr Surg*. 2015;136:58–66.
29. Sinno S, Chang J. Determining the safety and efficacy of gluteal augmentation: a systematic review of outcomes and complications. *Plast Reconstr Surg*. 2016;137:1151–1156.
30. Villanueva N, Del Vecchio D. Staying safe during gluteal fat transplantation. *Plast Reconstr Surg*. 2017;141:79–86.
31. Condé-Green A, Kotamarti V, Nini KT, et al. Fat grafting for gluteal augmentation: a systematic review of the literature and meta-analysis. *Plast Reconstr Surg*. 2016;138:437–446.
32. Fraipont MJ, Adamson GJ. Chronic exertional compartment syndrome. *J Am Acad Orthop Surg*. 2003;11:268–276.
33. Saladin, KS. *Anatomy and Physiology: The Unity of Form and Function*. New York, NY: McGraw Hill; 2012:315.
34. Andjelkov K, Atanasijevic TC, Popovic VM, et al. Safe composite calf augmentation: a staged procedure. *Aesthet Surg J*. 2021;41:NP26–NP35.