



POSTER PRESENTATION

Open Access

# The potential of CD127 as a prognostic and residual disease marker in chronic adult T cell leukaemia/lymphoma

Huseini Kagdi<sup>1\*</sup>, Anat Melamed<sup>1</sup>, Silva Hilburn<sup>1</sup>, Nicolas Gillet<sup>2</sup>, Andrew Hodson<sup>1</sup>, Maria A Demontis<sup>1</sup>, Aileen Rowan<sup>1</sup>, Lucy Cook<sup>1</sup>, Charles Bangham<sup>1</sup>, Graham Taylor<sup>1</sup>

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses  
Montreal, Canada. 26-30 June 2013

Adult T cell Leukaemia Lymphoma [ATL], a mature T-cell neoplasm has been classified into 4 subtypes: smouldering; chronic leukaemia; lymphoma and acute leukaemia. The diagnosis depends on clinical features, the immunophenotype, and demonstration of HTLV-1 infection & ideally of monoclonal proviral integration. The typical immunophenotype of ATL is not specific. The methods used to detect monoclonality are labour-intensive and/or expensive and are not widely available. We developed a flow cytometry assay for diagnosis and monitoring of ATL. We performed 11-colour immunophenotyping, HTLV-1 proviral quantification and proviral integration site [IS] analysis on 53 samples from 36 patients [25 non ATL HTLV-1-infected, 11 chronic/smouldering ATL], 3 uninfected individuals and 2 HTLV-1-immortalized cell lines. The non-ATL patients had CD127+ & CCR7-lo expression in CD4+CD25+CCR4+ cells, and a polyclonal distribution on IS analysis. Four ATL patients had CD127+ & CCR7-lo expression in CD4+CD25+CCR4+ cells and polyclonal distribution on IS analysis. These patients had an excellent outcome achieving remission with either PUVA or no therapy. Eight ATL patients had CD127-lo expression on CD4+ CD25+ CCR4+ cells with mono/oligoclonal distribution on IS analysis. One of nine patients with chronic ATLL had high CCR7 expression. Foxp3 expression was variable. All 8 patients required systemic ATL treatment and longitudinal study of 5 patients found the change in frequency of CD4+CD25+CCR4+ to correlate with PVL whilst CD127 expression correlated with IS analysis ( $p<0.005$ ) and disease remission status. CD127 expression

appears to be useful to identify patients needing treatment and for monitoring the treatment of chronic ATL.

**Authors' details**

<sup>1</sup>Infectious Diseases and Immunology, Faculty of Medicine, Imperial College, London, UK. <sup>2</sup>Molecular and Cellular epigenetics, Interdisciplinary Cluster for Applied Genoproteomics (GIGA) of University of Liège, Leige, Belgium.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P140

Cite this article as: Kagdi et al.: The potential of CD127 as a prognostic and residual disease marker in chronic adult T cell leukaemia/lymphoma. *Retrovirology* 2014 11(Suppl 1):P140.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: hkagdi@imperial.ac.uk

<sup>1</sup>Infectious Diseases and Immunology, Faculty of Medicine, Imperial College, London, UK

Full list of author information is available at the end of the article