

## RESEARCH ARTICLE

# Dominant bacteria and influencing factors of early intestinal colonization in very low birth weight infants: A prospective cohort study

Yanbo Lu<sup>1</sup> | Xiaohong Cai<sup>1</sup> | Yao Zheng<sup>2</sup> | Qin Lyv<sup>2</sup> | Junhua Wu<sup>2</sup> 

<sup>1</sup>School of Medicine, Ningbo University, Ningbo, China

<sup>2</sup>Ningbo Women and Children's Hospital, Ningbo, China

**Correspondence**

Junhua Wu, Liuting Street 339, Ningbo City, Zhejiang Province 315012, China.  
Email: wudata@163.com

**Funding information**

This research was funded by the Ningbo Clinical Research Center for Children's Health and Diseases (no. 2019A21002); the Zhejiang Provincial Medical and Health Project (no. 2020RC109) and the Zhejiang Provincial Medical and Health Project (no. 2020KY279)

**Abstract**

**Background:** The intestine of newborns is colonized by bacteria immediately after birth. This study explored dominant bacteria and influencing factors of early intestinal colonization in the early life of very low birth weight infants (VLBWI).

**Methods:** We enrolled 81 VLBWI and collected anal swabs at 24 h, 7th, 14th and 21st day after birth. We conducted bacterial culture for anal swabs, then selected the colony with obvious growth advantages in the plate for further culture and identification. Afterward, we analyzed the distribution and influencing factors of intestinal dominant microbiota combined with clinical data.

**Results:** A total of 300 specimens were collected, of which 62.67% (188/300) had obvious dominant bacteria, including 29.26% (55/188) Gram-positive bacteria and 70.74% (133/188) Gram-negative bacteria. The top five bacteria with the highest detection rates were *Klebsiella pneumoniae*, *Escherichia coli*, *Enterococcus faecium*, *Enterococcus faecalis* and *Serratia marcescens*. Meconium-stained amniotic fluid and chorioamnionitis were correlated with intestinal bacterial colonization within 24 h of birth. Mechanical ventilation and antibiotics were independent risk factors affecting colonization. Nosocomial infection of *K. pneumoniae* and *S. marcescens* were associated with intestinal colonization. The colonization rates of *K. pneumoniae*, *E. coli*, *E. faecium*, and *E. faecalis* increased with the birth time.

**Conclusions:** The colonization rate in the early life of VLBWI increased over time and the predominant bacteria were Gram-negative bacteria. Meconium-stained amniotic fluid and chorioamnionitis affect intestinal colonization in early life. Mechanical ventilation and antibiotics were independent risk factors for intestinal bacterial colonization. The nosocomial infection of some bacteria was significantly related to intestinal colonization.

**KEYWORDS**

intestinal bacterial colonization, neonatal intensive care unit, nosocomial infection, very low birth weight infants

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Journal of Clinical Laboratory Analysis* published by Wiley Periodicals LLC

## 1 | INTRODUCTION

Studies have shown that a large number of microorganisms are colonized in the human body, most of which exist in the gut.<sup>1</sup> There are about  $10^2$  bacteria in the gut, which is the most important, enormous and complex microbiota in the human body.<sup>3</sup> The gut microbiota plays a vital role in maintaining normal metabolism and physiological functions of the human body and is closely related to many diseases such as infectious diseases, digestive system diseases, nervous system diseases, metabolic diseases, autoimmune diseases and so on.<sup>4-8</sup>

Premature infants refer to newborns whose gestational ages are less than 37 weeks, and those whose birth weights are <1,500 g are very low birth weight infants (VLBWI). There are more clinical problems and higher mortality in preterm infants, especially VLBWI that are the focus of management of preterm infants.<sup>9</sup> In recent years, with the improvement of perinatal medicine and neonatal intensive care treatment, the survival rate of VLBWI has increased significantly.<sup>10</sup> The development of infants' gut microbiota is related to cognition, metabolism, immune system and nervous system, gradually forming a diversified, interrelated and relatively stable signal network.<sup>11,12</sup> This dynamic process is influenced by various factors, such as delivery mode, gestational age, feeding practices, environmental microbial exposures, diseases and pharmaceuticals.<sup>13</sup> VLBWI, as a special group of preterm newborns, may have a different intestinal microecological pattern from normal newborns due to the following complex factors: admission to neonatal intensive care unit (NICU) immediately after birth, isolation from mother, the delayed establishment of intestinal feeding, antibiotics and invasive operations.<sup>2,9,11,14</sup> Once the microecological imbalance occurs in the vulnerable, it will affect the development of the gut, brain and immune system and even lead to necrotizing enterocolitis (NEC), sepsis, white matter damage, neurodevelopmental disorders, metabolic syndrome and long-term atopic diseases.<sup>2,15-17</sup> These complications are related to the changes in gut microbiota.<sup>18-21</sup> Early life (including embryonic and infant years) is a critical period for the colonization and formation of intestinal flora.<sup>21</sup> For VLBWI, the occurrence of any of the above complications will cause a serious blow to them. Therefore, the study of bacterial colonization and influencing factors is of great importance. We collected four anal swabs from 81 VLBWI at 24 h, 7th, 14th and 21st day after birth and carried out bacterial culture to analyze the predominant colonization bacteria in the early life of VLBWI in NICU combination with clinical data.

## 2 | METHODS

### 2.1 | Participants

The VLBWI that met the inclusion criteria from 1 March 2020 to 29 February 2021 in the NICU of Ningbo Women's and Children's Hospital were selected as the research objects. Inclusion criteria:

those with birth weight less than 1500g were directly admitted to NICU for treatment after delivery. Exclusion criteria: severe congenital malformation, less than 72 h of hospital stay, giving up treatment or death within 2 weeks of hospitalization, family members refusing to study or quitting during hospitalization. All subjects had informed consent signed by their guardians. This study was approved by the medical ethics committee of Ningbo Women and Children's Hospital and is in line with the declaration of Helsinki.

### 2.2 | Anal swab collection, bacterial culture and identification

The personnel of NICU collected anal swabs 24 h, 7th, 14th and 21st day after admission. When the infant was critically ill or discharged automatically, we would stop the collection on that day. The method of collecting anal swabs is inserting a cotton swab 1-2 cm into the anus, rotating it gently for 2 weeks, then taking it out and putting it into a sterile test tube.

The collected samples were inoculated into the Blood Agar Plate (MacConkey Agar) by the zoned streak plate method (see below). The medium was placed in a 35°C incubator and incubated for 18-24 h. The colonies with the largest number and the best growth, or a single colony with good growth in zone 3 or 4, were selected for purification. The colonies were identified by the automatic microbial analyzer (Vitek2 Compact600, Bio-Meri, France). The whole process shall be operated in strict accordance with the national manual of clinical examination version 3.<sup>22</sup> If there were no dominant bacteria on the culture dish or no single colony in zone 3 and zone 4, we designated it as "negative." Anaerobic culture was not performed.

The method of zoned scribe separation: The experimenter shall first pick out the specimen with the inoculation ring, apply it to zone 1 of the plate (accounting for 1/4 of the total area of the medium) and draw several streaks, and then draw in zones 2, 3 and 4 in turn. After the streaking of one area, the inoculation ring shall be heated and sterilized, and the streaking of the next area shall be carried out after cooling. Each area shall be contacted 1-3 times from the streaking of the previous area.

### 2.3 | Diagnostic criteria

The diagnosis of chorioamnionitis was a combined clinical picture and histopathology of the placenta and umbilical cord.<sup>23,24</sup> Nosocomial infection refers to the infection that occurred 48 h after admission.<sup>25</sup> When infants had symptoms or signs of infection, we collected corresponding samples for bacterial cultures, such as sputum, urine and peritoneal drainage. All amniotic fluids were cultured for bacteria. As for the factor of antibiotics, if antibiotics are used on the day of collection, it is considered "yes"; otherwise it is "no."

## 2.4 | Statistical analyses

SPSS 20.0 software was used for data analysis. The Chi-square test was used for comparison, while Fisher's exact probability method was used when the Chi-square test was not applicable.  $p < 0.05$  was considered statistically significant.

## 3 | RESULTS

### 3.1 | Basic characteristics

In total, 81 VLBWI were registered whose information is summarized in Table 1. The length of hospital stay of all VLBWI was  $50.40 \pm 16.49$  days. It should be noted that we did not collect all samples of 81 infants at 4-time points, due to critical illness, automatic discharge or carelessness of the staff and so on. The specific collection process is shown in Figure 1. A total of 300 anal swabs were collected, of which 188 had obvious dominant bacteria after inoculating on the culture dish.

### 3.2 | Distribution of dominant bacteria

All bacterial culture results at four-time points are shown in Table 2. We conducted a linear-trend Chi-square test for the positive rates ( $\chi^2 = 125.53, p < 0.001$ ). It indicated that the probability of colonization of dominant bacteria in the intestine in VLBW infants increased significantly with time (Figure 2).

TABLE 1 Basic information for all infants

Gender (cases)	Male	46
	Female	35
Mode of delivery (cases)	Cesarean section	51
	Vaginal delivery	30
Birth weight (g)		$1238.35 \pm 169.49$
Gestational age (weeks)		$29.44 \pm 1.77$
Feeding mode (cases)	Hydrolyzed cow's milk	20
	premature infant formula	61
Mechanical ventilation (cases)	Yes	50
	No	31
Antibiotic (cases)	Yes	78
	No	3
Chorioamnionitis (cases)	Yes	14
	No	67
Amniotic fluid bacterial culture (cases)	Positive	5
	Negative	76
Nosocomial infection (cases)	Yes	26
	No	55
Necrotizing enterocolitis (cases)	Yes	6
	No	75

Note: Birth weight and gestational age are expressed in the form of mean  $\pm$  standard deviation, while other items represent the number of infants.

The bacterial distribution of 188 positive anal swabs was summarized in Table 3. A total of 19 genera were identified. The top five were *K. pneumoniae*, *E. coli*, *E. faecium*, *E. faecalis* and *Serratia marcescens*. Gram-positive ( $G^+$ ) bacteria detected included *E. faecium*, *E. faecalis*, *S. epidermidis*, *S. agalactiae*, *L. salivarius*, *Gemella morbillorum* and *Staphylococcus haemolyticus*; Gram-negative ( $G^-$ ) bacteria included *K. pneumoniae*, *E. coli*, *S. marcescens*, *E. cloacae*, *E. aerogenes*, *Acinetobacter baumannii*, *Raoultella planticola*, *Kirschner Citrobacter*, *Enterobacter gergoviae*, *Klebsiella oxytoca*, *Comamonas testosterone* and *Stenotrophomonas maltophil*.  $G^+$  bacteria accounted for 29.26% (55/188) and  $G^-$  bacteria accounted for 70.74% (133/188). The colonization rates of  $G^+$  bacteria ( $\chi^2 = 22.97, p < 0.001$ ) and  $G^-$  bacteria ( $\chi^2 = 51.49, p < 0.001$ ) increased over time (Table 4). However, there were some differences. Both colonization rates of  $G^+$  and  $G^-$  were low within 24 h of admission, but  $G^-$  rapidly increased and was significantly higher than  $G^+$  after 7 days of birth, and continued to stabilize at a high level, while  $G^+$  bacterial colonization rate increased gradually (Figure 3).

### 3.3 | Influence of non-infectious factors

We analyzed the non-infective factors including gender, mode of delivery (vaginal delivery or cesarean section), gestational age, birth weight, feeding mode and mechanical ventilation. The results related to all non-infectious factors are shown in Table 5. From the results, mechanical ventilation is the influencing factor of intestinal bacterial colonization.




FIGURE 1 The number of anal swabs collected 4 times

Result	24h	D7	D14	D21
Negative (cases)	73	25	8	6
Positive (cases)	6	49	66	67
Total number (cases)	79	74	74	73
Positive rate (%)	7.59 (6/79)	66.22 (49/74)	89.19 (66/74)	91.78 (67/73)

TABLE 2 Results of bacterial culture of anal swabs

Note: Positive represented the bacteria with obvious dominant growth in the culture dish. Negative did not mean that there were no bacteria, but bacteria without obvious dominant growth.




FIGURE 2 The detection rate of intestinal dominant colonization bacteria at different time

### 3.3.1 | Mode of delivery

We analyzed the impact of different modes of delivery on the colonization of intestinal dominant bacteria four times. There was no statistical difference between the two groups and the specific results are shown in Table 6.

### 3.3.2 | Feeding mode

It should be explained that all subjects began enteral nutrition on the second day after admission. The feeding regimen is divided into

premature infant formula and hydrolyzed cow's milk. Hydrolyzed cow's milk has lower osmotic pressure and calories than premature infant formula. Different feeding methods had no effect on the detection rate of intestinal dominant bacteria and the data were shown in Table 7.

### 3.3.3 | Mechanical Ventilation

The detection rate of intestinal dominant bacteria with mechanical ventilation was 18.75% (9/48) while it without mechanical ventilation was 71.03% (179/252). Mechanical ventilation is one of the factors affecting bacterial colonization in VLBWI ( $\chi^2 = 47.11, p < 0.001$ ) (Table 5). The duration of the mechanical ventilation time was  $5.00 \pm 5.64$  days. In addition, we analyzed the relationship between the duration of mechanical ventilation and the colonization of dominant bacteria. The result showed that there was no correlation ( $p = 0.435$ ).

## 3.4 | Influence of infectious factors

The infection factors we analyzed included meconium-stained amniotic fluid (MSAF), maternal chorioamnionitis, nosocomial infection, NEC and antibiotics. The results are shown in Table 8.

### 3.4.1 | Meconium-stained amniotic fluid

The total detection rate of dominant bacteria of VLBWI was statistically different from MSAF (Table 8). Then we examined the influence of MSAF at different times. Within 24 h after admission, the detection rate in the MSAF group was 60.00% (3/5), while the non-MSAF group was 4.05% (3/74). MSAF has a significant effect on intestinal colonization in VLBWI within 24 h of birth ( $p < 0.05$ ) (Table 9). There was no statistical significance between the detection of intestinal colonization bacteria on the 7<sup>th</sup>, 14<sup>th</sup> and 21<sup>st</sup> day ( $p > 0.05$ ) (Table 9). MSAF is an independent risk factor affecting early intestinal colonization.

### 3.4.2 | Chorioamnionitis

Although there was no significant difference in the total detection rate of intestinal bacterial colonization between maternal intrauterine


TABLE 3 Distribution of dominant bacteria in intestinal colonization

Dominant bacteria	24h	D7	D14	D21	Total	Detection rate (%)	Composition ratio (%)
<i>Klebsiella pneumoniae</i>	1	20	20	23	64	21.33 (64/300)	34.04 (64/188)
<i>Escherichia coli</i>	2	6	14	14	36	12.00 (36/300)	19.15 (36/188)
<i>Enterococcus faecium</i>	0	3	11	12	26	8.67 (26/300)	13.83 (26/188)
<i>Enterococcus faecalis</i>	0	4	7	8	19	6.33 (19/300)	10.11 (19/188)
<i>Serratia marcescens</i>	0	5	5	0	10	3.33 (10/300)	5.32 (10/188)
<i>Enterobacter cloacae</i>	0	6	1	2	9	3.00 (9/300)	4.79 (9/188)
<i>Staphylococcus epidermidis</i>	1	2	0	2	5	1.67 (5/300)	2.66 (5/188)
<i>Kirschner citrobacter</i>	0	0	2	1	3	1.00 (3/300)	1.60 (3/188)
<i>Enterobacter aerogenes</i>	0	0	1	2	3	1.00 (3/300)	1.60 (3/188)
<i>Acinetobacter baumannii</i>	0	1	0	1	2	0.67 (2/300)	1.06 (2/188)
<i>Raoultella planticola</i>	0	0	2	0	2	0.67 (2/300)	1.06 (2/188)
<i>Streptococcus agalactiae</i>	2	0	0	0	2	0.67 (2/300)	1.06 (2/188)
<i>Lactobacillus salivarius</i>	0	1	0	0	1	0.33 (1/300)	0.53 (1/188)
<i>Gemella morbillorum</i>	0	0	1	0	1	0.33 (1/300)	0.53 (1/188)
<i>Enterobacter gergoviae</i>	0	0	1	0	1	0.33 (1/300)	0.53 (1/188)
<i>Klebsiella oxytoca</i>	0	0	0	1	1	0.33 (1/300)	0.53 (1/188)
<i>Staphylococcus haemolyticus</i>	0	0	0	1	1	0.33 (1/300)	0.53 (1/188)
<i>Comamonas testosterone</i>	0	1	0	0	1	0.33 (1/300)	0.53 (1/188)
<i>Stenotrophomonas maltophilia</i>	0	0	1	0	1	0.33 (1/300)	0.53 (1/188)
Total	6	49	66	67	188	62.67 (188/300)	100.00 (188/188)

TABLE 4 Distribution of G<sup>+</sup> and G<sup>-</sup> bacteria

Result	24h	D7	D14	D21	$\chi^2$	p Value
G <sup>+</sup>	3	10	19	23	22.97	<0.001
Non-G <sup>+</sup>	76	64	55	50		
G <sup>-</sup>	3	39	47	44	51.49	<0.001
Non-G <sup>-</sup>	76	35	27	29		

Abbreviations; G<sup>+</sup>: Gram-positive bacteria, non-G<sup>+</sup>: non-Gram-positive bacteria, G<sup>-</sup>: Gram-negative bacteria, non-G<sup>-</sup>: non-Gram-negative bacteria.

FIGURE 3 The detection rate of G<sup>+</sup> and G<sup>-</sup> bacteria at different time

infection (Table 8). Comparisons at different times showed that the detection rate in the maternal chorioamnionitis group was significantly higher within 24 h after admission ( $p < 0.05$ ) (Table 10). Chorioamnionitis is an independent risk factor for early intestinal colonization.

### 3.4.3 | Nosocomial infection

We performed statistics of VLBWI intestinal bacterial colonization with nosocomial infection, and the results showed no statistical difference ( $p > 0.05$ ) (Table 8). Therefore, we analyzed the relationship between the common four nosocomial infection bacteria and the detection of intestinal dominant bacteria, including *K. pneumoniae*, *S. marcescens*, *E. coli* and *E. cloacae* (Table 11). The detection rate of *K. pneumoniae* was 100% (5/5) in the *K. pneumoniae*-infected group while 43.42% (33/76) in the non-*K. pneumoniae*-infected group. The probability of intestinal colonization of *K. pneumoniae* in the *K. pneumoniae*-infected group was significantly higher than that in the non-*K. pneumoniae*-infected group ( $p < 0.05$ ) (Table 11).

The colonization rate of *S. marcescens* in VLBWI infected *S. marcescens* was 100% (3/3), and non-infected was 7.69% (6/78). The colonization rate of VLBWI infected *S. marcescens* was significantly higher than the non-infected group ( $p < 0.05$ ) (Table 11).

TABLE 5 Detection results related to non-infectious factors

Result	Gender		Delivery		Birth weight (g)			Gestational age(weeks)			Feeding mode		Mechanical ventilation	
	Male	Female	Cesarean section	Vaginal delivery	<1000	1000-1250	1251-1500	<28	28-32	>32	Hydrolysed cow's milk	Premature infant formula	Yes	No
Positive (cases)	112	76	116	72	17	80	91	21	142	25	45	143	9	73
Negative (cases)	57	55	72	40	12	44	56	17	78	17	27	85	39	179
$\chi^2$	2.15		0.2		0.42			1.4			0.001		47.11	
p Value	0.14		0.65		0.81			0.5			0.97		<0.001	

Nosocomial infections such as *K. pneumoniae* and *S. marcescens* were significantly associated with intestinal colonization in VLBWI. However, the infection of *E. coli* and *E. cloacae* were not significantly associated with intestinal colonization ( $p > 0.05$ ). Therefore, we believed that bacterial colonization is a potential risk factor for bacterial infection.

#### 3.4.4 | Necrotizing Enterocolitis

The detection rate of intestinal dominant bacteria was 63.63% (14/22) in the NEC group and 62.59% (174/278) in the non-NEC group. NEC had no obvious correlation with dominant colonization bacteria ( $p = 0.92$ ).

#### 3.4.5 | Antibiotic

The detection rate of intestinal colonization bacteria was 27.82% (37/133) during antibiotic use, while 92.07% (151/164) during non-using of antibiotics. The detection rate in the period without antibiotics was significantly higher than that in the period with antibiotics ( $\chi^2 = 124.01$ ,  $p < 0.01$ ) (Table 8). Antibiotics are a strong factor affecting the colonization of intestinal bacteria.

### 3.5 | Distribution of main colonization bacteria detected at different time

We compared the colonization detection rate of the top five dominant colonizing bacteria at different times (Figure 4). It was obvious that the colonization rate of *K. pneumoniae* increased over time. *K. pneumoniae* always maintained a high detection rate within 7-21 days. In addition, we found that the colonization rates of *E. coli*, *Enterococcus faecium* and *E. faecalis* also increased over time. However, the detection rate of *S. marcescens* did not increase linearly with time (Table 12).

## 4 | DISCUSSION

Our results showed that Gram<sup>-</sup> bacteria were the main colonization bacteria in the early stage of VLBWI. The colonization rate of intestinal bacteria increased over time. In addition, the colonization rate of G<sup>-</sup> bacteria increased rapidly on the 7th day after birth and maintained a high level. However, Gram<sup>+</sup> bacteria increased slowly and were significantly lower than G<sup>-</sup> bacteria. Korpela found that the total abundance of intestinal microorganisms in preterm infants increased with age at birth.<sup>26</sup> Our results are consistent with the research reported that the intestinal microflora of premature infants less than 33 weeks old are characterized by G<sup>+</sup> cocci such as *Streptococcus* and *Staphylococcus*, but are rapidly replaced by G<sup>-</sup> bacteria (*Proteus* and *Enterobacteriaceae*).<sup>27</sup>

TABLE 6 Results of intestinal dominant bacteria culture related to delivery mode

Result	24 h		D7		D14		D21	
	Cesarean section	Vaginal delivery	Cesarean section	Vaginal delivery	Cesarean section	Vaginal delivery	Cesarean section	Vaginal delivery
Positive (cases)	4	2	30	19	41	25	41	26
Negative (cases)	46	27	15	10	6	2	5	1
$\chi^2$	-		0.01		-		-	
p value	<sup>a</sup> 0.61		0.92		<sup>a</sup> 0.70		<sup>a</sup> 0.40	

<sup>a</sup>Fisher's exact probability method.

TABLE 7 Results of intestinal dominant bacteria culture related to feeding mode

Result	24 h		D7		D14		D21	
	Hydrolyzed cow's milk*	Premature infant Formula*	Hydrolyzed cow's milk	Premature infant formula	Hydrolyzed cow's milk	Premature infant formula	Hydrolyzed cow's milk	Premature infant formula
Positive (cases)	2	4	14	35	14	52	15	52
Negative (cases)	18	55	6	19	2	6	1	5
$\chi^2$	-		0.18		-		-	
p Value	<sup>a</sup> 0.64		0.68		1		<sup>a</sup> 1.00	

Note: \*subjects were fasting at 24 h.

<sup>a</sup>Fisher's exact probability method.

TABLE 8 Detection results related to infectious factors

Infectious factors	MSAF		Chorioamnionitis			Nosocomial infection		NEC		Antibiotic	
	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	
Positive (cases)	14	174	31	157	59	129	14	174	37	151	
Negative (cases)	2	110	16	96	37	75	8	104	96	16	
$\chi^2$	4.46		0.26		0.09		0.01		124.01		
p Value	0.03		0.61		0.77		0.92		<0.001		

Abbreviations: MSAF, meconium-stained amniotic fluid, NEC, necrotizing enterocolitis.

TABLE 9 Effect of MSAF on the detection of intestinal dominant bacteria in VLBWI at different time

MSAF	24 h		D7		D14		D21	
	Yes	No	Yes	No	Yes	No	Yes	No
Positive (cases)	3	3	4	45	4	62	3	64
Negative (cases)	2	71	0	25	0	8	0	6
p Value	<sup>a</sup> 0.002		<sup>a</sup> 0.29		<sup>a</sup> 1.00		<sup>a</sup> 1.00	

<sup>a</sup>Fisher's exact probability method.

TABLE 10 Influence of maternal chorioamnionitis at different time

Intrauterine infection	24 h		D7		D14		D21	
	Yes	No	Yes	No	Yes	No	Yes	No
Positive (cases)	4	3	5	44	12	54	11	56
Negative (cases)	10	62	4	21	0	8	1	5
$\chi^2$	5.49		0.12		0.66		-	
p	0.02		0.73		0.42		<sup>a</sup> 1.00	

<sup>a</sup>Fisher's exact probability method.

TABLE 11 Detection of common nosocomial infection bacteria in intestinal colonization of VLBWI

Common bacteria	<i>Klebsiella pneumoniae</i>		<i>Serratia marcescens</i>		<i>Escherichia coli</i>		<i>Enterobacter cloacae</i>	
	Infected	Non-infected	Infected	Non-infected	Infected	Non-infected	Infected	Non-infected
Homologous bacterial colonization (cases)	5	33	3	6	3	20	1	7
Non-homologous bacterial colonization (cases)	0	43	0	72	2	56	1	72
p Value	<sup>a</sup> 0.02		<sup>a</sup> <.001		<sup>a</sup> 0.14		<sup>a</sup> 0.19	

<sup>a</sup>Fisher's exact probability method.




FIGURE 4 Broken line chart of colonization rate of main bacteria at different times

Afterward, we analyzed the non-infectious factors that may affect the colonization of intestinal dominant bacteria on VLBWI, including gender, mode of delivery, gestational age, birth weight, mechanical ventilation and feeding mode. It is worth noting that the detection rate of intestinal dominant colonization bacteria is only 7.59% (6/79) when VLBWI are fasting within 24 h of birth. This does not mean that the rectal swab within 24 h is completely sterile, but there is no obvious vigorous growth of bacteria. This may be related to the low bacterial diversity in meconium of VLBWI.<sup>28</sup> It has long been considered that the fetus grows in a sterile environment and bacterial colonization begins at rupturing of the fetal membrane. More recent careful research have shown that the amniotic fluid is not sterile, suggesting that colonization of the gut begins in utero.<sup>29</sup> The analysis of placenta obtained during full-term cesarean section without rupture of membranes showed that the microbiota of placenta, amniotic fluid and meconium were similar.<sup>30</sup> The development of the intestinal microbiota of newborns is usually driven by the colostrum. However, VLBWI admitted to NICU immediately after birth are usually fed with hydrolyzed milk and formula, which seem to have little effect on the colonization of intestinal dominant bacteria. The study has shown that diet has a lesser impact on the gut microbiota in extremely preterm infants than is seen in term infants.<sup>31</sup> In addition, the detection rate of intestinal bacteria without mechanical ventilation was significantly higher than that on VLBWI with mechanical ventilation (Table 5), which may be due to the severe condition, large dosage and frequent use of antibiotics in VLBWI with mechanical ventilation affected the normal colonization of intestinal bacteria.

The analysis of intestinal flora of preterm infants based on 16S pyrophosphate sequencing found that gestational age and mode of delivery were important influencing factors.<sup>32,33</sup> However, these differences were not found in this study. The possible reason is that our method based on bacterial culture has no way to compare its flora diversity and composition. From the dominant bacteria identification, gestational age and delivery mode have no effect on the colonization of intestinal dominant bacteria.

Among the infectious factors, we found MSAF and maternal chorioamnionitis were closely related to the bacterial colonization in the early stage of VLBWI, and the influence became smaller with time.



**TABLE 12** Distribution of main colonization bacteria detected at different time

Bacterium		24 h	D7	D14	D21	$\chi^2$	p Value
<i>Klebsiella pneumoniae</i>	Positive	1	20	20	23	19.12	<0.001
	Negative	78	54	54	50		
<i>Escherichia coli</i>	Positive	2	6	14	14	13.28	<0.001
	Negative	77	68	60	59		
<i>Enterococcus faecium</i>	Positive	0	3	11	12	17.25	<0.001
	Negative	79	71	63	61		
<i>Enterococcus faecium</i>	Positive	0	4	7	8	8.78	0.003
	Negative	79	70	67	65		
<i>Serratia marcescens</i>	Positive	0	5	5	0	0.007	0.932
	Negative	79	69	69	73		

MSAF and maternal chorioamnionitis may be one of the causes of premature birth. Early maternal exposure to inflammation can lead to a series of events, including prostaglandin increase, promoting uterine contraction, degradation of chorioamniotic extracellular matrix and cervical ripening.<sup>34</sup> Fetuses may be exposed to pathogenic microorganisms, resulting in a decrease in the diversity of intestinal microbiota and a relatively large number of pathogenic bacteria in infants.<sup>35</sup> It is well known that infection during pregnancy can change the vaginal flora of bacterial vaginosis, or increase the susceptibility of other infections with altered maternal and infant microbiota, thus interrupting normal pregnancy.<sup>36</sup> In this study, NEC and intestinal colonization bacteria detection on VLBWI had no obvious correlation. NEC in preterm infants is not only associated with a single pathogen. There may be a reason for insufficient sample size, and its correlation may need larger samples and more scientific research program to confirm.

There is no doubt that antibiotic is a strong influencing factor. We found that the rate was only 27.82% (37/133) during the use of antibiotics, while it was as high as 92.07% (151/164) during the non-use of antibiotics. The use of antibiotics markedly reduced the colonization rate of intestinal bacteria. VLBWI are almost universally treated with antibiotics while in NICU. Arboleya found a decrease in fecal diversity among premature infants who received long-term antibiotic treatment.<sup>37</sup> Vangay proposed four biological disorders induced by antibiotics: the loss of keystone taxa, the loss of diversity, the changes of metabolic capacity and the proliferation of pathogens.<sup>38</sup>

In the VLBWI enrolled in this study, the hospital infection rate was as high as 32.10% (26/81) which is much higher than usual. However, this data are authentic. Generally, the nosocomial infection rate of the neonatal department is 4.5%–5.0%. The vulnerability of VLBWI, long hospitalization and many invasive procedures may lead to this result. Of course, we can not rule out the effect of cross infection. We analyzed the influence of common bacteria in a nosocomial infection on VLBWI's intestinal colonization. Five children infected *K. pneumoniae* and three children with *S. marcescens* had corresponding bacterial colonization. The detection rates of *K. pneumoniae* and *S. marcescens* ranked first and fifth with a high colonization rate in this study. It is worth noting that late-onset septicemia

caused by Gram-negative bacteria is one of the main causes of incidence rate and mortality in the neonatal intensive care unit, and it is especially significant for premature infants with small gestational age.<sup>39–41</sup> Zaidi et al.<sup>42</sup> pointed out that *K. pneumoniae* is one of the most common pathogens of nosocomial respiratory tract infection, and is also the second-largest factor causing nosocomial sepsis and urinary tract infection. In this study, the detection rate of *K. pneumoniae* was as high as 21.33% (64/300) and increased over time. Epidemiological studies have suggested that *K. pneumoniae* host-to-host transmission requires close contact and usually occurs through the fecal-oral route.<sup>43</sup> Moles et al.<sup>28</sup> indicated that the presence of *S. marcescens* was strongly related to a higher degree of immaturity and hospital-related parameters, including antibiotic and mechanical ventilation. Although *S. marcescens* displays relatively low virulence, it can lead to nosocomial infections and outbreaks in immunocompromised or critically ill patients, particularly in settings such as NICU.<sup>44</sup> However, the contaminated hands of medical staff are considered to be a major medium of transmission. In NICU, colonized or infected newborns are the main potential source of pathogens, especially in the respiratory apparatus, but also in the gastrointestinal tract. To curb the spread of infection, it is necessary to identify colonized or infected patients as soon as possible and prompt implement infection control measures, particularly strict hand hygiene and contact precautions.

Garcia Hernandez pointed out that *E. coli* is the most common pathogen of nosocomial and community-acquired infections.<sup>45</sup> *E. cloacae* is responsible for nosocomial outbreaks in susceptible patients in NICU.<sup>46</sup> In this study, we did not find a significant correlation between intestinal colonization of *E. coli* and *E. cloacae* and nosocomial infection. The reasons may be that the sample size is insufficient or the two bacteria are not identified as the most dominant bacteria. This could be due to insufficient sample size or the failure of these two bacteria to be identified as the most dominant bacteria.

In this study, we adopted an anal swab bacterial culture to select the most dominant bacteria, and then analyzed the distribution and influencing factors of intestinal colonization from this perspective. Although flora sequencing can usually obtain more bacterial composition and diversity, it cannot distinguish living bacteria from dead

bacteria, let alone judge the vitality of bacteria. Of course, our study has some limitations. Firstly, we did not carry out anaerobic culture. Secondly, we did not set up a control group. For further research, we have collected feces from VLBWIs and term infants. After that, we will perform 16S pyrophosphate sequencing to obtain more information about the gut microbiota of VLBWI.

### CONFLICT OF INTEREST

The authors declare that they have no competing interests.

### CONSENT

All subjects had informed consent signed by their guardians.

### DATA AVAILABILITY STATEMENT

The data used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### ORCID

Junhua Wu  <https://orcid.org/0000-0003-3355-6550>

### REFERENCES

- Altveş S, Yildiz HK, Vural HC. Interaction of the microbiota with the human body in health and diseases. *Biosci Microbiota Food Health*. 2020;39(2):23-32.
- Desorcy-Scherer K, Bendixen MM, Parker LA. Determinants of the very low-birth-weight infant's intestinal microbiome: a systematic review. *J Perinat Neonatal Nurs*. 2020;34(3):257-275.
- Power SE, O'Toole PW, Stanton C, Ross RP, Fitzgerald GF. Intestinal microbiota, diet and health. *Br J Nutr*. 2014;111(3):387-402.
- Kataoka K. The intestinal microbiota and its role in human health and disease. *J Med Invest*. 2016;63(1-2):27-37.
- Zeevi D, Korem T, Godneva A, et al. Structural variation in the gut microbiome associates with host health. *Nature*. 2019;568(7750):43-48.
- Salvucci E. The human-microbiome superorganism and its modulation to restore health. *Int J Food Sci Nutr*. 2019;70(7):781-795.
- Lynch SV, Pedersen O. The human intestinal microbiome in health and disease. *N Engl J Med*. 2016;375(24):2369-2379.
- De Luca F, Shoenfeld Y. The microbiome in autoimmune diseases. *Clin Exp Immunol*. 2019;195(1):74-85.
- Hahn WH, Chang JY, Chang YS, Shim KS, Bae CW. Recent trends in neonatal mortality in very low birth weight Korean infants: in comparison with Japan and the USA. *J Korean Med Sci*. 2011;26(4):467-473.
- Chung SH, Bae CW. Improvement in the survival rates of very low birth weight infants after the establishment of the Korean neonatal network: comparison between the 2000s and 2010s. *J Korean Med Sci*. 2017;32(8):1228-1234.
- Gibson MK, Crofts TS, Dantas G. Antibiotics and the developing infant gut microbiota and resistome. *Curr Opin Microbiol*. 2015;27:51-56.
- Niemark HJ, De Meij TG, van Ganzwinkel CJ, et al. Necrotizing enterocolitis, gut microbiota, and brain development: Role of the brain-gut axis. *Neonatology*. 2019;115(4):423-431.
- Henderickx JGE, Zwittink RD, van Lingen RA, Knol J, Belzer C. The preterm gut microbiota: an inconspicuous challenge in nutritional neonatal care. *Front Cell Infect Microbiol*. 2019;2(9):85.
- Unger S, Stintzi A, Shah P, Mack D, O'Connor DL. Gut microbiota of the very-low-birth-weight infant. *Pediatr Res*. 2015;77(1-2):205-213.
- Neu J, Pammi M. Pathogenesis of NEC: impact of an altered intestinal microbiome. *Semin Perinatol*. 2017;41(1):29-35.
- Adlerberth I. Factors influencing the establishment of the intestinal microbiota in infancy. *Nestle Nutr Workshop Ser Pediatr Program*. 2008;62:13-29. discussion 29-33.
- Adlerberth I, Wold AE. Establishment of the gut microbiota in Western infants. *Acta Paediatr*. 2009;98(2):229-238.
- Rich BS, Dolgin SE. Necrotizing enterocolitis. *Pediatr Rev*. 2017;38(12):552-559.
- Dong Y, Speer CP. Late-onset neonatal sepsis: recent developments. *Arch Dis Child Fetal Neonatal Ed*. 2015;100(3):F257-F263.
- Fanaro S. Feeding intolerance in the preterm infant. *Early Hum Dev*. 2013;89(Suppl 2):S13-S20.
- Zhuang L, Chen H, Zhang S, Zhuang J, Li Q, Feng Z. Intestinal microbiota in early life and its implications on childhood health. *Genomics Proteomics Bioinformatics*. 2019;17(1):13-25.
- Ye Y, Wang Y, Shen Z. *The National Manual of Clinical Examination Verion3 [M]*. Southeast University Press; 2006:11.
- Greenberg MB, Anderson BL, Schulkin J, Norton ME, Aziz N. A first look at chorioamnionitis management practice variation among US obstetricians. *Infect Dis Obstet Gynecol*. 2012;2012:628362.
- Hui YunZ. *Diagnostic pathology of Obstetrics and Gynecology [M]*. Peking University Medical Press; 2007:1089.
- New Medicine*. Ministry of health of the people's Republic of China. Excerpt of diagnostic criteria for nosocomial infection (Trial) (1). 2005;36(8):1.
- Korpela K, Blakstad EW, Moltu SJ, et al. Intestinal microbiota development and gestational age in preterm neonates. *Sci Rep*. 2018;8(1):2453.
- Berkhout DJC, Niemarkt HJ, de Boer NKH, Benninga MA, de Meij TGJ. The potential of gut microbiota and fecal volatile organic compounds analysis as early diagnostic biomarker for necrotizing enterocolitis and sepsis in preterm infants. *Expert Rev Gastroenterol Hepatol*. 2018;12(5):457-470.
- Moles L, Gómez M, Heilig H, et al. Bacterial diversity in meconium of preterm neonates and evolution of their fecal microbiota during the first month of life. *PLoS One*. 2013;8(6):e66986.
- Wassenaar TM, Panigrahi P. Is a foetus developing in a sterile environment? *Lett Appl Microbiol*. 2014;59(6):572-579.
- Collado MC, Rautava S, Aakko J, Isolauri E, Salminen S. Human gut colonisation may be initiated in utero by distinct microbial communities in the placenta and amniotic fluid. *Sci Rep*. 2016;22(6):23129.
- Schwartz A, Gruhl B, Löbnitz M, Michel P, Radke M, Blaut M. Development of the intestinal bacterial composition in hospitalized preterm infants in comparison with breast-fed, full-term infants. *Pediatr Res*. 2003;54(3):393-399.
- Underwood MA, Sohn K. The Microbiota of the extremely preterm infant. *Clin Perinatol*. 2017;44(2):407-427.
- Dominguez-Bello MG, Costello EK, Contreras M, et al. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci USA*. 2010;107(26):11971-11975.
- Kemp MW. Preterm birth, intrauterine infection, and fetal inflammation. *Front Immunol*. 2014;1(5):574.
- Chernikova DA, Koestler DC, Hoen AG, et al. Fetal exposures and perinatal influences on the stool microbiota of premature infants. *J Matern Fetal Neonatal Med*. 2016;29(1):99-105.
- Redelinghuys MJ, Ehlers MM, Dreyer AW, Kock MM. Normal flora and bacterial vaginosis in pregnancy: an overview. *Crit Rev Microbiol*. 2016;42(3):352-363.
- Arbolea S, Sánchez B, Milani C, et al. Intestinal microbiota development in preterm neonates and effect of perinatal antibiotics. *J Pediatr*. 2015;166(3):538-544.
- Vangay P, Ward T, Gerber JS, Knights D. Antibiotics, pediatric dysbiosis, and disease. *Cell Host Microbe*. 2015;17(5):553-564.

39. Smith A, Saiman L, Zhou J, Della-Latta P, Jia H, Graham PL 3rd. Concordance of gastrointestinal tract colonization and subsequent bloodstream infections with gram-negative bacilli in very low birth weight infants in the neonatal intensive care unit. *Pediatr Infect Dis J*. 2010;29(9):831-835.
40. Alshaikh B, Yusuf K, Sauve R. Neurodevelopmental outcomes of very low birth weight infants with neonatal sepsis: systematic review and meta-analysis. *J Perinatol*. 2013;33(7):558-564.
41. Vergnano S, Menon E, Kennea N, et al. Neonatal infections in England: the NeonIN surveillance network. *Arch Dis Child Fetal Neonatal Ed*. 2011;96(1):F9-F14.
42. Zaidi AK, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. Hospital-acquired neonatal infections in developing countries. *Lancet*. 2005;365(9465):1175-1188.
43. Young TM, Bray AS, Nagpal RK, Caudell DL, Yadav H, Zafar MA. Animal model to study klebsiella pneumoniae gastrointestinal colonization and host-to-host transmission. *Infect Immun*. 2020;88(11):e00071-e20.
44. Cristina ML, Sartini M, Spagnolo AM. *Serratia marcescens* Infections in Neonatal Intensive Care Units (NICUs). *Int J Environ Res Public Health*. 2019;16(4):610.
45. Álvarez J, Rojas Á, Carvajal C, et al. Evaluation of susceptibility and response to therapy with piperacillin-tazobactam in patients with infections caused by *Escherichia coli* with extended-Spectrum  $\beta$ -Lactamase (ESBL) CTX-M. *Rev Chilena Infectol*, 2018;35(4):343-350. Spanish.
46. Ferry A, Plaisant F, Ginevra C, et al. Enterobacter cloacae colonisation and infection in a neonatal intensive care unit: retrospective investigation of preventive measures implemented after a multiclonal outbreak. *BMC Infect Dis*. 2020;20(1):682.

**How to cite this article:** Lu Y, Cai X, Zheng Y, Lyv Q, Wu J. Dominant bacteria and influencing factors of early intestinal colonization in very low birth weight infants: A prospective cohort study. *J Clin Lab Anal*. 2022;36:e24290. doi:[10.1002/jcla.24290](https://doi.org/10.1002/jcla.24290)