



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

^a Unidad de Estimulación Cardíaca de UCI, Servicio de Cuidados Críticos y Urgencias, Hospital Virgen de la Victoria de Málaga, Málaga, España

^b Unidad Coronaria, Servicio de Medicina Intensiva, Hospital Regional de Málaga, Málaga, España

^c Servicio de Medicina Intensiva, Hospital Quironsalud Miguel Domínguez, Pontevedra, España

^d Servicio Medicina Intensiva, Hospital Universitario de Torrejón, Torrejón de Ardoz (Madrid), España

* Autor para correspondencia.

Correos electrónicos: cristinasweb@gmail.com (C. Salazar Ramírez), rociogmzlpz@gmail.com (R. Gómez-López).

<https://doi.org/10.1016/j.medin.2020.08.014>
0210-5691/ © 2020 Publicado por Elsevier España, S.L.U.

Innate cell response in severe SARS-CoV-2 infection in children: Expression analysis of CD64, CD18 and CD11a



Respuesta de celular innata en infección pediátrica grave por SARS-CoV-2: análisis de expresión de CD64, CD18 y CD11a

Dear Editor,

In January 2020, a new coronavirus known as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was described in Wuhan, China. The virus, which produces coronavirus disease 2019 (COVID-19), has been declared a global health emergency and pandemic by the World Health Organization. Spain is one of the more severely affected countries.¹

The immune response to SARS-CoV-2 infection appears to be a critical factor in the development and prognosis of COVID-19 patients.² In children, severe forms of the disease like the pediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 appears to be related with some immune dysregulation.³ So, increase knowledge about the innate cellular immune response to SARS-CoV-2 is of great interest. To this, the study by flow cytometry (FC) may provide critical data and further understanding of this novel disease.³

In this paper, we study three molecules which are part of the innate cellular response to infection: CD64, CD18 and CD11a. The CD64 is a type I high-affinity receptor for the Fc fraction of the immunoglobulin G. It is located on monocytes, macrophages, dendritic cells, and neutrophils. The CD64 density on the cell surface is related to the stimulation received by inflammatory cytokines⁴. The CD18, also known as integrin beta-2, participates in leukocyte adhesion and signaling. The CD11a associates with CD18 to form the lymphocyte function-associated antigen 1, or LFA-1. This LFA-1 on leukocytes plays a central role in leukocyte cell-cell interactions and lymphocyte stimulation.

We study in this report three children with severe SARS-CoV-2 infection. Also, we compare them with a healthy control, a case of severe influenza infection and a case of Neisseria meningitidis sepsis. All cases included had SARS-CoV-2 infection confirmed by real-time reverse transcriptase-polymerase chain reaction (RT-PCR) on nasopharyngeal swab samples. The cases trajectories,

complementary tests, and therapy approaches are summarized in [Table 1](#). The children were studied after informed consent was obtained. One 0.5 ml sample of peripheral blood was extracted on admission to the pediatric intensive care unit (PICU). The samples obtained were collected in sterile EDTA at room temperature or refrigerated at 4 °C, after which they were used for CD45+ cell-marker studies and analyzed by FC within 24 h. Cell surface expression of CD64, CD18, and CD11a was measured by BD FACS Canto II flow cytometer (Becton Dickinson, New York, USA). CD64 (clone 10.1), CD18 (clone CBR LFA-1/2), and CD11a (clone HI111) monoclonal antibodies were obtained from Biolegend® (San Diego, CA, USA). Expressions were measured in monocytes, neutrophils, and lymphocytes. Cell viability was confirmed by 7-AAD staining. At least 10,000 events were recorded for each sample. Flow-cytometric settings and samples were prepared according to manufacturer instructions. Neutrophils, monocytes and lymphocytes were identified on a dot-plot and gated ([Fig. 1](#)). The intensity of CD64, CD18, and CD11a surface expression was measured as mean fluorescence intensity in arbitrary units (MFI, [Fig. 1B](#)).⁵ The FC was performed on PICU admission in all cases. All patients received methylprednisolone prior to FC.

As results, we provide the description of CD64, CD18, and CD11a expression on neutrophils, monocytes and lymphocytes in children with severe SARS-CoV-2 disease. This expression appears to be higher compared to other infections and may point to an exacerbated cellular innate response in these children.³⁻⁶

The cytopathic effects of SARS-CoV-2 combined with the host immune response may play a major role in disease severity. A dysregulated immune response may result in inflammation and clinical worsening in patients with COVID-19. Elevated CD64 expression have been previously described in infectious and noninfectious diseases.⁷ Our group carried out CD64 expression studies in acute bronchiolitis and severe viral and bacterial infections.⁵ As can be seen in [Fig. 1](#) children with SARS-CoV-2 show levels of CD64 expression that are higher than in previous published reports of bacterial or viral infections or autoinflammatory diseases.⁵ Regarding the CD11a and CD18 complex or LFA-1, it is known that plays a key role in migration. Through these, leukocytes are mobilized from the bloodstream into tissues. One of the main findings in COVID-19 patients is the presence of lymphopenia.⁸ It can be seen in our cases. This may be linked to the migration of CD8+ lymphocytes to the infected tissues. As seen in [Fig. 1](#), the CD11a upregulation in CD8+ is clear and could be linked to this process. The LFA-1 is also

Table 1 Epidemiologic characteristics, clinical features, radiologic findings, and management of children admitted for pediatric critical care due to *Influenza B*, *Neisseria meningitidis* and SARS-CoV-2 infection.

	Influenza B	Neisseria meningitidis	Case 1	Case 2	Case 3
Age in years	4	9	12	11	7
Sex	Male	Male	Male	Male	Female
Referring department	Emergency department, 1 day of symptoms	Emergency department, 1 day of symptoms	Emergency department, 3 days of symptoms	Emergency department, 2 days of symptoms	Pediatric ward, 4 days of symptoms (one day of admission)
Previous diseases	No	No	No	No	No
Signs and symptoms prior to PICU admission	Tachipnea, hypoxemia, fever	Tachycardia, hypotension, fever	Fever, nausea, vomiting, diarrhea	Fever, nausea, diarrhea, adenopathy	Fever, abdominal pain
Cause of PICU admission	Respiratory instability	Hemodynamic instability	Hemodynamic instability	Hemodynamic instability	Hemodynamic instability
PRIMS III	0	3	4	4	7
Total leukocytes/ μ L	9640	5710	11,410	7880	3820
Neutrophils/ μ L	4360	3280	10,510	7150	3160
Lymphocytes/ μ L	4450	2110	320	430	410
PCR mg/dl (0.01–1)	8.2	13	15.88	16.67	11
PCT ng/ml (0.1–0.5)	0.33	3.75	4.28	10.29	1.78
Ferritin ng/mL (7–140)	ND	ND	888	1110	1349
D-dimer mg/L (0–0.5)	ND	ND	3.85	4.22	7.37
IL-6 pg/ml (\leq 7)	ND	ND	63.2	11	85
Chest X-ray on PICU admission	Bilateral peribronchial thickening	No pathological findings	Bilateral pneumonia	No pathological findings	Bilateral pneumonia
Bilateral pneumonia developed while PICU treatment	No	No	Yes	No	Yes
Echocardiogram	Not done	Normal heart function	Normal heart function	Normal heart function	Normal heart function
Maximal respiratory support	BiPAP	Nassal cannula	HFNC	HFNC	HFNC
Inotropic support	No	No	No	No	Yes
Other support	No	No	No	No	No
Broad-spectrum antibiotics because of suspected bacterial coinfection	Yes	Yes	No	No	Yes
Azithromycin	No	No	Yes	Yes	Yes
Lopinavir/ritonavir	No	No	Yes	Yes	Yes
Remdesivir	No	No	No	No	No
Hydroxychloroquine	No	No	Yes	Yes	Yes
Steroids	No	No	Methylprednisolone (1 mg/kg/day)	Methylprednisolone (1 mg/kg/day)	Methylprednisolone (1 mg/kg/day)

Table 1 (Continued)

	Influenza B	Neisseria meningitidis	Case 1	Case 2	Case 3
Immunoglobulins	No	No	No	Yes	No
Tocilizumab	No	No	No	No	Yes
Heparin	No	No	Yes, prophylactic	Yes, prophylactic	Yes, prophylactic
Confirmed coinfection	No	No	No	No	No
Days of PICU admission	5	7	5	6	9

PICU: pediatric intensive care unit; HFNC: high flow nasal cannula; BiPAP: Bilevel Positive Airway Pressure; pSOFA: Pediatric Sequential Organ Failure Assessment; PRISM III: Pediatric Risk of Mortality Score; ND: not done.

CD64, CD11a and CD18 expression in neutrophils, monocytes, and CD8+ lymphocytes




Figure 1 (A) CD64 staining on granulocytes, monocytes, and lymphocytes in peripheral blood samples obtained on pediatric critical care unit (PICU) admission. From left to right, we can observe the CD64 expression. CD64 is expressed on monocytes and neutrophils but not lymphocytes (internal negative control). The positive CD64 region is located to the right of the dotted line. As can be seen, neutrophils gated in the control case are crossed by this line. (B) Mean fluorescence intensity (MFI) values for CD64, CD11a, and CD18 are given for each case in the form of a bar chart. As can be seen in all cases, CD64 and CD11a expression is higher in SARS-CoV-2 cases. This observation is clear for CD64. The CD11a expression on CD8+ lymphocytes is also upregulated compared to previous data.

involved in the process of cytotoxic T cell-mediated killing as well as antibody-mediated killing by granulocytes and monocytes.⁹ The upregulation of both leukocyte populations is also observed in our cases (Fig. 1).

Immunomodulatory treatment seems to have a great role in COVID19. Their use should be based on a risk-benefit analysis. In SARS-CoV-2 infections the cytokine storm theory coupled with analytical data are used to justify these approaches.^{2,10} Our FC results introduce a new approach to

analyzing the immune response to this new virus. We confirm the activation of the innate cellular response. Besides we observed that is different and maybe higher than in other infections. The description of this immune status using FC could individualize the diagnosis and optimize the therapies applied.

In summary, we describe the immunophenotype of three children with severe SARS-CoV-2 infection. We observed significant upregulation of CD64, CD18, and CD11a expression

on leukocytes. Compare to previous papers and to other types of infection it appears to be higher. This could inform about immune dysregulation triggered by SARS-CoV-2. The use of FC may lead to a better understanding of this response and optimize the therapies applies. Prospective studies with a higher number of cases should be conducted to confirm this observation.

Financial disclosure

The authors have no financial relationships relevant to this article to disclose.

Conflict of interest

The authors have no conflicts of interest to disclose.

Bibliografía

1. Tagarro A, Epalza C, Santos M, Sanz-Santaefemia FJ, Otheo E, Moraleda C, et al. Screening and Severity of Coronavirus Disease 2019 (COVID-19) in children in Madrid, Spain. *JAMA Pediatr.* 2020.
2. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of immune response in patients with Coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis.* 2020;71:762–8.
3. Carter MJ, Fish M, Jennings A, Doores KJ, Wellman P, Seow J, et al. Peripheral immunophenotypes in children with multisystem inflammatory syndrome associated with SARS-CoV-2 infection. *Nat Med.* 2020.
4. Hu BQ, Yang Y, Zhao CJ, Liu DF, Kuang F, Zhang LJ, et al. Accuracy of neutrophil CD64 expression in diagnosing infection in patients with autoimmune diseases: a meta-analysis. *Clin Rheumatol.* 2019;38:1319–28.
5. García-Salido A, de Azagra-Garde AM, García-Teresa MA, Caro-Patón GL, Iglesias-Bouzas M, Nieto-Moro M, et al. Accuracy of CD64 expression on neutrophils and monocytes in bacterial infection diagnosis at pediatric intensive care admission. *Eur J Clin Microbiol Infect Dis.* 2019;38:1079–85.
6. Gupta R, Gant VA, Williams B, Enver T. Increased Complement Receptor-3 levels in monocytes and granulocytes distinguish COVID-19 patients with pneumonia from those with mild symptoms. *Int J Infect Dis.* 2020.
7. de Jong E, de Lange DW, Beishuizen A, van de Ven PM, Girbes AR, Huisman A. Neutrophil CD64 expression as a longitudinal biomarker for severe disease and acute infection in critically ill patients. *Int J Lab Hematol.* 2016;38:576–84.
8. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet (London, England).* 2020;395:1054–62.
9. Hyun YM, Choe YH, Park SA, Kim M. LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18) distinctly regulate neutrophil extravasation through hotspots I and II. *Exp Mol Med.* 2019;51:1–13.
10. Zumla A, Hui DS, Azhar EI, Memish ZA, Maeurer M. Reducing mortality from 2019-nCoV: host-directed therapies should be an option. *Lancet (London, England).* 2020;395(10224):e35–6.

A. García-Salido^{a,*}, M.Á. García-Teresa^a, I. Leoz-Gordillo^a, A. Martínez de Azagra-Garde^a, M. Cabrero-Hernández^a, M. Ramirez-Orellana^b

^a *Pediatric Critical Care Unit, Hospital Infantil Universitario Niño Jesús, Madrid, Spain*

^b *Pediatric Oncohematology Unit, Flow Cytometry Laboratory, Hospital Infantil Universitario Niño Jesús, Madrid, Spain*

* Corresponding author.

E-mail address: citopensis@yahoo.es (A. García-Salido).

<https://doi.org/10.1016/j.medin.2020.09.003>

0210-5691/ © 2020 Elsevier España, S.L.U. y SEMICYUC. All rights reserved.

Restrictive or liberal transfusion for cardiac surgery: Spanish results of a randomized multicenter international parallel open-label clinical trial



Transfusión Restrictiva o liberal en cirugía cardíaca: Resultados españoles de un ensayo clínico aleatorizado, multicéntrico, internacional, paralelo y abierto

Dear Editor,

In cardiac surgery, red blood cell (RBC) transfusion is very frequent. Although transfusion is useful to treat anemia and avoid its complications, it represents a potential risk of acute kidney damage at 72 h after surgery, prolonged mechanical ventilation, need for hemodynamic support, increase-hospital mortality, as well as longer hospital stays.¹ Finding the optimal threshold of hemoglobin for indicating

the transfusion with maximal benefit and minimal risk is an aim of the clinical practice. However, a survey in 34 Spanish centers performed in 2007 showed that 70% centers did not have homeostasis protocols and 75% of patients undergoing cardiac surgery were transfused.²

Because of the wide variability in transfusion practices and high rates of transfusion in cardiac surgery in Spain, we participated in the Transfusion Requirements in Cardiac Surgery (TRICS) III trial. This is a randomized, multicentre, international, controlled, open-label clinical trial to assess whether a restrictive transfusion strategy, in which lower hemoglobin concentrations for RBC transfusion, applied throughout the perioperative period, would be non-inferior, in terms of major morbidities and mortality, to a liberal approach among 5243 patients undergoing cardiac surgery.^{3,4} This study was funded by national peer-review organizations from Australia, Canada, Spain (ISCIII and European Social Fund, CP15/00116) and New Zealand.

We present the results of the Spanish included patients followed up to 6 months after surgery.

Ethics approval was provided by each institutional review boards. All patients consented to participate in the clinical