



ORAL PRESENTATION

Open Access

Cancer occurrence during follow-up of the CAPP2 study -aspirin use for up to four years significantly reduces Lynch syndrome cancers for up to several years after completion of therapy

John Burn*, John C Mathers, Anne-Marie Gerdes, MarieLuise Bisgaard, Gareth Evans, Diana Eccles, Annika Lindblom, Findlay Macrae, Eamonn R Maher, Jukka-Pekka Mecklin, Gabriela Moslein, Sylviane Olschwang, Raj Ramesar, Hans FA Vasen, Juul Wijnen, Gail Barker, Faye Elliott, Henry Lynch, D Tim Bishop, the CAPP2 Consortium

From 13th Annual Meeting of the Collaborative Group of the Americas on Inherited Colorectal Cancer Honolulu, Hawaii, USA. 16-17 October 2009

Background/methods

The CAPP2 Study evaluated 600mg enteric coated aspirin and/or 30gms of Novelose (resistant starch) in a double blind factorial RCT in 1071 carriers of Lynch syndrome over a treatment period of 1 to 4 years, mean 29 months.

Results

The trial, reported in December 2008 [1], showed that there was no difference between the treatment and placebo groups for new colorectal neoplasia. Follow-up data for 667 participants for up to 120 months (mean 51m) is now available. Analysis reveals a striking reduction in subsequent cancers; overall, 102 participants have developed 110 Lynch syndrome cancers. Despite equal numbers being randomised to aspirin or placebo, cancer sufferers in the aspirin group are outnumbered 2 to 1. Lifetable analysis for time to first Lynch syndrome cancer reveals a hazard ratio of 0.62(0.41, 0.96) $p=0.03$. There is a clear effect of duration of treatment: <24months on treatment OR 0.90 (0.45, 1.81) $p=0.78$, treated >24 months OR 0.50 (0.28, 0.86) $p=0.01$.

Conclusions

All carriers of Lynch syndrome should consider aspirin chemoprevention. A dose finding study, CAPP3, is

under development. It will compare different doses of aspirin over a 5 to 10 year period.

Funding provided by the UK Medical Research Council, Cancer Research UK, EU Framework and Bayer Corp.

Published: 25 May 2010

Reference

1. Bum J, Bishop DT, Mecklin JP, *et al*: Effect of aspirin or resistant starch on colorectal neoplasia in the Lynch Syndrome. *NEJM* 2008, **359**(24):2567-78.

doi:10.1186/1897-4287-8-S1-O5

Cite this article as: Burn *et al.*: Cancer occurrence during follow-up of the CAPP2 study -aspirin use for up to four years significantly reduces Lynch syndrome cancers for up to several years after completion of therapy. *Hereditary Cancer in Clinical Practice* 2010 **8**(Suppl 1):O5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

