

Are Your Kidneys Ok? Detect Early to Protect Kidney Health



Joseph A. Vassalotti^{1,2,14}, Anna Francis^{3,14}, Augusto Cesar Soares dos Santos Jr⁴, Ricardo Correa-Rotter⁵, Dina Abdellatif⁶, Li-Li Hsiao⁷, Stefanos Roumeliotis⁸, Agnes Haris⁹, Latha A. Kumaraswami¹⁰, Siu-Fai Lui¹¹, Alessandro Balducci¹² and Vassilios Liakopoulos⁸; for the World Kidney Day Joint Steering Committee¹³

¹Mount Sinai Hospital, Department of Medicine-Renal Medicine, New York, New York, USA; ²National Kidney Foundation, Inc., New York, New York, USA; ³Queensland Children's Hospital, Department of Nephrology, South Brisbane, Queensland, Australia; ⁴Faculdade Ciências Médicas de Minas Gerais, Brazil, Hospital das Clínicas, Ebserh, Universidade Federal de Minas Gerais, Brazil; ⁵Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico; ⁶Department of Nephrology, Cairo University Hospital, Cairo, Egypt; ⁷Renal Division, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; ⁸Second Department of Nephrology, American Hellenic Educational Progressive Association (AHEPA) University Hospital Medical School, Aristotle University of Thessaloniki, Thessaloniki, Greece; ⁹Nephrology Department, Péterfy Hospital, Budapest, Hungary; ¹⁰Tamilnad Kidney Research (TANKER) Foundation, Chennai, India; ¹¹Division of Health System, Policy and Management, Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong; and ¹²Italian Kidney Foundation, Rome, Italy

Early identification of kidney disease can protect kidney health, prevent kidney disease progression and related complications, reduce cardiovascular disease risk, and decrease mortality. We must ask “Are your kidneys ok?” using serum creatinine to estimate kidney function and urine albumin to assess for kidney and endothelial damage. Evaluation for causes and risk factors for chronic kidney disease (CKD) includes testing for diabetes and measurement of blood pressure and body mass index. This World Kidney Day we assert that case-finding in high-risk populations, or even population level screening, can decrease the burden of kidney disease globally. Early-stage CKD is asymptomatic and simple to test for, and recent paradigm shifting CKD treatments such as sodium glucose co-transporter-2 inhibitors dramatically improve outcomes and favor the cost-benefit analysis for screening or case-finding programs. Despite this, numerous barriers exist, including resource allocation, health care funding, health care infrastructure, and health care professional and population awareness of kidney disease. Coordinated efforts by major kidney nongovernmental organizations to prioritize the kidney health agenda for governments and aligning early detection efforts with other current programs will maximize efficiencies.

Kidney Int Rep (2025) **10**, 629–636; <https://doi.org/10.1016/j.ekir.2025.01.033>

KEYWORDS: albuminuria; case finding; chronic kidney disease; prevention; proteinuria; screening

© 2025 World Kidney Day Steering Committee. Published by Elsevier Inc. on behalf of the International Society of Nephrology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Timely treatment is the primary strategy to protect kidney health, prevent kidney disease progression and related complications, reduce cardiovascular disease risk, and prevent premature

kidney-related and cardiovascular mortality.^{1–3} International population assessments show low awareness and low detection of kidney disease and substantial gaps in treatment.² People with kidney

failure universally express the preference for having been diagnosed early in their disease trajectory to allow more time for educational, lifestyle, and pharmacologic interventions.⁴ Therefore,

Correspondence: Anna Francis, Queensland Children's Hospital, Department of Nephrology, 501 Stanley Street, South Brisbane QLD 4101, Queensland, Australia. E-mail: anna.francis@uq.edu.au; or Stefanos Roumeliotis, Second Department of Nephrology, American Hellenic Educational Progressive Association (AHEPA) University Hospital, Medical School, Aristotle University of Thessaloniki, 1 St. Kyriakidi str, 54636 Thessaloniki, Greece. E-mail: st_roumeliotis@hotmail.com

This article is being published in *Kidney International* and reprinted concurrently in several journals. The articles cover identical concepts and wording, but vary in minor stylistic and spelling changes, detail, and length of manuscript in keeping with each journal's style. Any of these versions may be used in citing this article.

¹³The World Kidney Day Joint Steering Committee is listed in the [Appendix](#).

¹⁴JAV and AF are co-first authors.

Received 5 November 2024; revised 2 December 2024; accepted 11 December 2024

increasing knowledge and implementing sustainable solutions for early detection of kidney disease to protect kidney health are public health priorities.^{2,3}

Epidemiology and complications of kidney disease

Chronic kidney disease (CKD) is prevalent, affecting 10% of the world's population, or over 700 million people.⁵ Almost 80% of the population with CKD reside in low-income countries (LICs) and lower middle-income countries (LMICs), with approximately one-third of the known affected population living in China and India alone.^{5,6} Prevalence of CKD increased by 33% between 1990 and 2017.⁵ Increasing prevalence of CKD is driven by population growth, aging, and the obesity epidemic, resulting in higher prevalence of 2 major risk factors for CKD: type-2 diabetes (T2DM) and hypertension. In addition, risk factors for CKD beyond cardiometabolic conditions contribute to the rising burden of kidney disease, including social deprivation, pregnancy-related acute kidney injury, preterm birth, and increasing environmental threats (infections, toxins, climate change, air pollution).^{5,7} These threats disproportionately affect people in LICs and LMICs.⁸

Undetected and untreated CKD is more likely to progress to kidney failure and cause premature morbidity and mortality. Globally, more people died in 2019 of cardiovascular disease attributed to reduced kidney function (1.7 million people) than kidney disease alone (1.4 million).⁵ CKD is expected to rise to the fifth most common cause of years of life lost by 2040, surpassing type 2 diabetes, Alzheimer's disease, and road injuries.⁹ The rising mortality of kidney disease is remarkable in contrast to other noncommunicable

diseases, such as cardiovascular disease, stroke, and respiratory disease, which are projected to experience a decline in mortality.⁸ Even in early stage CKD, multi-system morbidity decreases quality of life. In particular, mild cognitive impairment is associated with early stage CKD and it is possible that early CKD detection and treatment could slow cognitive decline and reduce the risk of dementia.¹⁰ CKD in children has profound additional effects, threatening growth and cognitive development and with lifelong health and quality-of-life implications.^{11,12} The number of people on kidney failure replacement therapy—dialysis and transplantation—is anticipated to more than double to 5.4 million from 2010 to 2030.^{13,14} Kidney failure replacement therapy, especially hemodialysis, is unavailable or unaffordable to many in LICs and LMICs, contributing to millions of deaths annually. LICs and LMICs comprise 48% of the global population but account for only 7% of the treated kidney failure population.¹⁵

Who is at risk of kidney disease?

Testing people at high risk for kidney disease (case-finding) limits potential harms and false-positive test results compared with general population screening that should only be considered in high-income countries (HICs). Limiting testing to those at increased risk of CKD would still capture a large proportion of the global population. Moreover, targeted case-finding in patients at high risk of CKD, is not optimally performed even in HICs. About 1 in 3 people worldwide have diabetes and/or hypertension. There is a bidirectional relationship between cardiovascular disease and CKD, with each increasing the risk of the other. The American Heart

Association and European Society of Cardiology call for testing those with cardiovascular disease for CKD, as part of routine cardiovascular assessments.^{1,16}

Other CKD risk factors include family history of kidney disease (e.g., APOL1-mediated kidney disease common in people of West African ancestry), prior acute kidney injury, pregnancy-related kidney disease (e.g., preeclampsia), malignancy, autoimmune disorders (systemic lupus erythematosus, vasculitis), individuals born with low birth weight or preterm, obstructive uropathy, recurrent kidney stones, and congenital anomalies of the kidney and urinary tract (see Figure 1).³ The social determinants of health strongly affect CKD risk, both for individuals and at a country level. In LICs and LMICs, heat stress for agricultural workers is thought to cause CKD of unknown etiology, an increasingly recognized major global cause of CKD.¹⁷ In addition, envenomations, environmental toxins, traditional medicines, and infections (viral hepatitis B or C, HIV, and parasites) deserve consideration as risk groups, especially in endemic areas.^{18,19}

How can we check kidney health?

Conceptually, there are 3 levels of CKD prevention. Primary prevention reduces the incidence of CKD by treating risk factors; secondary prevention reduces progression and complications in those with detected CKD; and tertiary prevention improves outcomes in those with kidney failure by improving management, such as improved vaccination or optimal dialysis delivery.²⁰ Primary and secondary prevention strategies can use the 8 golden rules for kidney health promotion: healthy diet, adequate hydration, physical activity, blood pressure monitoring and control,




Figure 1. Risk factors for chronic kidney disease (CKD).

glycemic monitoring and control, avoidance of nicotine, avoidance of regular use of nonsteroidal anti-inflammatory drugs, and targeted testing for those with risk factors.²¹ Five of these are identical to “Life’s Essential 8” rules for improving and maintaining cardiovascular health that also include healthy weight, adequate sleep, and lipid management.²² Early detection focuses on secondary CKD prevention that involves protecting kidney health and reducing cardiovascular risk.

Are your kidneys okay?

Globally, early detection of CKD is rare, haphazard, and even less likely to occur in LICs or LMICs. Currently, only 3 countries have a national program to actively test for CKD in at-risk populations

and a further 17 countries actively test at-risk population during routine health encounters.²³ Even in HICs, albuminuria is not assessed in over one-half of people with T2DM and/or hypertension.^{24–26} Startlingly, in those with documented reduced kidney function, a diagnosis of CKD is often missing. A study in HICs showed absence of CKD diagnosis among 62%–96% of the population with laboratory evidence of CKD stage G3.²⁷

We recommend that health care professionals perform the following tests for all risk groups to assess kidney health (see [Figure 2](#)):

- Blood pressure measurements because hypertension is the most prevalent risk factor for kidney disease worldwide.^{3,28,29}

- Body mass index because obesity is epidemiologically associated with CKD risk indirectly through T2DM and hypertension and as an independent risk factor. Visceral adiposity contributes to monocyte microinflammation and cardiometabolic kidney risk.^{3,28,29}
- Testing for diabetes with glycosylated hemoglobin or fasting blood sugar or random glucose is part of kidney health assessment as T2DM is a common risk factor.^{3,28,29}
- Evaluating kidney function by using serum creatinine to determine the estimated glomerular filtration rate (eGFR) is recommended in all settings.³ GFR should be estimated with a validated, race-free equation appropriate for the country or




Figure 2. Conceptual framework of chronic kidney disease (CKD) testing, risk stratification, and treatment program. Reprinted with permission from *Kidney International*, Volume 99, Issue 1, Shlipak MG, Tummalaipalli SL, Boulware LE, et al., for Conference Participants. The case for early identification and intervention of chronic kidney disease: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference, 2021:34–47,³⁰ Copyright © 2020, Kidney Disease: Improving Global Outcomes (KDIGO). Published by Elsevier Inc. on behalf of the International Society of Nephrology, under the terms of a CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

region and age group.³ In general, the eGFR <60 ml/min per 1.73 m² is the threshold for CKD in adults and children, although a threshold of <90 ml/min per 1.73 m² can be flagged as “low” in children and adolescents over the age of 2 years.³ A limitation of creatinine-based eGFR is that creatinine is also a marker of nutrition and muscle mass. Therefore, states of malnutrition and frailty overestimate kidney function.^{3,30} Thus, eGFR using the combination of serum creatinine and cystatin C is generally more accurate than either biomarker alone in most clinical contexts. However, the feasibility of cystatin C use is mainly limited to HICs because of assay availability and cost relative to creatinine.^{3,30,31}

- Testing for kidney damage (albuminuria). In adults and children, a first morning sample is preferred for assessing albuminuria.³ In adults, quantitative urinary albumin-creatinine ratio (uACR) is preferred as the most sensitive test.³ Importantly, urinary albumin is in the process of being standardized analytically, which should ultimately facilitate worldwide uACR standardization.³² In children, both urinary protein-creatinine ratio and uACR should be tested to assess tubular proteinuria.³ Semiquantitative albuminuria testing allows for flexibility for point-of-care or home-based testing.³³ Semiquantitative or qualitative screening tests should be positive in >85% of individuals with quantitative uACR 30 mg/g or more to be useful.³⁴ In resource-constrained settings, urine protein dipstick testing may be used with a threshold of +2 proteinuria or greater to reduce false positive results for repeat confirmatory testing.³⁵

In specific populations, the following can be considered:

- Testing for hematuria is notable as the forgotten risk factor in recent clinical practice guidelines. It is particularly important for those at risk for glomerular disease, particularly IgA nephropathy.³⁶
- Baseline imaging in groups with signs or symptoms of structural abnormalities (e.g., pain and hematuria) to evaluate for kidney masses, cysts, stones, hydronephrosis, or urinary retention is important. Antenatal ultrasound can detect hydronephrosis and other congenital anomalies of the kidney and urinary tract.
- With increasing availability of genetic testing, family cascade CKD testing is indicated when there is a known genetic risk of kidney disease.³⁷
- In those who have an occupational risk of developing kidney disease, kidney testing should be offered as part of occupational health programs.
- Those who donate kidneys should be included in a post-donation surveillance program to assess kidney health over the long term.³⁸

Potential benefits of early detection

Screening for CKD fits with many of the Wilson-Jungner principles of the World Health Organization (WHO). Early stage CKD is asymptomatic, and effective treatments, including lifestyle modification, interdisciplinary care, and pharmacologic interventions, need to be implemented.^{2,3,30,35} WHO essential medicines that improve CKD outcomes should be widely available, including angiotensin converting enzyme inhibitors, angiotensin receptor blockers, statins, and sodium glucose co-transporter-2 inhibitors.^{2,39}

Sodium glucose co-transporter-2 inhibitors alone are estimated to decrease the risk of CKD progression by 37% in people with and without diabetes.⁴⁰ For a 50-year-old person with albuminuria and nondiabetic CKD, this could extend their future period of healthy kidney function from 9.6 to 17 years.⁴¹ These essential medicines reduce progression to more advanced CKD stages and limit cardiovascular hospitalization to provide short-term cost-effectiveness, especially for LICs. Where available and affordable, the range of new paradigm-shifting medications to slow CKD progression also includes glucagon-like peptide-1 receptor antagonists, nonsteroidal mineralocorticoid receptor antagonists, endothelin receptor antagonists, and specific disease-modifying drugs (e.g., complement-inhibitors) that herald an exciting new era for nephrology.

Considering the significant health care costs associated with CKD, particularly hospitalization and kidney failure, effective preventive measures offer clear economic benefits for both HICs and LICs. CKD confers enormous costs to the individual, their families, health care systems, and governments worldwide. In the United States, CKD costs Medicare over US\$85 billion annually.¹³ In many HICs and MICs, 2%–4% of the health budget is spent on kidney failure care alone. In Europe, health care costs associated with CKD are higher than those associated with cancer or diabetes.⁴² Reducing the burden of kidney care worldwide will also have profound environmental effects, because it will save water and plastic waste, especially associated with dialysis.⁴³ On an individual level, CKD costs are frequently catastrophic, particularly in LICs and LMICs, where the individual

largely bears the burden of payment. Only 13% of LICs and 19% of LMICs cover the costs of kidney failure replacement therapy for adults.¹⁵ CKD causes 188 million people in LICs and LMICs annually to be faced with catastrophic health care expenditures.⁴⁴

The most widely cited and studied incremental cost-effectiveness ratio threshold to assess screening is <US\$50,000 per quality-adjusted life year.⁴⁵ If the prevalence of CKD is high, a population-wide screening strategy should be considered in HIC.^{33,46} For example, in the United States, a recent Markov simulation model of population-wide screening for CKD, which included appropriate sodium glucose co-transporter-2 inhibitors treatment added to standard of care angiotensin converting enzyme inhibitors or angiotensin receptor blockers for adults age 35 to 75 years old with albuminuria, concluded that screening to identify CKD would be cost-effective.⁴⁶ In addition, an analysis of a home-based general population semi-quantitative albuminuria screening in Holland was also found to be cost-effective.³³ Case-finding to detect CKD in higher risk groups rather than mass or general population screening will reduce costs and other harms while increasing the true-positive rate of the screening tests.^{3,35,45} An alternate incremental cost-effectiveness ratio threshold proposed by WHO is <1–3 times the ratio of the gross domestic product per capita income per quality-adjusted life year and can be used to assess case-finding approaches in LICs and LMICs.⁴⁵ The recommended tests for detecting kidney disease are low cost and minimally invasive, facilitating their administration across diverse settings. Basic testing of eGFR and uACR are widely available, and using urine dipstick testing where quantitative proteinuria testing is

unavailable or unaffordable will drastically reduce testing costs.³¹

If coupled with effective intervention, early identification of people with kidney disease will benefit the individual, the health care system, governments, and the economy.⁴⁴ Health and quality-of-life benefits for the individual would lead to improved productivity, especially in the young with more working years ahead, and to developmental or educational (or both) improvements in children and young adults. Individuals would face fewer catastrophic health expenses; governments and health care systems will save money not only on CKD care, but also on cardiovascular disease costs; and economies will benefit from more worker participation. This is especially crucial for LICs, where the greatest burden of CKD exists and is cruelly coupled with the lowest ability for governments and individuals to afford kidney care.

Challenges and solutions for implementation

Structural barriers to widespread identification and treatment of people with CKD include cost, reliability of testing, and lack of health information systems to track CKD burden. These are underpinned by a lack of relevant government and health care policy, low health care professional knowledge and implementation, poor general population perceived kidney disease risk, and low patient CKD awareness. Solutions for implementing effective interventions include tying CKD identification to existing screening programs, educating the public and primary care professionals, and leveraging nongovernmental organization joint advocacy programs to focus health policy agendas on kidney disease. Any solutions must balance the potential benefits and harms of screening and case-finding

programs. Ethical implications for consideration include the availability of resources (such as health care workers and medicines), the affordability of testing and treatment, false positives or negatives, and anxiety for patients and their families.⁴⁷

Screening and case-finding programs require workforce capacity; health information systems; reliable testing equipment; and equitable access to medical care, medicines, vaccines, and medical technologies. Primary care is at the front lines of the battle to protect kidney health, particularly in LICs and LMICs. The tiny nephrology workforce, with a median global prevalence of 11.8 nephrologists per million population and an 80-fold difference between LICs and HICs, is inadequate to detect and manage the vast majority of CKD.²³ As for other chronic diseases, primary care clinicians and other frontline health workers are foundational to early detection of CKD.⁴⁸ Testing must be affordable, simple, and practical; point-of-care creatinine testing and urine dipsticks are useful in resource-limited settings.³¹ Educational efforts directed at primary care clinicians are key to integrating CKD detection into routine care, despite constrained time and resources.^{49–51} Automated clinical decision support could leverage electronic health records to identify people with CKD or who are at high risk of CKD and recommend appropriate actions to clinicians (Figure 2).

Currently, few countries have CKD registries, limiting our ability to highlight the disease burden to governments. Knowledge of CKD burden assists in prioritizing kidney health needs, which should then progressively expand to encompass the full spectrum of kidney care.⁵² A global survey revealed only a quarter of the countries (41 of 162) had a nation-specific CKD strategy and fewer than a third (48 of 162) recognized

CKD as a public health priority.²³ WHO's recognition of CKD as a major driver of noncommunicable diseases mortality would be impactful in increasing awareness, improving local surveillance, monitoring to implement clinical practice guidelines, and improving resource allocation.²

Programs for the early detection of CKD will require extensive coordination and engagement of stakeholders, including governments, health systems, and insurers. International and national kidney organizations, such as the International Society of Nephrology, already advocate to the WHO and individual governments for the prioritization of kidney disease. We must continue this work, collaborating to streamline early detection program planning and implementation. Connection to existing community interventions (e.g., cardiovascular disease prevention) in LMICs and HICs can decrease cost and maximize efficiencies by integrating into existing programs. Such programs will need to be adapted to the local context and can be held in a variety of settings, such as individual health care practices, hospitals, as well as regional or national health care facilities or as outreaches in rural communities. Depending on local regulations and resources, screening and case-finding can also take place outside of medical settings such as town halls, churches, or markets. Volunteers in the community can also assist with community-based screening and case-finding efforts.

In conjunction with reorienting the clinical practice of health care professionals to a greater focus on timely detection of CKD, we must focus on general population perceived risk education and health promotion activities, as well as education programs aimed at patient awareness and

Box 1. Are your kidneys okay? Person perspectives on CKD awareness, detection, and treatment from the literature, see Guha *et al.*⁴ and Tuot *et al.*⁵⁷

I actually didn't fully understand because nobody had actually given me the full information of what I had in a way that I could kind of go, "Well this is what I've got [CKD], and this is why I've got it."

[The clinicians] they can answer those [kidney health] questions, . . . but it's all very jargonistic.

I didn't know what it [CKD] meant so I couldn't really share it with other people.

I may not know what my [kidney health] numbers are, but I do know what the tests are, and I do know that I've had them done before.

Well, let me put this way: I'm now well aware now of the significance of the kidneys and about what the issues are here. And I would definitely consider . . . When I go to the doctor, I would say to him, "Now, listen. You did the blood tests. But how are my kidneys doing? What are the numbers?"

I know that they have done urine tests in the past, and I know protein and sugar was in my urine.

I went from never taking a tablet to taking 22 tablets. What going on here? I didn't know what they were. But I just number them and that did help me a lot because I realized what was going on but some of them, every time I went there [to see the doctor], I'd get another tablet. I knew that I had to take it because they knew what they were doing, the doctors that I went to see.

This [CKD] is something new, so immediately I was like, just another thing to be concerned about. But then I felt kind of empowered, and like I really do want to get ahead of this thing. I feel like I do want to have a conversation with my primary care physician.

What I would be mostly interested in is what is happening, why is it happening, and what can I do to slow it [CKD] down?

CKD, chronic kidney disease.

empowerment. General population awareness of kidney disease is poor, with 9 of 10 people with CKD unaware they are affected.⁵³ Coverage of kidney disease is missing from the mainstream conversation, with an analysis of lay press showing kidney disease was 11-times underrepresented in discussed compared to the actual cause of death.⁵⁴ A number of national and international organizations have developed public-facing quizzes on risk of kidney disease, supported by a regional study that showed socially vulnerable patients with hypertension do not understand their kidney risks.^{21,55–57} Online and direct education for health care professionals can improve consumer health literacy. Patient activation, engagement, and shared decision-making are downstream impacts of awareness. Awareness education is nuanced for CKD, including detection and risk stratification to inform and empower rather than frighten regarding the timing and extent of interventions (see [Box 1](#)).^{4,27,57} Getting the balance right will optimize self-efficacy and patient, family, and caregiver engagement.

Conclusion: a call to action

We call on all health care professionals to check the kidney health of their patients at risk of kidney disease. In tandem, we must work with public health organizations to improve the general population's perceived risk of kidney disease and empower people at risk to seek kidney health checks. To ensure this change can be delivered, we must work with health care systems, governments, and the WHO to prioritize kidney disease and create effective and efficient early detection programs for kidney disease. Only then will the paradigm-shifting benefits of lifestyle change and pharmacologic treatments translate to better kidney and overall health for people all around the world.

APPENDIX

The World Kidney Day Joint Steering Committee includes Valerie A. Luyckx, Marcello Tonelli, Ifeoma Ulasi, Vivekanand Jha, Marina Wainstein, Siddiq Anwar, Daniel O'Hara, Elliot K. Tannor, Jorge Cerda, Elena Cervantes, and María Carlota González.

DISCLOSURE

All the authors declared no competing interests.

ACKNOWLEDGMENTS

We thank Valerie A. Luyckx, Marcello Tonelli, Ifeoma Ulasi, Vivekanand Jha, Marina Wainstein, Siddiq Anwar, Daniel O'Hara, Elliot K. Tan-nor, Jorge Cerda, Elena Cervantes, and María Carlota González Bedat for their invaluable feedback on this paper.

REFERENCES

- Ndumele CE, Neeland IJ, Tuttle KR, et al. A synopsis of the evidence for the science and clinical management of cardiovascular-kidney-metabolic (CKM) syndrome: a scientific statement from the American Heart Association. *Circulation*. 2023;148:1636–1664.
- Luyckx VA, Tuttle KR, Abdellatif D, et al. World Kidney Day Joint Steering Committee. Mind the gap in kidney care: translating what we know into what we do. *Kidney Int*. 2024;105:406–417.
- Stevens PE, Ahmed SB, Carrero JJ, et al. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int*. 2024;105:S117–S314.
- Guha C, Lopez-Vargas P, Ju A, et al, PAVE-CKD Workshop Investigators. Patient needs and priorities for patient navigator programmes in chronic kidney disease: a workshop report. *BMJ Open*. 2020;10:e040617.
- GBD Chronic Kidney Disease Collaboration. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2020;395:709–733.
- Cojuc-Konigsberg G, Guijosa A, Moscona-Nissan A, et al. Representation of low- and middle-income countries in CKD drug trials: a systematic review. *Am J Kidney Dis*. 2025;85:55–66.e1.
- Hsiao LL, Shah KM, Liew A, et al, World Kidney Day Joint Steering Committee. Kidney health for all: preparedness for the unexpected in supporting the vulnerable. *Kidney Int*. 2023;103:436–443.
- Francis A, Harhay MN, Ong ACM, et al. Chronic kidney disease and the global public health agenda: an international consensus. *Nat Rev Nephrol*. 2024;20:473–485.
- Foreman KJ, Marquez N, Dolgert A, et al. Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes of death: reference and alternative scenarios for 2016–40 for 195 countries and territories. *Lancet*. 2018;392:2052–2090.
- Viggiano D, Wagner CA, Martino G, et al. Mechanisms of cognitive dysfunction in CKD. *Nat Rev Nephrol*. 2020;16:452–469.
- Chen K, Didsbury M, van Zwielen A, et al. Neurocognitive and educational outcomes in children and adolescents with CKD: a systematic review and meta-analysis. *Clin J Am Soc Nephrol*. 2018;13:387–397.
- Francis A, Didsbury MS, van Zwielen A, et al. Quality of life of children and adolescents with chronic kidney disease: a cross-sectional study. *Arch Dis Child*. 2019;104:134–140.
- United States Renal Data System. 2023 USRDS Annual Data Report: Epidemiology of Kidney Disease in the United States. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases; 2023.
- Liyanage T, Ninomiya T, Jha V, et al. Worldwide access to treatment for end-stage kidney disease: a systematic review. *Lancet*. 2015;385:1975–1982.
- Bello AK, Levin A, Tonelli M, et al. Assessment of global kidney health care status. *JAMA*. 2017;317:1864–1881.
- Ortiz A, Wanner C, Gansevoort R, ERA Council. Chronic kidney disease as cardiovascular risk factor in routine clinical practice: a position statement by the Council of the European Renal Association. *Nephrol Dial Transplant*. 2023;38:527–531.
- Johnson RJ, Wesseling C, Newman LS. Chronic kidney disease of unknown cause in agricultural communities. *N Engl J Med*. 2019;380:1843–1852.
- McCulloch M, Luyckx VA, Cullis B, et al. Challenges of access to kidney care for children in low-resource settings. *Nat Rev Nephrol*. 2021;17:33–45.
- Stanifer JW, Muir A, Jafar TH, Patel UD. Chronic kidney disease in low- and middle-income countries. *Nephrol Dial Transplant*. 2016;31:868–874.
- Levey AS, Schoolwerth AC, Burrows NR, et al. Comprehensive public health strategies for preventing the development, progression, and complications of CKD: report of an expert panel convened by the Centers for Disease Control and Prevention. *Am J Kidney Dis*. 2009;53:522–535.
- International Society of Nephrology. World Kidney Day 2025. Accessed January 11, 2025. <https://www.worldkidneyday.org/about-kidney-health/>
- Lloyd-Jones DM, Allen NB, Anderson CAM, et al. Life's Essential 8: updating and enhancing the American Heart Association's Construct of Cardiovascular Health: a presidential advisory from the American Heart Association. *Circulation*. 2022;146:e18–e43.
- Bello AK, Okpechi IG, Levin A, et al, ISN-GKHA Group. An update on the global disparities in kidney disease burden and care across world countries and regions. *Lancet Glob Health*. 2024;12:e382–e395.
- Ferre S, Storfer-Isser A, Kinderknecht K, et al. Fulfillment and validity of the kidney health evaluation measure for people with diabetes. *Mayo Clin Proc Innov Qual Outcomes*. 2023;7:382–391.
- Alfego D, Ennis J, Gillespie B, et al. Chronic kidney disease testing among at-risk adults in the U.S. remains low: real-world evidence from a national laboratory database. *Diabetes Care*. 2021;44:2025–2032.
- Stempniewicz N, Vassalotti JA, Cuddeback JK, et al. Chronic kidney disease testing among primary care patients with type 2 diabetes across 24 U.S. health care organizations. *Diabetes Care*. 2021;44:2000–2009.
- Kushner PR, DeMeis J, Stevens P, et al. Patient and clinician perspectives: to create a better future for chronic kidney disease, we need to talk about our kidneys. *Adv Ther*. 2024;41:1318–1324.

28. Farrell DR, Vassalotti JA. Screening, identifying, and treating chronic kidney disease: why, who, when, how, and what? *BMC Nephrol.* 2024;25:34.
29. Tuttle KR. CKD screening for better kidney health: Why? Who? How? When? *Nephrol Dial Transplant.* 2024;39:1537–1539.
30. Shlipak MG, Tummalapalli SL, Boulware LE, et al. The case for early identification and intervention of chronic kidney disease: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int.* 2021;99:34–47.
31. Tummalapalli SL, Shlipak MG, Damster S, et al. Availability and affordability of kidney health laboratory tests around the globe. *Am J Nephrol.* 2020;51:959–965.
32. Seegmiller JC, Bachmann LM. Urine albumin measurements in clinical diagnostics. *Clin Chem.* 2024;70:382–391.
33. van Mil D, Kieneker LM, Heerspink HJL, Gansevoort RT. Screening for chronic kidney disease: change of perspective and novel developments. *Curr Opin Nephrol Hypertens.* 2024;33:583–592.
34. Sacks DB, Arnold M, Bakris GL, et al. Guidelines and recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus. *Clin Chem.* 2023;69(8):808–868.
35. Tonelli M, Dickinson JA. Early detection of CKD: implications for low-income, middle-income, and high-income countries. *J Am Soc Nephrol.* 2020;31:1931–1940.
36. Moreno JA, Martin-Cleary C, Gutierrez E, et al. Haematuria: the forgotten CKD factor? *Nephrol Dial Transplant.* 2012;27(1):28–34.
37. Franceschini N, Feldman DL, Berg JS, et al. Advancing genetic testing in kidney diseases: report from a National Kidney Foundation Working Group. *Am J Kidney Dis.* 2024;84:751–766.
38. Mjøs G, Hallan S, Hartmann A, et al. Long-term risks for kidney donors. *Kidney Int.* 2014;86:162–167.
39. Francis A, Abdul Hafidz MI, Ekrikpo UE, et al. Barriers to accessing essential medicines for kidney disease in low- and lower middle-income countries. *Kidney Int.* 2022;102:969–973.
40. Nuffield Department of Population Health Renal Studies Group. SGLT2 Inhibitor Meta-Analysis Cardio-Renal Trialists' Consortium. Impact of diabetes on the effects of sodium glucose co-transporter-2 inhibitors on kidney outcomes: collaborative meta-analysis of large placebo-controlled trials. *Lancet.* 2022;400(10365):1788–1801.
41. Vart P, Vaduganathan M, Jongs N, et al. Estimated lifetime benefit of combined RAAS and SGLT2 inhibitor therapy in patients with albuminuric CKD without diabetes. *Clin J Am Soc Nephrol.* 2022;17:1754–1762.
42. Vanholder R, Annemans L, Brown E, et al. European Kidney Health Alliance. Reducing the costs of chronic kidney disease while delivering quality health care: a call to action. *Nat Rev Nephrol.* 2017;13:393–409.
43. Berman-Parks N, Berman-Parks I, Gómez-Ruiz IA, et al. Combining patient care and environmental protection: a pilot program recycling polyvinyl chloride from automated peritoneal dialysis waste. *Kidney Int Rep.* 2024;9:1908–1911.
44. Essue BM, Laba M, Knaul F, et al. Economic burden of chronic ill health and injuries for households in low- and middle-income countries. In: Jamison DT, Gelband H, Horton S, Jha P, Laxminarayan R, Mock CN, et al., eds. *Disease Control Priorities: Improving Health and Reducing Poverty*. The International Bank for Reconstruction and Development; 2018 and The World Bank; 2017.
45. Yeo SC, Wang H, Ang YG, et al. Cost-effectiveness of screening for chronic kidney disease in the general adult population: a systematic review. *Clin Kidney J.* 2024;17:sfad137.
46. Cusick MM, Tisdale RL, Chertow GM, et al. Population-wide screening for chronic kidney disease : a cost-effectiveness analysis. *Ann Intern Med.* 2023;176:788–797.
47. Yadla M, John P, Fong VK, Anandh U. Ethical issues related to early screening programs in low resource settings. *Kidney Int Rep.* 2024;9:2315–2319.
48. Szczech LA, Stewart RC, Su HL, et al. Primary care detection of chronic kidney disease in adults with type-2 diabetes: the ADD-CKD Study (Awareness, Detection and Drug Therapy in Type 2 Diabetes and Chronic Kidney Disease). *PLoS One.* 2014;9:e110535.
49. Vassalotti JA, Centor R, Turner BJ, et al. National Kidney Foundation Kidney Disease Outcomes Quality Initiative. Practical approach to detection and management of chronic kidney disease for the primary care clinician. *Am J Med.* 2016;129(2):153–162.e7.
50. Thavarajah S, Knically DH, Choi MJ. CKD for primary care practitioners: can we cut to the chase without too many shortcuts? *Am J Kidney Dis.* 2016;67:826–829.
51. Vassalotti JA, Boucree SC. Integrating CKD Into US primary care: bridging the knowledge and implementation gaps. *Kidney Int Rep.* 2022;7:389–396.
52. Luyckx VA, Moosa MR. Priority setting as an ethical imperative in managing global dialysis access and improving kidney care. *Semin Nephrol.* 2021;41:230–241.
53. US Centers for Disease Control and Prevention. Chronic kidney disease in the United States 2023. Accessed January 11, 2025. <https://www.cdc.gov/kidney-disease/php/data-research/index.html>
54. Dattani S, Spooner F, Ritchie H, Roser M. Causes of death. 2023. Accessed January 11, 2025. <https://ourworldindata.org/causes-of-death>
55. Boulware LE, Carson KA, Troll MU, et al. Perceived susceptibility to chronic kidney disease among high-risk patients seen in primary care practices. *J Gen Intern Med.* 2009;24:1123–1129.
56. National Kidney Foundation. Kidney quiz 2024. Accessed January 11, 2025. <https://www.kidney.org/kidney-quiz/>
57. Tuot DS, Crowley ST, Katz LA, et al. Usability testing of the kidney score platform to enhance communication about kidney disease in primary care settings: qualitative think-aloud study. *JMIR Form Res.* 2022;6:e40001.