

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(1-Allyl-1H-indazol-5-yl)-4-methylbenzenesulfonamide

 Hakima Chicha,^a El Mostapha Rakib,^a Hafid Abderrafia,^{a*} Mohamed Saadi^b and Lahcen El Ammari^b
^aLaboratoire de Chimie Organique et Analytique, Université Sultan Moulay Slimane, Faculté des Sciences et Techniques, Béni-Mellal, BP 523, Morocco, and ^bLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed

V-Agdal, Avenue Ibn Battouta, BP. 1014, Rabat, Morocco

Correspondence e-mail: h_abderrafia@yahoo.fr


Received 12 November 2013; accepted 25 November 2013

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.042; wR factor = 0.120; data-to-parameter ratio = 16.4.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$, contains two independent molecules linked by an $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond. The molecules show different conformations. In the first molecule, the fused five- and six-membered ring system is almost perpendicular to the plane through the atoms forming the allyl group, as indicated by the dihedral angle of $85.1(4)^\circ$. The dihedral angle with the methylbenzenesulfonamide group is $78.8(1)^\circ$. On the other hand, in the second molecule, the dihedral angles between the indazole plane and the allyl and methylbenzenesulfonamide groups are $80.3(3)$ and $41.5(1)^\circ$, respectively. In the crystal, molecules are further linked by $\text{N}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For the biological activity of sulfonamides, see: Bouissane *et al.* (2006); El-Sayed *et al.* (2011); Mustafa *et al.* (2012). For similar compounds, see: Abbassi *et al.* (2012, 2013); Chicha *et al.* (2013).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$
 $M_r = 327.40$

 Triclinic, $P\bar{1}$
 $a = 8.8200(4)$ Å

 $b = 10.4769(5)$ Å

 $c = 19.7407(10)$ Å

 $\alpha = 80.211(1)^\circ$
 $\beta = 78.984(1)^\circ$
 $\gamma = 69.784(1)^\circ$
 $V = 1669.51(14)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.21$ mm⁻¹
 $T = 296$ K

 $0.42 \times 0.35 \times 0.28$ mm

Data collection

Bruker X8 APEX diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2009)

 $T_{\min} = 0.693$, $T_{\max} = 0.747$

31703 measured reflections

6791 independent reflections

 5368 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.120$
 $S = 1.02$

6791 reflections

415 parameters

H-atom parameters constrained

 $\Delta\rho_{\max} = 0.32$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N6}-\text{H6N}\cdots\text{O1}$	0.79	2.11	2.900 (2)	176
$\text{N3}-\text{H3N}\cdots\text{N5}^i$	0.80	2.19	2.983 (2)	175
$\text{C21}-\text{H21}\cdots\text{O4}^{ii}$	0.93	2.49	3.245 (2)	138
$\text{C7}-\text{H7}\cdots\text{O2}^{iii}$	0.93	2.48	3.358 (2)	158

 Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x, -y + 2, -z + 2$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: PLATON (Spek, 2009) and publCIF (Westrip, 2010).

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2445).

References

- Abbassi, N., Chicha, H., Rakib, E. M., Hannioui, A., Alaoui, M., Hajjaji, A., Geffken, D., Aiello, C., Gangemi, R., Rosano, C. & Viale, M. (2012). *Eur. J. Med. Chem.* **57**, 240–249.
- Abbassi, N., Rakib, E. M., Hannioui, A., Saadi, M. & El Ammari, L. (2013). *Acta Cryst.* **E69**, o190–o191.
- Bouissane, L., El Kazzouli, S., Léonce, S., Pfeiffer, B., Rakib, E. M., Khouili, M. & Guillaume, G. (2006). *Bioorg. Med. Chem.* **14**, 1078–1088.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chicha, H., Kouakou, A., Rakib, E. M., Saadi, M. & El Ammari, L. (2013). *Acta Cryst.* **E69**, o1353.
- El-Sayed, N. S., El-Bendary, E. R., El-Ashry, S. M. & El-Kerdawy, M. M. (2011). *Eur. J. Med. Chem.* **46**, 3714–3720.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Mustafa, G., Khan, I. U., Ashraf, M., Afzal, I., Shahzad, S. A. & Shafiq, M. (2012). *Bioorg. Med. Chem.* **20**, 2535–2539.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2013). E69, o1847 [doi:10.1107/S1600536813032091]

***N*-(1-Allyl-1*H*-indazol-5-yl)-4-methylbenzenesulfonamide**

Hakima Chicha, El Mostapha Rakib, Hafid Abderrafia, Mohamed Saadi and Lahcen El Ammari

1. Comment

Sulfonamides are an important class of compounds which are widely used in the design of diverse classes of drug candidates. These compounds exhibit a wide range of biological activities such as anticancer, anti-inflammatory, and antiviral functions (Bouissane, *et al.*, 2006; El-Sayed, *et al.*, 2011; Mustafa, *et al.* 2012). The present work is a continuation of the investigation of sulfonamides derivatives published recently by our team (Abbassi, *et al.*, 2012; Abbassi, *et al.*, 2013; Chicha, *et al.*, 2013).

The two independent molecules forming the asymmetric unit of the title compound are linked by a weak hydrogen bond (C13–H13···O4) and have different conformations as shown in Fig.1. Each molecule is built up from fused five- and six-membered rings linked to a methylbenzenesulfonamide and allyl groups. In the first molecule, the indazole ring system makes dihedral angles of 78.8 (1)° and 85.1 (4)° with the plane through the methylbenzenesulfonamide group and that through the allyl group, respectively. In the second molecule, the indazole system is almost perpendicular to the plane through the atoms forming the allyl group, as indicated by the dihedral angle of 80.3 (3)°. The dihedral angle between the indazole ring and the benzene ring belonging to the methylbenzenesulfonamide is 41.5 (1)°. In addition, the most important difference between the two conformations of molecules is the orientation of their allyl substituents (Fig.2). In the crystal, molecules are linked by N–H···N, N–H···O and C–H···O hydrogen bonds, forming a three-dimensional network (Table 1).

2. Experimental

A mixture of 1-allyl-5-nitroindazole (250 mg, 1.22 mmol) and anhydrous SnCl₂ (1.1 g, 6.1 mmol) in 25 ml of absolute ethanol was heated to 333 K for 6 h. After reduction, the starting material disappeared, and the solution was allowed to cool down. The pH was made slightly basic (pH 7–8) by addition of 5% aqueous potassium bicarbonate before extraction with ethyl acetate. The organic phase was washed with brine and dried over magnesium sulfate. The solvent was removed to afford the amine, which was immediately dissolved in pyridine (5 ml) and then reacted with 4-methylbenzenesulfonyl chloride (240 mg, 1.25 mmol) at room temperature for 24 h. After the reaction mixture was concentrated *in vacuo*, the resulting residue was purified by flash chromatography (eluted with ethyl acetate: hexane 1:9). The title compound was recrystallized from ethanol at room temperature (yield: 75%, m.p. 374 K).


3. Refinement

H atoms were located in a difference map and treated as riding with N–H = 0.89 Å, C–H = 0.96 Å (methyl), C–H = 0.97 Å (methylene) and C–H = 0.93 Å (aromatic CH, terminal =CH₂), respectively. All hydrogen atoms were refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$ for methyl groups and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$ for all other hydrogen atoms.

Values of atomic displacements of C1, C2, C14 and C15 which belong to the first molecule (S1, O1, O2, N1 - N3, C1 - C17) and are larger than those observed in the second molecule for corresponding carbons.

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

**Figure 1**

The two molecules building the asymmetric unit of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small circles.




Figure 2

Automatic fit of the two crystallographically independent molecules.

N-(1-Allyl-1*H*-indazol-5-yl)-4-methylbenzenesulfonamide

Crystal data

$C_{17}H_{17}N_3O_2S$

$M_r = 327.40$

Triclinic, $P\bar{1}$

Hall symbol: $-p\ 1$

$a = 8.8200$ (4) Å

$b = 10.4769$ (5) Å

$c = 19.7407$ (10) Å

$\alpha = 80.211$ (1)°

$\beta = 78.984$ (1)°

$\gamma = 69.784$ (1)°

$V = 1669.51$ (14) Å³

$Z = 4$

$F(000) = 688$

$D_x = 1.303$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6791 reflections

$\theta = 2.4$ – 26.4 °

$\mu = 0.21$ mm⁻¹

$T = 296$ K

Block, colourless

$0.42 \times 0.35 \times 0.28$ mm

Data collection

Bruker X8 APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2009)

$T_{\min} = 0.693$, $T_{\max} = 0.747$

31703 measured reflections

6791 independent reflections

5368 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 26.4$ °, $\theta_{\min} = 2.4$ °

$h = -10 \rightarrow 11$

$k = -13 \rightarrow 13$

$l = -24 \rightarrow 24$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.120$

$S = 1.02$

6791 reflections

415 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0594P)^2 + 0.4278P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against all reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on all data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.5331 (4)	1.4103 (5)	1.1048 (2)	0.1376 (14)
H1A	-0.4762	1.3460	1.1375	0.165*
H1B	-0.5362	1.5009	1.1007	0.165*
C2	-0.6054 (3)	1.3750 (3)	1.06626 (16)	0.0943 (8)
H2	-0.6610	1.4419	1.0341	0.113*
C3	-0.6085 (3)	1.2319 (3)	1.06826 (13)	0.0859 (7)
H3A	-0.5292	1.1694	1.0966	0.103*
H3B	-0.7156	1.2276	1.0894	0.103*
C4	-0.6237 (3)	1.1553 (3)	0.90373 (15)	0.0899 (8)
H4	-0.6796	1.1473	0.8701	0.108*
C5	-0.4527 (3)	1.1281 (2)	0.89489 (11)	0.0631 (5)
C6	-0.4237 (3)	1.1507 (2)	0.95842 (10)	0.0567 (5)
C7	-0.2677 (3)	1.1323 (2)	0.97123 (10)	0.0626 (5)
H7	-0.2500	1.1460	1.0140	0.075*
C8	-0.1409 (3)	1.0934 (2)	0.91886 (9)	0.0561 (5)
H8	-0.0355	1.0816	0.9260	0.067*
C9	-0.1670 (2)	1.07082 (18)	0.85424 (9)	0.0483 (4)
C10	-0.3201 (3)	1.0860 (2)	0.84226 (10)	0.0623 (5)
H10	-0.3363	1.0688	0.8000	0.075*
C11	0.0714 (2)	0.7588 (2)	0.81912 (10)	0.0537 (4)
C12	-0.0059 (3)	0.7344 (2)	0.77026 (12)	0.0680 (6)
H12	-0.0317	0.7982	0.7317	0.082*
C13	-0.0443 (3)	0.6161 (3)	0.77871 (15)	0.0810 (7)
H13	-0.0977	0.6011	0.7460	0.097*
C14	-0.0053 (3)	0.5185 (2)	0.83494 (16)	0.0806 (7)
C15	0.0682 (3)	0.5455 (3)	0.88345 (15)	0.0874 (8)

H15	0.0935	0.4814	0.9219	0.105*
C16	0.1063 (3)	0.6655 (2)	0.87706 (12)	0.0717 (6)
H16	0.1543	0.6826	0.9111	0.086*
C17	-0.0423 (4)	0.3862 (3)	0.8436 (2)	0.1211 (12)
H17A	-0.1549	0.4054	0.8394	0.182*
H17B	-0.0214	0.3383	0.8886	0.182*
H17C	0.0258	0.3305	0.8084	0.182*
C18	-0.0239 (3)	0.5553 (3)	0.40309 (16)	0.0876 (8)
H18A	-0.1103	0.5665	0.4393	0.105*
H18B	0.0332	0.4688	0.3894	0.105*
C19	0.0165 (3)	0.6589 (2)	0.37214 (12)	0.0632 (5)
H19	0.1036	0.6438	0.3362	0.076*
C20	-0.0659 (2)	0.80224 (19)	0.38962 (10)	0.0522 (4)
H20A	-0.1273	0.8559	0.3527	0.063*
H20B	-0.1423	0.8019	0.4320	0.063*
C21	0.2282 (2)	0.96925 (19)	0.36555 (9)	0.0511 (4)
H21	0.2911	1.0194	0.3383	0.061*
C22	0.2391 (2)	0.91991 (17)	0.43648 (8)	0.0428 (4)
C23	0.3346 (2)	0.92338 (17)	0.48490 (8)	0.0442 (4)
H23	0.4118	0.9684	0.4729	0.053*
C24	0.3107 (2)	0.85819 (16)	0.55050 (8)	0.0419 (4)
C25	0.1910 (2)	0.79282 (18)	0.56925 (9)	0.0484 (4)
H25	0.1768	0.7507	0.6143	0.058*
C26	0.0952 (2)	0.78944 (18)	0.52319 (9)	0.0488 (4)
H26	0.0158	0.7467	0.5361	0.059*
C27	0.1216 (2)	0.85298 (16)	0.45583 (8)	0.0407 (4)
C28	0.5359 (2)	0.58393 (18)	0.63513 (9)	0.0463 (4)
C29	0.5396 (2)	0.5001 (2)	0.58778 (10)	0.0552 (5)
H29	0.5773	0.5188	0.5411	0.066*
C30	0.4874 (3)	0.3886 (2)	0.60989 (11)	0.0619 (5)
H30	0.4889	0.3330	0.5775	0.074*
C31	0.4327 (2)	0.3571 (2)	0.67903 (11)	0.0590 (5)
C32	0.4335 (3)	0.4411 (2)	0.72592 (11)	0.0726 (6)
H32	0.3994	0.4207	0.7728	0.087*
C33	0.4837 (3)	0.5539 (2)	0.70484 (10)	0.0665 (6)
H33	0.4825	0.6095	0.7371	0.080*
C34	0.3732 (3)	0.2362 (3)	0.70195 (15)	0.0841 (7)
H34A	0.2562	0.2675	0.7115	0.126*
H34B	0.4174	0.1867	0.7432	0.126*
H34C	0.4081	0.1772	0.6658	0.126*
N1	-0.5718 (2)	1.18978 (19)	0.99908 (10)	0.0694 (5)
N2	-0.6939 (2)	1.1936 (2)	0.96546 (12)	0.0855 (6)
N3	-0.03312 (19)	1.03964 (16)	0.79882 (7)	0.0523 (4)
H3N	-0.0594	1.0518	0.7611	0.063*
N4	0.05029 (18)	0.86568 (15)	0.39871 (7)	0.0470 (3)
N5	0.1175 (2)	0.93510 (16)	0.34324 (7)	0.0521 (4)
N6	0.40966 (19)	0.85977 (15)	0.60082 (7)	0.0487 (4)
H6N	0.3615	0.8700	0.6388	0.058*
O1	0.22363 (18)	0.91011 (16)	0.73713 (7)	0.0677 (4)

O2	0.20026 (17)	0.91093 (16)	0.86333 (7)	0.0688 (4)
O3	0.67127 (17)	0.72657 (15)	0.53792 (8)	0.0677 (4)
O4	0.6511 (2)	0.76466 (16)	0.66010 (8)	0.0748 (4)
S1	0.12954 (6)	0.90722 (5)	0.80483 (2)	0.05223 (14)
S2	0.58344 (6)	0.73592 (5)	0.60662 (2)	0.05151 (14)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.092 (2)	0.163 (4)	0.149 (3)	-0.015 (2)	0.001 (2)	-0.071 (3)
C2	0.0788 (17)	0.105 (2)	0.0888 (19)	-0.0081 (15)	0.0001 (15)	-0.0433 (17)
C3	0.0829 (16)	0.099 (2)	0.0613 (14)	-0.0196 (14)	0.0163 (12)	-0.0202 (13)
C4	0.0640 (14)	0.117 (2)	0.0809 (17)	-0.0096 (14)	-0.0194 (13)	-0.0214 (16)
C5	0.0589 (12)	0.0669 (13)	0.0564 (12)	-0.0073 (10)	-0.0145 (9)	-0.0089 (10)
C6	0.0649 (12)	0.0509 (11)	0.0464 (10)	-0.0117 (9)	-0.0036 (9)	-0.0046 (8)
C7	0.0738 (13)	0.0749 (14)	0.0397 (10)	-0.0214 (11)	-0.0072 (9)	-0.0148 (9)
C8	0.0629 (11)	0.0668 (12)	0.0418 (10)	-0.0212 (10)	-0.0109 (8)	-0.0104 (9)
C9	0.0598 (11)	0.0455 (10)	0.0361 (9)	-0.0112 (8)	-0.0090 (8)	-0.0054 (7)
C10	0.0628 (12)	0.0757 (14)	0.0446 (10)	-0.0091 (10)	-0.0179 (9)	-0.0141 (9)
C11	0.0500 (10)	0.0561 (11)	0.0461 (10)	-0.0064 (8)	-0.0039 (8)	-0.0082 (8)
C12	0.0773 (14)	0.0674 (14)	0.0587 (12)	-0.0190 (11)	-0.0152 (11)	-0.0090 (10)
C13	0.0810 (16)	0.0753 (16)	0.0906 (18)	-0.0244 (13)	-0.0089 (14)	-0.0253 (14)
C14	0.0586 (13)	0.0602 (14)	0.109 (2)	-0.0098 (11)	0.0117 (13)	-0.0202 (14)
C15	0.0765 (16)	0.0666 (15)	0.0922 (19)	-0.0075 (13)	-0.0019 (14)	0.0186 (13)
C16	0.0676 (13)	0.0738 (15)	0.0627 (13)	-0.0123 (11)	-0.0157 (11)	0.0060 (11)
C17	0.096 (2)	0.0656 (17)	0.186 (4)	-0.0242 (15)	0.022 (2)	-0.0225 (19)
C18	0.0931 (18)	0.0565 (14)	0.113 (2)	-0.0224 (13)	-0.0220 (16)	-0.0035 (13)
C19	0.0685 (13)	0.0618 (13)	0.0663 (13)	-0.0244 (10)	-0.0145 (10)	-0.0131 (10)
C20	0.0522 (10)	0.0506 (10)	0.0589 (11)	-0.0191 (8)	-0.0206 (9)	-0.0003 (8)
C21	0.0652 (11)	0.0579 (11)	0.0378 (9)	-0.0308 (9)	-0.0112 (8)	0.0015 (8)
C22	0.0522 (9)	0.0408 (9)	0.0368 (8)	-0.0165 (7)	-0.0081 (7)	-0.0034 (7)
C23	0.0528 (10)	0.0440 (9)	0.0401 (9)	-0.0199 (8)	-0.0102 (7)	-0.0032 (7)
C24	0.0502 (9)	0.0379 (8)	0.0354 (8)	-0.0087 (7)	-0.0091 (7)	-0.0069 (6)
C25	0.0555 (10)	0.0491 (10)	0.0356 (9)	-0.0142 (8)	-0.0053 (7)	0.0016 (7)
C26	0.0500 (10)	0.0489 (10)	0.0469 (10)	-0.0193 (8)	-0.0063 (8)	0.0025 (8)
C27	0.0449 (9)	0.0374 (8)	0.0393 (8)	-0.0107 (7)	-0.0093 (7)	-0.0050 (7)
C28	0.0499 (10)	0.0464 (9)	0.0412 (9)	-0.0119 (8)	-0.0129 (7)	-0.0018 (7)
C29	0.0680 (12)	0.0533 (11)	0.0410 (9)	-0.0129 (9)	-0.0123 (9)	-0.0054 (8)
C30	0.0750 (13)	0.0516 (11)	0.0620 (12)	-0.0168 (10)	-0.0230 (10)	-0.0072 (9)
C31	0.0508 (11)	0.0517 (11)	0.0723 (13)	-0.0131 (9)	-0.0212 (10)	0.0055 (10)
C32	0.0875 (16)	0.0749 (15)	0.0500 (12)	-0.0296 (13)	0.0003 (11)	0.0029 (10)
C33	0.0954 (16)	0.0643 (13)	0.0425 (10)	-0.0284 (12)	-0.0078 (10)	-0.0098 (9)
C34	0.0725 (15)	0.0758 (16)	0.109 (2)	-0.0343 (13)	-0.0311 (14)	0.0193 (14)
N1	0.0679 (11)	0.0688 (12)	0.0599 (11)	-0.0126 (9)	0.0029 (9)	-0.0103 (9)
N2	0.0631 (12)	0.0942 (16)	0.0848 (15)	-0.0095 (11)	-0.0043 (11)	-0.0130 (12)
N3	0.0643 (9)	0.0581 (9)	0.0312 (7)	-0.0139 (8)	-0.0103 (7)	-0.0054 (6)
N4	0.0537 (8)	0.0483 (8)	0.0432 (8)	-0.0204 (7)	-0.0141 (6)	0.0000 (6)
N5	0.0657 (10)	0.0584 (9)	0.0375 (8)	-0.0261 (8)	-0.0145 (7)	0.0008 (7)
N6	0.0615 (9)	0.0488 (8)	0.0349 (7)	-0.0134 (7)	-0.0111 (6)	-0.0074 (6)
O1	0.0662 (9)	0.0856 (10)	0.0476 (8)	-0.0247 (8)	0.0051 (6)	-0.0126 (7)

O2	0.0656 (9)	0.0885 (11)	0.0535 (8)	-0.0167 (8)	-0.0229 (7)	-0.0129 (7)
O3	0.0603 (8)	0.0687 (9)	0.0640 (9)	-0.0187 (7)	0.0032 (7)	0.0018 (7)
O4	0.0865 (11)	0.0710 (10)	0.0846 (11)	-0.0340 (8)	-0.0475 (9)	0.0018 (8)
S1	0.0529 (3)	0.0649 (3)	0.0371 (2)	-0.0147 (2)	-0.00791 (19)	-0.0086 (2)
S2	0.0554 (3)	0.0524 (3)	0.0496 (3)	-0.0189 (2)	-0.0165 (2)	-0.0002 (2)

Geometric parameters (Å, °)

C1—C2	1.252 (4)	C20—N4	1.451 (2)
C1—H1A	0.9300	C20—H20A	0.9700
C1—H1B	0.9300	C20—H20B	0.9700
C2—C3	1.502 (4)	C21—N5	1.317 (2)
C2—H2	0.9300	C21—C22	1.417 (2)
C3—N1	1.450 (3)	C21—H21	0.9300
C3—H3A	0.9700	C22—C23	1.402 (2)
C3—H3B	0.9700	C22—C27	1.404 (2)
C4—N2	1.317 (3)	C23—C24	1.372 (2)
C4—C5	1.417 (3)	C23—H23	0.9300
C4—H4	0.9300	C24—C25	1.409 (2)
C5—C6	1.400 (3)	C24—N6	1.448 (2)
C5—C10	1.406 (3)	C25—C26	1.367 (2)
C6—N1	1.364 (3)	C25—H25	0.9300
C6—C7	1.388 (3)	C26—C27	1.398 (2)
C7—C8	1.368 (3)	C26—H26	0.9300
C7—H7	0.9300	C27—N4	1.359 (2)
C8—C9	1.410 (2)	C28—C29	1.376 (3)
C8—H8	0.9300	C28—C33	1.384 (3)
C9—C10	1.365 (3)	C28—S2	1.7592 (18)
C9—N3	1.435 (2)	C29—C30	1.375 (3)
C10—H10	0.9300	C29—H29	0.9300
C11—C16	1.380 (3)	C30—C31	1.382 (3)
C11—C12	1.384 (3)	C30—H30	0.9300
C11—S1	1.762 (2)	C31—C32	1.383 (3)
C12—C13	1.370 (3)	C31—C34	1.503 (3)
C12—H12	0.9300	C32—C33	1.375 (3)
C13—C14	1.383 (4)	C32—H32	0.9300
C13—H13	0.9300	C33—H33	0.9300
C14—C15	1.369 (4)	C34—H34A	0.9600
C14—C17	1.506 (4)	C34—H34B	0.9600
C15—C16	1.390 (4)	C34—H34C	0.9600
C15—H15	0.9300	N1—N2	1.355 (3)
C16—H16	0.9300	N3—S1	1.6219 (16)
C17—H17A	0.9600	N3—H3N	0.7979
C17—H17B	0.9600	N4—N5	1.363 (2)
C17—H17C	0.9600	N6—S2	1.6396 (16)
C18—C19	1.280 (3)	N6—H6N	0.7925
C18—H18A	0.9300	O1—S1	1.4337 (14)
C18—H18B	0.9300	O2—S1	1.4253 (14)
C19—C20	1.493 (3)	O3—S2	1.4283 (15)
C19—H19	0.9300	O4—S2	1.4260 (14)

C2—C1—H1A	120.0	N5—C21—H21	124.3
C2—C1—H1B	120.0	C22—C21—H21	124.3
H1A—C1—H1B	120.0	C23—C22—C27	120.20 (15)
C1—C2—C3	124.6 (4)	C23—C22—C21	135.59 (16)
C1—C2—H2	117.7	C27—C22—C21	104.21 (15)
C3—C2—H2	117.7	C24—C23—C22	117.89 (15)
N1—C3—C2	111.3 (2)	C24—C23—H23	121.1
N1—C3—H3A	109.4	C22—C23—H23	121.1
C2—C3—H3A	109.4	C23—C24—C25	121.15 (15)
N1—C3—H3B	109.4	C23—C24—N6	118.47 (15)
C2—C3—H3B	109.4	C25—C24—N6	120.37 (14)
H3A—C3—H3B	108.0	C26—C25—C24	122.00 (16)
N2—C4—C5	111.7 (2)	C26—C25—H25	119.0
N2—C4—H4	124.2	C24—C25—H25	119.0
C5—C4—H4	124.2	C25—C26—C27	117.11 (16)
C6—C5—C10	119.11 (19)	C25—C26—H26	121.4
C6—C5—C4	104.1 (2)	C27—C26—H26	121.4
C10—C5—C4	136.8 (2)	N4—C27—C26	131.53 (16)
N1—C6—C7	131.43 (19)	N4—C27—C22	106.84 (14)
N1—C6—C5	106.56 (19)	C26—C27—C22	121.63 (15)
C7—C6—C5	122.02 (18)	C29—C28—C33	120.04 (18)
C8—C7—C6	117.81 (18)	C29—C28—S2	120.13 (14)
C8—C7—H7	121.1	C33—C28—S2	119.67 (14)
C6—C7—H7	121.1	C30—C29—C28	119.62 (18)
C7—C8—C9	121.28 (19)	C30—C29—H29	120.2
C7—C8—H8	119.4	C28—C29—H29	120.2
C9—C8—H8	119.4	C29—C30—C31	121.56 (19)
C10—C9—C8	120.84 (17)	C29—C30—H30	119.2
C10—C9—N3	119.27 (16)	C31—C30—H30	119.2
C8—C9—N3	119.79 (17)	C30—C31—C32	117.79 (19)
C9—C10—C5	118.93 (18)	C30—C31—C34	120.7 (2)
C9—C10—H10	120.5	C32—C31—C34	121.5 (2)
C5—C10—H10	120.5	C33—C32—C31	121.6 (2)
C16—C11—C12	120.0 (2)	C33—C32—H32	119.2
C16—C11—S1	120.87 (17)	C31—C32—H32	119.2
C12—C11—S1	119.12 (16)	C32—C33—C28	119.35 (19)
C13—C12—C11	119.9 (2)	C32—C33—H33	120.3
C13—C12—H12	120.0	C28—C33—H33	120.3
C11—C12—H12	120.0	C31—C34—H34A	109.5
C12—C13—C14	121.3 (3)	C31—C34—H34B	109.5
C12—C13—H13	119.4	H34A—C34—H34B	109.5
C14—C13—H13	119.4	C31—C34—H34C	109.5
C15—C14—C13	118.1 (2)	H34A—C34—H34C	109.5
C15—C14—C17	120.2 (3)	H34B—C34—H34C	109.5
C13—C14—C17	121.7 (3)	N2—N1—C6	111.58 (18)
C14—C15—C16	122.1 (2)	N2—N1—C3	120.2 (2)
C14—C15—H15	119.0	C6—N1—C3	128.1 (2)
C16—C15—H15	119.0	C4—N2—N1	106.1 (2)

C11—C16—C15	118.6 (2)	C9—N3—S1	123.01 (12)
C11—C16—H16	120.7	C9—N3—H3N	114.5
C15—C16—H16	120.7	S1—N3—H3N	110.5
C14—C17—H17A	109.5	C27—N4—N5	111.13 (13)
C14—C17—H17B	109.5	C27—N4—C20	128.27 (15)
H17A—C17—H17B	109.5	N5—N4—C20	120.10 (14)
C14—C17—H17C	109.5	C21—N5—N4	106.45 (14)
H17A—C17—H17C	109.5	C24—N6—S2	118.67 (11)
H17B—C17—H17C	109.5	C24—N6—H6N	114.2
C19—C18—H18A	120.0	S2—N6—H6N	108.4
C19—C18—H18B	120.0	O2—S1—O1	118.84 (9)
H18A—C18—H18B	120.0	O2—S1—N3	109.12 (8)
C18—C19—C20	124.7 (2)	O1—S1—N3	104.33 (8)
C18—C19—H19	117.6	O2—S1—C11	108.08 (9)
C20—C19—H19	117.6	O1—S1—C11	107.75 (9)
N4—C20—C19	111.89 (16)	N3—S1—C11	108.31 (9)
N4—C20—H20A	109.2	O4—S2—O3	120.18 (10)
C19—C20—H20A	109.2	O4—S2—N6	105.67 (9)
N4—C20—H20B	109.2	O3—S2—N6	106.80 (8)
C19—C20—H20B	109.2	O4—S2—C28	108.23 (9)
H20A—C20—H20B	107.9	O3—S2—C28	108.43 (9)
N5—C21—C22	111.35 (15)	N6—S2—C28	106.79 (8)
C1—C2—C3—N1	-132.5 (3)	C29—C30—C31—C34	178.9 (2)
N2—C4—C5—C6	1.2 (3)	C30—C31—C32—C33	1.3 (3)
N2—C4—C5—C10	-179.8 (3)	C34—C31—C32—C33	-178.2 (2)
C10—C5—C6—N1	-179.90 (19)	C31—C32—C33—C28	-0.5 (4)
C4—C5—C6—N1	-0.6 (2)	C29—C28—C33—C32	-1.0 (3)
C10—C5—C6—C7	-0.3 (3)	S2—C28—C33—C32	174.52 (18)
C4—C5—C6—C7	178.9 (2)	C7—C6—N1—N2	-179.6 (2)
N1—C6—C7—C8	-179.2 (2)	C5—C6—N1—N2	0.0 (2)
C5—C6—C7—C8	1.3 (3)	C7—C6—N1—C3	4.1 (4)
C6—C7—C8—C9	-0.8 (3)	C5—C6—N1—C3	-176.4 (2)
C7—C8—C9—C10	-0.6 (3)	C2—C3—N1—N2	-100.4 (3)
C7—C8—C9—N3	175.55 (18)	C2—C3—N1—C6	75.7 (3)
C8—C9—C10—C5	1.6 (3)	C5—C4—N2—N1	-1.2 (3)
N3—C9—C10—C5	-174.58 (18)	C6—N1—N2—C4	0.8 (3)
C6—C5—C10—C9	-1.1 (3)	C3—N1—N2—C4	177.5 (2)
C4—C5—C10—C9	179.9 (3)	C10—C9—N3—S1	-124.00 (18)
C16—C11—C12—C13	-1.5 (3)	C8—C9—N3—S1	59.8 (2)
S1—C11—C12—C13	176.62 (18)	C26—C27—N4—N5	-179.05 (18)
C11—C12—C13—C14	-0.9 (4)	C22—C27—N4—N5	1.07 (19)
C12—C13—C14—C15	2.2 (4)	C26—C27—N4—C20	-7.3 (3)
C12—C13—C14—C17	-178.1 (2)	C22—C27—N4—C20	172.87 (17)
C13—C14—C15—C16	-1.0 (4)	C19—C20—N4—C27	-81.8 (2)
C17—C14—C15—C16	179.2 (2)	C19—C20—N4—N5	89.4 (2)
C12—C11—C16—C15	2.6 (3)	C22—C21—N5—N4	1.4 (2)
S1—C11—C16—C15	-175.51 (17)	C27—N4—N5—C21	-1.5 (2)
C14—C15—C16—C11	-1.3 (4)	C20—N4—N5—C21	-174.08 (16)

C18—C19—C20—N4	129.8 (2)	C23—C24—N6—S2	-90.15 (17)
N5—C21—C22—C23	178.45 (19)	C25—C24—N6—S2	90.84 (18)
N5—C21—C22—C27	-0.7 (2)	C9—N3—S1—O2	-56.27 (17)
C27—C22—C23—C24	0.9 (2)	C9—N3—S1—O1	175.74 (14)
C21—C22—C23—C24	-178.2 (2)	C9—N3—S1—C11	61.16 (16)
C22—C23—C24—C25	-1.6 (3)	C16—C11—S1—O2	-4.08 (19)
C22—C23—C24—N6	179.38 (14)	C12—C11—S1—O2	177.84 (16)
C23—C24—C25—C26	0.9 (3)	C16—C11—S1—O1	125.51 (17)
N6—C24—C25—C26	179.90 (16)	C12—C11—S1—O1	-52.57 (18)
C24—C25—C26—C27	0.5 (3)	C16—C11—S1—N3	-122.18 (17)
C25—C26—C27—N4	178.88 (17)	C12—C11—S1—N3	59.74 (18)
C25—C26—C27—C22	-1.3 (3)	C24—N6—S2—O4	-179.54 (13)
C23—C22—C27—N4	-179.55 (15)	C24—N6—S2—O3	51.40 (15)
C21—C22—C27—N4	-0.22 (19)	C24—N6—S2—C28	-64.46 (14)
C23—C22—C27—C26	0.6 (3)	C29—C28—S2—O4	-150.75 (16)
C21—C22—C27—C26	179.88 (16)	C33—C28—S2—O4	33.78 (19)
C33—C28—C29—C30	1.6 (3)	C29—C28—S2—O3	-18.86 (18)
S2—C28—C29—C30	-173.83 (15)	C33—C28—S2—O3	165.67 (17)
C28—C29—C30—C31	-0.8 (3)	C29—C28—S2—N6	95.90 (16)
C29—C30—C31—C32	-0.6 (3)	C33—C28—S2—N6	-79.58 (17)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N6—H6N \cdots O1	0.79	2.11	2.900 (2)	176
N3—H3N \cdots N5 ⁱ	0.80	2.19	2.983 (2)	175
C21—H21 \cdots O4 ⁱⁱ	0.93	2.49	3.245 (2)	138
C7—H7 \cdots O2 ⁱⁱⁱ	0.93	2.48	3.358 (2)	158

Symmetry codes: (i) $-x, -y+2, -z+1$; (ii) $-x+1, -y+2, -z+1$; (iii) $-x, -y+2, -z+2$.