



The Acorn Honey Used for a Published Study was Mistakenly Listed as *Quercus pyrenaica* and Should Be Corrected to *Quercus* species

Integrative Cancer Therapies
Volume 21: 1–2
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15347354221110818
journals.sagepub.com/home/ict


Abdurrahim Kocyigit, MD, PhD^{1,2} , Gokhan Aydogdu, MSc³,
Ezgi Balkan, MSc¹, Vildan Betül Yenigun, PhD¹, Eray Metin Guler, PhD¹,
Huri Bulut, PhD¹, Fatmanur Koktasoglu, PhD¹,
Ahmet Ceyhan Gören, PhD¹, and Ali Timucin Atayoglu, MD, PhD⁴

Submitted June 14, 2022; revised May 30, 2022; accepted June 15, 2022

A critical review letter¹ was published concerning the article “*Quercus pyrenaica* honeydew honey effects on gastric adenocarcinoma cells” published in *Integrative Cancer Therapies* in 2019.² According to the letter, the authors reported that the *Quercus pyrenaica* mentioned in the study was neither found in Mount Ida nor within the borders of Turkey as stated in the study, therefore the results obtained in the study may belong to another *Quercus* species.

In fact, within the scope of this project, 14 different types of honey obtained from different regions of Turkey were studied. The total phenol and flavonoid contents of each were analyzed. According to the results, 2 honey species with the highest and lowest phenolic content were selected to research other biological activities on cancer cells. One of them was acorn honey which was obtained from Mount Ida, and the other one is a multifloral honey which was obtained from the same region. At that time, unfortunately, we did not have the opportunity to investigate the flora of the region where each honey was produced in detail. When verbally asked about the origin of acorn honey to the producer, they reported that the origin of honey is *Quercus pyrenaica*. Therefore, this genus was mentioned as *Quercus pyrenaica* in the article. In fact, our main aim with this study was to compare the total phenolic and flavonoid contents of different honey samples and their cytotoxic, genotoxic, apoptotic, and pro-oxidant activities on cancer cells. Therefore, we focused on the phenolic compounds and biological activities of the honeys rather than the flora of the region where the honeys are produced. After the letter, we focused on the flora of the region where the acorn honey is produced. The genus *Quercus*, with high diversity and number of species, dominates many regions of temperate, tropical, and subtropical forests of Europe and North America.³ The genus contains more than 500 species distributed in the temperate zone in the Northern Hemisphere.⁴

Turkey is among the richest countries in the number of *Quercus* taxa, with 18 species belonging to 3 sections, *Quercus*, *Cerris*, and *Ile*,⁵ and 11 of them are known to be distributed in the Mount Ida region.⁵ When we asked the Forest Regional Directorate of the region where honey is produced with an official letter, they replied that there are 3 different *Quercus* species, *Quercus frainetto*, *Quercus cerris*, and *Quercus petraea* in the region where honey is produced (letter available from the authors on reasonable request).

In conclusion, our main focus with this study was to investigate the phenolic contents of honeys and the effects of honey with the highest and lowest phenolic content on the genotoxic, cytotoxic, apoptotic, and reactive oxygen species (ROS) production capacities on cancer cells, rather than the characteristics of the flora from which 14 different honey species are produced. Information on the origin of the honey with the highest phenolic content was requested from the producer, and according to the response received, it was incorrectly mentioned as *Quercus pyrenaica* in the article. Therefore, we think that it would be appropriate to publish a corrigendum to correct the errors in our article.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

¹Bezmailem Vakif University, Istanbul, Turkey

²Advanced Research and Application Center of Traditional and Complementary Medicine, Istanbul, Turkey

³Uludag University, Bursa, Turkey

⁴Istanbul Medipol University, Istanbul, Turkey

Corresponding Author:

Abdurrahim Kocyigit, Department of Medical Biochemistry, Medical Faculty, Bezmailem Vakif University, Istanbul 34093, Turkey.
Email: abdurrahimkocyigit@yahoo.com



Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Abdurrahim Kocyigit  <https://orcid.org/0000-0003-2335-412X>

References

1. Battal F, Kaya S, Köseoğlu M. The Potential Error on the *Quercus pyrenaica* Honeydew Honey Identifications. *Integr Cancer Ther.* 2022;21:15347354221096809.
2. Kocyigit A, Aydogdu G, Balkan E, et al. *Quercus pyrenaica* Honeydew Honey with high phenolic contents cause DNA damage, apoptosis, and cell death through generation of reactive oxygen species in gastric adenocarcinoma cells. *Integr Cancer Ther.* 2019;18:1534735419876334.
3. Ardi M, Rahmani F, Siami A. Genetic variation among Iranian oaks (*Quercus* spp.) using random amplified polymorphic DNA (RAPD) markers. *Afr J Biotechnol.* 2012;11:10291-10296.
4. Yücedağ C, Gailing O. Morphological and genetic variation within and among four *Quercus petraea* and *Q. robur* natural populations. *Turk J Botany.* 2013;37:619-629.
5. Güner A, Aslan S. *Türkiye bitkileri listesi:(damarlı bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi Yayınları; 2012.