

# Sequence variations of *GRM6* in patients with high myopia

Xiaoyu Xu, Shiqiang Li, Xueshan Xiao, Panfeng Wang, Xiangming Guo, Qingjiong Zhang

State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China

**Purpose:** Mutations in the glutamate receptor metabotropic 6 gene (*GRM6*) have been identified in patients with congenital stationary night blindness (CSNB1B). High myopia is usually observed in CSNB1B patients. This study tested if any mutations in *GRM6* were solely responsible for high myopia.

**Methods:** DNA was prepared from the venous leukocytes of 96 Chinese patients with high myopia (refraction of spherical equivalent of at least -6.00 diopters [D]) and 96 controls (refraction of spherical equivalent between -0.50 D and +2.00 D with normal visual acuity). The coding regions and adjacent intronic sequence of *GRM6* were amplified by a polymerase chain reaction (PCR) and then analyzed by cycle sequencing. Detected variations were evaluated in normal controls by heteroduplex-single-strand-conformation (SSCP) polymorphism analysis or restriction fragment polymorphism (RFLP).

**Results:** Four novel variations predicted to have potential functional changes were identified: c.67-82delCAGGCGGGCCTGGCGCinsT (p.Gln23\_Arg28delinsCys), c.858-5a>g (r.spl?), c.1172G>A (p.Arg391Gln), and c.1537G>A (p.Val513Met). Except for c.1172G>A, the other three were not detected in the 96 controls. In addition, five rare variations—(c.72G>A, c.504+10g>t, c.726-50g>c, c.1359C>T, and c.1383C>T)—and one common variation (c.2437-6g>a) without predicted functional consequences and nine known single nucleotide polymorphisms (SNPs) were also detected.

**Conclusion:** Three novel variations with potential functional consequences were identified in the *GRM6* of patients with high myopia, suggesting a potential role in the development of myopia in rare cases.

The glutamate receptor metabotropic 6 gene (*GRM6*; OMIM 604096), mapped to 5q35, contained 10 exons and encoded an 877 amino acid protein, mGluR6. As a member of Group III mGluRs (mGluR4, 6, 7, and 8), it contains a signal peptide, a large bi-lobed extracellular NH<sub>2</sub>-terminal domain containing the glutamate binding site, seven G protein-coupled receptor (GPCR) transmembrane domains and an intracellular COOH-terminal domain [1-3]. In the rat retina, mGluR6 is specifically expressed in ON bipolar cells at the postsynaptic site [4]. In the retinal neural network, an increase in light reduced the release of glutamate from cones and rods, while a decrease in light increases its release, which acted as inputs from photoreceptors to bipolar cells. The visual signals of light and dark were segregated into parallel ON and OFF pathways through ON and OFF bipolar cells. The ON bipolar cells utilize a metabotropic glutamate receptor (mGluR6) for a light-activated depolarization, whereas the OFF bipolar cells utilized ionotropic glutamate receptors (iGluRs) for a light-activated hyperpolarization. ON and OFF bipolar cells made synaptic contacts with ON and OFF ganglion cells and transmitted visual signals to the brain [5-8].

Functional defects involving retinal ON-pathways have been demonstrated by retinal electrophysiology studies in the complete form of congenital stationary night blindness,

including CSNB1A (OMIM 310500) and CSNB1B (OMIM 257270) [7,9-12]. Mutations in *NYX* or *GRM6* are responsible for CSNB1A or CSNB1B, respectively [10,11,13-15]. Besides night blindness, high myopia is also frequently documented as a typical sign in CSNB1A patients with *NYX* mutations [13,14,16-18]. A mouse model with *Nyx* mutation and retinal ON-pathway defect has high susceptibility to experimental myopia [19]. Recently, mutations in *NYX* have been reported to associate with high myopia alone without night blindness [20]. Similarly, moderate to high myopia is also a common sign in CSNB1B patients with *GRM6* mutations [10,21]. A mouse model lacking the *GRM6* gene showed a loss of ON response, but an unchanged OFF response to light, demonstrating its essential role in ON synaptic transmission [4].

Congenital stationary night blindness (CSNB) can be caused by mutations in genes *GNATI*, *PDE6B*, *RHO*, *CABP4*, *GRK1*, *GRM6*, *RDH5*, *SAG*, *CACNA1F*, and *NYX* (RetNet). Myopia is not always associated with CSNB, except in cases resulting from mutations in *NYX* and *GRM6* [10,16], suggesting a gene-specific phenotype rather than association with night blindness. As mutations in *GRM6* resulted in phenotypes extremely similar to those in *NYX*, this suggested that *GRM6* might be a candidate susceptibility gene for isolated high myopia. In this study, we analyzed the genomic sequence of *GRM6* in 96 Chinese patients with high myopia.

## METHODS

**Subjects:** Ninety-six unrelated probands with high myopia and 96 unrelated normal controls were collected from

Correspondence to: Qingjiong Zhang, M.D., Ph.D., State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, 54 South Xianlie Road, Guangzhou, 510060, China; Phone: (+86)-20-87330422; FAX: (+86)-20-87333271; email: qingjiongzhang@yahoo.com

TABLE 1. PRIMERS USED FOR POLYMERASE CHAIN REACTION AMPLIFICATION AND DIRECT SEQUENCING OF *GRM6*.

Exons	Primer sequence(5'-3')	Product length (bp)	Annealing temperature (°C)	Note
1	F-CCGAGCGCCTTCTCCCCAGGAC R-TGCGTCGGGCTCAACTGACAGG	785	68	
2	F-TGTGCCCCCGTCCGTGTCC R-TGAGGCAGGGCTGAGTCGTC	481	63	
3	F-GGGCGGGACCTCTTTGTT R-GCTATTCAGTCTGGGCTTGTG	521	65	
4~7	F-CCGGCCCCGTTTTTCCTG R-CACCTGGGACGCACAAAACAC	4132	69	LA Taq DNA Polymerase
	F1-GCCATGCCTAGTTCACCT	/	/	Sequencing primer
	F2-ATTTGCACGTCCCTTATGAGC	/	/	Sequencing primer
	R1-CTTTCCTCGCACGCCATCC	/	/	Sequencing primer
	R2-TCAGCCTCACCCAGCCCTTCC	/	/	Sequencing primer
8	F-TCAACGAGAACGGAGATGC R-CCCTTTGGCTTTGTAACG	900	61	
9	F-CTGAGAGCCACCACGAAGAG R-GCAGCCAGATACGGGATAAA	567	65	
10	F-GGGCCCCTGCTCATACTGTT R-CTCCCTGCCACTGACTGTTC	460	65	

In the "Primer sequence" column, "F" indicates the forward primer and "R" indicates the reverse primer. GC1 buffer (Takara Biotechnology Dalian CO. Ltd, Liaoning, China) is designed for amplification of templates having complex secondary structure or high GC content. LA Taq DNA Polymerase (Takara Biotechnology Dalian CO. Ltd, Liaoning, China) was designed for amplification of long targets. The amplicon from exon 4 to 7 was sequenced with the additional primers listed as F1, F2, R1 and R2.

Zhongshan Ophthalmic Center. Informed consent conforming to the tenets of the Declaration of Helsinki was obtained from the participants prior to the study. This study was approved by the Institutional Review Boards of Zhongshan Ophthalmic Center. Ophthalmological examinations were performed by ophthalmologists (Drs. Q.Z. and X.G.). The diagnostic criteria for high myopia were as we previously described [22]: 1) bilateral refraction of  $-6.00D$  or lower (spherical equivalent) and 2) no other known ocular or systemic diseases associated with high myopia. Normal controls met the following criteria: 1) bilateral refraction between  $-0.50 D$  and  $+2.00 D$  with normal visual acuity, 2) no family history of high myopia, and 3) exclusion of known ocular or systemic diseases. The refractive error of all eyes was measured with cycloplegic autorefractometry after mydriasis (Mydrin<sup>®</sup>-P, a compound tropicamide; Santen Pharmaceutical Co. Ltd., Osaka, Japan). Genomic DNA was prepared from venous blood.

**Variation analysis:** Seven pairs of primers (Table 1) were used to amplify the 10 coding exons and the adjacent intronic sequence of *GRM6* (human genome build 36.2 [NC\\_000005.8](#) for gDNA, [NM\\_000843.3](#) for mRNA, and [NP\\_000843.2](#) for protein). DNA fragments from individual exons were amplified by touchdown PCR where higher annealing temperatures were set for the first five cycles, followed by moderate annealing temperature for the next five cycles and finally by a lower annealing temperature as listed in Table 1 for the remaining 23 cycles. The procedures for sequencing

and variation detection were basically the same as previously described [22]. Potential mutations detected in *GRM6* of patients were further evaluated in the 96 controls by using either heteroduplex-single-strand-conformational polymorphism (HA-SSCP) [23] or polymerase chain reaction combined with restriction fragment length polymorphism (PCR-RFLP) analysis [24]. Extra pairs of primers were designed for HA-SSCP or PCR-RFLP analysis (Table 2). The 102 bp amplicons with the c.1172G>A variation were cut into 26 bp and 76 bp as the variation created a new BstNI site, while the wild amplicons remained unchanged. The 136 bp amplicons with the c.1537G>A variation could not be digested by Bsp1286I since the variation erased the Bsp1286I site that presented in wild type amplicons. The variation of c.1172G>A was genotyped and statistically analyzed using continuity correction of Pearson's Chi-Square test with a significance level of 0.05.

**Database and online tools:** Polymorphism Phenotyping (PolyPhen) [25-27] and Sorting Intolerant From Tolerant (SIFT) [28] were used to evaluate the potential pathogenicity of sequence alterations at the protein level. Automated Splice Site Analysis (ASSA) [29] and Berkeley Drosophila Genome Project (BDGP) [30] were used for predicting the alterations of splicing sites. National Center for Biotechnology Information (NCBI), Conserved Domain Database (CDD), PSORT II, WoLF PSORT [31,32], Simple Modular Architecture Research Tool (SMART) [33,34], and

TABLE 2. PRIMERS USED FOR EVALUATING THE VARIATIONS BY HA-SSCP OR PCR-RFLP ANALYSIS OF *GRM6*.

Variations	Primer sequence (5'-3')	Product length (bp)	Annealing temperature (°C)	Note
c.68-82delAGGCGG GCCTGGCGCinsT	F-CCGAGCGCCTTCTCCCCAGGAC R-GGTGTCCCGCGAGCAGGTG	430	71	
c.726-50g>c	F-GGGCGGGACCTCTTTGTT R-CCTGGGAATCTTGATAGACTGG	214	68-66-64	Touchdown PCR
c.858-5a>g	F-GGAAGGTGCCCTAAATGC R-AAGTGGCCGGTCAGGTTGG	202	65	
c.1172G>A	F-TAGGACAGGCGAGGAACG R-TGGAGGGCGTGGGCAATG	102	70-68-66	Touchdown PCR
c.1537G>A	F-GGATCCCGGAGGCAGACG R-CCATCTTCTCCGCTCCC	136	70-68-66	Touchdown PCR

pTARGET [35,36] were used for analyzing structures and functions of protein domains and predicting protein topology and subcellular localization. Search Tool for the Retrieval of Interacting Genes/Proteins (STRING) was used for showing and predicting protein interactions [37].

## RESULTS

Upon complete analysis of *GRM6* in 96 patients with high myopia, four novel variations predicted to have potential functional consequence were identified: c.67-82delCAGGCGGGCCTGGCGCinsT (p.Gln23\_Arg28delinsCys), c.858-5a>g (r.spl?), c.1172G>A (p.Arg391Gln), and c.1537G>A (p.Val513Met; Figures 1A,B; Table 3). The c.67-82delCAGGCGGGCCTGGCGCinsT (p.Gln23\_Arg28delinsCys) variation was further determined by cloning sequencing (Figure 1A). Of the four variations, three were not present in the 96 controls but the other one (c.1172G>A) was detected in four of the 96 controls. In addition, five rare variations (c.72G>A, c.504+10g>t, c.726-50g>c, c.1359C>T, and c.1383C>T) and a common variation (c.2437-6g>a) without predicted functional consequence by the abovementioned online tools [25-30] were also detected in patients with high myopia, but the presence of these variations in controls was not examined. Furthermore, nine reported SNPs (rs2645329, rs2071246, rs2645339, rs4701014, rs2067011, rs11746675, rs2071247, rs2071249, and rs17078853) were also detected in patients (Table 3).

The c.67-82delCAGGCGGGCCTGGCGCinsT (p.Gln23\_Arg28delinsCys) variation was detected in one patient in a heterozygous status, but not in the 96 controls. It was a del\_ins variation without frame shifting, resulting in the deletion of six residues and the addition of cysteine at protein. Conserved domain analysis showed the affected oligopeptide was not conserved among different species or different members of Group III mGluRs (Figures 1D, E). Functional domain prediction revealed that amino acid residues one to 24

of mGluR6 would act as the signal peptide, and the following 25 to 585 amino acid residues would form an extracellular N-terminal domain related to periplasmic ligand binding. The NH<sub>2</sub>-terminal domain is vital in glutamate binding and the activation/inactivation of mGluR6 [38,39].

The c.858-5a>g variation was detected in one patient, but not in the 96 controls. Analysis of the corresponding splicing site by both the ASSA server and the BDGP server revealed a comparatively weak leaky effect (from 9.6 to 9.3), resulting in a -1.2 fold decrease on the natural splicing acceptor site.

The c.1172G>A variation resulted in a substitution of arginine (a strongly basic residue) to glutamine (a small polar amino acid; p.Arg391Gln), with a residue weight of one on Blosum 62 and "benign" or "tolerated" effect by PolyPhen and SIFT, respectively. The arginine in this position was conserved between different species except in *Rattus norvegicus* and *Mus musculus*, whereas it varied in different members of Group III mGluRs (Figures 1D, E). The residue with variation is in the NH<sub>2</sub>-terminal domain and nearby one of the key positions of the glutamate binding pocket p.Lys400 [40], presumably and, therefore, is presumed to affect the receptor family ligand binding region according to domain analyzing databases. However, this variation was found in two patients and four controls, with no statistical significance on continuity correction of Pearson's  $\chi^2$  test (Table 3).

The c.1537G>A variation was found in three high myopia patients, where two were heterozygous and one homozygous. It resulted in a substitution of a hydrophobic valine by a highly conserved sulfur-containing and polar methionine (p.Val513Met), with a Blosum 62 score of one and a "benign" or "tolerated" effect by the abovementioned online tools. Residues 513 to 564 were predicted to form a highly conserved extracellular domain of family 3 GPCR forming disulphide bridges (Figures 1D, E). The function of this region remains uncertain but may play a role in calcium sensing and proper trafficking of proteins [41,42]. Residue 524 is predicted to be one of the nuclear localization signals of




Figure 1. Four novel variations were detected in *GRM6* of 96 unrelated subjects with isolated high myopia. **A**: Sequences with c.67-82delCAGGCGGGCCTGGCGCinsT p.Gln23\_Arg28delinsCys variation, normal control, as well as cloning sequences of both forward and reverse directions. **B**: Sequences showing the rest three novel variations as well as normal controls. The substitute bases are highlighted by an arrow, with protein consequence marked below. **C**: Variations were not observed in normal controls (NC) by SSCP or RFLP, with the exception of c.1172G>A p.Arg391Gln. Plus signs indicate samples with variation; minus signs indicate normal controls. **D** and **E**: Conservation analysis of the variations between different species (**D**) or between different members of Group III mGluRs (**E**).

mGluR6. Residue 513 may play a role in nuclear localization and translocation of mGluR6.

### DISCUSSION

High myopia usually occurs alone (nonsyndromic) but, in rare cases, may present as a sign in a number of syndromes (syndromic). Genetic factors have been demonstrated to play

an important role in the development of high myopia [43,44]. The genes responsible for nonsyndromic high myopia are mostly unknown although identification of such genes has been sought by using various approaches, including linkage analysis, case-control association study, and sequence analyzing of candidate genes [45]. Of these, a number of studies have tested the association of nonsyndromic high

TABLE 3. SEQUENCE VARIATIONS IN *GRM6* DETECTED IN 96 HIGH MYOPIA PATIENTS.

Variation	Location	Effect	High myopia (n=96)	Normal control (n=96)
# <b>c.72G&gt;A</b>	<b>Exon 1</b>	<b>synonymous</b>	<b>1</b>	<b>ND</b>
<b>1 c.67-82delCAGGCGGGCCTGGCGCinsT</b>	<b>Exon 1</b>	<b>p.Gln23_Arg28delinsCys</b>	<b>1</b>	<b>0</b>
2 c.176A>C	Exon 1	rs2645329	76	ND
<b>3 c.504+10g&gt;t</b>	<b>Intron 1</b>	<b>No splicing site changed</b>	<b>1</b>	<b>ND</b>
<b>4 c.726-50g&gt;c</b>	<b>Intron 2</b>	<b>Splicing acceptor unaffected</b>	<b>1</b>	<b>0</b>
<b>5 c.858-5a&gt;g</b>	<b>Intron 3</b>	<b>Decrease of splicing acceptor</b>	<b>1</b>	<b>0</b>
6 c.1131C>T	Exon 5	rs2071246	87	ND
<b>7 c.1172G&gt;A</b>	<b>Exon 6</b>	<b>p.Arg391Gln</b>	<b>2</b>	<b>4*</b>
8 c.1227C>T	Exon 6	rs2645339	84	ND
9 c.1308T>C	Exon 6	rs4701014	96	ND
10 c.1353T>C	Exon 6	rs2067011	84	ND
<b>11 c.1359C&gt;T</b>	<b>Exon 7</b>	<b>synonymous</b>	<b>2</b>	<b>ND</b>
<b>12 c.1383C&gt;T</b>	<b>Exon 7</b>	<b>synonymous</b>	<b>1</b>	<b>ND</b>
13 c.1392A>G	Exon 7	rs11746675	75	ND
<b>14 c.1537G&gt;A</b>	<b>Exon 8</b>	<b>p.Val513Met</b>	<b>3</b>	<b>0</b>
15 c.2196G>A	Exon 9	rs2071247	52	ND
<b>16 c.2437-6g&gt;a</b>	<b>Intron 9</b>	<b>No splicing site changed</b>	<b>19</b>	<b>ND</b>
17 c.2457G>A	Exon 10	rs2071249	19	ND
18 c.2634+56a>t	Intron 10	rs17078853	19	ND

The novel variations are highlighted in bold. An asterisk indicates that there is no statistical difference between patients and controls. ND: no detected.

myopia with common SNPs in genes responsible for syndromic high myopia but, unfortunately, the results are inconclusive and controversial in general [40,46-49]. This approach is based on the understanding that the genetic basis of complex diseases is associated with common variant (common disease, common variant [CDCV]). However, routine association study of common SNPs may not discover rare variations that might contribute to complex diseases (common disease, rare variant [CDRV]) [50-52]. Sequencing the functional regions of the target gene, as in this study, would detect such rare variations as well as mutations, which should be a preferable approach for a disease like high myopia where both Mendelian traits and complex traits are involved.

Here in this study, the entire coding and adjacent intronic regions of *GRM6* were sequenced for 96 patients with nonsyndromic high myopia. Consequently, four novel variations were detected in *GRM6* of the Chinese patients with high myopia, where three were only present in patients but not among the 96 controls. These variations were predicted to affect the functions of the encoded proteins if the mutant allele is expressed. This was the first sequencing evaluation of *GRM6* in a group of patients with high myopia. The information obtained based on the current case-control sequence analyzing may not lead to a definite conclusion; however, it not only expands our understanding of *GRM6* variations in human beings but also may suggest a potential role of *GRM6* rare variations in the development of myopia in rare cases. Further studies in other ethnic populations may provide useful information to verify our findings.

#### ACKNOWLEDGMENTS

The authors thank all patients and normal controls for their participation. This study was supported in part by grants 30772390 from the National Natural Science Foundation of China, Grant 7001571 and 8251008901000020 from the National Science Foundation of Guangdong Province. Prof. Qingjiong Zhang is a recipient of the National Science Fund for Distinguished Young Scholars (30725044).

#### REFERENCES

1. Nakajima Y, Iwakabe H, Akazawa C, Nawa H, Shigemoto R, Mizuno N, Nakanishi S. Molecular characterization of a novel retinal metabotropic glutamate receptor mGluR6 with a high agonist selectivity for L-2-amino-4-phosphonobutyrate. *J Biol Chem* 1993; 268:11868-73. [PMID: 8389366]
2. Hashimoto T, Inazawa J, Okamoto N, Tagawa Y, Bessho Y, Honda Y, Nakanishi S. The whole nucleotide sequence and chromosomal localization of the gene for human metabotropic glutamate receptor subtype 6. *Eur J Neurosci* 1997; 9:1226-35. [PMID: 9215706]
3. Weng K, Lu C, Daggett LP, Kuhn R, Flor PJ, Johnson EC, Robinson PR. Functional coupling of a human retinal metabotropic glutamate receptor (hmGluR6) to bovine rod transducin and rat Go in an in vitro reconstitution system. *J Biol Chem* 1997; 272:33100-4. [PMID: 9407094]
4. Masu M, Iwakabe H, Tagawa Y, Miyoshi T, Yamashita M, Fukuda Y, Sasaki H, Hiroi K, Nakamura Y, Shigemoto R, Takadac M, Nakamura K, Nakaog K, Katsukig M, Nakanishi S. Specific deficit of the ON response in visual transmission by targeted disruption of the mGluR6 gene. *Cell* 1995; 80:757-65. [PMID: 7889569]

5. Kew JN, Kemp JA. Ionotropic and metabotropic glutamate receptor structure and pharmacology. *Psychopharmacology (Berl)* 2005; 179:4-29. [PMID: 15731895]
6. Maddox DM, Vessey KA, Yarbrough GL, Invergo BM, Cantrell DR, Inayat S, Balannik V, Hicks WL, Hawes NL, Byers S, Smith RS, Hurd R, Howell D, Gregg RG, Chang B, Naggert JK, Troy JB, Pinto LH, Nishina PM, McCall MA. Allelic variance between GRM6 mutants, Grm6nob3 and Grm6nob4 results in differences in retinal ganglion cell visual responses. *J Physiol* 2008; 586:4409-24. [PMID: 18687716]
7. McCall MA, Gregg RG. Comparisons of structural and functional abnormalities in mouse b-wave mutants. *J Physiol* 2008; 586:4385-92. [PMID: 18653656]
8. Snellman J, Kaur T, Shen Y, Nawy S. Regulation of ON bipolar cell activity. *Prog Retin Eye Res* 2008; 27:450-63. [PMID: 18524666]
9. Khan NW, Kondo M, Hiriyanna KT, Jamison JA, Bush RA, Sieving PA. Primate Retinal Signaling Pathways: Suppressing ON-Pathway Activity in Monkey With Glutamate Analogues Mimics Human CSNB1-NYX Genetic Night Blindness. *J Neurophysiol* 2005; 93:481-92. [PMID: 15331616]
10. Dryja TP, McGee TL, Berson EL, Fishman GA, Sandberg MA, Alexander KR, Derlacki DJ, Rajagopalan AS. Night blindness and abnormal cone electroretinogram ON responses in patients with mutations in the GRM6 gene encoding mGluR6. *Proc Natl Acad Sci USA* 2005; 102:4884-9. [PMID: 15781871]
11. Zeitz C, van Genderen M, Neidhardt J, Luhmann UF, Hoeben F, Forster U, Wycisk K, Matyas G, Hoyng CB, Riemslag F, Meire F, Cremers FP, Berger W. Mutations in GRM6 cause autosomal recessive congenital stationary night blindness with a distinctive scotopic 15-Hz flicker electroretinogram. *Invest Ophthalmol Vis Sci* 2005; 46:4328-35. [PMID: 16249515]
12. Kabanarou SA, Holder GE, Fitzke FW, Bird AC, Webster AR. Congenital stationary night blindness and a "Schubert-Bornschein" type electrophysiology in a family with dominant inheritance. *Br J Ophthalmol* 2004; 88:1018-22. [PMID: 15258017]
13. Pusch CM, Zeitz C, Brandau O, Pesch K, Achatz H, Feil S, Scharfe C, Maurer J, Jacobi FK, Pinckers A, Andreasson S, Hardcastle A, Wissinger B, Berger W, Meindl A. The complete form of X-linked congenital stationary night blindness is caused by mutations in a gene encoding a leucine-rich repeat protein. *Nat Genet* 2000; 26:324-7. [PMID: 11062472]
14. Bech-Hansen NT, Naylor MJ, Maybaum TA, Sparkes RL, Koop B, Birch DG, Bergen AA, Prinsen CF, Polomeno RC, Gal A, Drack AV, Musarella MA, Jacobson SG, Young RS, Weleber RG. Mutations in NYX, encoding the leucine-rich proteoglycan nyctalopin, cause X-linked complete congenital stationary night blindness. *Nat Genet* 2000; 26:319-23. [PMID: 11062471]
15. Abramowicz MJ, Ribai P, Cordonnier M. Congenital stationary night blindness: report of an autosomal recessive family and linkage analysis. *Am J Med Genet A* 2005; 132A:76-9. [PMID: 15551339]
16. Xiao X, Jia X, Guo X, Li S, Yang Z, Zhang Q. CSNB1 in Chinese families associated with novel mutations in NYX. *J Hum Genet* 2006; 51:634-40. [PMID: 16670814]
17. Zeitz C, Minotti R, Feil S, Matyas G, Cremers FP, Hoyng CB, Berger W. Novel mutations in CACNA1F and NYX in Dutch families with X-linked congenital stationary night blindness. *Mol Vis* 2005; 11:179-83. [PMID: 15761389]
18. Zito I, Allen LE, Patel RJ, Meindl A, Bradshaw K, Yates JR, Bird AC, Erskine L, Cheetham ME, Webster AR, Poopalasundaram S, Moore AT, Trump D, Hardcastle AJ. Mutations in the CACNA1F and NYX genes in British CSNBX families. *Hum Mutat* 2003; 21:169. [PMID: 12552565]
19. Pardue MT, Faulkner AE, Fernandes A, Yin H, Schaeffel F, Williams RW, Pozdeyev N, Iuvone PM. High susceptibility to experimental myopia in a mouse model with a retinal on pathway defect. *Invest Ophthalmol Vis Sci* 2008; 49:706-12. [PMID: 18235018]
20. Zhang Q, Xiao X, Li S, Jia X, Yang Z, Huang S, Caruso RC, Guan T, Sergeev Y, Guo X, Hejtmancik JF. Mutations in NYX of individuals with high myopia, but without night blindness. *Mol Vis* 2007; 13:330-6. [PMID: 17392683]
21. O'Connor E, Allen LE, Bradshaw K, Boylan J, Moore AT, Trump D. Congenital stationary night blindness associated with mutations in GRM6 encoding glutamate receptor mGluR6. *Br J Ophthalmol* 2006; 90:653-4. [PMID: 16622103]
22. Li T, Xiao X, Li S, Xing Y, Guo X, Zhang Q. Evaluation of EGR1 as a candidate gene for high myopia. *Mol Vis* 2008; 14:1309-12. [PMID: 18636116]
23. Zhang Q, Minoda K. Detection of congenital color vision defects using heteroduplex-SSCP analysis. *Jpn J Ophthalmol* 1996; 40:79-85. [PMID: 8739504]
24. Deng GR. A sensitive non-radioactive PCR-RFLP analysis for detecting point mutations at 12th codon of oncogene c-Ha-ras in DNAs of gastric cancer. *Nucleic Acids Res* 1988; 16:6231. [PMID: 2456524]
25. Ramensky V, Bork P, Sunyaev S. Human non-synonymous SNPs: server and survey. *Nucleic Acids Res* 2002; 30:3894-900. [PMID: 12202775]
26. Sunyaev S, Ramensky V, Bork P. Towards a structural basis of human non-synonymous single nucleotide polymorphisms. *Trends Genet* 2000; 16:198-200. [PMID: 10782110]
27. Sunyaev S, Ramensky V, Koch I, Lathe W 3rd, Kondrashov AS, Bork P. Prediction of deleterious human alleles. *Hum Mol Genet* 2001; 10:591-7. [PMID: 11230178]
28. Ng PC, Henikoff S. Predicting deleterious amino acid substitutions. *Genome Res* 2001; 11:863-74. [PMID: 11337480]
29. Nalla VK, Rogan PK. Automated splicing mutation analysis by information theory. *Hum Mutat* 2005; 25:334-42. [PMID: 15776446]
30. Reese MG, Eeckman FH, Kulp D, Haussler D. Improved splice site detection in Genie. *J Comput Biol* 1997; 4:311-23. [PMID: 9278062]
31. Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K. WoLF PSORT: protein localization predictor. *Nucleic Acids Res* 2007; 35:W585-7. [PMID: 17517783]

32. Nakai K, Horton P. Computational prediction of subcellular localization. *Methods Mol Biol* 2007; 390:429-66. [PMID: 17951705]
33. Schultz J, Milpetz F, Bork P, Ponting CP. SMART, a simple modular architecture research tool: identification of signaling domains. *Proc Natl Acad Sci USA* 1998; 95:5857-64. [PMID: 9600884]
34. Letunic I, Doerks T, Bork P. SMART 6: recent updates and new developments. *Nucleic Acids Res* 2009; 37:D229-32. [PMID: 18978020]
35. Guda C, Subramaniam S. pTARGET [corrected] a new method for predicting protein subcellular localization in eukaryotes. *Bioinformatics* 2005; 21:3963-9. [PMID: 16144808]
36. Guda C. pTARGET: a web server for predicting protein subcellular localization. *Nucleic Acids Res* 2006; 34:W210, 3. [PMID: 16844995]
37. von Mering C, Jensen LJ, Kuhn M, Chaffron S, Doerks T, Kruger B, Snel B, Bork P. STRING 7--recent developments in the integration and prediction of protein interactions. *Nucleic Acids Res* 2007; 35:D358-62. [PMID: 17098935]
38. Tsuchiya D, Kunishima N, Kamiya N, Jingami H, Morikawa K. Structural views of the ligand-binding cores of a metabotropic glutamate receptor complexed with an antagonist and both glutamate and Gd<sup>3+</sup>. *Proc Natl Acad Sci USA* 2002; 99:2660-5. [PMID: 11867751]
39. Kunishima N, Shimada Y, Tsuji Y, Sato T, Yamamoto M, Kumasaka T, Nakanishi S, Jingami H, Morikawa K. Structural basis of glutamate recognition by a dimeric metabotropic glutamate receptor. *Nature* 2000; 407:971-7. [PMID: 11069170]
40. Wang P, Li S, Xiao X, Jia X, Jiao X, Guo X, Zhang Q. High myopia is not associated with the SNPs in the TGIF, lumican, TGFB1, and HGF genes. *Invest Ophthalmol Vis Sci* 2009; 50:1546-51. [PMID: 19060265]
41. Zeitz C, Forster U, Neidhardt J, Feil S, Kalin S, Leifert D, Flor PJ, Berger W. Night blindness-associated mutations in the ligand-binding, cysteine-rich, and intracellular domains of the metabotropic glutamate receptor 6 abolish protein trafficking. *Hum Mutat* 2007; 28:771-80. [PMID: 17405131]
42. Jo J, Heon S, Kim MJ, Son GH, Park Y, Henley JM, Weiss JL, Sheng M, Collingridge GL, Cho K. Metabotropic glutamate receptor-mediated LTD involves two interacting Ca(2+) sensors, NCS-1 and PICK1. *Neuron* 2008; 60:1095-111. [PMID: 19109914]
43. Feldkamper M, Schaeffel F. Interactions of genes and environment in myopia. *Dev Ophthalmol* 2003; 37:34-49. [PMID: 12876828]
44. Young TL, Metlapally R, Shay AE. Complex trait genetics of refractive error. *Arch Ophthalmol* 2007; 125:38-48. [PMID: 17210850]
45. Tang WC, Yap MK, Yip SP. A review of current approaches to identifying human genes involved in myopia. *Clin Exp Optom* 2008; 91:4-22. [PMID: 18045248]
46. Mutti DO, Cooper ME, O'Brien S, Jones LA, Marazita ML, Murray JC, Zadnik K. Candidate gene and locus analysis of myopia. *Mol Vis* 2007; 13:1012-9. [PMID: 17653045]
47. Metlapally R, Li YJ, Tran-Viet KN, Abbott D, Czaja GR, Malecaze F, Calvas P, Mackey D, Rosenberg T, Paget S, Zayats T, Owen MJ, Guggenheim JA, Young TL. COL1A1 and COL2A1 genes and myopia susceptibility: evidence of association and suggestive linkage to the COL2A1 locus. *Invest Ophthalmol Vis Sci* 2009; 50:4080-6. [PMID: 19387081]
48. Inamori Y, Ota M, Inoko H, Okada E, Nishizaki R, Shiota T, Mok J, Oka A, Ohno S, Mizuki N. The COL1A1 gene and high myopia susceptibility in Japanese. *Hum Genet* 2007; 122:151-7. [PMID: 17557158]
49. Liang CL, Hung KS, Tsai YY, Chang W, Wang HS, Juo SH. Systematic assessment of the tagging polymorphisms of the COL1A1 gene for high myopia. *J Hum Genet* 2007; 52:374-7. [PMID: 17273809]
50. Schork NJ, Murray SS, Frazer KA, Topol EJ. Common vs. rare allele hypotheses for complex diseases. *Curr Opin Genet Dev* 2009; 19:212-9. [PMID: 19481926]
51. Nejentsev S, Walker N, Riches D, Egholm M, Todd JA. Rare variants of IFIH1, a gene implicated in antiviral responses, protect against type 1 diabetes. *Science* 2009; 324:387-9. [PMID: 19264985]
52. Bodmer W, Bonilla C. Common and rare variants in multifactorial susceptibility to common diseases. *Nat Genet* 2008; 40:695-701. [PMID: 18509313]