

# Exploring the Molecular Mechanisms of Pterygium by Constructing lncRNA–miRNA–mRNA Regulatory Network

Nuo Xu,<sup>1</sup> Yi Cui,<sup>2</sup> Jiaxing Dong,<sup>3</sup> and Li Huang<sup>2</sup>

<sup>1</sup>Department of Ophthalmology, Fujian Provincial Hospital, Shengli Clinical Medical College of Fujian Medical University, Fuzhou City, Fujian Province, China

<sup>2</sup>Department of Ophthalmology, Fujian Medical University Union Hospital, Fuzhou City, Fujian Province, China

<sup>3</sup>Xinxiang Medical University, Xinxiang City, Henan Province, China

Correspondence: Yi Cui, Department of Ophthalmology, Fujian Medical University Union Hospital, No. 29, Xinquan Road, Gulou District, Fuzhou, Fujian Province, 350001, China; [oph\\_cy@163.com](mailto:oph_cy@163.com).

NX, YC, and JD contributed equally to this work.

**Received:** March 13, 2020

**Accepted:** June 4, 2020

**Published:** July 9, 2020

Citation: Xu N, Cui Y, Dong J, Huang L. Exploring the molecular mechanisms of pterygium by constructing lncRNA–miRNA–mRNA regulatory network. *Invest Ophthalmol Vis Sci.* 2020;61(8):12. <https://doi.org/10.1167/iovs.61.8.12>

**PURPOSE.** This research explores the aberrant expression of the long non-coding RNA (lncRNA), microRNA (miRNA), and messenger RNA (mRNA) in pterygium. A competitive endogenous RNA (ceRNA) network was constructed to elucidate the molecular mechanisms in pterygium.

**METHODS.** We obtained the differentially expressed mRNAs based on three datasets (GSE2513, GSE51995, and GSE83627), and summarized the differentially expressed miRNAs (DEmiRs) and differentially expressed lncRNAs (DELS) data by published literature. Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway, protein-protein interaction (PPI), and gene set enrichment analysis (GSEA) analysis were performed. DEmiRs were verified in GSE21346, and the regulatory network of hub mRNAs, DELs, and DEmiRs were constructed.

**RESULTS.** Overall, 40 upregulated and 40 downregulated differentially expressed genes (DEGs) were obtained. The KEGG enrichment showed the DEGs mainly involved in extracellular matrix (ECM)-receptor interaction, focal adhesion, and PI3K-Akt signaling pathway. The GSEA results showed that cornification, keratinization, and cornified envelope were significantly enriched. The validation outcome confirmed six upregulated DEmiRs (miR-766-3p, miR-184, miR-143-3p, miR-138-5p, miR-518b, and miR-1236-3p) and two downregulated DEmiRs (miR-200b-3p and miR-200a-3p). Then, a ceRNA regulatory network was constructed with 22 upregulated and 15 downregulated DEmiRs, 4 downregulated DELs, and 26 upregulated and 33 downregulated DEGs. The network showed that lncRNA SNHG1/miR-766-3p/FOS and some miRNA-mRNA axes were dysregulated in pterygium.

**CONCLUSIONS.** Our study provides a novel perspective on the regulatory mechanism of pterygium, and lncRNA SNHG1/miR-766-3p/FOS may contribute to pterygium development.

**Keywords:** pterygium, bioinformatics analysis, competing endogenous RNA

Pterygium is a prevalent ocular surface disease that occurs most frequently in tropical equatorial areas.<sup>1,2</sup> It comprises a wing-shaped progressively growing fibrovascular tissue usually located on the nasal side, and could lead to visual impairment, astigmatism, and dry eye.<sup>3</sup> Over the years, various medical measures, such as anti-inflammatory eye drops and chemotherapeutic agents have been used in the treatment of pterygium. Surgical removal can be performed if the patient desires symptomatic or cosmetic improvement, but recurrence remains the main complication. The recurrence rate of different surgical techniques ranges from 0% to 88%.<sup>4–6</sup> Hence, elucidating the pathogenesis and molecular mechanisms of pterygium is crucial for improving surgical outcomes and decreasing the risk of recurrence.

Previous clinical and laboratory studies showed that immunologic mechanisms, extracellular matrix (ECM) modulation under ultraviolet radiation exposure, cell proliferation and hyperplasia, inflammation, angiogenesis, chole-

sterol metabolism modification, and hereditary factors attributed to the pathogenesis of pterygium.<sup>7,8</sup> Moreover, in recent years, the discovery of non-coding RNAs (regulatory RNAs) made the molecular etiology of pterygium more complex.

MicroRNA (miRNA) is a group of single-stranded noncoding RNA (ncRNA) that downregulates gene expression at the post-transcriptional level by inhibiting translation or promoting degradation of target messenger RNA (mRNAs). Long non-coding RNA (lncRNA) also belongs as a member of the noncoding RNA family, which is longer than 200 nt in length and has little or no protein-coding ability. MiRNAs and lncRNAs are involved in many physiological and pathological conditions. Recent studies have demonstrated that lncRNAs can work as competitive endogenous RNA (ceRNA) with miRNAs to compete with mRNAs for binding with miRNAs, thus affecting gene expression.<sup>9</sup> However, the research of core RNAs and lncRNA-miRNA-mRNA regulatory

networks in pterygium using bioinformatics analysis was still devoid.

In the present study, we utilized mRNA microarray dataset from Gene Expression Omnibus (GEO) to obtain and analyze differentially expressed genes (DEGs) and pooled the expression profiling of differentially expressed miRNAs (DEmiRs) and lncRNAs (DELS) between pterygium and normal conjunctiva tissues. Afterward, gene ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis, protein-protein interaction (PPI) network, and gene set enrichment analysis (GSEA) were performed. Furthermore, lncRNA-miRNA-mRNA regulatory networks were explored using StarBase and DIANA-LncBase, which helped us understand the pathogenesis and molecular mechanisms of pterygium.

## METHODS

### Subjects and Gene Information

The GEO is a national center for genetic information database, including a large number of gene chips, methylation, and sequencing data.<sup>10</sup> The including criteria: 1. Samples were obtained from Homo sapiens. 2. The chip data included both pterygium and normal conjunctiva tissues. 3. Chip data belonged to different independent studies, and all chip data did not contain each other. In this study, three gene expression profiles (GSE2513, GSE51995, and GSE83627) were searched and selected from the GEO database. In brief, GSE2513 consisted of 4 conjunctiva samples and 8 pterygium samples, which were harvested from 7 Chinese and 7 non-Chinese with age distribution of 42 to 57 years. In GSE51995, four primary nasal pterygium and four uninvolved conjunctiva tissues were collected from the superior temporal quadrant of the same eye. GSE83627 contained four donor-matched pterygium and conjunctiva tissues without mentioning the other clinical information.

### Data Analysis and DEGs Screening

We used the GEO mirror of R packages to get the expression matrix of GSE2513, GSE51995, and GSE83627 from the GEO Dataset. Then the expression matrixes were normalized and differential genes were screened by limma package (<http://www.bioconductor.org/packages/release/bioc/html/limma.html>)<sup>11</sup> between pterygium and conjunctiva samples. The  $|\log_2 \text{Fold change}| > 1.5$  and adjust  $P$  value  $< 0.05$  were used as the selection criteria.

### Enrichment Analysis of Gene Functions

GO<sup>12</sup> enrichment analysis and KEGG<sup>13</sup> enrichment analysis are the two most widely used analysis strategies for gene functions. The basic unit of GO is "term," which can be used for identifying cellular component (CC), molecular function (MF), and biological process (BP). The analysis of KEGG enrichment can show the main enrichment pathways of DEGs. In order to identify the GO annotations and pathways in which relevant DEGs were enriched, GO term and KEGG pathway enrichment analyses were performed with the Org.Hs.eg.db packages (<http://www.bioconductor.org/packages/release/data/annotation/html/org.Hs.eg.db.html>). The adjusted  $P$  value  $< 0.05$  was served to distinguish significant enriched genes.

## PPI Network Visualization

PPI network of DEGs was simulated to screen out hub proteins, which played a key role in the progress of pterygium. The Search Tool for the Retrieval of Interacting Genes (STRING) database<sup>14</sup> and Cytoscape<sup>15</sup> were utilized for the visualization of PPI network. The STRING database covers a large number of information about proteins, including the results of protein interaction, three-dimensional structure, and related functional enrichment. Through this database, the network structure of multiple DEGs can be constructed.

## Analysis of GSEA

GSEA<sup>16</sup> analysis was conducted in order to avoid missing the genes that actually play a crucial part during the process of screening out DEGs among three sets of data. The normal conjunctiva tissue was class A, and pterygium tissue was class B. Gene set permutations were performed 1000 times for each analysis. Absolute value of normalized enrichment score (NES)  $> 1$  and nominal  $P$  value  $< 0.05$  were considered as the threshold for statistical significance.

## Construction of lncRNA-miRNA-mRNA Regulatory Network

We identified miRNA through miRBase (<http://www.mirbase.org>),<sup>17</sup> and constructed miRNA-mRNA regulatory network by screening Targetscan ([http://www.targetscan.org/vert\\_72/](http://www.targetscan.org/vert_72/))<sup>18</sup> and miRDB (<http://mirdb.org/>).<sup>19</sup> Both Targetscan and miRDB provided a wide-range of information on the interaction between miRNA and mRNA. Then, predicted lncRNAs interacted with miRNAs were constructed in downloaded databases StarBase version 2.0<sup>20</sup> and DIANA-LncBase version 2.0,<sup>21</sup> both of which provided the experimentally validated lncRNA - miRNA interaction effect. DELS and DEmiRs were acquired by searching PubMed database for recent studies on pterygium.

## Validation of Key DEmiRs

We made a systemic search on PubMed and summarized the DEmiRs that met the criteria of  $P$  value  $< 0.05$  and  $|\log_2 \text{FC}| > 1$ . The validation procedure proceeded in GSE21346, which consisted of three pterygium samples and three matched conjunctiva samples from patients diagnosed with primary pterygium. The dataset was based on GPL7723, from which we could match and compare the DEmiRs with those previously reported. The DEmiRs were considered as significant difference with  $P < 0.05$ .

## RESULTS

### Screening of DEGs

A total of 422, 1374, and 420 DEGs were identified from the GSE2513, GSE51995, and GSE83627 datasets, respectively. The specific filtering results of each dataset are shown in Table 1. A total of 40 upregulated genes were found in all 3 datasets and 40 downregulated genes were found in 2 datasets (Figs. 1A, 1B). The heatmap of 80 DEGs in GSE51995 was shown in Figure 2.

TABLE 1. Specific Filtering Results of Each Data Set

GEO Dataset	Up	Down	All
GSE2513	272	150	422
GSE51995	860	514	1374
GSE83627	420	0	420

### GO and KEGG Pathway Enrichment Analysis

GO analysis of individual DEGs and KEGG pathway enrichment analysis were performed by R software to obtain more insightful details into the diverse functions of particular DEGs. The main MF with significant enrichment involved with all DEGs were ECM structural constituent, peptidase regulator activity, serine-type endopeptidase inhibitor activity, sulfur compound binding, heparin binding, and endopeptidase regulator activity (Supplementary Table S1), and the main BP were response to steroid hormone, skin development, epidermis development, keratinocyte differentiation, and epidermal cell differentiation, etc. (Supplementary Table S2). The results of GO enrichment analysis of upregulated and downregulated DEGs were shown in Figures 3A, 3B. The upregulated genes were mainly related to ECM-receptor interaction, focal adhesion, PI3K-Akt signaling pathway, regulation of actin cytoskeleton, primary bile acid biosynthesis, and one carbon pool by folate, and the downregulated genes were mainly related to osteoclast differentiation, glycerolipid metabolism, and IL-17 signaling pathway (Table 2, Supplementary Tables S3A, S3B).

### PPI Network Visualization

The 80 DEGs were introduced into the STRING online tool, and the proteins that did not interact with any other protein were removed to obtain the final protein interaction diagram (Fig. 4). A total of 58 nodes and 128 edges were selected to plot the PPI network, which consisted of 31 upregulated genes and 27 downregulated genes. Subsequently, 20 hub DEGs were screened out with degree  $\geq 5$ , which might play a meaningful role in the development of pterygium (Table 3). With the help of MCODE plug-in, the top three sub-networks with most importance were analyzed (Figs. 5A–5C).

### GSEA Enrichment of all mRNAs in GSE2513

GSEA was applied to analyze the main gene sets enriched on pterygium. The results showed that cornification, keratinization, cornified envelope, peptide cross linking, extracellular structure organization, collagen fibril organization, ECM structural constituent conferring tensile strength, complex of collagen trimers, ECM component, and vascular smooth muscle contraction were top 10 gene sets with the largest NES, and the details were reported in Figure 6 and Supplementary Table S4.

### DEmiRs and DELs in Pterygium

Through searching the results of 9 studies<sup>22–30</sup>, there were 49 DEmiRs that met the criteria of  $P$  value  $< 0.05$  and  $|\log_2FC| > 1$ , all of these DEmiRs' information was shown in Table 4. A total of 26 DEmiRs were upregulated and 23 DEmiRs were




FIGURE 1. Venn diagram DEGs were selected with a fold change  $> 1.5$  and adjust  $P$ -value  $< 0.05$  among the mRNA expression profiling datasets. (A) 40 up-regulated genes shared among GSE83627, GSE51995 and GSE2513. (B) Forty downregulated genes shared among GSE51995 and GSE2513.






FIGURE 2. The heatmap of 80 DEGs in GSE2513, GSE51995, and GSE83627.

A



B




**FIGURE 3.** Gene ontology analysis of the DEGs. (A) GO enrichment of downregulated DEGs. (B) GO enrichment analysis of upregulated DEGs. BP, biological process; CC, cellular component; MF, molecular function. The X axis represents the number of DEGs involved in GO terms.

**TABLE 2.** Mainly Pathways Involved With all DEGs

ID	Pathways, All Genes	Counts	P Value	Gene
hsa04512	ECM-receptor interaction	4	1.30E-03	COL1A1, FN1, SPP1, TNC
hsa04510	Focal adhesion	5	4.33E-03	ACTN1, COL1A1, FN1, SPP1, TNC
hsa04380	Osteoclast differentiation	3	3.26E-02	FOS, FOSB, JUNB
hsa04210	Apoptosis	3	3.80E-02	CTSV, FOS, PMAIP1
hsa04371	Apelin signaling pathway	3	3.87E-02	SPP1, EGRI, PPARGC1A
hsa04151	PI3K-Akt signaling pathway	5	4.27E-02	COL1A1, FN1, NTRK2, SPP1, TNC
hsa00561	Glycerolipid metabolism	2	4.39E-02	CEL, LIPC

**TABLE 3.** Top 20 Hub Genes With Degree  $\geq 5$

Gene Symbol	Degree	LogFC (GSE2513)	LogFC (GSE51995)	LogFC (GSE83627)
FN1	22	1.904340196	1.919574909	1.501942764
EGRI	14	-2.759612281	-2.594288111	N/A
FOS	13	-5.825731498	-3.270838601	N/A
SPP1	9	1.787503117	2.209450969	1.588880771
ATF3	9	-2.96106942	-1.73277313	N/A
PI3	9	0.984055795	3.690047454	3.052663221
BTG2	8	-0.814282557	-1.385548881	N/A
COL1A1	8	1.493514951	2.011230519	1.536520954
JUNB	7	-0.7886312	-1.316569032	N/A
FOSB	7	-5.160410507	-2.97242355	N/A
SPRR1B	7	2.425475595	3.373383589	2.403416753
DUSP1	7	-1.749318839	-0.634950586	N/A
DSG3	7	0.800044612	1.86852976	1.421462526
DSC2	7	0.651243446	1.515902315	1.077400291
LTBP1	6	0.696185299	1.793949203	1.264300642
SPRR3	6	2.950888463	3.831094018	2.647137921
SPRR2B	6	1.191061582	2.802224378	1.883683433
ZFP36	6	-1.45025058	-1.476260495	N/A
GRP	5	0.659772976	6.281685003	5.023794136
MCAM	5	1.503220261	0.93347822	0.655456146



**FIGURE 4.** Protein-protein interaction network of DEGs. Note: A circle represents a protein, and an edge represents a degree. The higher the degree is, the more important the protein is in the network structure.

downregulated. From the research of Liu, 20 DELs were acquired (Table 5), among which 10 DELs were upregulated and 10 DELs were downregulated.

**lncRNA-miRNA-mRNA Regulatory Network**

In the light of the DEMiR-DEL and DEMiR-DEG interactive pairs, the pterygium related lncRNA-miRNA-mRNA network was established in Figure 7, including 22 upregulated and 15 downregulated DEMiRs, 4 downregulated DELs, and 26 upregulated and 33 downregulated DEGs.

**Validation of Key DEMiRs**

In total, 6 upregulated DEMiRs (miR-766-3p, miR-184, miR-143-3p, miR-138-5p, miR-518b, and miR-1236-3p) and 2 downregulated DEMiRs (miR-200b-3p and miR-200a-3p) were verified in GSE21346, with a significant difference ( $P < 0.05$ ; Fig. 8). The expression trends of these eight DEMiRs are consistent with our ceRNA network results.

**DISCUSSION**

Pterygium is one of the most common ocular surface diseases. Epidemiological observations have suggested




FIGURE 5. Top three sub-networks in PPI.




FIGURE 6. Enrichment plots by GSEA.

TABLE 4. DE miRNAs of 9 Studies

	Annotation	Log <sub>2</sub> (Fold change)	P Value	
İçme G2019 <sup>22</sup>	miR-182-5p	4.37	<0.0001	
	miR-183-5p	4.15	0.01	
	miR-184	3.00	0.01	
	miR-221-3p	-1.41	0.02	
	miR-143-3p	2.4	N/A	
Lee 2016 <sup>23</sup>	miR-181a-2-3p	3.4	N/A	
	miR-377-5p	2.1	N/A	
	miR-411-5p	3.9	N/A	
	miR-145-3p	2.1	N/A	
	miR-145-5p	1.28	NA	
Teng 2018 <sup>24</sup>	miR-143-3p	UP	0.017	
	miR-145-5p	UP	0.028	
	miR-138-5p	3.024437	0.019	
Lan 2015 <sup>25</sup>	miR-215-3p	-1.8617594	0.028	
	miR-518b	1.7451285	0.032	
	miR-1236-3p	1.6934906	0.047	
Han 2019 <sup>26</sup>	miR-766-3p	1.5764362	0.017	
	miR-218-5p	DOWN	<0.01	
Li 2018 <sup>27</sup>	miR-21-5p	UP	<0.01	
	miR-1246	4.5	0.001	
	miR-486-3p	4.4	0.004	
	miR-451	4.1	0.010	
	miR-3175	3.3	<0.001	
	miR-1972	3.0	<0.001	
	miR-143-3p	2.7	0.008	
	miR-211-5p	2.7	0.030	
	miR-665	2.3	0.010	
	miR-1973	2.2	0.040	
	Engelsvold 2013 <sup>28</sup>	miR-18a-5p	2.1	0.004
		miR-143-5p	2.0	0.006
		miR-663b	2.0	0.020
		miR-675-5p	-2.0	0.005
		miR-200b-3p	-2.1	0.002
		miR-200b-5p	-2.3	<0.001
		miR-200a-3p	-2.7	0.002
		miR-200a-5p	-2.3	<0.001
		miR-29b-3p	-2.3	0.005
		miR-210-3p	-2.4	<0.001
Wu 2014 <sup>29</sup>	miR-141-3p	-2.5	<0.001	
	miR-31-5p	-2.6	0.020	
	miR-934	-3.0	<0.001	
	miR-375	-3.7	0.030	
	miR-221	DOWN	<0.0001	
	miR-1298-5p	1.36	0.019	
	miR-122-3p	-3.87	0.005	
	miR-122-5p	-2.88	0.037	
	miR-192-3p	-1.74	0.006	
	miR-192-5p	-2.29	<0.001	
Cui 2016 <sup>30</sup>	miR-194-5p	-2.51	0.015	
	miR-302f	-1.50	0.006	
	miR-802	-2.03	0.039	
	miR-1973	-1.27	0.036	
	miR-5000-3p	-1.3	0.04	

TABLE 5. DELs of 1 Research

	Annotation	Fold Change
Liu 2016 <sup>31</sup>	FOXD2-AS1	92.68
	RP11-78F17.1	88.76
	RP11-702F3.4	82.01
	RP5-963E22.4	75.08
	RP11-611E13.2	70.63
	AF196972.9	66.01
	KIAA0664L3	63.47
	LOC283761	61.28
	RP3-416J7.2	58.06
	LOC100130264	56.05
	LINC00638	0.0088
	WI2-2373I1.2	0.0096
	ARL6IP6	0.011
	CLCA4	0.015
TECR	0.017	
RP11-398J10.2	0.018	
RP11-61L23.2	0.021	
RP11-420A23.1	0.022	
RP11-217B1.2	0.023	
RP3-434O14.8	0.026	

In this study, we identified a total of 80 DEGs (40 upregulated and 40 downregulated mRNAs), and hub-genes with 3 subnetworks with most importance in PPI, including FN1, PI3, ERG1, SPRR1B, FOS, and FOSB, suggesting they may play a very important role in the pathogenesis of pterygium. SPRR1B belongs to keratinocyte protein, and their mRNA transcripts in conjunctival tissues increased in response to desiccating stress, which is associated with pterygium recurrence.<sup>33</sup> FOS and FOSB encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. The encoded proteins have been implicated as regulators of cell proliferation, and experimental reports describe a tumor-suppressive function in various tumors.<sup>34,35</sup> EGR1 was found to suppress cell survival, proliferation, and activates expression of p53 and TGF-beta,<sup>36</sup> which was proved to play a pivotal role in the occurrence of pterygium.<sup>8,37,38</sup> FN1 and COL1A1 were key molecules in EMT, and were demonstrated to be involved in the pathogenic mechanism of pterygium.<sup>28</sup> Further studies were required to elucidate the complex interaction with these genes and clinical features.

In this study, the results of functional enrichment analysis indicated that ECM-receptor interaction, focal adhesion, apoptosis, and PI3K-Akt signaling pathway were involved with significant DEGs. These results were in agreement with previous proteomics study by Hou,<sup>39</sup> in which they compared the protein expression from the conditioned medium of paired pterygium and normal conjunctival fibroblast cells from the same patients by iTRAQ-based proteomics strategy, and they found the differenced protein might serve as extracellular ligands to activate intracellular pathways. Aberrant ECM expressions were found to be a major characteristic feature of pterygium. ECM and its receptors, including fibronectin, versican, collagen III, and SPARC, were shown to be upregulated and remodeled in pterygium,<sup>40-42</sup> and they have turned out to be regulated by matrix metalloproteinase (MMP), an essential enzyme in local proteolysis of the extracellular matrix.<sup>43</sup> UV irradiation may lead to the imbalance of MMPs and its inhibitors TIMP, which enable the pterygial cells to dissolve corneal epithelium

that environmental alteration, like ultraviolet radiation, is the most important factor contributing to this disease.<sup>32</sup> However, its exact pathogenesis remains unknown. The development of high throughput microarray technology with high efficiency and high accuracy combined with bioinformatic algorithm allows us to identify key genes, which could provide a deeper understanding of molecular mechanisms on pterygium.




FIGURE 7. LncRNA-miRNA-mRNA regulatory network. Note: *Circles* represent DEGs, *triangles* represent DEmiRs, and *chevrans* represent DEs. *Red* means upregulated and *blue* means downregulated.




FIGURE 8. Validation of key DEmiRs in GSE21346. Note: *Green* represents the control group, and *purple* represents the pterygium group. \*  $P < 0.05$ .

and Bowman's layer and invade the corneal stroma.<sup>44–46</sup> Besides, impression cytology specimens of pterygium show fewer apoptotic markers, and PI3K-Akt inhibitor has been proved to reduce the TGF- $\beta$ -induced synthesis of FN in human pterygium fibroblasts.<sup>47</sup> Moreover, our GSEA-based GO analysis confirmed keratinization and cornification pathway participated in the pathogenesis of pterygium. Previous reports have confirmed that the expression of keratin markers is upregulated in the epithelium of pterygium, and takes effect in the epithelial abnormal differentiation.<sup>48–50</sup> Liu constructed pterygium-related mRNA libraries by using microarray and found upregulated mRNAs were closely related to proliferation and differentiation.<sup>51</sup> If our results were combined with publications mentioned above, we could surmise the underlying mechanisms of pterygium that dysregulation of ECM proteins caused by increased expression of MMP might activate intracellular PI3K-Akt signaling

pathway, leading to anti-apoptosis of fibroblast, abnormal matrix protein deposition, and hyperproliferation and keratinization of pterygial cells.

It has been discovered that a group of miRNAs, including miR-200,<sup>28</sup> miR-145,<sup>24,52</sup> miR-122,<sup>50</sup> and miR-21,<sup>27</sup> were shown to contribute to the development of pterygium. In the current study, by summarizing 10 relevant studies in recent years, we found a total of 26 upregulated and 23 downregulated DEmiRs. Moreover, by establishing the miRNA-mRNA network, we found miR-215-3p significantly downregulated and directly regulated seven DEGs, including the hub gene FN. A previous research has shown that miR-215-3p takes part in regulating cell cycling and inhibiting proliferation of primary fibroblast cells from the ocular surface, and was downregulated in pterygium.<sup>25</sup> Because FN has been reported to be a key gene of epithelial mesenchymal transition (EMT) and was also identified as a target of



miR-200b,<sup>28</sup> we speculated that downregulated miR-215-3p, along with miR-200b, might play important roles in pterygium by regulating EMT.

Previous studies have identified the roles of various lncRNAs in regulating pterygium proliferation and apoptosis,<sup>31</sup> but no study was reported about lncRNA-associated ceRNA network based on bioinformatics analysis in pterygium. Combining the results of DEmiR-DEL, DEmiR-DEG interactive network, and hub genes produced by the STRING, we found that the expression of lncRNA SNHG1, FOS, and FOSB were downregulated, whereas the expression of has-miR-766-3p was upregulated, which was consistent with the mechanism ceRNA hypothesis. lncRNA SNHG1 is a marker of tumor progression,<sup>53</sup> and it has been proved to negatively regulate p53, which is a tumor suppressor protein that could induce apoptosis and show strong expression in pterygium.<sup>38</sup> Downregulation of lncRNA SNHG1 could explain p53 reactivation in pterygium. MiR-766-3p has also been reported to act as a key gene in different tumors and immune diseases, and its expression was elevated in hepatocellular carcinoma, inflamed pulp, and acute promyelocytic leukemia.<sup>53-55</sup> This miRNA contributes to anti-inflammatory responses, cell proliferation, and apoptosis by targeting different downstream genes and signaling pathways, but the precise post-transcriptional mechanisms of miR-766-3p remain to be explored in pterygium. Further work will be needed with multiple clinical samples to clarify the ceRNA hypothesis that lncRNA SNHG1 regulates FOS and FOSB to act on the progression of pterygium through sponging miR-766-3p.

There were also limitations in our study. The sample size of gene expression profile was not large, and no microarray data of dysregulated lncRNAs in pterygium was found in GEO and Array Express databases. In addition, the platform of GSE21346 was published 10 years ago with limited probe annotation for miRNA profile. All these may prevent us from finding a comprehensive noncoding RNA expression profile and related regulatory network. Although we proved that the expression of matched miRNAs on GSE21346 were consistent with previous studies, the mechanism and validation of these ncRNAs in pterygium still need further research in clinical and molecular biology experiments.

To sum up, we identified core genes, and related crucial pathways, especially PI3K-Akt, keratinization, and cornification pathway involved in the pathogenesis of pterygium. Furthermore, we summarized all the studies on the miRNA and lncRNA related to pterygium from PubMed database and established lncRNA-miRNA-mRNA regulatory network. This study might deepen the understanding of potential molecular mechanism underlying pterygium and provide some new insights for use in further identification and development of new therapeutic targets for pterygium.

### Acknowledgments

Supported by the Fujian Provincial Natural Science Foundation (Grant numbers 2018J01238; 2018J01310; and 2019J01152), the Joint Funds for the Innovation of Science and Technology of Fujian province (Grant number 2018Y9035), the Startup Fund for Scientific Research of Fujian Medical University (Grant number 2017XQ1022), and the Youth Foundation of Fujian Provincial Health Commission (Grant number 2019-1-31).

Disclosure: **N. Xu**, None; **Y. Cui**, None; **J. Dong**, None; **L. Huang**, None

### References

- Durkin SR, Abhary S, Newland HS, Selva D, Aung T, Casson RJ. The prevalence, severity and risk factors for pterygium in central Myanmar: the Meiktila Eye Study. *Br J Ophthalmol*. 2008;92:25-29.
- McCarty CA, Fu CL, Taylor HR. Epidemiology of pterygium in Victoria, Australia. *Br J Ophthalmol*. 2000;84:289-292.
- Liu C, Song Y, Wang X, et al. The key role of VEGF in the cross talk between pterygium and dry eye and its clinical significance. *Ophthalmic Res*. 2020;63:320-331.
- Frucht-Pery J, Raiskup F, Ilsar M, Landau D, Orucov F, Solomon A. Conjunctival autografting combined with low-dose mitomycin C for prevention of primary pterygium recurrence. *Am J Ophthalmol*. 2006;141:1044-1050.
- Chen PP, Ariyasu RG, Kaza V, LaBree LD, McDonnell PJ. A randomized trial comparing mitomycin C and conjunctival autograft after excision of primary pterygium. *Am J Ophthalmol*. 1995;120:151-160.
- Clearfield E, Muthappan V, Wang X, Kuo IC. Conjunctival autograft for pterygium. *Cochrane Database Syst Rev*. 2016;2:CD011349.
- Bradley JC, Yang W, Bradley RH, Reid TW, Schwab IR. The science of pterygia. *Br J Ophthalmol*. 2010;94:815-820.
- Chu WK, Choi HL, Bhat AK, Jhanji V. Pterygium: new insights. *Eye (Lond)*. 2020;34:1047-1050.
- Cesana M, Cacchiarelli D, Legnini I, et al. A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell*. 2011;147:358-369.
- Barrett T, Suzek TO, Troup DB, et al. NCBI GEO: mining millions of expression profiles—database and tools. *Nucleic Acids Res*. 2005;33:D562-D566.
- Ritchie ME, Phipson B, Wu D, et al. Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*. 2015;43:e47.
- Ashburner M, Ball CA, Blake JA, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25:25-29.
- Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. New approach for understanding genome variations in KEGG. *Nucleic Acids Res*. 2019;47:D590-D595.
- Szklarczyk D, Franceschini A, Wyder S, et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res*. 2015;43:D447-D452.
- Shannon P, Markiel A, Ozier O, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*. 2003;13:2498-2504.
- Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102:15545-15550.
- Kozomara A, Birgaoanu M, Griffiths-Jones S. miRBase: from microRNA sequences to function. *Nucleic Acids Res*. 2019;47:D155-D162.
- Agarwal V, Bell GW, Nam JW, Bartel DP. Predicting effective microRNA target sites in mammalian mRNAs. *eLife*. 2015;4:e05005.
- Wong N, Wang X. miRDB: an online resource for microRNA target prediction and functional annotations. *Nucleic Acids Res*. 2015;43:D146-D152.
- Li JH, Liu S, Zhou H, Qu LH, Yang JH. starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Nucleic Acids Res*. 2014;42:D92-D97.
- Paraskevopoulou MD, Vlachos IS, Karagkouni D, et al. DIANA-LncBase v2: indexing microRNA targets on non-coding transcripts. *Nucleic Acids Res*. 2016;44:D231-D238.
- Ice G, Yilmaz A, Dinc E, Gorur A, Fidanci SB, Tamer L. Assessment of miR-182, miR-183, miR-184, and miR-221

- expressions in primary pterygium and comparison with the normal conjunctiva. *Eye Contact Lens*. 2019;45:208–211.
23. Lee JH, Jung SA, Kwon YA, Chung JL, Kim US. Expression of microRNAs in fibroblast of pterygium. *Int J Ophthalmol*. 2016;9:967–972.
  24. Teng Y, Yam GH, Li N, et al. MicroRNA regulation of MDM2-p53 loop in pterygium. *Exp Eye Res*. 2018;169:149–156.
  25. Lan W, Chen S, Tong L. MicroRNA-215 regulates fibroblast function: insights from a human fibrotic disease. *Cell Cycle*. 2015;14:1973–1984.
  26. Han S, Chen Y, Gao Y, Sun B, Kong Y. MicroRNA-218-5p inhibit the migration and proliferation of pterygium epithelial cells by targeting EGFR via PI3K/Akt/mTOR signaling pathway. *Exp Eye Res*. 2019;178:37–45.
  27. Li X, Dai Y, Xu J. MiR-21 promotes pterygium cell proliferation through the PTEN/AKT pathway. *Mol Vis*. 2018;24:485–494.
  28. Engelsen DH, Utheim TP, Olstad OK, et al. miRNA and mRNA expression profiling identifies members of the miR-200 family as potential regulators of epithelial-mesenchymal transition in pterygium. *Exp Eye Res*. 2013;115:189–198.
  29. Wu CW, Cheng YW, Hsu NY, et al. MiRNA-221 negatively regulated downstream p27Kip1 gene expression involvement in pterygium pathogenesis. *Mo Vis*. 2014;20:1048–1056.
  30. Cui YH, Li HY, Gao ZX, et al. Regulation of apoptosis by miR-122 in pterygium via targeting Bcl-w. *Invest Ophthalmol Vis Sci*. 2016;57:3723–3730.
  31. Liu J, Ding X, Yuan L, Zhang X. Identification of pterygium-related long non-coding RNAs and expression profiling by microarray analysis. *Int J Mol Med*. 2016;38:529–536.
  32. Ang M, Li X, Wong W, et al. Prevalence of and racial differences in pterygium: a multiethnic population study in Asians. *Ophthalmology*. 2012;119:1509–1515.
  33. Tan J, Vollmer-Conna U, Tat L, Coroneo M. Dry-eye disease in recurrent pterygium. *Ophthalmic Res*. 2019;61:199–203.
  34. Tang C, Jiang Y, Shao W, et al. Abnormal expression of FOSB correlates with tumor progression and poor survival in patients with gastric cancer. *Int J Oncol*. 2016;49:1489–1496.
  35. Durchdewald M, Angel P, Hess J. The transcription factor Fos: a Janus-type regulator in health and disease. *Histol Histopathol*. 2009;24:1451–1461.
  36. Liu C, Rangnekar VM, Adamson E, Mercola D. Suppression of growth and transformation and induction of apoptosis by EGR-1. *Cancer Gene Ther*. 1998;5:3–28.
  37. Di Girolamo N, Chui J, Coroneo MT, Wakefield D. Pathogenesis of pterygia: role of cytokines, growth factors, and matrix metalloproteinases. *Prog Retin Eye Res*. 2004;23:195–228.
  38. Weinstein O, Rosenthal G, Zirkin H, Monos T, Lifshitz T, Argov S. Overexpression of p53 tumor suppressor gene in pterygia. *Eye (Lond)*. 2002;16:619–621.
  39. Hou A, Law KP, Tin MQ, Lim YP, Tong L. In vitro secretomics study of pterygium-derived fibroblasts by iTRAQ-based quantitative proteomics strategy. *Exp Eye Res*. 2016;153:14–22.
  40. Naib-Majani W, Eltohami I, Wernert N, et al. Distribution of extracellular matrix proteins in pterygia: an immunohistochemical study. *Graefes Arch Clin Exp Ophthalmol*. 2004;42:332–338.
  41. Seet LF, Tong L, Su R, Wong TT. Involvement of SPARC and MMP-3 in the pathogenesis of human pterygium. *Invest Ophthalmol Vis Sci*. 2012;53:587–595.
  42. John-Aryankalayil M, Dushku N, Jaworski CJ, et al. Microarray and protein analysis of human pterygium. *Mol Vis*. 2006;12:55–64.
  43. Nagase H, Woessner JF Matrix metalloproteinases. *J Biol Chem*. 1999;274:21491–21494.
  44. Tsai YY, Chiang CC, Yeh KT, Lee H, Cheng YW. Effect of TIMP-1 and MMP in pterygium invasion. *Invest Ophthalmol Vis Sci*. 2010;51:3462–3467.
  45. Ardan T, Nemcova L, Bohuslavova B, et al. Reduced levels of tissue inhibitors of metalloproteinases in UVB-irradiated corneal epithelium. *Photochem Photobiol*. 2016;92:720–727.
  46. Di Girolamo N, Coroneo M, Wakefield D. Epidermal growth factor receptor signaling is partially responsible for the increased matrix metalloproteinase-1 expression in ocular epithelial cells after UVB radiation. *Am J Pathol*. 2005;167:489–503.
  47. Chen K, Lai K, Zhang X, et al. Bromfenac inhibits TGF-beta1-induced fibrotic effects in human pterygium and conjunctival fibroblasts. *Invest Ophthalmol Vis Sci*. 2019;60:1156–1164.
  48. Wang G, Dong N, Wang Y, et al. Epithelial dysplasia in pterygium postoperative granuloma. *Exp Eye Res*. 2018;175:199–206.
  49. Peng J, Sha XY, Liu Y, Yang RM, Wen Y. Pterygium epithelium abnormal differentiation related to activation of extracellular signal-regulated kinase signaling pathway in vitro. *Int J Ophthalmol*. 2015;8:1118–1125.
  50. Zhang M, Liu Z, Xie Y. The study on the expression of keratin proteins in pterygial epithelium. *Yan Ke Xue Bao*. 2000;16:48–52.
  51. Liu J, Ding X, Yuan L, Zhang X. Identification of pterygium-related mRNA expression profiling by microarray analysis. *Eye (Lond)*. 2017;31:1733–1739.
  52. Chien KH, Chen SJ, Liu JH, et al. Correlation of microRNA-145 levels and clinical severity of pterygia. *Ocul Surf*. 2013;11:133–138.
  53. Yang C, Ma X, Guan G, et al. MicroRNA-766 promotes cancer progression by targeting NR3C2 in hepatocellular carcinoma. *FASEB J*. 2019;33:1456–1467.
  54. Zhong S, Zhang S, Bair E, Nares S, Khan AA. Differential expression of microRNAs in normal and inflamed human pulps. *J Endod*. 2012;38:746–752.
  55. Liang H, Li X, Wang L, et al. MicroRNAs contribute to promyelocyte apoptosis in As2O3-treated APL cells. *Cell Physiol Biochem*. 2013;32:1818–1829.