



# Ruxolitinib In The Treatment Of Polycythemia Vera: An Update On Health-Related Quality Of Life And Patient-Reported Outcomes

This article was published in the following Dove Press journal:  
*Journal of Blood Medicine*

Shashank Cingam <sup>1</sup>  
Lainey Flatow-Trujillo<sup>2</sup>  
Leslie A Andritsos<sup>1</sup>  
Cecilia Arana Yi <sup>1</sup>

<sup>1</sup>University of New Mexico Comprehensive Cancer Center, Albuquerque, NM, USA; <sup>2</sup>Department of Internal Medicine, University of New Mexico School of Medicine, Albuquerque, NM, USA

**Abstract:** Polycythemia vera (PV) is a rare myeloproliferative neoplasm (MPN) associated with significant impairment in quality of life (QoL) due to disease-related symptoms and complications. Assessment of disease burden constitutes standard monitoring of symptoms and response. Conventional treatments for MPN, such as hydroxyurea, phlebotomy, or interferon, have not shown a significant impact in QoL or patient-reported outcomes (PRO). Ruxolitinib (RUX) is a JAK2 inhibitor approved for patients intolerant or resistant to hydroxyurea (HA). We conducted a systematic review of clinical trials of RUX in patients with PV that incorporated PRO measures to evaluate the effects on PRO and QoL. Three randomized Phase 3 studies reported in four publications were relevant for analysis. Although the small number of trials and potential for treatment bias in the review, treatment with RUX was associated with improved QoL and PRO in PV patients intolerant or resistant to hydroxyurea.

**Keywords:** polycythemia vera, ruxolitinib, quality of life, patient-reported outcomes

## Introduction

Polycythemia vera (PV) is a rare myeloproliferative neoplasm (MPN) characterized by increased red blood cell mass, bone marrow panmyelosis, and Janus kinase 2 (JAK2) mutation.<sup>1,2</sup> Constitutional symptoms such as fatigue, weight loss, night sweats and pruritus (so-called “cytokine symptoms”) are present in approximately 61% of patients with PV and affect the quality of life (QoL), overall health and productivity.<sup>3–6</sup> Healthcare utilization and costs are also higher in patients with PV than in non-cancer controls.<sup>7</sup>

Patient-reported outcomes (PRO) measures are tools for the objective evaluation of symptoms, physical and social functioning, and mental health and well-being.<sup>8</sup> Despite their pivotal importance in the understanding of disease- and treatment-related adverse events, information regarding PRO in hematologic malignancies remains sparse.<sup>9</sup> However, growing recognition of the symptom burden in PV has led to the incorporation of PRO instruments for symptom assessment into the standard evaluation of MPN’s (NCCN guidelines, version 2.2019).

In clinical trial settings, PRO instruments evaluate therapeutic responses, quality of life, and prognosis.<sup>10</sup>

The goals of PV therapy are to decrease thrombotic events, manage symptoms, minimize adverse effects of treatment, and reduce the risk of transformation to myelofibrosis or leukemia.<sup>1,11,12</sup> Treatments for PV include phlebotomy, aspirin,

Correspondence: Cecilia Arana Yi  
University of New Mexico  
Comprehensive Cancer Center, 1201  
Camino De Salud NE, Albuquerque, NM  
87131, USA  
Email caranayi@salud.unm.edu

and cytoreductive therapies such as hydroxyurea (HA), anagrelide, and interferon. HA decreases the hematocrit, reduces the spleen size, and lowers thrombotic risk, but does not reduce symptom burden in all patients.<sup>13,14</sup> Intolerance of or resistance to HA occurs in approximately 25% of patients and is associated with poor outcomes.<sup>15</sup>

Ruxolitinib (RUX) is an oral JAK1/JAK2 inhibitor approved for the management of myelofibrosis and for PV patients who are intolerant or resistant to HA, per modified ELN criteria.<sup>4,16,17</sup> In myelofibrosis, treatment with RUX may result in the reduction of spleen size, decreased symptom burden, and improvement in QoL measures.<sup>18,19</sup> In PV, the phase 3 RESPONSE study demonstrated the superiority of RUX compared to the best available therapy in the reduction of hematocrit, spleen size, and PV symptoms.<sup>4</sup> Since then, additional studies have explored the impact of RUX on symptom burden and QoL. Herein we present a review of the PRO literature in PV patients treated with RUX and propose future concepts for research.

## Methods

### Search Criteria

We conducted a systematic review in PubMed, Medline, EMBASE, and Cochrane of English-language scientific

articles using the MeSH-terms “randomized clinical trials” “polycythemia vera” “ruxolitinib,” “quality of life” and “patient-reported outcome” as title or abstract terms from January 2000 to January 2018. We reviewed the bibliographies of all retrieved papers to identify randomized controlled trials that studied the effect of RUX on symptoms and quality of life as primary or secondary endpoints. The exclusion criteria were the following:

- Non-English publication
- Publication before 2000
- Patients are younger than 18 years.
- Non- PV myeloproliferative neoplasms.
- Only abstract available.

We synthesized the data according to PRISMA guidelines. (Figure 1).<sup>20</sup> The quality of the study was assessed using the Mixed Methods Appraisal Tool (MMAT).<sup>21</sup>

### Data Extraction And Analysis

We extracted the following data: Demographics, number of patients enrolled, symptoms, adverse effects, missing data and results, and instruments of QoL and PRO assessments. (Table 1)




Figure 1 PRISMA diagram of clinical trials of ruxolitinib in polycythemia vera.

**Table 1** Common Instruments Used In Myeloproliferative Neoplasms For Quantitatively Measuring Symptoms And Quality Of Life

Abbreviated QoL Instrument	Full title	Scoring	Description	References
MPN-SAF	Myeloproliferative Neoplasm Symptoms Assessment Form	14 individual symptoms scores, multiplied by 10 to achieve 0–100 scale.	3 symptom clusters: Mechanisms related to cytokines (TSS-C); Hyperviscosity (TSS-H), and splenomegaly (TSS-S); MPN SAF is validated in PV.	Emanuel et al <sup>28</sup>
				Johansson et al <sup>33</sup>
				Scherber et al <sup>24</sup>
PSIS	Pruritus Symptom Impact Scale	Five point questionnaire. 10 point scale; zero equals “no itching/not bothered at all” and 10 equals “bothered as bad as you can imagine/interfered as bad as you can.”	Evaluates severity, interruption of daily life and improvement or worsening of their itching/pruritus since the start of treatment.	Vannucchi et al <sup>4</sup>
EORTC-QLQ-C30	European Organization for Research and Treatment of Cancer Quality of Life Core Questionnaire-for all cancer patients.	Multi-item scales and single-item measures. Five functional scales, three symptom scales, a global health status/QoL scale, and six single items. All of the scales and single-item measures range in score from 0 to 100.	High score represents a high/healthy status/QoL. High score for a symptom represents a high level of symptomatology. Validated for all cancers but does not capture some hematological symptoms.	Aaronson et al <sup>34</sup>
EQ-5D-5L	EuroQOL Group non-disease specific QoL instrument	Measures health outcomes for a wide range of health conditions and treatments. Descriptive system and a visual analogue scale (EQ VAS). Five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression.	Endpoints are labeled “best imaginable health state” and “worst imaginable health state.” Scores can be summarized into a single index score that provides a simple measure of health for clinical and economic appraisal.	Herdman et al <sup>35</sup>
WPPI	Work Productivity and Activity Impairment Questionnaire	Six-item questionnaire that measures self-reported productivity loss during the past seven days	Questions about absence from work, hours spent at work, reduction in productivity at work, and reduction in productivity while performing regular activities.	Relly et al <sup>36</sup>
PGIC	Patient Global Impression of Change	Scale from 1 to 7 from “very much improved” to “very much worse”	Evaluates all aspects of patients’ health and assesses in there has been an improvement or decline in clinical status	Dorwkin et al <sup>37</sup>

## Ethical Considerations

This study evaluated the published data and did not require institutional ethics board approval.

## Results

The initial search in PubMed, Medline, EMBASE, and Cochrane resulted in 40 scientific publications. Nine records were eligible for review. Two files were excluded because one

study was retrospective, and another was a Phase 2 study. (Figure 1)

Three Phase 3 multicenter randomized trials of RUX in PV included QoL as a primary or secondary endpoint. The RESPONSE trial evaluated RUX (n = 110) versus standard therapy (SOC) (n = 112) in patients with PV and splenomegaly who were resistant or intolerant to HA.<sup>4</sup> The primary endpoints of the study were hematocrit

control and spleen volume; symptom reduction, QoL and safety were secondary endpoints, with PRO analysis in the same patient cohort. Patients crossed over to RUX at 32 weeks.<sup>11</sup>

The RESPONSE 2 evaluated RUX (n = 74) versus BAT (n = 75) in patients with PV without splenomegaly who were resistant to or intolerant of HA.<sup>22</sup> The primary endpoint was hematocrit control. Secondary endpoints were symptom reduction, quality of life (QoL), and safety. Patients crossed over to RUX at 28 weeks if the primary endpoint was not met.

The RELIEF study evaluated RUX (n=54) versus HA (n=56) in patients who had achieved disease control with a stable dose of HA but who continued to experience PV-related symptoms and impairment in QoL.<sup>23</sup> The primary endpoint was a 50% improvement from baseline in myeloproliferative neoplasm related symptom assessment, the total symptom score and the cytokine symptom cluster at Week 16. Patients could cross over at 16 weeks. (Table 2)

## Quality Of Life And PRO Instruments

We measured and reported the outcomes of these studies using standard instruments. Six different QoL and PRO instruments were applied: MPN-SAF, PSIS, EORTC-QLQ-C30, EQ-5D-5L, WPAI, and PGIC were applied. (Table 1) MPN-SAF is the only instrument for the assessment of the prevalence and symptoms of MPNs.<sup>24</sup>

## The Methodological Quality Of Studies

The overall quality score was 100% (4 out of 4 criteria met) in the three studies, as shown in Table 2. One study was missing more than 20% of PRO data, and another had more than a 20% dropout rate.

The RESPONSE trial did not consider missing patient assessments for the primary endpoint but did not mention strategies for handling missing data. PRO instruments reported an improvement of symptoms in the RUX arm, although not powered for statistical comparisons. The study did not prevent potential patient treatment bias and reported outcomes in less than 80% of patients.

RESPONSE 2 trial had a trial profile describing the number of eligible and randomized patients. PRO studies reported symptom improvements on RUX, but the follow up was short and not statistically comparable.

RELIEF study was a randomized, double-blind, double-dummy study. At the time of randomization, the discontinuation rate was 13% in RUX versus 10.7% in the HA arm. At 16 weeks, only 64.8% continue RUX versus 64.3% HA, leading to a higher than 20% dropout rate. PRO instruments showed a

trend toward symptom improvement in the RUX arm that was not statistically significant. In addition to the limited number of cases, the follow-up time was only for 16 weeks, and there was a higher proportion of female patients in the HA arm.

## Effect Of Ruxolitinib On Patient-Reported Outcomes Symptom Control

The primary symptoms evaluated at baseline in these studies were fatigue, insomnia, pain, dyspnea, pruritus, myalgias, night sweats, and sweats while awake (cytokine cluster), visual disturbance, dizziness, concentration difficulties, headache, numbness/tingling in hands or feet, tinnitus, skin redness (hyperviscosity cluster), abdominal discomfort and early satiety (splenomegaly symptom cluster).

In the RESPONSE trial, patients in the RUX arm reported significant symptom improvement compared to SOC at week 32. 49% (36/74) of patients on RUX reported more than 50% improvement from baseline in MPN-SAF total symptom score at week 32, versus 5% (4/81) in the standard treatment arm, 4% (2/49) with HA and 6% (2/32) in the non-HA patients. In the cytokine cluster, symptom improvement was achieved in 64% (47/74) on RUX, versus 11% (9/80) on SOC, 4% (2/48) on HA, and 22% (7/32) on non-HA respectively. In the hyperviscosity cluster, 37% (26/71) on (RUX, versus 13% (10/80) on standard treatment, 12% (6/49) on HA, and 13% (4/31) on non-HA). In the splenomegaly cluster, 62% (39/63) on RUX, versus 17% (12/71) on SOC, 14% (6/44) on HA, 22% (6/27) on non-HA. Patients randomized to the RUX arm also showed rapid improvements in all five components of the PSIS compared to SOC.

In RESPONSE 2, patients randomized to the RUX arm reported improvements in all individual symptoms with RUX compared to BAT where most symptoms worsened at week 28. At week 80, patients randomized initially to RUX showed an increase in the majority of individual symptom scores, except abdominal discomfort and fever. At week 28, 45.3% (29/64) of patients in the RUX group had a 50% improvement in MPN-SAF TSS vs 22.7% (5/22) on BAT. The median percentage change from baseline in MPN-SAF TSS was -45.3% in the RUX group versus 2.4% in the BAT group at week 28, where negative scores indicate improvement. MPN-SAF TSS of at least 20 at baseline, 17 (50%) of 34 patients treated with RUX achieved complete resolution of disease-related symptoms compared with two (8%) of 26 patients treated with BAT. At week 80, 45% of patients randomized to RUX demonstrated a higher than 50% reduction in the

**Table 2** Published Studies Of Ruxolitinib In PV And Quality Of Life And PRO

Topic	Authors	Year	Patient Group	QoL Instruments	No Of Patients	Outcome	Main Symptom Topics	Limitations	Quality Assessment
QoL PRO	Vannucchi et al <sup>4</sup>	2015	PV, intolerant/ resistant to HA on RUX	MPN-SAF EORTC-QLQ-C30 PSIS PGIC	222 RUX=110 SOC=112	At week 32, 36/74 had 50% reduction of MPN SAF symptom score. Reduction in score by PSIS. EORTC QLQ-C30 scores were higher compared to standard. PGIC comparable to SOC.	Pruritus Tiredness, Night sweats Headaches Dizziness/Vision problems Muscle aches Abdominal discomfort Early satiety	Open-label, potential treatment bias	****>20% missing PRO data
QoL PRO	Mesa et al <sup>11</sup>	2016	PV, intolerant/ resistant to HA on RUX	EORTC-QLQ-C3 0MPN- SAFPSISPGIC	222 RUX=110 SOC=112	At week 32, 44% on RUX achieved MID (minimally important difference) in GHS/ QoL compared to 9% MPN SAF 49% achieved better total symptom score compared to 5% PSIS Improvement in the 5 components of PSIS as early as week 4 PGIC 46% condition improved at week 4 versus 11%	Fatigue Insomnia Pain Dyspnea Appetite loss Diarrhea, Nausea and vomiting Pruritus Role functioning Cognitive functioning Social functioning Physical functioning	Open label Not powered for statistical comparisons	****
QoL PRO	Passamonti et al <sup>38</sup>	2017	PV without splenomegaly, intolerant/ resistant to HA	MPN-SAF EuroQoL-5D-5L PSIS WPAl PGIC	173 RUX=74 BAT=75	Median percentage change from baseline (-45.3%) in RUX vs BAT. At week 28, 29/64 had 50% or greater reduction in MPN-SAF score. EQ-5D-5L no problems in 5 dimensions and improvement in work productivity. Reduction in score by PSIS. Improvements in scores from PGIC and WPAl.	Pruritus Overall QoL, Fatigue Activity impairment	Open-label	****
QoL PRO	Mesa et al <sup>23</sup>	2016	PV on HA ≥ 4 weeks with cytokine symptom cluster (TSS-C)	MPN-SAF PGIC	110 RUX=54 HA=56	At week 16, ≥50% improvement from baseline TSS-C in 43.4% of RUX no statistically significant	Tiredness Muscle aches, Night sweats Itching	Did not anticipate high proportion of patients achieving endpoint on HA arm	** Drop-out rate >20%

**Notes:** \*\*\*\*Represents a quality assessment of 80%, and \*\*\*\*\*Represents a quality assessment of 100%.

MPN-SAF TSS. It was also reflected in the sustained improvement (indicated by a negative mean change) in the total symptom scores observed in the RUX arm, with a mean (standard deviation) change of  $-9.0$  (13.52) at week 80, consistent with what was observed at week 28 ( $-10.46$  [14.25]). PSIS recorded rapid improvements in the severity of pruritus in patients treated with RUX at week 28 compared to patients receiving BAT, who had a worsening in pruritus symptom severity at most assessments. At week 80, 71% of patients treated with RUX showed improvement in pruritus.

In the RELIEF trial, the proportion of patients who achieved a  $\geq 50\%$  improvement in the MPN-SAF TSS was similar in both groups. In the RUX arm, there was a more significant proportion of patients who achieved higher than 50% improvement from baseline at Week 16 (RUX- 43.4% vs HC - 29.6%; however, the difference was not statistically significant (OR, 1.82; 95% CI, 0.82–4.04;  $P = 0.139$ )). A more substantial proportion of patients in the RUX arm achieved higher than 50% improvement from baseline in the individual TSS-C symptoms compared with the HA arm, and only the difference in itchiness was statistically significant (OR, 2.51; 95% CI, 1.10–5.71;  $P = 0.027$ ). The median percentage change from baseline symptoms was in favor of RUX at Week 16 and showed continued benefit at week 24 weeks and 48 weeks. The proportion of patients who achieved more than 50% improvement in the individual hyperviscosity or splenomegaly-related symptoms was similar between the treatment arms. The patients had no splenomegaly (0 in both the groups) and had well-controlled hematocrit (mean hematocrit in the RUX arm - 42.1% vs HA 43.7%).

### Work Productivity And Activity Impairment

In RESPONSE 2, utilizing the WPAI scale, patients reported improvement in work impairment, productivity, and days missed due to symptoms compared to the BAT where they experienced worsening of the same.

### Adverse Events Potentially Affecting QoL

The most common adverse events (AE) associated with RUX were a headache (9–16%), fatigue (7–20%), dizziness, diarrhea, constipation, pruritus, and weight increase. An increased incidence of Herpes zoster infection occurred in patients on the RUX arm (7 (6.4%) vs 0 in the RESPONSE trial, 1(1%) vs 0 RESPONSE- 2 trial, 1(0.5%) vs 0 in RELIEF trial). Overall, the infection rate was also slightly higher in the RUX arm. In the RESPONSE trial, the rate of infections of any grade was 41.8% in the RUX group and


36.9% in the standard therapy group. In RESPONSE-2, 19 (25.6%) patients had an infection of any grade (cystitis, influenza, nasopharyngitis, respiratory infection) compared to 15 (20%) patients (influenza, nasopharyngitis, respiratory infection) on BAT. Four patients in the RUX group and two patients in the standard therapy group had newly diagnosed non-melanoma skin cancer (basal cell or squamous-cell carcinoma) in the RESPONSE trial; all patients but one on standard treatment had a history of skin cancer. Two patients in the RELIEF trial developed squamous cell carcinoma while on RUX. Finally, patients on RUX reported mild elevations of total cholesterol and triglycerides.

Hematological AE was mostly graded 1–2 in all three studies. Hematologic laboratory abnormalities primarily consisted of mild anemia and thrombocytopenia with RUX. Anemia and thrombocytopenia of any category were less common in the RESPONSE-2 trial (14% and 3% respectively) compared to the RESPONSE (43.6% and 24.5% respectively) and RELIEF trials (37% and 9% respectively) trials. Thromboembolic events occurred in one patient in the RUX group versus six patients in the SOC group. In the RELIEF trial, thromboembolic events occurred in two patients on the RUX arm and in two patients on the HA arm. In the RESPONSE trial, three patients on RUX progressed to myelofibrosis, and one patient received a diagnosis of AML at day 56 after randomization. One patient in the SOC arm evolved to myelofibrosis on day 101. Also, two patients assigned to standard treatment received a diagnosis of myelofibrosis on days 308 and 378 after crossover, and one progressed to AML. In RELIEF, one patient in the RUX arm developed MF transformation (on day 211, which was 24 days after the final dose of RUX) and AML (on day 216, or 29 days after the last RUX treatment).

### Effect Of Ruxolitinib On QoL In PV

In the RESPONSE trial, according to the PGIC scale patients on the RUX arm (67%) reported that their condition was “much improved” or “very much improved” at Week 32 compared to the 13% on standard therapy.

On the EORTC QLQ-C30 scale, at Week 32 patients who received RUX experienced improvements from baseline compared to significant worsening in the SOC arm. Patients who received RUX showed improvement in social, physical, role, and cognitive functional scales from baseline at Week 32, whereas patients who received standard therapy experienced a worsening of these measurements. In the emotional functioning subscale, the RUX group had a higher proportion of patients who showed improvement. A more significant



**Figure 2** Histograms comparing ruxolitinib with standard therapy or hydroxyurea alone in RESPONSE at 32 weeks, RESPONSE at 28 weeks and RELIEF at 16 weeks. Graph 1 depicts proportion of patients who achieved a  $\geq 50\%$  reduction in the MPN-SAF TSS. The RELIEF study only reported the TSS- C (Cytokine symptoms). Graph 2 depicts the proportion of patients who reported "very much" or "much improved" on the PGIC scale.

**Abbreviations:** TSS, Total Symptom Score; PGIC, Patient Global Impression of Change.

proportion of patients in the RUX arm compared with the standard therapy arm (44% vs 9% respectively) achieved a minimally significant difference (MID;  $\geq$ a 10-point improvement from baseline) in global health status/QoL from baseline at each post-baseline study visit through Week 32.

In the RESPONSE-2 trial, according to the PGIC scale patients on the RUX arm, 83% reported that their condition was "much improved" or "very much improved" at Week 32 compared to 14% on standard therapy.

At Week 80 PGIC scores remained similar in patients originally randomized to RUX. On EQ-5D-5L, a higher proportion of patients in the RUX group reported having no problems in all five dimensions (mobility 52.7% vs 18.7%, self-care 74.3% vs 29.3%, usual activities 59.5% vs 17.3%, pain/discomfort 52.7% vs 12%, anxiety/depression 56.8% vs 18.7%). EQ-5D-5L scores did not change at 80 weeks of RUX.

In the RELIEF trial, 48.1% of patients in the RUX arm reported that their PV-related symptoms were "very much improved" or "much improved" at week 16 compared to 30.4% in HA arm. (OR, 2.13; 95% CI, 0.98–4.65). Patients in the HA arm were more likely to describe their symptoms as "minimally improved" or "no change" (HA, 55.4%; RUX, 33.3%) (Figure 2).

## Correlation Of PRO With JAK2 Allele Burden

The RESPONSE trial evaluated the JAK2 allele burden. At week 32, there was a  $-12.2\%$  mean change compared to a baseline (mean difference of  $-34.7\%$ ). The interpretation of this finding is limited, given that the allele burden in the study was used for exploratory analysis and not for a specific biomarker-related hypothesis.<sup>4</sup>

## Discussion

Three RCTs have evaluated the impact of RUX on QoL and PRO in adult patients with PV. Although symptom burden is a very well recognized feature of MPNs, the first publication of QoL and PRO in PV was not reported until 2007.<sup>25</sup> Since then, multiple efforts have been made to develop instruments to assess symptoms and functionality in patients with MPNs.<sup>24,26,27</sup>

The MPN-SAF is the instrument validated for QoL and PRO in myeloproliferative neoplasms. Originally consisting of a 20-item tool, the instrument subsequently incorporated 27-item tools and evaluates ten symptoms. (MPN-SAF TSS; MPN-10).<sup>28</sup> This instrument is part of the regular assessment of MPNs (NCCN guidelines V1.2019).

In PV, constitutively activated JAK2 recruits signal transducers and activators of transcription (STATs) to cytokine receptors and hematopoietic factors leading to chronic inflammation.<sup>29</sup> Symptoms such as fatigue, early satiety, night sweats, and itching belong to the cluster of “cytokine symptoms” on MPN-SAF.<sup>24</sup>

Ruxolitinib is a potent kinase inhibitor of the JAK1/2-STAT pathways leading to decreased expression of cytokines and cell growth factors necessary for hematopoiesis.<sup>18,30</sup> Other JAK2 inhibitors share similar mechanisms of action and are under evaluation in other myeloproliferative neoplasms.

In myelofibrosis, RUX is effective in reducing spleen size and significantly improving the total symptom score according to the modified Myelofibrosis Symptom Assessment Form (MFSAF).<sup>31</sup> More recently, fedratinib, a selective JAK2 inhibitor, received FDA approval for the management of primary or secondary myelofibrosis.<sup>32</sup> Although fedratinib improved symptom responses per TSS from baseline up to 24 weeks,<sup>33</sup> no ongoing studies are evaluating its efficacy in PV.

An open-label, randomized Phase 2 study evaluated momelotinib, a JAK1/JAK2 inhibitor in PV;<sup>34</sup> the study was terminated because of limited efficacy; momelotinib did not change symptom burden per MPN-SAF TSS.

RESPONSE and RESPONSE 2 demonstrated the efficacy of RUX in decreasing symptom burden in PV. (Table 1) The authors proposed that the results from the studies have real-life applications since the comparison group was the best available therapy. However, an important caveat is that the open-label design of RESPONSE and RESPONSE 2 could not determine statistical differences in PROs between arms. Additionally, the evaluation of PRO in PV patients treated with RUX, who do not have splenomegaly, will require further study. The effect of RUX on PRO needs to consider other variables such as age, gender, and leukemic transformation. The role of PROs as a prognostic marker in PV and their correlation with cytokine activity are also exciting options to consider for further analyses.

Conversely, RELIEF did not show statistical differences in PRO in the RUX arm compared to HA.<sup>23</sup> Despite the design limitations of the three studies, the current data strongly supports the efficacy of RUX in PRO compared to other conventional PV therapies (Figure 1).

The progressive incorporation of PROs in RCTs in oncology is ongoing.<sup>35</sup> Current clinical trials in PV include PRO data in the efficacy assessment of MPNs (Table 3).

**Table 3** Ongoing Clinical Trials In Polycythemia Vera Incorporating PRO In Efficacy Assessment

Title	Clinical Trial Identifier	PRO Instruments	Phase	Recruitment Status
TGR-1202 + Ruxolitinib in Subjects With Myelofibrosis, MDS/MPN, or Polycythemia Vera Resistant to Hydroxyurea	NCT02493530	MPN-SAF TSS	Phase I	Recruiting
Expanded Treatment Protocol (ETP) of Ruxolitinib in Patients With Polycythemia Vera Who Are Hydroxyurea Resistant or Intolerant and for Whom no Treatment Alternatives Are Available.	NCT02292446	MPN-SAF	Phase III	Completed
SAR302503 in Patients With Polycythemia Vera or Essential Thrombocythemia	NCT01420783	MPN-SAF	Phase II	Completed
Low Dose Interferon Alpha Versus Hydroxyurea in Treatment of Chronic Myeloid Neoplasms (DALIAH)	NCT01387763	EORTC QLQ C-30 and MPN-SAF	Phase III	Active, not recruiting
The Benefit/Risk Profile of AOP2014 in Low-risk Patients With PV (Low-PV)	NCT03003325	Functional Assessment of cancer Therapy-Anaemia (FACT-An) and MPN-SAF TSS	Phase II	Recruiting
Idasanutlin Monotherapy in Patients With Hydroxyurea-Resistant/Intolerant Polycythemia Vera	NCT03287245	MPN-SAF TSS EORTC QLQ-C30 PGIC	Phase II	Recruiting
KRT-232 Compared to Ruxolitinib in Patients With Phlebotomy-Dependent Polycythemia Vera	NCT03669965	MPN-SAF TSS EORTC-QLQ-C30	Phase II	Recruiting



The data regarding the benefit of RUX versus standard treatment in HA intolerance/resistant PV is limited. The effect of MPN treatments in PRO and QoL needs confirmation with large-scale and long-term studies. Interventions directed to improve PRO in MPNs would provide a better understanding of symptoms, optimize therapy selection, and increase patient survival.

## Conclusion

RUX in PV patients who are intolerant or resistant to HA is associated with significantly decreased symptom burden. HA is an effective cytoreductive agent, but its effect PRO is primarily limited to pruritus control. PRO and QoL assessments are an integral part of the evaluation and treatment of MPNs, and patients with PV on RUX will experience improvements in PRO in addition to improving hematological parameters.

## Disclosure

CAY reports involvement in advisory boards with JAZZ in 2018/2019 and with Pfizer in 2017. The authors report no other conflicts of interest in this work.

## References

- Vannucchi AM, Guglielmelli P. What are the current treatment approaches for patients with Polycythemia vera and essential thrombocythemia? *Hematology*. 2017;(1):480–488. doi:10.1182/asheducation-2017.1.480
- Tefferi A. Primary myelofibrosis: 2019 update on diagnosis, risk-stratification and management. *Am J Hematol*. 2018;93:1551–1560. doi:10.1002/ajh.25230
- Mesa R, Boccia RV, Grunwald MR, et al. Patient-reported outcomes data from REVEAL at the Time of Enrollment (Baseline): a prospective observational study of patients with Polycythemia vera in the United States. *Clin Lymphoma, Myeloma Leuk*. 2018;18:590–596. doi:10.1016/j.clml.2018.05.020
- Vannucchi AM, Kiladjian JJ, Griesshammer M, et al. Ruxolitinib versus standard therapy for the treatment of Polycythemia vera. *N Engl J Med*. 2015;372:426–435. doi:10.1038/mt.2009.265
- Mesa R, Miller CB, Thyne M, et al. Myeloproliferative neoplasms (MPNs) have a significant impact on patients' overall health and productivity: the MPN Landmark survey. *BMC Cancer*. 2016;16:167. doi:10.1186/s12885-016-2208-2
- Abelsson J, Andréasson B, Samuelsson J, et al. Patients with polycythemia vera have worst impairment of quality of life among patients with newly diagnosed myeloproliferative neoplasms. *Leuk Lymphoma*. 2013;54(10):2226–2230.
- Stein BL, Moliterno AR, Tiu RV. Polycythemia vera disease burden: contributing factors, impact on quality of life, and emerging treatment options. *Ann Hematol*. 2014;93:1965–1976. doi:10.1007/s00277-014-2205-y
- Nelson EC, Eftimovska E, Lind C, et al. Patient reported outcome measures in practice. *BMJ*. 2015;350:g7818–g7818. doi:10.1136/bmj.g7818
- Buckley SA, Kirtane K, Walter RB, et al. Patient-reported outcomes in acute myeloid leukemia: where are we now? *Blood Rev*. 2018;32:81–87. doi:10.1016/j.blre.2017.08.010
- Bryant AL, Drier SW, Lee S, Bennett AV. A systematic review of patient reported outcomes in phase II or III clinical trials of myelodysplastic syndromes and acute myeloid leukemia. *Leuk Res*. 2018;70:106–116. doi:10.1016/j.leukres.2018.06.006
- Mesa R, Verstovsek S, Kiladjian JJ, et al. Changes in quality of life and disease-related symptoms in patients with polycythemia vera receiving ruxolitinib or standard therapy. *Eur J Haematol*. 2016;97:192–200. doi:10.1111/ejh.12707
- Blum S, Martins F, Alberio L. Ruxolitinib in the treatment of polycythemia vera: patient selection and special considerations. *J Blood Med*. 2016;7:205–215. doi:10.2147/JBM.S102471
- Mesa RA, Miller CB, Mascarenhas JO, et al. Hydroxyurea treatment history and quality of life in patients with polycythemia vera: results from the MPN landmark survey in the United States. *Blood*. 2015;126(23):4077.
- Scherber R, Dueck A, Kiladjian J, et al. Conventional therapeutic options have limited impact on MPN symptoms: insights from a prospective analysis of the MPN-SAF TSS. *Haematologica*. 2012.
- Alvarez-Larrán A, Pereira A, Cervantes F, et al. Assessment and prognostic value of the European leukemiaNet criteria for clinico-hematologic response, resistance, and intolerance to hydroxyurea in polycythemia vera. *Blood*. 2012;119:1363–1369. doi:10.1182/blood-2011-10-387787
- Raessler LA. Jakafi (Ruxolitinib): first FDA-approved medication for the treatment of patients with Polycythemia vera. *Am Heal Drug Benefits*. 2015;8:75–79.
- Barosi G, Mesa R, Finazzi G, et al. Revised response criteria for polycythemia vera and essential thrombocythemia: an ELN and IWG-MRT consensus project. *Blood*. 2013;121(23):4778–4781. doi:10.1182/blood-2013-01-478891
- Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. 2012;366(9):799–807. doi:10.1056/NEJMoa1110557
- Harrison C, Kiladjian J, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. 2012;366(9):787–798.
- Moher D, Liberati A, Tetzlaff J, Altman D. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(6):e1000097. doi:10.1371/journal.pmed1
- Hong QN, Gonzalez-Reyes A, Pluye P. Improving the usefulness of a tool for appraising the quality of qualitative, quantitative and mixed methods studies, the Mixed Methods Appraisal Tool (MMAT). *J Eval Clin Pract*. 2018;24(3):459–467. doi:10.1111/jep.12884
- P F, G M, P F, et al. Ruxolitinib for the treatment of inadequately controlled polycythaemia vera without splenomegaly (RESPONSE-2): a randomised, open-label, phase 3b study. *Lancet Oncol*. 2017;18(1):p. 88–99. doi:10.1016/S1470-2045(16)30558-7
- Mesa R, Vannucchi AM, Yacoub A, et al. The efficacy and safety of continued hydroxycarbamide therapy versus switching to ruxolitinib in patients with polycythaemia vera: a randomized, double-blind, double-dummy, symptom study (RELIEF). *Br J Haematol*. 2017;176(1):76–85. doi:10.1111/bjh.14382
- Scherber R, Dueck AC, Johansson P, et al. The Myeloproliferative Neoplasm Symptom Assessment Form (MPN-SAF): international prospective validation and reliability trial in 402 patients. *Blood*. 2011;118(2):401–408. doi:10.1182/blood-2011-01-328955
- Mesa RA, Niblack J, Wadleigh M, et al. The burden of fatigue and quality of life in myeloproliferative disorders (MPDs): an international internet-based survey of 1179 MPD patients. *Cancer*. 2007;109(1):68–76. doi:10.1002/encr.22365

26. Mesa RA, Schwager S, Radia D, et al. The Myelofibrosis Symptom Assessment Form (MFSAF): an evidence-based brief inventory to measure quality of life and symptomatic response to treatment in myelofibrosis. *Leuk Res.* 2009;33(9):1199–1203. doi:10.1016/j.leukres.2009.01.035
27. Mesa RA, Gotlib J, Gupta V, et al. Effect of ruxolitinib therapy on myelofibrosis-related symptoms and other patient-reported outcomes in COMFORT-I: a randomized, double-blind, placebo-controlled trial. *J Clin Oncol.* 2013;31(10):1285–1292. doi:10.1200/JCO.2012.44.4489
28. Emanuel RM, Dueck AC, Geyer HL, et al. Myeloproliferative neoplasm (MPN) symptom assessment form total symptom score: prospective international assessment of an abbreviated symptom burden scoring system among patients with MPNs. *J Clin Oncol.* 2012;30(33):4098–4103. doi:10.1200/JCO.2012.42.3863
29. Pourcelot E, Trocme C, Mondet J, et al. Cytokine profiles in polycythemia vera and essential thrombocythemia patients: clinical implications. *Exp Hematol.* 2014;42(5):360–368. doi:10.1016/j.exphem.2014.01.006
30. Verstovsek S, Kantarjian H, Mesa RA, et al. Safety and Efficacy of INCB018424, a JAK1 and JAK2 Inhibitor, in Myelofibrosis. *N Engl J Med.* 2010;363(12):1117–1127. doi:10.1056/NEJMoa1002028
31. Mesa RA, Kantarjian H, Tefferi A, et al. Evaluating the serial use of the myelofibrosis symptom assessment form for measuring symptomatic improvement: performance in 87 myelofibrosis patients on a JAK1 and JAK2 inhibitor (INCB018424) clinical trial. *Cancer.* 2011;117(21):4869–4877. doi:10.1002/cncr.26129
32. Fedratinib becomes new option in myelofibrosis. *Cancer Discov.* 2019;9(10):1332. doi:10.1158/2159-8290.CD-NB2019-102
33. Pardanani A, Harrison C, Cortes JE, et al. Safety and efficacy of fedratinib in patients with primary or secondary myelofibrosis: A randomized clinical trial. *JAMA Oncol.* 2015;1(5):643–51. doi:10.1001/jamaoncol.2015.1590
34. Verstovsek S, Courby S, Griesshammer M, et al. A phase 2 study of momelotinib, a potent JAK1 and JAK2 inhibitor, in patients with polycythemia vera or essential thrombocythemia. *Leuk Res.* 2017. doi:10.1016/j.leukres.2017.05.002
35. Trotti A, Colevas AD, Setser A, Basch E. Patient-reported outcomes and the evolution of adverse event reporting in oncology. *J Clin Oncol.* 2007;25(32):5121–5127.

## Journal of Blood Medicine

Dovepress

### Publish your work in this journal

The Journal of Blood Medicine is an international, peer-reviewed, open access, online journal publishing laboratory, experimental and clinical aspects of all aspect pertaining to blood based medicine including but not limited to: Transfusion Medicine; Blood collection, Donor issues, Transmittable diseases, and Blood banking logistics; Immunohematology; Artificial and alternative blood based

therapeutics; Hematology; Biotechnology/nanotechnology of blood related medicine; Legal aspects of blood medicine; Historical perspectives. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/journal-of-blood-medicine-journal>