



ORIGINAL RESEARCH

Open Access

Evolution of female urinary continence after physical therapy and associated factors

Caroline Baldini Prudencio¹, Guilherme Thomaz de Aquino Nava², Marco Aurélio Cardoso³,
Rafaela Bresciani Marreto⁴, Érica Almeida Sousa⁴, Vitor E Valenti³ and Angélica Mércia Pascon Barbosa^{1*}

Abstract

Background: Urinary incontinence (UI) is defined as any involuntary loss of urine that can influence the quality of life, personal hygiene and social interaction. The types of UI that most affect women are stress urinary incontinence, urge incontinence and mixed urinary incontinence. There are several risk factors that result in specific treatments. We aimed to investigate the evolution of female urinary continence after physical therapy intervention and its associated factors.

Method: A retrospective cross-sectional study was conducted with 71 participants who were discharged from physiotherapy sector from August 2006 to April 2012 and met the inclusion criteria.

Results: Among the studied variables, the number of sessions and completion of home pelvic floor exercises showed a significant association. The urinary continence appeared in 43.7% of the cases, and factors, performance of home exercises, and number of sessions showed a significant association.

Conclusion: The number of sessions and completion of home pelvic floor exercises showed a significant relationship with each other.

Keywords: Urinary incontinence, Patient outcome assessment, Urogynecology, Women's health

Background

Among the main complaints of women, especially after menopause, we may include urinary incontinence (UI) [1]. The UI is defined as involuntary loss of urine according to the International Continence Society [2], leading to impairment in quality of life, personal hygiene and social interaction [1]. The main types of urinary incontinence are stress urinary incontinence (SUI) that is characterized by involuntary loss when the intra-abdominal pressure exceeds urethral pressure in the absence of detrusor muscle contraction [3], since urgency urinary incontinence (UUI) is the complaint of involuntary leakage accompanied by or immediately preceded by urgency and when there is a junction of both the mixed urinary incontinence (MUI) [4].

The UI is caused by multifactorial causes, from factors related to biomechanical efforts and others associated to comorbidities that predisposes to the UI. Knowledge of

these factors is important for avoidance. The change of life style, weight control and proper nutrition reduce the risk of developing diseases which are associated with loss [5,6].

The treatments for this condition is conservative in the literature and recent studies highlighted their effectiveness, whereas approximately 72% of women undergoing treatment become continent [7]. Therapeutic intervention will depend on the type of UI. For conservative treatment of SUI it is used stretching, strengthening muscle groups accessories, exercises pelvic floor contraction (PF) alone, with the help of cone and/or biofeedback, the latter is related to proprioceptive stimulus [8-10]. In UUI therapeutic interventions are stretching, proprioception [11], electrical stimulation of the posterior tibial nerve [12-14], guidance reeducation bladder, bowel and fluid intake [9,10].

Enhancement of life expectancy increased the concern about public health. In this context, it is necessary studies on the main bouts that affect subjects with UI, preventing them from living with impaired quality of life. Thus, this investigation was undertaken to evaluate the

* Correspondence: pasconbarbosa@uol.com.br

¹Post Graduate Program in Women's Health, USP, Sao Paulo, Brazil
Full list of author information is available at the end of the article

evolution of female urinary continence after physical therapy intervention and its risks factors associated.

Method

It is a cross-sectional study approved by the Ethics Committee in Research of the Faculty of Philosophy and Science (FFC) UNESP Marília campus (n° 1898/2010). All participants signed an informed consent letter that provided information on the procedures.

Women included in the study were treated in the physiotherapy sector UBS New Marília from August 2006 to April 2012. There were one to two weekly sessions lasting 50 minutes in which interventions made were suitable to their type of UI. After the patient report continence for 30 days and present contraction of pelvic floor grade II during the service we granted pre-discharge, in which the patient should remain continent, to perform PF exercises and follow the directions at home. 30 days after the pre-discharge the patient returned to the industry and should report being continent, presenting the maintenance of the degree of contraction of the pre-discharge physical examination and be reoriented regarding care reported in the pre-high, with the caveat that they should be performed continuously and daily, to then be granted discharge.

Data collection was conducted in approximately 412 medical records of patients who underwent evaluation and physical therapy in Basic Health Unit neighborhood in New Marília Marília-SP. Among all patients, 121 were selected for the study by meeting the inclusion criteria.

Inclusion criteria were being continent, having been discharged more than six months at the time of collection by phone. We excluded participants who refused to answer the questions, who did not attend the calls or the phone was not on the list of subscribers. At the end 71 participants were included in the study.

In the medical records we selected data to the personal data of the participant and their state of health at the time of evaluation. By phone we collected information about the current state of health-related UI.

The results concerning the characterization of the sample are presented as mean \pm standard deviation (minimum-maximum) {median} for quantitative variables, such as age, BMI, height, weight and number of sessions, and by percentage qualitative variables, race, marital status, group belonging (continent/incontinent), type of UI presented before treatment. To analyze the association of UI with qualitative variables of hypertension (HBP), smoking, diabetes, exercise performance of PF and presence of normal delivery was applied the chi-square test, and linear regression was used to associate the quantitative variables related to BMI, age and number of sessions associated with the UI. We considered $p < 0.05$ for statistical significance. The statistical program used was a Biostat 2009 Professional for Windows 5.8.4.

Results

The age of the 71 study participants was 57.8 ± 10.7 (22–84) {59} years. Regarding ethnic 74.65% (53) reported that were white, 16.9% (12) caboclo, 7.04% (5) black and 1.41% (1) yellow. Regarding marital status 2.82% (2) were single, 73.24% (52) married, 14.08% (10) divorced and 9.86% (7) widows. We observed that 11.27% (8) had UUI, 15.49% (11) SUI and 73.24% (52) presented MUI. The body mass index (BMI) found for the sample was 28.15 ± 5.30 (18–48) {28} kg/cm². The weight of the participants was 69.10 ± 14.59 (42–142) {67} kg, height was 1.57 ± 0.07 (1.44 to 1.75) {1.55} m. The total number of sessions of the participants was 19.97 ± 12.70 (4–79) {17}.

It was observed that 43.7% (31) of the participants remained continents, 56.3% (40) presented incontinence again.

The group of incontinent participants after physiotherapy was composed by 40 women who had an average age of 59.62 ± 8.43 (47–78) {58.5} years old, with a predominance of Caucasians and 70% (28), puma 17 5% (7), 10% black (4) 2.5% yellow (1). Most participants reported being married and 67.5% (27) 17.5% (7) divorced, 12.5% (5) widowed and 2.5% (1) single.

The group of continent participants after therapy were 31 participants aged 55.55 ± 12.67 (22–84) {59} years old, mostly Caucasians 80.6% (25), followed by caboclo 16.1% (5), 3.3% black (1), none of which declared being yellow. Regarding marital status of the participants 80.6% (25) stated that they were married, 9.7% (3) divorced, 6.4% (2) widows and 3.3% (1) single.

Anthropometric variables and the number of sessions are presented in Table 1.

The IU subdivision prior to physical therapy protocol separately in continent and incontinent groups are displayed in Table 2.

Among the patients who continued continents, 83.9% (26) followed the recommendations of the physiotherapist and the execution of PF exercises three times weekly household. It did not occur in the incontinent group, and only 37.5% (15) continued to follow the proposed exercises.

Table 1 Anthropometric data and number of sessions participants continent and incontinent

Variable	Continents	Incontinents
BMI	$27,504 \pm 6,707$ (18,1-48,5) {25,2}	$28,650 \pm 3,909$ (21-37) {29}
Weight	$69,254 \pm 18,998$ (42-142) {67,5}	$68,99 \pm 10,223$ (49-100) {67,5}
Height	$1,584 \pm 0,0767$ (1,44-1,75) {1,58}	$1,551 \pm 0,053$ (1,47-1,7) {1,5}
Number of sessions	$16,580 \pm 8,369$ (8-44) {15}	$22,6 \pm 14,819$ (4-79) {20}

Table 2 Urinary incontinence and incontinence continents in subjects prior to treatment

Variable	Continents n (%)	Incontinents n (%)
Kind of UI	UUI	7 (22,6%)
	SUI	6 (19,3%)
	MUI	18 (58,1%)
		1 (2,5%)
		5 (12,5%)
		34 (85%)

Table 3 shows BMI, age and number of sessions associated with the variable incontinence. The number of sessions was significantly associated with incontinence, however, BMI and age showed no significant association.

The association of hypertension (HBP), smoking, diabetes, execution of pelvic floor exercises and the presence of normal delivery with urinary incontinence are described in Table 4. There was a significant association between urinary incontinence and PF exercise household.

Discussion

In this study we aimed to analyze the evolution of urinary continence in women who received physical therapy and associated factors. Among the main findings of this study we reported that 43.7% of participants remained continents and there was a significant association between the number of sessions and the PF exercises performed at home, the other variables did not present significant difference.

The main type of incontinence presented by participants before the physical therapy protocol was the MUI (73.2% of cases), this finding contrast with the literature that showed that the most frequent type is SUI [15,16]. Conversely, previous studies indicated that the urodynamic examination [17], which is the gold standard in the diagnosis of UI is more sensitive to distinguish them, Sartori et al. demonstrated difference between the clinical and urodynamic study, and the number of women with MUI [17].

The data regarding the number of participants who remained continent was 43.7% in contrast to the study of Krüger et al. [3] which observed that the amount of continent patients reached 60%. This difference may be noticed by the difference between the number of study participants, in addition to cultural difference and educational level of the patients. Another investigation showed that continents participants represented 46.2%, similar our findings, this slight increase in the percentage of continents

Table 3 Association between UI and quantitative variables by linear regression

Variable	P	R	R ²
BMI	0,370	0,11	0,01
Number of sessions	0,031	0,25	0,06
Age	0,108	0,19	0,04

Table 4 Association between UI and qualitative variables by chi-square test

Variable	P
SAH	0,64
Smoker	0,30
Diabetes	0,22
PF exercise	<0,0001
Vaginal delivery previous	0,25

can be explained by the data regarding continence that were collected immediately after the end of the sessions of physiotherapy [18].

In relation to age we noted that the incontinent patients had a mean age of 59.6 years old, however, this variable was not associated with UI. In the study of Hannestad et al. [19] they observed that the higher the age the higher the severity of UI, which agrees with the results of Higa and Lopes that urinary incontinence is associated with increased age [20].

The same was observed with BMI, which did not present significant association with IU. It is in contrast to studies showing that the probability of UI can be reduced by 3% for every pound lost and that there is an association between incontinence and BMI in women aged like this study [6,21]. However, these data still show up with conflicting studies that showed no association between BMI and IU [20,22].

Regarding chronic diseases and their association with the UI, we reported that hypertension and diabetes presented no significant association. In this sense, the study of Higa and Lopes [20] displayed significant associations between hypertension and UI. On the other hand, the same was not observed with diabetes. Blanco et al. reported no significant association in both diseases and UI [23].

Cigarette smoking showed no significant association in our study. The data found in the study of Tampakoudis et al. is that smokers have a greater association with the UI when compared to non smokers [24]. Bump and McClisch demonstrated in incontinent groups smoking and a non smoker, in which groups incontinent smoking were associated with UI due to cough this group be more vigorous [25]. Higa and Lopes and found no association between smoking and UI data that is opposed to previously presented by other authors and confirming the data found in this study [20].

In participants who underwent previous vaginal delivery, there was no significant association with IU, contradicting the literature that demonstrated an association between this type of delivery and IU [20,26,27].

Based on our study, it is important to note that the participants who remained continents performed an average of 16.5 sessions and 83.9% of them continued to perform PF exercises in domicile. Patients who have already returned

to complain UI performed an average of 22.6 sessions, however, only 37.5% of them continued with home exercises.

The lower amount of sessions in patients continents can be explained by the fact that they followed the same guidelines and conducted PF exercises in domicile correctly, aiding rapid evolution and maintenance of the results. Berquó presented a review that corroborates our finding [28].

The opposite is also true when it comes to incontinent patients, they required more sessions to achieve the same goal, perhaps because this group did not properly follow the guidelines, the data that leads us to hypothesize that this is only 37.5% of patients continued to follow the guidelines.

There are some points in our study that are worth to be raised, the phone records was not updated or was not in the list of more subscribers and by some participants who had hearing problems to understand the questionnaire.

Conclusion

The female urinary continence was maintained at less than half the studied group even after six months or more. The number of sessions and the implementation of home exercises were associated with continence.

Competing interests

The authors declare no conflict of interests.

Authors' contributions

All authors participated in the design of the project. CBP, GTAN, RBM and EAS conducted the interviews and discussion. MAC and VEV performed the statistical analysis. ABMP revised the final version. All authors read and approved the final manuscript.

Acknowledgements

We thank the Municipal Health and staff of the Basic Health Unit of Marília by the space to collect the data and interviews. This study received financial support from FAPESP.

Author details

¹Post Graduate Program in Women's Health, USP, Sao Paulo, Brazil. ²Post Graduate Program in Functional Rehabilitation of Posture and Movement, USP, Sao Paulo, Brazil. ³Post Graduate Program in Physiotherapy, Faculty of Sciences and Technology, UNESP, Presidente Prudente, SP, Brazil.

⁴Departament of Physical Therapy and Occupational Therapy, Faculty of Philosophy Sciences, UNESP, Av. Hygino Muzzi Filho 747 Marília, SP, Brazil.

Received: 10 September 2013 Accepted: 17 April 2014

Published: 12 May 2014

References

1. Abreu N: **Qualidade de vida na perspectiva de idosas com incontinência urinária.** *Rev Bras Fisioter* 2007, **11**(6):429–436.
2. Haylen BT, de Ridder D, Freeman RM, Swift SE, Berghmans B, Lee J, Monga A, Petri E, Rizk DE, Sand PK, Schaer GN: **An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on the terminology for female pelvic floor dysfunction.** *Int Braz J Urol* 2010, **36**:116.
3. Krüger AP, Luz SCT, Virtuoso JF: **Home exercises for pelvic floor in continent women one year after physical therapy treatment for urinary incontinence: an observational study.** *Rev Bras Fisioter* 2011, **15**(5):351–356.
4. Abrams P, Andersson KE, Birder L, Brubaker L, Cardozo L, Chapple C, Cottenden A, Davila W, de Ridder D, Dmochowski R, Drake M, Dubeau C, Fry C, Hanno P, Smith JH, Herschorn S, Hosker G, Kelleher C, Koelbl H, Khoury S, Madoff R, Milsom I, Moore K, Newman D, Nitti V, Norton C, Nygaard I, Payne C, Smith A, Staskin D, Tekgul S, Thuroff J, Tubaro A, Vodusek D, Wein A, Wyndaele JJ, Members of Committees; Fourth International Consultation on Incontinence: **Fourth International Consultation on Incontinence Recommendations of the International Scientific Committee: Evaluation and treatment of urinary incontinence, pelvic organ prolapse, and fecal incontinence.** *Neurourol Urodyn* 2010, **29**(1):213–240.
5. Sievert K: **Can we prevent incontinence?** *Neurourol Urodyn* 2012, **31**(3):390–399.
6. Phelan S, Kanaya AM, Subak LL, Hogan PE, Espeland MA, Wing RR, Burgio KL, DiLillo V, Gorin AA, West DS, Brown JS, Look AHEAD Research Group: **Weight loss prevents urinary incontinence in women with type 2 diabetes: results from the Look AHEAD trial.** *J Urol* 2012, **187**(3):939–944.
7. Bernardes NO: **Métodos de Tratamento Utilizados na Incontinência Urinária de Esforço Genuína: um Estudo Comparativo entre Cinesioterapia e Eletroestimulação Endovaginal Pacientes e Métodos.** *Rev Bras Ginecol Obstet* 2000, **22**(1):49–54.
8. Silva AMN, Oliva LMP: **Exercícios de Kegel associados ao uso de cones vaginais no tratamento da incontinência urinária: estudo de caso.** *Sci Med* 2011, **10**:120–125.
9. Sociedade brasileira de urologia: **Projeto Diretrizes Incontinência Urinária de Esforço : Tratamento Não Cirúrgico e Não Farmacológico Projeto Diretrizes.** São Paulo: Guanabara Coogan; 2006:1–10.
10. Vasconcelos MMA: **Eficácia de um programa de reeducação miccional em crianças e adolescentes com distúrbio funcional do trato urinário inferior: cinesioterapia vs biofeedback do assoalho pélvico um estudo clínico aleatorizado.** PhD Thesis. Department of Health Sciences. Universidade Federal de Minas Gerais. 2005.
11. Arruda RM, Sousa GB, Castro RA, Sartori MG, Baracat EC, Girão MJB: **Hiperatividade do detrusor: comparação entre oxibutina, eletroestimulação funcional do assoalho pélvico e exercícios perineais. Estudo randomizado.** *Rev Bras Ginecol Obstet* 2007, **29**:452–458.
12. Franco MM, Souza FO, Vasconcelos ECLM, Freitas MMS, Ferreira CHJ: **Avaliação da qualidade de vida e da perda urinária de mulheres com bexiga hiperativa tratadas com eletroestimulação transvaginal ou do nervo tibial.** *Fisio & Pesq* 2011, **18**(2):145–150.
13. Alves FK, Florencio AC, Ricceto C, Carvalho RL: **Efeito da Electroestimulação do Nervo Tibial Posterior na Hiperatividade do Detrusor Neurogênico: Revisão de Literatura.** *Acta Urol* 2011, **1**:23–30.
14. Haddad JM: **Reabilitação do Assoalho Pélvico – Nas disfunções urinárias e anorretais.** 2ath edition. São Paulo: Segmento Farma; 2007.
15. Berquó MS: **Qualidade de vida de mulheres com incontinência urinária antes e após o tratamento fisioterapêutico.** Accessed in: [http://www.sbpnet.org.br/livro/63ra/conpeex/doutorado/trabalhos-doutorado/doutorado-marcela-souza.pdf] Acesso em: 21 nov. 2012.
16. Roussennq KR: **Atendimento fisioterapêutico na incontinência urinária: Resultados e Vivência prática.** Accessed in: [http://www.udesc.br/arquivos/id_submenu/797/artigo_cefid_11.pdf] Acesso em: 21 nov. 2012.
17. Sartori JP: **Distúrbios Urinários no Climatério: Avaliação Clínica e Urodinâmica.** *Rev Bras Ginecol Obstet* 1999, **21**:43–47.
18. Rett MT: **Qualidade de vida em mulheres após tratamento da incontinência urinária de esforço com fisioterapia.** *Rev Bras Ginecol Obstet* 2007, **3**:110–116.
19. Hannestad YS, Rortveit G, Sandvik H, Hunskaar S, Norwegian EPINCONT study, Epidemiology of Incontinence in the County of Nord-Trøndelag: **A community-based epidemiological survey of female urinary incontinence: The Norwegian EPINCONT Study.** *J Clin Epidemiol* 2000, **53**:1150–1157.
20. Higa R, Lopes MHBM: **Fatores associados com a incontinência urinária na mulher.** *Rev Bras Enferm* 2005, **10**:901–909.
21. Müller LM, Lose G, Jürgensen T: **Risk factors of lower urinary tract symptoms in women aged 40–60 years.** *Ugeskr Laeger* 2001, **163**:6598–6601.
22. Oliveira E: **Influência do índice de massa corporal na incontinência urinária feminina.** *Rev Bras Ginecol Obstet* 2010, **10**:111–118.
23. Blanco EN, Pérez JC, Alvarez VD, García MPL, Bejar PM, Lorente MP, Molina LS, Redondo BF: **Epidemiología e impacto de la incontinencia urinaria en mujeres de 40 a 65 años de edad en un área sanitaria de Madrid.** *Atención Primaria* 2003, **32**(7):410–414.

24. Tampakoudis P, Tantanassis T, Grimbizis G, Papaletsos M, Mantalenakis S: **Cigarette smoking and urinary incontinence in women—a new calculative method of estimating the exposure to smoke.** *Eur J Obstet Gynecol Reprod Biol* 1995, **63**(1):27–30.
25. Bump RC, McClish DM: **Cigarette smoking and pure genuine stress incontinence of urine: a comparison of risk factors and determinants between smokers and nonsmokers.** *Am J Obstet Gynecol* 1994, **170**:579–582.
26. Oliveira E: **Avaliação dos fatores relacionados à ocorrência da incontinência urinária feminina.** *Rev Assoc Med Bras* 2010, **56**:688–690.
27. Scarpa KP: **Sintomas do trato urinário inferior três anos após o parto: estudo prospectivo.** *Rev Bras Ginecol Obstet* 2008, **8**:733–741.
28. Berquó MS, Ribeiro MO, Amaral RG: **Fisioterapia no tratamento da incontinência urinária feminina.** *FEMINA* 2009, **5**:221–227.

doi:10.1186/1755-7682-7-24

Cite this article as: Prudencio *et al.*: Evolution of female urinary continence after physical therapy and associated factors. *International Archives of Medicine* 2014 **7**:24.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

