

# Complex characterization of oat (*Avena sativa* L.) lines obtained by wide crossing with maize (*Zea mays* L.)

Edyta Skrzypek<sup>1</sup>, Tomasz Warzecha<sup>2</sup>, Angelika Noga<sup>1</sup>, Marzena Warchoł<sup>1</sup>, Ilona Czyczyło-Mysza<sup>1</sup>, Kinga Dziurka<sup>1</sup>, Izabela Marcińska<sup>1</sup>, Kamila Kapłoniak<sup>1</sup>, Agnieszka Sutkowska<sup>2</sup>, Zygmunt Nita<sup>3</sup>, Krystyna Werwińska<sup>3</sup>, Dominika Idziak-Helmcke<sup>4</sup>, Magdalena Rojek<sup>4</sup> and Marta Hosiawa-Barańska<sup>4</sup>

<sup>1</sup> Department of Biotechnology, Polish Academy of Sciences, The Franciszek Górski Institute of Plant Physiology, Kraków, Poland

<sup>2</sup> Department of Plant Breeding and Seed Science, University of Agriculture, Kraków, Polska

<sup>3</sup> Plant Breeding Strzelce Ltd., PBAI Group, Strzelce, Polska

<sup>4</sup> Department of Plant Anatomy and Cytology, University of Silesia in Katowice, Katowice, Polska

## ABSTRACT

**Background.** The oat × maize addition (OMA) lines are used for mapping of the maize genome, the studies of centromere-specific histone (CENH3), gene expression, meiotic chromosome behavior and also for introducing maize C4 photosynthetic system to oat. The aim of our study was the identification and molecular-cytogenetic characterization of oat × maize hybrids.

**Methods.** Oat DH lines and oat × maize hybrids were obtained using the wide crossing of *Avena sativa* L. with *Zea mays* L. The plants identified as having a *Grande-1* retrotransposon fragment, which produced seeds, were used for genomic *in situ* hybridization (GISH).

**Results.** A total of 138 oat lines obtained by crossing of 2,314 oat plants from 80 genotypes with maize cv. Waza were tested for the presence of maize chromosomes. The presence of maize chromatin was indicated in 66 lines by amplification of the PCR product (500 bp) generated using primers specific for the maize retrotransposon *Grande-1*. Genomic *in situ* hybridization (GISH) detected whole maize chromosomes in eight lines (40%). All of the analyzed plants possessed full complement of oat chromosomes. The number of maize chromosomes differed between the OMA lines. Four OMA lines possessed two maize chromosomes similar in size, three OMA—one maize chromosome, and one OMA—four maize chromosomes. In most of the lines, the detected chromosomes were labeled uniformly. The presence of six 45S rDNA loci was detected in oat chromosomes, but none of the added maize chromosomes in any of the lines carried 45S rDNA locus. Twenty of the analyzed lines did not possess whole maize chromosomes, but the introgression of maize chromatin in the oat chromosomes. Five of 66 hybrids were shorter in height, grassy type without panicles. Twenty-seven OMA lines were fertile and produced seeds ranging in number from 1–102 (in total 613). Sixty-three fertile DH lines, out of 72 which did not have an addition of maize chromosomes or chromatin, produced seeds in the range of 1–343 (in total 3,758). Obtained DH and OMA lines were fertile and produced seeds.

**Discussion.** In wide hybridization of oat with maize, the complete or incomplete chromosomes elimination of maize occur. Hybrids of oat and maize had a complete

Submitted 24 March 2018

Accepted 6 June 2018

Published 25 June 2018

Corresponding author

Edyta Skrzypek, e.skrzypek@ifr-pan.edu.pl

Academic editor

Sheila McCormick

Additional Information and  
Declarations can be found on  
page 21

DOI 10.7717/peerj.5107

© Copyright  
2018 Skrzypek et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

set of oat chromosomes without maize chromosomes, and a complete set of oat chromosomes with one to four retained maize chromosomes.

**Subjects** Agricultural Science, Biotechnology, Genetics, Plant Science

**Keywords** Genomic *in situ* hybridization (GISH), Fertility, Oat × maize hybrids, Doubled haploids (DH)

## INTRODUCTION

The fusion of genomes from various genetic backgrounds through wide hybridization is essential for plant breeders seeking new crop cultivars, especially in the context of climate changes. On the other hand, unlikely hybrids with genomes from both the parental species are not always obtained. Uniparental chromosome elimination, the removal of the one parent chromosomes from the wide hybrids, might be a benefit to the breeders for fast genetic improvement of the crop cultivars ([Chaudhary et al., 2013](#)).

In crosses between Panicoideae and Pooideae, in hybrids e.g., wheat × sorghum, wheat × pearl millet, wheat × maize, oat × maize or oat × pearl millet, either Panicoideae chromosomes are eliminated shortly after fertilization at the beginning of embryogenesis, or incomplete chromosomes elimination occurred ([Laurie & Bennett, 1986](#); [Laurie & Bennett, 1988](#); [Laurie, 1989](#); [Rines & Dahleen, 1990](#)). In the case of maize crosses, the elimination of maize chromosomes was due to their limited mobility for the duration of mitotic metaphase and anaphase as a result of the failure of spindle fiber attachment to the centromeres ([Laurie & Bennett, 1989](#)).

Chromosome elimination of the alien genome after fertilization of the different species can take place in various distant hybrids ([Dunwell, 2010](#)). Interspecific hybridization was the first observed in the genus *Hordeum*, during crossing cultivated barley (*Hordeum vulgare* L.) as the female with wild species *Hordeum bulbosum* L. ([Kasha & Kao, 1970](#)). In an early stage of development of a hybrid embryo, chromosomes of the female are eliminated and developed haploid embryos contain only chromosomes of cultivated barley ([Liu et al., 2014](#)). Chromosome elimination has been described in crosses of *Triticum aestivum* and Triticeae species (*H. vulgare* and *H. bulbosum*) as well as more distantly related species (*Zea mays*, *Pennisetum glaucum*, *Sorghum bicolor*, *Coix lacryma-jobi*, *Imperata cylindrica*) ([Barclay, 1975](#); [Laurie & Bennett, 1986](#); [Laurie & Bennett, 1988](#); [Laurie, 1989](#); [Inagaki & Mujeeb-Kazi, 1995](#); [Mochida & Tsujimoto, 2001](#); [Gernand et al., 2005](#); [Komeda et al., 2007](#); [Chaudhary et al., 2013](#)).

A new situation was observed in the crosses between oat (*Avena sativa* L.) and maize (*Zea mays* L.) where oat × maize hybrid embryos developed into euhaploid plants with the whole oat chromosomes without maize chromosomes and into aneuhaploid plants with the complete set of oat chromosome and added a different numbers of maize chromosomes ([Kynast et al., 2012](#)). In the aneuhaploid plants numerous maize chromosomes were not removed during embryogenesis, and the preserved maize chromosomes ultimately were stable and act as oat chromosomes in mitosis. The consequence of it was obtaining oat × maize addition (OMA) lines.

The first sweet maize line used to obtain OMA was the Seneca 60. In oat was identified a fragment of Seneca 60 chromosome 3 containing a neocentromere (Topp *et al.*, 2009). Over the past few years, fertile disomic OMA of diverse oat cultivars have been reported, in which *Zea mays* was the recipient of chromosomes 1–7 and 9, the short arm of 10, as well as a monosomic OMA with chromosome 8 (Rines *et al.*, 2009). Amongst maize crosses with other cereals, only oat × maize hybrids formed embryos which are able to germinate and grow into partially fertile plants. In these OMA lines the partial fertility is a result of meiotic restitution, which improved the production of unreduced gametes. The observation of haploid oat microsporocytes (with or without maize chromosomes) displayed that in microsporogenesis karyokinesis is irregular, mainly in meiosis (Kynast *et al.*, 2012).

Ishii *et al.* (2013) reported that in crosses of oat × pearl millet (*Pennisetum glaucum* L.), chromosome and marker analyses showed that the seedlings were hybrids that had all oat and some pearl millet chromosomes. Twenty-one embryos in this study germinated and showed only shoot growth which turned to the scutellum and finally died under light regime.

The OMA lines are irreplaceable and valuable genetic materials, where the heterologous organism is required to investigate the role and structure of single maize chromosome, to ascribe and localize molecular markers and genes on maize chromosomes, or to study the dynamic of chromosome pairing. Dissecting the maize chromosomes by OMA lines can deliver plants with maize subchromosomal fragments. As OMA lines, subchromosomal materials are used in a series of genomics and chromosome studies (Kynast & Riera-Lizarazu, 2011).

The OMA lines were used in the cytogenetic mapping system for maize genome (Riera-Lizarazu, Rines & Phillips, 1996; Maquieira, 1997; Kynast *et al.*, 2001; Kynast *et al.*, 2002; Okagaki *et al.*, 2001; Koumbaris & Bass, 2003), gene expression (Jin *et al.*, 2004), meiotic chromosome arrangements (Bass *et al.*, 2000), investigation of conservation of gene expression of prolamin storage protein (Garcia *et al.*, 2015) as well as investigation of genes transcriptional regulation in new genomic environments (Dong *et al.*, 2018). It was shown that the majority of maize genes displayed maize-specific transcription in the oat genomic environment. Oat × maize hybrids were also used for identification (Topp *et al.*, 2009) and analyzing the size and location of maize chromosomes centromeres transferred into oat by immunoprecipitation for the histone CENH3 (Wang *et al.*, 2014). The mutation of CENH3 offers promising opportunities for application in a wide range of crop species for haploid production (Ravi & Chan, 2010; Ishii *et al.*, 2015; Karimi-Ashtiyani *et al.*, 2015; Maheshwari *et al.*, 2015; Britt & Kuppu, 2016; Kelliher *et al.*, 2016). A single-point amino acid exchange in the centromere-targeting domain of CENH3 leads to reduced centromere loading of CENH3 in barley, sugar beet, and *Arabidopsis thaliana*. Haploids were obtained when *cenh3*-null mutants were crossed with desired plants. The OMA lines were also used in the examination for expression of the C<sub>4</sub> photosynthetic system (Kynast *et al.*, 2001; Kowles *et al.*, 2008; Dong *et al.*, 2018).

The aim of the study was the identification and molecular-cytogenetic characterization of oat × maize hybrids. The paper also presents vigor of DH and OMA lines and their effectiveness in seeds production.

## MATERIAL AND METHODS

### Oat wide crossing with maize

The experiments were done on 80 oat genotypes (listed in [Table 1](#)) obtained from Plant Breeding Strzelce Ltd., PBAI Group, Strzelce, Łódź Voivodeship, Poland. Oat plants and maize cultivar Waza were sown and cultivated according to [Noga et al. \(2016\)](#). Oat haploid plants were obtained using the wide crossing method by pollination with maize cv. Waza, as described by [Marcińska et al. \(2013\)](#). Three weeks after floret pollination, enlarged ovaries were sterilized in 70% (v/v) ethanol (1 min), in a 2.5% (w/v) calcium hypochlorite (8 min), and washed three times with sterile water. Then, haploid embryos were isolated from caryopses and transferred to the 190-2 medium ([Zhuang & Xu, 1983](#)) with 9% (w/v) maltose, 0.6% (w/v) agar, 0.5 mg L<sup>-1</sup> kinetin (KIN) and 0.5 mg L<sup>-1</sup> α-naphthaleneacetic acid (NAA) and pH 6.0. Haploid embryos were germinated in an *in vitro* chamber with a 16 h photoperiod, light intensity of 100 μmol m<sup>2</sup> s<sup>-1</sup> and temperature 21 °C. The germination capacity of haploid embryos was evaluated under a stereomicroscope (SMZ 1500, Nikon, Tokyo, Japan) and photographs were taken using a digital CCD camera (DS-Ri1, Nikon, Tokyo, Japan).

The plants developed from haploid embryos were transferred to MS medium ([Murashige & Skoog, 1962](#)) with 0.6% (w/v) agar. Then, the well rooted plants were acclimated to natural conditions by transferring them subsequently to wet perlite and soil. For chromosome doubling, the roots of haploid plants were treated by colchicine solution and the ploidy level of the plants was evaluated using a MACSQuant flow cytometer (MACSQuant, Miltenyi Biotec GmbH, Bergisch Gladbach, Germany) as described by [Noga et al. \(2016\)](#).

Next, the plants were tested for the presence of the maize retrotransposon *Grande-1* to indicate maize chromatin in oat. Oat cv. Stoper and maize cv. Waza were used as a negative and positive control of the presence of the maize retrotransposon *Grande-1*. Leaves of studied lines, oat cv. Stoper, and maize cv. Waza (about 200 mg) were lyophilized in high vacuum at 40 μbar, coil temperature -52 °C (lyophiliser FreeZone 6L; Labconco, Kansas City, MO, USA). Then, the tissue was homogenized using ball mill (MM400; Retsch, Haan, Germany) for 5 min at frequency 25 Hz.

### DNA extraction and PCR analyses

DNA was extracted from 20 mg of lyophilized leaves, using the Genomic Mini AX Plant kit (A&A Biotechnology, Gdynia, Poland) according to the producer recommendations. After precipitation, the DNA pellet was washed twice in 70% (v/v) ethanol and dried in a concentrator (Concentrator Plus; Eppendorf, Hamburg, Germany) for 10 min. under 20 hPa. The precipitate was suspended in sterile distilled water. DNA concentration was measured at 260 nm with a spectrophotometer NanoDrop 2000c (Thermo Scientific, USA).

Two Grande 1F (5'-AAA GAC CTC ACG AAA GGC CCA AGG-3'), Grande1R (5'-AAA TGG TTC ATG CCG ATT GCA CG-3') primers (GenBank accession number [X97604](#);

**Table 1** The influence of oat (*Avena sativa* L.) genotype on the efficiency of haploid embryos, plants and seeds production using wide crossing with maize (*Zea mays* L.).

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/florets (%)	Germinated embryos/florets (%)	Acclimated plants/florets (%)	Lines/florets (%)	No. of seeds	No. of maize chromosome/chromatin added to oat genome
1	STH 4.403/1	27	858	11	0	0	0	1.28	0	0	0	0 <sup>d</sup>	0
2	STH 4.403/2	29	856	23	3	1	1	2.69	0.35	0.12	0.12	105 <sup>bc</sup>	0
3	STH 4.403/3	32	984	10	3	1	1	1.02	0.30	0.10	0.10	35 <sup>d</sup>	2
4	STH 4.403/4	23	763	12	3	2	2	1.57	0.39	0.26	0.26		
							STH 4.403/4 a					128 <sup>bc</sup>	0
							STH 4.403/4 b					9 <sup>d</sup>	0
5	STH 4.4576	26	439	21	9	2	1	4.78	2.05	0.46	0.23	10 <sup>d</sup>	chromatin
6	STH 4.4577	33	559	23	6	1	0	4.11	1.07	0.18	0	0 <sup>d</sup>	0
7	STH 4.4586	40	674	45	21	3	1	6.68	3.12	0.45	0.15	0 <sup>d</sup>	n/a*
8	STH 4.4595	40	523	21	13	5	0	4.03	2.49	0.96	0	0 <sup>d</sup>	0
9	STH 4.4606	39	832	60	34	8	2	7.22	4.09	0.96	0.24		
							STH 4.4606 a					36 <sup>cd</sup>	chromatin
							STH 4.4606 b					0 <sup>d</sup>	n/a
10	STH 4.4690	45	1,094	106	62	20	16	9.69	5.67	1.83	1.46		
							STH 4.4690 a					19 <sup>d</sup>	0
							STH 4.4690 b					119 <sup>bc</sup>	0
							STH 4.4690 c					23 <sup>d</sup>	0
							STH 4.4690 d					10 <sup>d</sup>	chromatin
							STH 4.4690 e					12 <sup>d</sup>	0
							STH 4.4690 f					2 <sup>d</sup>	chromatin
							STH 4.4690 g					200 <sup>bc</sup>	0
							STH 4.4690 h					2 <sup>d</sup>	0
							STH 4.4690 i					6 <sup>d</sup>	0
							STH 4.4690 j					1 <sup>d</sup>	0
							STH 4.4690 k					1 <sup>d</sup>	0
							STH 4.4690 l					0 <sup>d</sup>	0
							STH 4.4690 m					0 <sup>d</sup>	n/a
							STH 4.4690 n					0 <sup>d</sup>	n/a
							STH 4.4690 o					0 <sup>d</sup>	n/a
							STH 4.4690 p					1 <sup>d</sup>	4
11	STH 4.4729	36	927	33	4	2	1	3.56	0.43	0.22	0.11	0 <sup>d</sup>	n/a

(continued on next page)

Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/ florets (%)	Germinated embryos/ florets (%)	Acclimated plants/ florets (%)	Lines/ florets (%)	No. of seeds	No. of maize chromosome/ chromatin added to oat genome	
12	STH 4.4731	35	776	33	13	5	4	4.25	1.68	0.64	0.52	91 <sup>bc</sup>	0	
													14 <sup>d</sup>	0
													14 <sup>d</sup>	0
													6 <sup>d</sup>	0
13	STH 4.4740	33	717	38	10	3	2	5.30	1.39	0.42	0.28	20 <sup>d</sup>	0	
													57 <sup>bcd</sup>	0
14	STH 4.4742	22	647	34	17	7	4	5.26	2.63	1.08	0.62	77 <sup>bc</sup>	chromatin	
													222 <sup>b</sup>	0
													5 <sup>d</sup>	0
													1 <sup>d</sup>	chromatin
15	STH 4.8402	36	626	31	8	1	0	4.95	1.28	0.16	0	0 <sup>d</sup>	0	
16	STH 4.8411	28	623	27	9	1	1	4.33	1.44	0.16	0.16	165 <sup>bc</sup>		
17	STH 4.8417	26	648	28	10	4	4	4.32	1.54	0.62	0.62	0 <sup>d</sup>	n/a	
													0 <sup>d</sup>	n/a
													0 <sup>d</sup>	n/a
													10 <sup>d</sup>	0
18	STH 4.8432	34	681	26	4	1	1	3.82	0.59	0.15	0.15	0 <sup>d</sup>	0	
19	STH 4.8435	35	653	29	5	1	0	4.44	0.77	0.15	0	0 <sup>d</sup>	0	
20	STH 4.8437	30	602	35	10	5	2	5.81	1.66	0.83	0.33	4 <sup>d</sup>	0	
													0 <sup>d</sup>	n/a
21	STH 4.8442	33	826	45	9	3	1	5.45	1.09	0.36	0.12	0 <sup>d</sup>	n/a	
22	STH 4.8448	29	518	33	11	2	1	6.37	2.12	0.39	0.19	0 <sup>d</sup>	n/a	
23	STH 4.8456/1	51	1,019	29	3	2	2	2.85	0.29	0.20	0.20	7 <sup>d</sup>	0	
													6 <sup>d</sup>	0
24	STH 4.8456/2	45	1,043	27	10	7	7	2.59	0.96	0.67	0.67	80 <sup>bc</sup>	0	
													39 <sup>cd</sup>	0
													17 <sup>d</sup>	0

(continued on next page)

Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/ florets (%)	Germinated embryos/ florets (%)	Acclimated plants/ florets (%)	Lines/ florets (%)	No. of seeds	No. of maize chromosome/ chromatin added to oat genome
							STH 4.8456/2 d					11 <sup>d</sup>	0
							STH 4.8456/2 e					53 <sup>cd</sup>	chromatin
							STH 4.8456/2 f					1 <sup>d</sup>	chromatin
							STH 4.8456/2 g					0 <sup>d</sup>	0
25	STH 4.8457/1	42	887	30	1	1	1	3.38	0.11	0.11	0.11	53 <sup>cd</sup>	0
26	STH 4.8457/2	47	1,141	47	8	5	5	4.12	0.70	0.44	0.44		
							STH 4.8457/2 a					105 <sup>bc</sup>	0
							STH 4.8457/2 b					0 <sup>d</sup>	0
							STH 4.8457/2 c					0 <sup>d</sup>	n/a
							STH 4.8457/2 d					0 <sup>d</sup>	n/a
							STH 4.8457/2 e					0 <sup>d</sup>	n/a
27	STH 4.8459	20	286	13	2	1	0	4.55	0.70	0.35	0	0 <sup>d</sup>	0
28	STH 5.451/1	25	627	10	2	0	0	1.59	0.32	0	0	0 <sup>d</sup>	0
29	STH 5.451/2	23	614	17	5	3	2	2.77	0.81	0.49	0.33		
							STH 5.451/2					0 <sup>d</sup>	n/a
							STH 5.451/2					0 <sup>d</sup>	n/a
30	STH 5.451/3	21	509	6	4	1	1	1.18	0.79	0.20	0.20	0 <sup>d</sup>	0
31	STH 5.451/4	21	518	9	1	0	0	1.74	0.19	0	0	0 <sup>d</sup>	0
32	STH 5.8046	2	77	2	0	0	0	2.60	0	0	0	0 <sup>d</sup>	0
33	STH 5.8421	35	998	32	8	7	7	3.21	0.80	0.70	0.70		
							STH 5.8421 a					183 <sup>bc</sup>	0
							STH 5.8421 b					24 <sup>d</sup>	0
							STH 5.8421 c					20 <sup>d</sup>	0
							STH 5.8421 d					102 <sup>bc</sup>	chromatin
							STH 5.8421 e					0 <sup>d</sup>	n/a
							STH 5.8421 f					0 <sup>d</sup>	n/a
							STH 5.8421 g					0 <sup>d</sup>	n/a
34	STH 5.8422/1	15	620	7	0	0	0	1.13	0	0	0	0 <sup>d</sup>	0
35	STH 5.8422/2	15	487	5	2	0	0	1.03	0	0	0	0 <sup>d</sup>	0
36	STH 5.8423	39	1,014	47	8	3	2	4.64	0.79	0.30	0.20		
							STH 5.8423 a					1 <sup>d</sup>	0
							STH 5.8423 b					0 <sup>d</sup>	n/a
37	STH 5.8424/1	22	455	13	5	2	1	2.86	1.10	0.44	0.22	21 <sup>d</sup>	0
38	STH 5.8424/2	30	874	44	6	2	1	5.03	0.69	0.23	0.11	0 <sup>d</sup>	n/a

(continued on next page)

Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/florets (%)	Germinated embryos/florets (%)	Acclimated plants/florets (%)	Lines/florets (%)	No. of seeds	No. of maize chromosome/chromatin added to oat genome											
39	STH 5.8425	40	1,284	84	21	9	5	6.54	1.64	0.70	0.39	3 <sup>d</sup>	chromatin											
							STH 5.8425 a						1 <sup>d</sup>	chromatin										
							STH 5.8425 b						0 <sup>d</sup>	n/a										
							STH 5.8425 c						0 <sup>d</sup>	n/a										
							STH 5.8425 d						0 <sup>d</sup>	n/a										
40	STH 5.8426	26	738	33	5	2	1	4.47	0.68	0.27	0.14	28 <sup>d</sup>	2											
							STH 5.8426 e						0 <sup>d</sup>	n/a										
							STH 5.8427 a						211 <sup>bc</sup>	0										
							STH 5.8427 b						179 <sup>bc</sup>	0										
							STH 5.8427 c						158 <sup>bc</sup>	0										
41	STH 5.8427	31	1,145	46	11	6	5	4.02	0.96	0.52	0.44	65 <sup>bcd</sup>	chromatin											
							STH 5.8427 d						1 <sup>d</sup>	chromatin										
							STH 5.8427 e						0 <sup>d</sup>	0										
							STH 5.8428/1						3	61	0	0	0	0	0	0	0	0	0 <sup>d</sup>	0
							STH 5.8428/2						35	690	11	2	1	1	1.59	0	0.14	0.14	4 <sup>d</sup>	chromatin
44	STH 5.8429	34	1,253	64	22	7	7	5.11	1.76	0.56	0.56	343 <sup>a</sup>	0											
							STH 5.8429 a						8 <sup>d</sup>	0										
							STH 5.8429 b						4 <sup>d</sup>	chromatin										
							STH 5.8429 c						0 <sup>d</sup>	n/a										
							STH 5.8429 d						0 <sup>d</sup>	n/a										
							STH 5.8429 e						0 <sup>d</sup>	n/a										
							STH 5.8429 f						0 <sup>d</sup>	n/a										
							STH 5.8429 g						0 <sup>d</sup>	n/a										
45	STH 5.8430/1	18	351	2	0	0	0	0.57	0	0	0	0 <sup>d</sup>	0											
46	STH 5.8430/2	17	307	17	3	1	1	5.54	0.98	0.33	0.33	130 <sup>bc</sup>	0											
47	STH 5.8432	32	698	29	4	1	1	4.15	0.57	0.14	0.14	58 <sup>cd</sup>	0											
48	STH 5.8436	39	799	24	3	2	2	3.00	0.38	0.25	0.25	58 <sup>cd</sup>	0											
							STH 5.8436 a						8 <sup>d</sup>	2										
							STH 5.8436 b						0 <sup>d</sup>	n/a										
49	STH 5.8440	35	817	43	7	2	2	5.26	0.86	0.24	0.24	96 <sup>bcd</sup>	0											
							STH 5.8440 a						51 <sup>cd</sup>	chromatin										
							STH 5.8440 b						0 <sup>d</sup>	n/a										
50	STH 5.8449	38	844	21	4	2	2	2.49	0.47	0.24	0.24													

(continued on next page)



Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/florets (%)	Germinated embryos/florets (%)	Acclimated plants/florets (%)	Lines/florets (%)	No. of seeds	No. of maize chromosome/chromatin added to oat genome
							STH 5.8449 a					70 <sup>bcd</sup>	0
							STH 5.8449 b					55 <sup>cd</sup>	chromatin
51	STH 5.8450	30	628	27	0	0	0	4.30	0	0	0	0 <sup>d</sup>	0
52	STH 5.8458	34	727	27	4	2	2	3.71	0.55	0.28	0.28		
							STH 5.8458 a					57 <sup>cd</sup>	0
							STH 5.8458 b					35 <sup>d</sup>	2
53	STH 5.8460	25	511	16	1	0	0	3.13	0	0	0	0 <sup>d</sup>	0
54	STH 5.8504	22	840	25	9	4	2	2.98	1.07	0.48	0.24		
							STH 5.8504 a					2 <sup>d</sup>	0
							STH 5.8504 b					5 <sup>d</sup>	1
55	STH 5.8504/1	25	808	34	15	7	3	4.21	1.86	0.87	0.37		
							STH 5.8504/1 a					2 <sup>d</sup>	0
							STH 5.8504/1 b					0 <sup>d</sup>	n/a
							STH 5.8504/1 c					0 <sup>d</sup>	n/a
56	STH 5.8505	27	946	33	10	4	3	3.49	1.06	0.42	0.32		
							STH 5.8505 a					19 <sup>d</sup>	0
							STH 5.8505 b					0 <sup>d</sup>	n/a
							STH 5.8505 c					0 <sup>d</sup>	n/a
57	STH 5.8505/1	27	707	16	5	2	1	2.26	0.71	0.28	0.14	3 <sup>d</sup>	0
58	STH 5.8506	22	644	30	13	4	3	4.66	2.02	0.62	0.47		
							STH 5.8506 a					84 <sup>bcd</sup>	0
							STH 5.8506 b					12 <sup>d</sup>	1
							STH 5.8506 c					0 <sup>d</sup>	n/a
59	STH 5.8506/1	24	644	19	6	2	1	2.95	0.93	0.31	0.16	99 <sup>bcd</sup>	0
60	STH 5.8507	25	663	25	14	6	4	3.77	2.11	0.90	0.60		
							STH 5.8507 a					11 <sup>d</sup>	1
							STH 5.8507 b					0 <sup>d</sup>	n/a
							STH 5.8507 c					0 <sup>d</sup>	n/a
							STH 5.8507 d					0 <sup>d</sup>	0
61	STH 5.8508	23	593	33	9	2	1	5.56	1.52	0.34	0.17	63 <sup>bcd</sup>	0
62	STH 5.8509	23	644	13	4	2	1	2.02	0.62	0.31	0.16	0 <sup>d</sup>	0
63	STH 5.8512	28	1092	28	8	2	1	2.56	0.73	0.18	0.09	41 <sup>cd</sup>	0
64	STH 5.8513	27	727	13	3	1	0	1.79	0.41	0.14	0	0 <sup>d</sup>	0
65	STH 5.8514	30	726	17	4	0	0	2.34	0.55	0	0	0 <sup>d</sup>	0
66	STH 5.8518	29	728	24	3	0	0	3.30	0.41	0	0	0 <sup>d</sup>	0

(continued on next page)

Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/ florets (%)	Germinated embryos/ florets (%)	Acclimated plants/ florets (%)	Lines/ florets (%)	No. of seeds	No. of maize chromosome/ chromatin added to oat genome
67	STH 5.8518/1	23	582	30	10	6	4	5.15	1.72	1.03	0.69		
							STH 5.8518/1 a					61 <sup>cd</sup>	0
							STH 5.8518/1 b					51 <sup>cd</sup>	0
							STH 5.8518/1 c					14 <sup>d</sup>	0
							STH 5.8518/1 d					2 <sup>d</sup>	0
68	STH 5.8522	25	672	12	3	2	0	1.79	0.45	0.30	0	0 <sup>d</sup>	0
69	STH 5.8522/1	32	733	8	2	1	1	1.09	0.27	0.14	0.14	1 <sup>d</sup>	chromatin
70	STH 5.8523	31	742	9	1	1	0	1.21	0.13	0.13	0	0 <sup>d</sup>	0
71	STH 5.8525	23	692	39	12	5	0	5.64	1.73	0.72	0	0 <sup>d</sup>	0
72	STH 5.8526	33	682	16	5	0	0	2.35	0.73	0	0	0 <sup>d</sup>	0
73	STH 5.8528	23	575	17	2	0	0	2.96	0.35	0	0	0 <sup>d</sup>	0
74	STH 5.8529	29	818	15	0	0	0	1.83	0.00	0	0	0 <sup>d</sup>	0
75	STH 5.8530	20	532	19	9	2	2	3.57	1.69	0.38	0.38		
							STH 5.8530 a					0 <sup>d</sup>	0
							STH 5.8530 b					0 <sup>d</sup>	n/a
76	STH 5.8530/1	22	576	20	7	3	2	3.47	1.22	0.52	0.35		
							STH 5.8530/1 a					74 <sup>bcd</sup>	0
							STH 5.8530/1 b					56 <sup>cd</sup>	0
77	STH 5.8535	35	784	39	9	2	2	4.97	1.15	0.26	0.26		
							STH 5.8535 a					19 <sup>d</sup>	0
							STH 5.8535 b					1 <sup>d</sup>	chromatin
78	STH 5.8536	31	835	35	10	2	1	4.19	1.20	0.24	0.12	0 <sup>d</sup>	n/a
79	STH 5.8536/1	25	774	32	7	2	2	4.13	0.90	0.26	0.26		
							STH 5.8536/1 a					1 <sup>d</sup>	chromatin
							STH 5.8536/1 b					0 <sup>d</sup>	0

(continued on next page)

Table 1 (continued)

No	Genotype	No. of plants	No. of emasculated florets	No. of isolated embryos	No. of germinated embryos	No. of acclimated plants	No. of lines	Embryos/ florets (%)	Germinated embryos/ florets (%)	Acclimated plants/ florets (%)	Lines/ florets (%)	No. of seeds	No. of maize chromosome/ chromatin added to oat genome
80	STH 5.8540	29	908	22	5	1	1	2.42	0.55	0.11	0.11	0 <sup>d</sup>	0
	Total	2,314	57,515	2,129	601	210	138					4371	
	Average							3.58	0.99	0.34	0.21		
	Min.							0	0	0	0		
	Max.							9.69	2.63	1.83	1.46		

**Notes.**

Significant differences between oat lines according to Duncan's test,  $p \leq 0.05$ , are marked with different letters.

*Ananiev, Phillips & Rines, 1998*) were used for the PCR reaction, which in successive cycles enabled the amplification of the 500 bp retrotransposon region of *Grande-1* and detection of the presence of maize chromatin in oat plants. For the PCR, the thermostable Taq polymerase (Fermentas, USA) was used.

After DNA isolation and dilution (50  $\mu\text{g}/\mu\text{l}$  DNA), 5  $\mu\text{l}$  of DNA was taken and introduced into reaction mixture using 0.5  $\mu\text{l}$  of each two primers. The reaction mixture was placed in a thermocycler (2720 Thermal Cycler; Applied Biosystems, Foster City, CA, USA) and used the following thermal program: 1. Initial denaturation  $-94\text{ }^{\circ}\text{C}$ , 5 min.; 2. Cyclic denaturation:  $94\text{ }^{\circ}\text{C}$ –30 s; 3. Starter connection— $58\text{ }^{\circ}\text{C}$ , 30 s; 4. Polymerization— $72\text{ }^{\circ}\text{C}$ —30 s; 5. Final Polymerization— $72\text{ }^{\circ}\text{C}$ , 5 min. Stages 2-4 were repeated cyclically 25 times. In the last stage the samples were cooled to  $4\text{ }^{\circ}\text{C}$  until removed from the amplification apparatus (*Ananiev, Phillips & Rines, 1998*).

The obtained products were mixed with 2.5  $\mu\text{L}$  DNA Gel Loading Dye buffer (Thermo Fisher Scientific, Waltham, MA, USA) and then separated in 1.5% agarose gel with ethidium bromide (Sigma-Aldrich, St. Louis, MO, USA) in TBE buffer, under 90 V for 90 min. DNA markers of 100 bp to 1000 bp and concentration of 0.5  $\mu\text{g}/\mu\text{l}$  (GeneRuler 100bp; Fermentas, Waltham, MA, USA) were used to estimate the length of PCR products. The image of electrophoretic separation was archived using the Imagemaster VDS gel reader (Amersham, Pharmacia Biotech, Piscataway, NJ, USA) and the Liscap Capture Application ver. 1.0. Electrophoretic gel analysis was performed using GelScan ver. 1.45, (Kucharczyk Electrophoretic Techniques, Warsaw, Poland). The plants identified as having a *Grande-1* retrotransposon fragment, which were also fertile and produced seeds, were used for genomic *in situ* hybridization (GISH).

### Cytogenetic analysis

The seeds of oat, maize, and oat-maize chromosome addition and DH lines were germinated for 2–3 days in Petri dishes on a wet filter paper. Seedlings with the root tips of 1.5–2 cm long were immersed in ice-cold water and incubated for 24 h at  $4\text{ }^{\circ}\text{C}$ . The procedure allowed to stop the cell division at metaphase through the spindle degradation, and to increase chromosome condensation. After 24 h, the root tips were fixed in a mixture of pure methanol and glacial acetic acid at a 3:1 ratio. The material was kept at  $4\text{ }^{\circ}\text{C}$  for 24 h, and stored in  $-20\text{ }^{\circ}\text{C}$  until used.

Excised root tips (about 5 mm in length) were washed in citrate buffer (pH 4.8) for 15–20 min and then digested in a solution of 20% pectinase (Sigma, St Louis, MO, USA), 1% cellulose (Calbiochem, San Diego, CA, USA) and 1% cellulase 'Onozuka R-10' (Serva, Heidelberg, Germany) in a citrate buffer at  $37\text{ }^{\circ}\text{C}$  for 2 h. The meristems were dissected from the root tips and squashed in a drop of 45% acetic acid. Good quality preparations were frozen on dry ice. After freezing, the coverslips were removed and the slides were air-dried and stored at  $4\text{ }^{\circ}\text{C}$  until required.

In order to discriminate maize chromatin introgressions, the total genomic DNA of the maize cv. Waza was used as probe for GISH. Maize DNA was isolated from plants grown under greenhouse conditions ( $21\text{ }^{\circ}\text{C}$ , photoperiod 8 h/16 h). Young leaves were frozen with liquid nitrogen and homogenized. DNA isolation was performed using the C-TAB method

(Doyle & Doyle, 1987). The concentration and purity of the isolated DNA was measured by NanoDrop 2000c (Thermo Scientific, Waltham, MA, USA) and the length of the DNA sequence fragments was checked on agarose gel.

Maize genomic DNA and 25S rDNA sequences that were used as additional chromosome markers were labeled by nick-translation using the digoxigenin-11-dUTP and rhodamine-5-dUTP (Roche), respectively. The GISH procedure was performed according to the protocol of Hasterok *et al.* (2006). In order to verify the specificity of the gDNA probe and appropriate hybridization conditions, maize and oat root tips preparations were also used as positive and negative control in GISH, respectively. Hybridization mixture was prepared by adding appropriate volumes of 50% formamide, 10% dextran sulfate,  $20 \times$  SSC, water, SDS, and labeled DNA probes ( $\sim 500$  ng/slide). Due to the high phylogenetic distance between oat and maize and related to that low degree of DNA sequence homology, the use of total genomic oat DNA to block non-specific hybridization was not necessary. Posthybridization stringent washes were carried out in  $10\% \times$  SSC at  $42^\circ\text{C}$ , which is equivalent to 79% stringency. Immunodetection of digoxigenin-labeled probes was performed according to the standard protocols using FITC-conjugated anti-digoxigenin antibodies (Roche). The rhodamine probes were visualized directly. The preparations were mounted in VectaShield antifade (Vector Laboratories, Inc. Burlingame, CA, USA) containing  $2.5 \mu\text{m ml}^{-1}$  DAPI (Serva, Heidelberg, Germany). Hybridization signals were visualized and captured using fluorescence microscope Axio Imager Z2 (ZEISS, Germany) equipped with monochromatic camera AxioCam MRm (ZEISS, Oberkochen, Germany) and archived by ZEN blue program (ZEISS, Oberkochen, Germany). All images were processed uniformly and superimposed using Photoshop CS3 (Adobe).

## STATISTICS

Statistical analysis of embryo formation, DH and OMA lines as well as seeds production was performed using analysis of variance at  $p \leq 0.05$  using the STATISTICA 12.0 software package (Stat-Soft, Inc., Tulsa, OK). For comparison of means for seeds number, Duncan's multiple test at the significance level of  $p \leq 0.05$  was used. Correlation coefficient for seeds number dependently on maize chromosome/chromatin addition to oat genome was also calculated.

## RESULTS

### Oat DH lines and oat $\times$ maize hybrids formation

Analysis of variance indicated no significant differences between oat genotypes in the production of embryos per emasculated florets and lines per pollinated florets (Table 2). The efficiency of oat lines production by wide crossing with maize is shown in Table 1. Total of 57,515 florets from 80 oat genotypes were pollinated with maize that resulted in development of 2,129 embryos (Fig. 1A). Average embryo formation frequency per pollinated florets was 3.58%. Six hundred and one embryos germinated within three weeks of *in vitro* culturing (28.23% of all embryos) (Fig. 1B). Number of embryos produced by various genotypes varied from 0 to 106 (9.69% embryos per pollinated florets). Only one

**Table 2** Analysis of variance of haploid embryos formation and lines production depending on oat genotype as well as seeds production depending on oat hybrids.

Trait	SS	df	MS	MS for residual	F	p
Embryos formation	217.13	79	2.895	34.464	0.084	0.751 ns
Lines production	6.27	79	0.084	0.024	3.457	0.116 ns
DH/OMA seeds production	49,347.9	1	49,347.9	2,607.8	18.923	0.000*

**Notes.**

SS, sum of squares; df, degrees of freedom; MS, mean squares; ns, not significant  $p \leq 0.05$ .

\*Significant at  $p \leq 0.05$ .


genotype out of eighty examined did not formed haploid embryos (STH 5.8428). Most embryos per pollinated floret were obtained from the genotypes: STH 4.4690 (9.69%), STH 4.4606 (7.22%), STH 4.4586 (6.68%), STH 5.8425 (6.54%) and STH 4.8448 (6.37%). Even if the genotype did not have the significant effect on the haploid plants production, 15 of 80 genotypes did not develop haploid plants. Total of 210 plants survived the acclimatization to the natural condition (Figs. 1C–1E). The procedure of chromosome doubling using colchicine reduced the number of obtained lines to 138. The highest number of lines (16) was obtained from the STH 4.4690 genotype. Plants from seven genotypes completely failed this process. The ploidy of the plants was measured by flow cytometry before and after application of colchicine (Fig. 2). Most plants doubled chromosomes number, except plants STH 4.4690 n, STH 4.4690 o, STH 5.8505 b, STH 4.8417 c and STH 5.8540.

### Identification of oat × maize hybrids

A total of 138 oat lines obtained by crossing with maize were tested for the presence of maize chromosomes. The presence of maize chromatin was indicated by amplification of the PCR product (500 bp) generated using primers specific for the maize retrotransposon *Grande-1* (GenBank accession number X97604; Ananiev, Phillips & Rines, 1998). Copies of the *Grande-1* are densely dispersed and located on each of the maize chromosomes. The 500 bp PCR product was identified in 66 oat plants. Figure 3 shows the exemplary agarose gel with 500 bp product corresponding to the maize retrotransposon *Grande-1*. Out of the 66 OMA lines containing maize chromatin, only 27 were fertile and produced seeds (Table 1). The 20 OMA lines that produced the highest number of seeds were then selected for cytogenetic analysis.

The seeds of two out of 20 OMA lines did not germinate. Among the remaining 20 lines, 8 lines (40%) had the character of additive lines, since GISH has detected one or more maize chromosomes in their cells. All of the analyzed plants possessed full complement of oat chromosomes. The number of detected maize chromosomes differed between the OMA lines (Table 1). They were usually smaller than the oat chromosomes and easy to discriminate (Fig. 4). In the plant STH 4.4690f, two pairs of chromosomes of varying size were found (Fig. 4A), whereas in STH 5.8504b only one maize chromosome was present (Fig. 4B). The majority of OMA lines possessed two maize chromosomes similar in size. In most of the lines, the detected chromosomes were labeled uniformly (Fig. 4C); however, in some, a banding pattern was visible that can potentially be used for the identification of maize addition (Fig. 4D). The banding pattern on both chromosomes in a given line was






**Figure 1** Oat haploid embryo formed after crossing with maize. (A); germinated haploid embryo on 190-2 medium (B); haploid plant on MS medium (C); acclimatization of haploid plant in perlite (D); DH plants in the greenhouse (E); panicles of OMA line STH5.8536/1, some of immature panicles are senesced (F); grassy OMA line STH5.8429 without panicles (G). Photo credit: Edyta Skrzypek.

Full-size [DOI: 10.7717/peerj.5107/fig-1](https://doi.org/10.7717/peerj.5107/fig-1)

similar, which might indicate that the added chromosomes are homologs. The presence of six 45S rDNA loci was detected in oat chromosomes (red hybridization signals, Fig. 4), but none of the added maize chromosomes in any of the lines carried 45S rDNA locus.

Twenty of the analyzed lines did not possess whole chromosomes of maize, but the introgression of maize chromatin was found in the oat chromosomes. A total of 12–14 hybridization signals for the maize gDNA probe were detected (Fig. 5). In all these lines, six of the maize gDNA hybridization signals colocalized with 25S rDNA sequence hybridization signals. In another pair of chromosomes, two strong signals were present in the pericentromeric positions on each of the chromosomes. Additionally, weaker maize gDNA signals were observed in one or two pairs of chromosomes of the lines STH 4.4690d (Fig. 5A), and STH 4.4606a (Fig. 5C), depending on the metaphase plate, and in one chromosome pair in the case of the line STH 4.4576 (Fig. 5B).



**Figure 2** Flow cytometry histograms of oat plants; (A) control 2n, (B) doubled haploid 2n, (C) haploid 1n and (D) mixoploid.

Full-size [DOI: 10.7717/peerj.5107/fig-2](https://doi.org/10.7717/peerj.5107/fig-2)


The colocalization of the gDNA and 25S rDNA signals may be related to the hybridization of the 25S rDNA sequence present in the maize genomic DNA to the same sequence in the oat genome. These sequences are highly conservative, hence the presence of these signals in the chromosomes of the analyzed lines. Additional hybridization signals found in 2–3 pairs of chromosomes, depending on the analyzed line, are presumably maize DNA sequences that have been incorporated into the oat chromosomes. Accurate determination of the origin of these sequences requires further analysis.

### Vigor of DH and OMA lines in their effectiveness in seeds production

The progeny of oat-maize hybrids showed some differences in morphology. Five of 66 hybrids were shorter in height, grassy type without panicles (Fig. 1G). Three of them had only a few short shoots with small panicles (Fig. 1F). Leaves' blades color and panicle shape were similar to their oat parent.

Analysis of variance showed significant variation in fertility in dependence on the addition of maize chromosomes to the oat genome (Table 1). OMA plants produced significantly less seeds than DH lines. Negative correlation coefficient ( $-0.279$ ) for seeds





**Figure 3 Identification of oat × maize F1 plants.** PCR products of genomic DNA of oat, maize, and a selection of 22 oat × maize F1 plants shown after electrophoresis in 1.5% (w/v) agarose gel. Bands represent 500 bp DNA fragments that were amplified with marker *Grande-1*. Marker leader is shown in the first line. Maize cv. Waza specificity is shown by product presence in maize DNA (positive control) and absence in oat cv. Stoper DNA (negative control). The presence of retained maize chromosomes is indicated in 14 out of the 22 F<sub>1</sub> plant DNAs shown. Photo credit: Tomasz Warzecha.

Full-size  DOI: [10.7717/peerj.5107/fig-3](https://doi.org/10.7717/peerj.5107/fig-3)


production dependently on the maize chromosomes/chromatin added to the oat genome was observed.

Sixty-three fertile DH lines out of 72 which did not have an addition of maize chromosomes or chromatin produced seeds (Table 2). Nine DH lines belonging to various genotypes did not produce seeds. The number of seeds was from 1 to 343, dependently on the genotype. In total, 3,758 seeds were produced. The highest number of seeds (343 in total) was produced by the STH 5.8429 genotype.

Twenty-seven OMA lines (from 66 identified) were fertile and produced seeds ranging in number from 1–102 (Table 2). In total, 613 seeds were produced by OMA lines. As for DH lines, the number of seeds produced by OMA lines depended on genotype, however nearly 41% of OMA plants were sterile. The differences in the seed production were not dependent on the number of chromosomes added to the oat genome or introgression of chromosome fragments to the oat chromosomes. OMA lines with the addition of whole chromosomes produced seeds as well as the ones with the introgression of fragments of maize chromosomes. All DH and OMA lines were fertile and produced seeds in the subsequent generation.

## DISCUSSION

In wide hybridization in Poaceae, the complete or incomplete chromosomes elimination of one of the parents occurs (Laurie & Bennett, 1986; Laurie & Bennett, 1988; Laurie, 1989; Rines & Dahleen, 1990). There are several reports concerning the mechanisms of this process. In wheat × maize hybrids, Mochidaetal2004 found the partial addition of the spindle microtubules to the centromeres of maize. In wheat × maize crosses, haploid




**Figure 4** Visualization of added maize chromosomes in oat genome by genomic *in situ* hybridization (GISH). (A) STH 4.4690f plant with tetrasomic addition of maize; (B) STH 5.8504b plant with monosomic addition of maize chromosome, (C) STH 5.8436b plant with disomic addition of maize chromosome. The maize chromosomes are labelled uniformly, (D) STH 4.4576 plant with disomic addition of maize chromosome. A banding pattern is visible on additional chromosomes. Maize gDNA is labeled with digoxigenin and detected anti-dig FITC (green fluorescence), rhodamine-5-dUTP—labeled 25S rDNA (red fluorescence) is used as an internal control of hybridization efficiency. Chromosomes are stained with DAPI (blue fluorescence). Scale bar: 10  $\mu$ m. Photo credit: Dominika Idziak-Helmcke.

Full-size  DOI: [10.7717/peerj.5107/fig-4](https://doi.org/10.7717/peerj.5107/fig-4)

wheat plants are obtained as a effect of complete elimination of the maize chromosomes prior to embryo rescue. In a wheat  $\times$  pearl millet hybrids cells, [Gernand et al. \(2005\)](#) detected delaying pearl millet chromosomes in anaphase and nuclear discarding of pearl millet chromatin in interphase. In the same wheat  $\times$  pearl millet hybrids, [Ishii et al. \(2010\)](#) noticed chromosomes breaking and lack of consistency of the pearl millet chromosomes.

The wide-cross combinations between various species of Triticeae as the female and a wide variety of Panicoideae species including sorghum and pearl millet as the male, showed a similar pattern of hybrid zygote formation with uniparental loss of the donor pollen chromosomes during subsequent embryo development ([Rines et al., 2003](#)). Chromosomes elimination was reported in crosses of wheat (female) and *Hordeum bulbosum* L. with maize (*Zea mays*), pearl millet (*Pennisetum glaucum*), sorghum (*Sorghum bicolor*), *Coix lacryma-jobi*, and *Imperata cylindrica* ([Barclay, 1975](#); [Laurie & Bennett, 1986](#); [Laurie & Bennett, 1988](#); [Laurie & Bennett, 1989](#); [Laurie, 1989](#); [Inagaki & Mujeeb-Kazi, 1995](#); [Mochida & Tsujimoto, 2001](#); [Komeda et al., 2007](#); [Ishii et al., 2010](#)). The cells of *H. bulbosum* fertilize



**Figure 5** Visualization of added fragments of maize chromosomes in oat genome by genomic *in situ* hybridization (GISH). (A) STH 4.4690d; (B) STH 4.4576; (C) STH 4.4606 F1 plants. The arrows point to maize introgressions into oat chromosomes (green fluorescence). Yellow signals result from colocalization of hybridization signals for maize gDNA (green fluorescence) and 25S rDNA (red fluorescence). Chromosomes are stained with DAPI (blue fluorescence). Scale bar: 10  $\mu$ m. Photo credit: Dominika Idziak-Helmcke.

Full-size  DOI: [10.7717/peerj.5107/fig-5](https://doi.org/10.7717/peerj.5107/fig-5)

the cells of wheat typically, but in hybrid embryogenesis the chromosomes of *H. bulbosum* are progressively extruded from the hybrid nucleus in the anaphase–telophase transition. The process is preceded by micronuclei and condensed chromosomes formation (Zenktelek & Straub, 1979).

Genomic *in situ* hybridization (GISH) is a modification of fluorescence *in situ* hybridization (FISH) in which the total genomic DNA of a given species is used as a probe to hybridize with the complementary DNA in a cytogenetic preparation (Schwarzacher, 2003). This technique allows for the genome-specific chromosome labeling in the cells of allopolyploid or hybrid species. It has been widely used in plant studies to identify putative parental genomes in the interspecific or intergeneric hybrids or allopolyploids (for the review see Chester *et al.*, 2010; Silva & Souza, 2013), to study the evolution of polyploid genomes (Song *et al.*, 1995; Gaeta *et al.*, 2007; Majka *et al.*, 2018; Tan *et al.*, 2017), or to detect the introgression of alien chromatin (Schneider *et al.*, 2005; Tan *et al.*, 2005). It has also become a valuable tool to analyze the meiotic recombination and behavior of chromosomes in the hybrids (Zwierzykowski *et al.*, 2008) as well as translocation (Kruppa *et al.*, 2013), addition (Molnar-Lang *et al.*, 2000; Ji & Chetelat, 2007), or substitution (Pan *et al.*, 2017) lines.

In the GISH-based studies, an excess of unlabeled blocking genomic DNA derived from the other parental species is often necessary in order to avoid non-specific cross-hybridization. Alternatively, the genomic DNAs of all parental species can be labeled differentially and hybridized simultaneously to the chromosomes of a hybrid/allopolyploid. In the studies of the genus *Paphiopedilum*, various ratios of blocking DNA has been applied in order to distinguish between the component genomes of the analyzed hybrid species, depending on the phylogenetic distance between the putative parents (Lee, Chang & Chung, 2011). In the case of a very close phylogenetic relationship between the parental species, even very high ratios of the blocking DNA are often not sufficient to distinguish

the chromosomes of each donor genome (*Xiong et al., 2006; Lee, Chang & Chung, 2011*). On the other hand, in the intergeneric hybrids and allopolyploids, GISH can be applied successfully even without blocking DNA (*Tang et al., 2011*). In our studies, the large genetic distance between maize and oat allowed us to specifically label and distinguish maize chromatin with no blocking DNA from oat. It should be noted, though, that the ease of discriminating the genomes of genetically distant species is usually coupled with difficulty in generating the crosses between them.

Although the GISH results allow us to unambiguously detect maize chromatin, the information provided by this method is not sufficient to identify which particular maize chromosome was added to the oat genome. The observation that some chromosomes in were labeled uniformly, whereas other displayed clearly a banding hybridization pattern, indicates that different maize chromosomes were retained in various OMA lines. The banding pattern probably reflects varying distribution of highly repetitive sequences in the chromosomes of maize since these sequences usually constitute the most of genomic DNA used as a probe. In the studies by *Rines et al. (2009)*, the identification of the maize chromosomes in the produced OMA lines was achieved using maize chromosome-specific simple sequence repeat (SSR) markers. An alternative, cytogenetic approach would require performing a FISH experiment with several different types of repetitive sequences as probes, labeled with different fluorophores and hybridized simultaneously in order to create unique banding pattern on added maize chromosomes. Comparing the results with the FISH on maize as a control would allow to discover the identity of the chromosome addition. Such multicolor-FISH method utilizing tandemly repeated DNA sequences has been applied before to maize chromosomes and permitted to distinguish each of the 10 chromosomes (*Kato, Lamb & Birchler, 2004*).

Oat (*Avena sativa* L.) haploid plants can be induced by hybridization either with maize (*Rines et al., 1996; Sidhu et al., 2006; Marcińska et al., 2013; Nowakowska et al., 2015; Warchoła et al., 2016*) or with pearl millet (*Rines et al., 1997; Ishii et al., 2013*). In oat and maize crosses, sometimes it happens that some maize chromosomes are not extruded during embryogenesis, and they ultimately stabilized and acted as oat chromosomes in mitosis (*Okagaki et al., 2001*). About one-third of the plants obtained from oat × maize crosses contain one or more maize chromosomes added to a complete set of oat chromosomes (*Rines et al., 2003; Rines et al., 2009*). Over the past few years, all ten maize chromosomes have been recovered as single chromosome additions to a haploid oat complement.

*Riera-Lizarazu, Rines & Phillips (1996)* after oat pollination by maize obtained 30 plants (one third of all plants) with retained one to four maize chromosomes, as a result of partial maize chromosomes elimination. Hybrid plants, in which some cells contained maize chromosomes and other did not, also were found. Authors explained that maize chromosomes elimination in crosses between oat and maize is more gradual than e.g., in wheat and maize. In the study by *Kynast et al. (2012)*, hybrids of oat and maize were euhaploids with a complete set of oat chromosomes without maize chromosomes, and aneuhaploids with a complete set of oat chromosomes and various number of retained maize chromosomes. Some of these haploids formed seeds to fifty percent of their spikelets. However, little is known about hybrids fertility *Kynast et al. (2012)* noticed that the

total seeds of F<sub>1</sub> hybrids generation with maintained maize chromosomes seemed to be meaningfully influenced by the number of retained maize chromosomes as well as by interaction of maize chromosomes with oat chromosomes. Also, the environmental conditions during F<sub>1</sub> hybrids cultivation appeared to had an impact on fertility. Based on chromosome observations and marker analyses in F<sub>1</sub> generation of oat × pearl millet crosses, *Ishii et al. (2015)* showed that the seedlings were full hybrids with all oat and pearl millet chromosomes. Although the hybrids of oat with pearl millet developed, they became necrotic in light condition.

In our experiment we found 53% of haploid plants and 47% of hybrids obtained after oat × maize crosses. The hybrids retained one to four maize chromosomes or fragments of maize chromosomes added to the oat chromosomes. Nearly 41% of hybrids was infertile. In the study by *Kynast & Riera-Lizarazu (2011)*, offspring of oat × maize consisted of 50–66% haploid oat plants and 34–50% partial hybrids. Haploid oat and partial hybrids with one to three maize chromosomes were incompletely fertile. Partial hybrids with more than three maize chromosomes had were smaller than haploid plants and sterile.

In our experiment seeds production depended on oat genotype and retained maize chromosomes in oat genome. Among 72 oat DH lines, nine did not produce seeds, whereas in the group of 66 hybrids—39 were infertile. Seeds produced even hybrid with retained four maize chromosomes. The hybrids also showed differences in morphology. Some of them were visibly shorter than maternal and DH plants. In contrast to *Kynast et al. (2001)*, they did not show differences in the color of leaf blade and panicle shape.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The research was funded by the National Centre for Research and Development project no. PBS3/B8/17/2015. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

The National Centre for Research and Development: PBS3/B8/17/2015.

### Competing Interests

Zygmunt Nita and Krystyna Werwińska are employed by Plant Breeding Strzelce Ltd., PBAI Group.

### Author Contributions

- Edyta Skrzypek and Tomasz Warzecha conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Angelika Noga performed the experiments, analyzed the data, approved the final draft.

- Marzena Warchoł performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Ilona Czyczyło-Mysza, Kinga Dziurka, Izabela Marcińska, Kamila Kapłoniak, Agnieszka Sutkowska, Krystyna Werwińska, Magdalena Rojek and Marta Hosiawa-Barańska performed the experiments, approved the final draft.
- Zygmunt Nita performed the experiments, contributed reagents/materials/analysis tools, approved the final draft.
- Dominika Idziak-Helmcke performed the experiments, analyzed the data, authored or reviewed drafts of the paper, approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in a [Supplemental File](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5107#supplemental-information>.

## REFERENCES

- Ananiev EV, Phillips RL, Rines HW. 1998.** Complex structure of knob DNA on maize chromosome 9: retrotransposon invasion into heterochromatin. *Genetics* **149**:2025–2037.
- Barclay IR. 1975.** High frequencies of haploid production in wheat (*Triticum aestivum*) by chromosome elimination. *Nature* **256**:410–411 DOI [10.1038/256410a0](https://doi.org/10.1038/256410a0).
- Bass HW, Riera-Lizarazu O, Ananiev EV, Bordoli SJ, Rines HW, Phillips RL, Sedat JW, Agard DA, Cande WZ. 2000.** Evidence for the coincident initiation of homolog pairing and synapsis during the telomere-clustering (bouquet) stage of meiotic prophase. *Journal of Cell Science* **113**:1033–1042.
- Britt AB, Kupp S. 2016.** CenH3: an emerging player in haploid induction technology. *Frontiers in Plant Science* **7**:357 DOI [10.3389/fpls.2016.00357](https://doi.org/10.3389/fpls.2016.00357).
- Chaudhary HK, Tayeng T, Kaila V, Rather SA. 2013.** Use of asynchrony in flowering for easy and economical polyhaploid induction in wheat following *Imperata cylindrica*—mediated chromosome elimination approach. *Plant Breeding* **132**(2):155–158 DOI [10.1111/pbr.12036](https://doi.org/10.1111/pbr.12036).
- Chester M, Leitch AR, Soltis PS, Soltis DE. 2010.** Review of the application of modern cytogenetic methods (FISH/GISH) to the study of reticulation (polyploidy/hybridisation). *Genes* **1**:166–192 DOI [10.3390/genes1020166](https://doi.org/10.3390/genes1020166).
- Dong Z, Yu J, Li H, Huang W, Xu L, Zhao Y, Zhang T, Xu W, Jiang J, Su Z, Jin W. 2018.** Transcriptional and epigenetic adaptation of maize chromosomes in Oat-Maize addition lines. *Nucleic Acids Research* **46**(10):5012–5028 DOI [10.1093/nar/gky209](https://doi.org/10.1093/nar/gky209).
- Doyle JJ, Doyle JL. 1987.** A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* **19**:11–15.



- Dunwell JM. 2010. Haploids in flowering plants: origins and exploitation. *Plant Biotechnology Journal* 8:377–424 DOI 10.1111/j.1467-7652.2009.00498.x.
- Gaeta RT, Pires JC, Iniguez-Luy F, Leon E, Osborn TC. 2007. Genomic changes in resynthesized *Brassica napus* and their effect on gene expression and phenotype. *The Plant Cell* 19:3403–3417 DOI 10.1105/tpc.107.054346.
- Garcia N, Zhang W, Wu Y, Messing J. 2015. Evolution of gene expression after gene amplification. *Genome Biology and Evolution* 7(5):1303–1312 DOI 10.1093/gbe/evv075.
- Gernand D, Bruss C, Houben A, Kumlehn J, Matzk F, Prodanovic S, Rubtsova M, Rutten T, Varshney A. 2005. Uniparental chromosome elimination at mitosis and interphase in wheat and pearl millet crosses involves micronucleus formation, progressive heterochromatinization, and DNA fragmentation. *The Plant Cell* 17:2431–2438 DOI 10.1105/tpc.105.034249.
- Hasterok R, Marasek A, Donnison IS, Armstead I, Thomas A, King IP, Wolny E, Idziak D, Draper J, Jenkins G. 2006. Alignment of the genomes of *Brachypodium distachyon* and temperate cereals and grasses using bacterial artificial chromosome landing with fluorescence in situ hybridization. *Genetics* 173:349–362 DOI 10.1534/genetics.105.049726.
- Inagaki M, Mujeeb-Kazi A. 1995. Comparison of polyhaploid production frequencies in crosses of hexaploid wheat with maize, pearl millet and sorghum. *Breeding Science* 45:157–161.
- Ishii T, Sunamura N, Matsumoto A, Eltayeb AE, Tsujimoto H. 2015. Preferential recruitment of the maternal centromere-specific histone H3 (CENH3) in oat (*Avena sativa* L.) × pearl millet (*Pennisetum glaucum* L.) hybrid embryos. *Chromosome Research* 23(4):709–718 DOI 10.1007/s10577-015-9477-5.
- Ishii T, Tanaka H, Eltayeb AE, Tsujimoto H. 2013. Wide hybridization between oat and pearl millet belonging to different subfamilies of Poaceae. *Plant Reproduction* 26:25–32 DOI 10.1007/s00497-012-0205-4.
- Ishii T, Ueda T, Tanaka H, Tsujimoto H. 2010. Chromosome elimination by wide hybridization between Triticeae or oat plant and pearl millet: pearl millet chromosome dynamics in hybrid embryo cells. *Chromosome Research* 18:821–831 DOI 10.1007/s10577-010-9158-3.
- Ji Y, Chetelat RT. 2007. GISH analysis of meiotic chromosome pairing in *Solanum lycopersicoides* introgression lines of cultivated tomato. *Genome* 50:825–833 DOI 10.1139/G07-069.
- Jin W, Melo JR, Nagaki K, Talbert PB, Henikoff S, Dawe RK, Jiang J. 2004. Maize centromeres: organization and functional adaptation in the genetic background of oat. *The Plant Cell* 16:571–581 DOI 10.1105/tpc.018937.
- Karimi-Ashtiyani R, Ishii T, Niessen M, Stein N, Heckmann S, Gurushidze M, Banaei-Moghaddam AM, Fuchs J, Schubert V, Koch K, Weiss O, Demidov D, Schmidt K, Kumlehn J, Houben A. 2015. Point mutation impairs centromeric CENH3 loading and induces haploid plants. *Proceedings of the National Academy of Sciences of the United States of America* 112(36):11211–11216 DOI 10.1073/pnas.1504333112.

- Kasha KJ, Kao KN. 1970.** High frequency haploid production in barley (*Hordeum vulgare* L.). *Nature* **225**:874–876 DOI [10.1038/225874a0](https://doi.org/10.1038/225874a0).
- Kato A, Lamb JC, Birchler JA. 2004.** Chromosome painting using repetitive DNA sequences as probes for somatic chromosome identification in maize. *Proceedings of the National Academy of Sciences of the United States of America* **101**:13554–13559 DOI [10.1073/pnas.0403659101](https://doi.org/10.1073/pnas.0403659101).
- Kelliher T, Starr D, Wang W, McCuiston J, Zhong H, Nuccio ML, Martin B. 2016.** Maternal haploids are preferentially induced by CENH3-tailswap transgenic complementation in maize. *Frontiers in Plant Science* **7**:414 DOI [10.3389/fpls.2016.00414](https://doi.org/10.3389/fpls.2016.00414).
- Komeda N, Chaudhary HK, Suzuki G, Mukai Y. 2007.** Cytological evidence for chromosome elimination in wheat × *Imperata cylindrica* hybrids. *Genes & Genetic Systems* **82**:241–248 DOI [10.1266/ggs.82.241](https://doi.org/10.1266/ggs.82.241).
- Koumbaris GL, Bass HW. 2003.** A new single-locus cytogenetic mapping system for maize (*Zea mays* L.): overcoming FISH detection limits with marker-selected sorghum (*S. propinquum* L.) BAC clone. *The Plant Journal* **35**:647–659 DOI [10.1046/j.1365-313X.2003.01829.x](https://doi.org/10.1046/j.1365-313X.2003.01829.x).
- Kowles RV, Walch MD, Minnerath JM, Bernacchi CJ, Stec AO, Rines HW, Phillips RL. 2008.** Expression of C4 photosynthetic enzymes in oat–maize chromosome addition lines. *Maydica* **53**:69–78.
- Kruppa K, Sepsi A, Szakacs E, Roder MS, Molnar-Lang M. 2013.** Characterization of a 5HS-7DS.7DL wheat-barley translocation line and physical mapping of the 7D chromosome using SSR markers. *Journal of Applied Genetics* **54**:251–258 DOI [10.1007/s13353-013-0152-2](https://doi.org/10.1007/s13353-013-0152-2).
- Kynast RG, Davis DW, Phillips RL, Rines HW. 2012.** Gamete formation via meiotic nuclear restitution generates fertile amphiploid F1 (oat × maize) plants. *Sexual Plant Reproduction* **25**:111–122 DOI [10.1007/s00497-012-0182-7](https://doi.org/10.1007/s00497-012-0182-7).
- Kynast RG, Okagaki RJ, Rines HW, Phillips RL. 2002.** Maize individualized chromosome and derived radiation hybrid lines and their use in functional genomics. *Functional and Integrative Genomics* **2**:60–69 DOI [10.1007/s10142-002-0052-8](https://doi.org/10.1007/s10142-002-0052-8).
- Kynast RG, Riera-Lizarazu O. 2011.** Development and use of oat-maize chromosome additions and radiation hybrids. In: Birchler JA, ed. *Plant chromosome engineering: methods and protocols, methods in molecular biology*, vol. 701, New York: Humana Press, c/o Springer Science + Business Media, 259–284.
- Kynast RG, Riera-Lizarazu O, Vales MI, Okagaki RJ, Maquieira SB, Chen G, Ananiev EV, Odland WE, Russell CD, Stec AO, Livingston SM, Zaia HA, Rines HW, Phillips RL. 2001.** A complete set of maize individual chromosome additions to the oat genome. *Plant Physiology* **125**:1216–1227 DOI [10.1104/pp.125.3.1216](https://doi.org/10.1104/pp.125.3.1216).
- Laurie DA. 1989.** The frequency of fertilization in wheat × pearl millet crosses. *Genome* **32**:1063–1067 DOI [10.1139/g89-554](https://doi.org/10.1139/g89-554).
- Laurie DA, Bennett MD. 1986.** Wheat × maize hybridization. *Canadian Journal of Genetics and Cytology* **28**:313–316 DOI [10.1139/g86-046](https://doi.org/10.1139/g86-046).



- Laurie DA, Bennett MD. 1988. Cytological evidence for fertilization in hexaploid wheat × sorghum crosses. *Plant Breeding* **100**:73–82  
DOI [10.1111/j.1439-0523.1988.tb00220.x](https://doi.org/10.1111/j.1439-0523.1988.tb00220.x).
- Laurie DA, Bennett MD. 1989. The timing of chromosome elimination in hexaploid wheat × maize crosses. *Genome* **32**:953–961 DOI [10.1139/g89-537](https://doi.org/10.1139/g89-537).
- Lee YI, Chang FC, Chung MC. 2011. Chromosome pairing affinities in interspecific hybrids reflect phylogenetic distances among lady's slipper orchids (*Paphiopedilum*). *Annals of Botany* **108**:113–121 DOI [10.1093/aob/mcr114](https://doi.org/10.1093/aob/mcr114).
- Liu D, Zhang H, Zhang L, Yuan Z, Hao M, Zheng Y. 2014. Distant hybridization: a tool for interspecific manipulation of chromosomes. In: Pratap A, Kumar J, eds. *Alien gene transfer in crop plants, volume 1: innovations, methods and risk assessment*. New York: Springer Science + Business Media, 25–42.
- Maheshwari S, Tan EH, West A, Franklin FCH, Comai L, Chan SWL. 2015. Naturally occurring differences in CENH3 affect chromosome segregation in zygotic mitosis of hybrids. *PLOS Genetics* **11**:e1004970 DOI [10.1371/journal.pgen.1004970](https://doi.org/10.1371/journal.pgen.1004970).
- Majka J, Zwierzykowski Z, Majka M, Kosmala A. 2018. Karyotype reshufflings of *Festuca pratensis* × *Lolium perenne* hybrids. *Protoplasma* **255**(2):451–458  
DOI [10.1007/s00709-017-1161-5](https://doi.org/10.1007/s00709-017-1161-5).
- Maquieira SB. 1997. Production and characterization of plants from oat × maize and oat × pearl millet. MS thesis, University of Minnesota, St. Paul.
- Marcińska I, Nowakowska A, Skrzypek E, Czyczyło-Mysza I. 2013. Production of double haploids in oat (*Avena sativa* L.) by pollination with maize (*Zea mays* L.). *Central European Journal of Biology* **8**:306–313 DOI [10.2478/s11535-013-0132-2](https://doi.org/10.2478/s11535-013-0132-2).
- Mochida K, Tsujimoto H. 2001. Production of wheat double haploids by pollination with Job's Tears (*Coix lacrymajobi* L.). *Heredity* **92**:81–83  
DOI [10.1093/jhered/92.1.81](https://doi.org/10.1093/jhered/92.1.81).
- Molnar-Lang M, Linc G, Logojan A, Sutka J. 2000. Production and meiotic pairing behaviour of new hybrids of winter wheat (*Triticum aestivum*) × winter barley (*Hordeum vulgare*). *Genome* **43**:1045–1054 DOI [10.1139/g00-079](https://doi.org/10.1139/g00-079).
- Murashige T, Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum* **15**:473–497  
DOI [10.1111/j.1399-3054.1962.tb08052.x](https://doi.org/10.1111/j.1399-3054.1962.tb08052.x).
- Noga A, Skrzypek E, Warchoń M, Czyczyło-Mysza I, Dziurka K, Marcińska I, Juzoń K, Warzecha T, Sutkowska A, Nita Z, Werwinska K. 2016. Conversion of oat (*Avena sativa* L.) haploid embryos into plants in relation to embryo developmental stage and regeneration media. *In Vitro Cellular & Developmental Biology-Plant* **52**:590–597  
DOI [10.1007/s11627-016-9788-z](https://doi.org/10.1007/s11627-016-9788-z).
- Nowakowska A, Skrzypek E, Marcińska I, Czyczyło-Mysza I, Dziurka K, Juzoń K, Cyganek K, Warchoń M. 2015. Application of chosen factors in the wide crossing method for the production of oat doubled haploids. *Open Life Sciences* **10**:112–118  
DOI [10.1515/biol-2015-0014](https://doi.org/10.1515/biol-2015-0014).

- Okagaki RJ, Kynast RG, Livingston SM, Russell CD, Rines HW, Phillips RL. 2001.** Mapping maize sequences to chromosome using oat-maize chromosome addition materials. *Plant Physiology* **125**:1228–1235 DOI [10.1104/pp.125.3.1228](https://doi.org/10.1104/pp.125.3.1228).
- Pan C, Li Q, Lu Y, Zhang J, Yang X, Li X, Li L, Liu W. 2017.** Chromosomal localization of genes conferring desirable agronomic traits from *Agropyron cristatum* chromosome 1P. *PLOS ONE* **12**:e0175265 DOI [10.1371/journal.pone.0175265](https://doi.org/10.1371/journal.pone.0175265).
- Ravi M, Chan SWL. 2010.** Haploid plants produced by centromere-mediated genome elimination. *Nature* **464**:615–618 DOI [10.1038/nature08842](https://doi.org/10.1038/nature08842).
- Riera-Lizarazu O, Rines HW, Phillips RL. 1996.** Cytological and molecular characterization of oat × maize partial hybrids. *Theoretical and Applied Genetics* **93**:123–135 DOI [10.1007/BF00225737](https://doi.org/10.1007/BF00225737).
- Rines HW, Dahleen LS. 1990.** Haploid oat plants produced by application of maize pollen to emasculated oat florets. *Crop Science* **30**:1073–1078 DOI [10.2135/cropsci1990.0011183X003000050023x](https://doi.org/10.2135/cropsci1990.0011183X003000050023x).
- Rines HW, Phillips RL, Kynast RG, Okagaki RJ, Galatowitsch MW, Huettl PA, Stec AO, Jacobs MS, Suresh J, Porter HL, Walch MD, Cabral CB. 2009.** Addition of individual chromosomes of maize inbreds B73 and Mo17 to oat cultivars Starter and Sun II: maize chromosome retention, transmission, and plant phenotype. *Theoretical and Applied Genetics* **119**:1255–1264 DOI [10.1007/s00122-009-1130-2](https://doi.org/10.1007/s00122-009-1130-2).
- Rines HW, Phillips RL, Kynast RG, Okagaki RJ, Odland WE, Stec AO, Jacobs MS, Granath SR. 2003.** Maize chromosome additions and radiation hybrids in oat and their use in dissecting the maize genome. In: Tuberosa R, Phillips RL, Gale M, eds. *Proceedings of the international congress “in the wake of the double helix: from the green revolution to the gene revolution,” 27–31 May 2003, Bologna, Italy*, 427–441.
- Rines HW, Riera-Lizarazu O, Maquieira SB, Phillips RL. 1996.** Wide crosses for haploids. In: Scoles G, Rossnagel B, eds. *Proceedings of the fifth international oat conference and eighth international barley genetic symposium, Saskatoon, Saskatchewan, Canada*, 207–212.
- Rines HW, Riera-Lizarazu O, Nunez VM, Davis DM, Phillips RL. 1997.** Oat haploids from anther culture and from wide hybridizations. In: Jain SM, Sopory SK, Veilleux RE, eds. *In vitro production of haploids in higher plants 4*. Dordrecht: Kluwer Academic Publishers, 205–221.
- Schneider A, Linc G, Molnar I, Molnar-Lang M. 2005.** Molecular cytogenetic characterization of *Aegilops biuncialis* and its use for the identification of 5 derived wheat-*Aegilops biuncialis* disomic addition lines. *Genome* **48**:1070–1082 DOI [10.1139/g05-062](https://doi.org/10.1139/g05-062).
- Schwarzacher T. 2003.** DNA, chromosomes, and in situ hybridization. *Genome* **46**:953–962 DOI [10.1139/g03-119](https://doi.org/10.1139/g03-119).
- Sidhu PK, Howes NK, Aung T, Zwer PK, Davies PA. 2006.** Factors affecting oat haploid production following oat × maize hybridization. *Plant Breeding* **125**:243–247 DOI [10.1111/j.1439-0523.2006.01206.x](https://doi.org/10.1111/j.1439-0523.2006.01206.x).
- Silva GS, Souza MM. 2013.** Genomic in situ hybridization in plants. *Genetics and Molecular Research* **12**:2953–2965 DOI [10.4238/2013.August.12.11](https://doi.org/10.4238/2013.August.12.11).

- Song K, Lu P, Tang K, Osborn TC. 1995.** Rapid genome change in synthetic polyploids of *Brassica* and its implications for polyploid evolution. *Proceedings of the National Academy of Sciences of the United States of America* **92**:7719–7723 DOI [10.1073/pnas.92.17.7719](https://doi.org/10.1073/pnas.92.17.7719).
- Tan C, Cui C, Xiang Y, Ge X, Li Z. 2017.** Development of *Brassica oleracea-nigra* monosomic alien addition lines: genotypic, cytological and morphological analyses. *Theoretical and Applied Genetics* **130**:2491–2504 DOI [10.1007/s00122-017-2971-8](https://doi.org/10.1007/s00122-017-2971-8).
- Tan G, Jin H, Li G, He R, Zhu L, He G. 2005.** Production and characterization of a complete set of individual chromosome additions from *Oryza officinalis* to *Oryza sativa* using RFLP and GISH analyses. *Theoretical and Applied Genetics* **111**:1585–1595 DOI [10.1007/s00122-005-0090-4](https://doi.org/10.1007/s00122-005-0090-4).
- Tang F, Wang H, Chen S, Chen F, Liu Z, Fang W. 2011.** Intergeneric hybridization between *Dendranthema nankingense* and *Tanacetum vulgare*. *Scientia Horticulturae* **132**:1–6 DOI [10.1016/j.scienta.2011.10.003](https://doi.org/10.1016/j.scienta.2011.10.003).
- Topp CN, Okagaki RJ, Melo JR, Kynast RG, Phillips RL, Dawe RK. 2009.** Identification of a maize neocentromere in an oat-maize addition line. *Cytogenetic and Genome Research* **124**:228–238 DOI [10.1159/000218128](https://doi.org/10.1159/000218128).
- Wang K, Wu Y, Dawe K, Jiang J. 2014.** Maize centromeres expand and adopt a uniform size in the genetic background of oat. *Genome Research* **24**:107–116 DOI [10.1101/gr.160887.113](https://doi.org/10.1101/gr.160887.113).
- Warchoń M, Skrzypek E, Nowakowska A, Barcińska I, Czyczyło-Mysza I, Dziurka K, Juzoń K, Cyganek K. 2016.** The effect of auxin and genotype on the production of *Avena sativa* L. doubled haploids lines. *Plant Growth Regulation* **78**:155–165 DOI [10.1007/s10725-015-0082-6](https://doi.org/10.1007/s10725-015-0082-6).
- Xiong ZY, Tan GX, He GY, He GC, Song YC. 2006.** Cytogenetic comparisons between A and G genomes in *Oryza* using genomic in situ hybridization. *Cell Research* **16**:260–266 DOI [10.1038/sj.cr.7310033](https://doi.org/10.1038/sj.cr.7310033).
- Zenkteler M, Straub J. 1979.** Cytoembryological studies on the process of fertilization and the development of haploid embryos of *Triticum aestivum* L. ( $2n = 47$ ) after crossing with *Hordeum bulbosum* ( $2n = 14$ ). *Zeitschrift für Pflanzenzüchtung* **82**:32–44.
- Zhuang JJ, Xu J. 1983.** Increasing differentiation frequencies in wheat pollen callus. In: Hu H, Vega MR, eds. *Cell and tissue culture techniques for cereal crop improvement*. Beijing: Science Press 431.
- Zwierzykowski Z, Zwierzykowska E, Taciak M, Jones N, Kosmala A, Krajewski P. 2008.** Chromosome pairing in allotetraploid hybrids of *Festuca pratensis* x *Lolium perenne* revealed by genomic in situ hybridization (GISH). *Chromosome Research* **16**:575–585 DOI [10.1007/s10577-008-1198-6](https://doi.org/10.1007/s10577-008-1198-6).