

Research article

Open Access

Angiomotin and angiomotin like proteins, their expression and correlation with angiogenesis and clinical outcome in human breast cancer

Wen G Jiang*¹, Gareth Watkins¹, Anthony Douglas-Jones², Lars Holmgren³ and Robert E Mansel¹

Address: ¹Metastasis and Angiogenesis Research Group, Wales College of Medicine, Cardiff University, Cardiff, UK, ²Department of Pathology, Wales College of Medicine, Cardiff University, Cardiff, UK and ³Department of Oncology, Karolinska Institutet, Stockholm, Sweden

Email: Wen G Jiang* - jiangw@cf.ac.uk; Gareth Watkins - watkinsg1@cf.ac.uk; Anthony Douglas-Jones - douglas-jones@cf.ac.uk; Lars Holmgren - lars.holmgren@cck.ki.se; Robert E Mansel - manselRE@cf.ac.uk

* Corresponding author

Published: 23 January 2006

Received: 07 July 2005

BMC Cancer 2006, 6:16 doi:10.1186/1471-2407-6-16

Accepted: 23 January 2006

This article is available from: <http://www.biomedcentral.com/1471-2407/6/16>

© 2006 Jiang et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Angiomotin is a newly discovered molecule that regulates the migration and tubule formation of endothelial cells. It therefore has been implicated in the control of angiogenesis under physiological and pathological conditions. This study examined the expression of angiomotin and its analogues, angiomotin-like 1 (L1) and -like 2 (L2) in breast tumour tissues, and analysed their correlation with angiogenesis and clinical outcomes.

Methods: Human breast tissues (normal $n = 32$ and tumours $n = 120$) were used. The levels of expression of angiomotin, L1 and L2 were determined using reverse transcription PCR. Microvessels were stained using antibodies against PECAM, von Willebrand factor (factor 8, or vWF) and VE-cadherin. The transcript levels of angiomotin and its analogues were assessed against the clinical and pathological background, including long term survival (120 months).

Results: Breast cancer tissues expressed significantly higher levels of angiomotin transcript, compared with normal mammary tissues (33.1 ± 11 in normal versus 86.5 ± 13.7 in tumour tissues, $p = 0.003$). Both L1 and L2 were seen at marginally higher levels in tumour than normal tissues but the difference was not statistically significant. Levels of angiomotin were at significantly higher levels in grade 2 and grade 3 tumours compared with grade 1 ($p < 0.01$ and $p = 0.05$ respectively). The levels of angiomotin in tumours from patients who had metastatic disease were also significantly higher than those patients who remained disease free ($p = 0.03$). Multivariate analysis indicated that angiomotin transcript was an independent prognostic factor ($p = 0.031$). No significant correlations were seen between angiomotin-L1 and L2 with the clinical outcome. Furthermore, high levels of angiomotin transcript were associated with shorter overall survival ($p < 0.05$). There was a high degree of correlation between levels of vW factor and that of angiomotin ($p < 0.05$), but not angiomotin-L1 and angiomotin-L2.

Conclusion: Angiomotin, a putative endothelial motility factor, is highly expressed in human breast tumour tissues and linked to angiogenesis. It links to the aggressive nature of breast tumours and the long term survival of the patients. These data point angiomotin as being a potential therapeutic target.

Background

Angiogenesis is essential in the development and vascular spread of cancer, by providing nutrients, oxygen, and passages for the departing cancer cells [1-3]. The angiogenic process is regulated by a carefully maintained balance between angiogenic factors and angiogenic inhibitors (angiogenic factors). In cancer, the pro-angiogenic factors frequently gain the 'upper hand', which stimulate vascular endothelial cells to growth, migrate and forming new vascular/capillary tubules. Most angiogenic factors are growth factors that increase the proliferation of vascular endothelial cells. Some factors, however, are strongly involved in the migration and morphogenesis of endothelial cells, such as hepatocyte growth factor. These factors are mainly produced by stromal cells and act via a paracrine pathway. Angiogenic factors, their receptors or molecules specific to vascular endothelial cells have been used to assess angiogenesis. Notable ones include von Willebrand factor (factor-8 or vWF), PE-CAM, VE-Cadherin, VEGF-receptors [4-6].

Angiomotin (AF286598) is a molecule recently identified from its ability to bind to angiostatin using a yeast two hybrid screen [7]. Angiomotin exerts a strong effect in inducing the migration and tubule formation from endothelial cells and promotes angiogenesis. The effect appears to be via its interaction with and subsequent inhibition of angiostatin, an angiogenesis inhibitor. However, other mechanism(s) may also operate, including possible interaction with integrins. Angiostatin is known to inhibit angiogenesis and metastasis in solid tumours [8]. Angiomotin belongs to a new protein family with which its members share sequence and structural similarities. Two other known members in the family include angiomotin-like-1 and angiomotin-like-2 proteins [9]. Angiomotin-like-1 is also known as junction-enriched and -associated protein (JEAP) that is highly located at tight junctions and co-localised with JEAP [10]. Angiomotin-like-2 has, however, no known functions identified.

The potential pro-motility function of angiomotin has suggested an important role of the molecule in angiogen-

esis. Indeed, it has been shown that transfection of micro-capillary endothelial cells with angiomotin expression vector increases the migration and tubule forming of the cells [7]. Angiomotin deficient mice died in their early days due to a migration defect during their development, further indicating the potential role of angiomotin in cell motility [11]. The important biological role of angiomotin and its analogues indicates that it may play an important role in angiogenesis in tumours. Indeed, angiomotin has been found to be expressed highly in vessels of Kaposi sarcoma, and weakly in vessels from normal tissues in an early study [7]. However, the expression, distribution pattern and the clinical implications of angiomotin in other tumour types are yet to be explored.

Breast cancer is the leading female cancer in U.S. and U.K. The metastatic spread of the tumour is the primary cause of death of the patients. In the past decade, angiogenesis has been shown to be an important biological marker in predicting prognosis and clinical outcome of patients with breast cancer. Traditionally, angiogenesis has been assessed using markers including von Willebrand factor (vWF), VE-cadherin (also known as cadherin-5) and PE-CAM (CD31), and has been found to be increased in tumour tissues compared with normal tissues [12-15]. Micro-vessel density (MVD) has been used as a mean to calculate angiogenesis in these studies and has been shown to be associated with the progress and metastasis of breast cancer [16,17]. A number of the angiogenic factors, such as VEGF has also shown to be linked to prognosis in patients with breast cancer.

We examined the expression of angiomotin and its analogue molecules angiomotin-like-1 and like-2 in a cohort of breast tumours against the clinical information. Here, we report for the first time the aberrant expression of angiomotin in breast tumours, its correlation with angiogenesis and association with metastatic disease in patients with breast cancer.

Table 1: Clinical and pathological information of the study cohort. Shown are number of samples in each group

| | | | | | | |
|------------------|---------------|-----------------|-------------------|-----------------------|----------------------------|--------|
| Node status | Node negative | Node positive | | | | |
| n= | 65 | 55 | | | | |
| Grade | Grade 1 | Grade 2 | Grade 3 | | | |
| n= | 23 | 41 | 56 | | | |
| Histology | Ductal | Lobular | Medullary | Tubular | Mucinous | Others |
| n= | 94 | 14 | 2 | 2 | 4 | 4 |
| TNM staging | TNM 1 | TNM 2 | TNM 3 | TNM 4 | | |
| n= | 69 | 40 | 7 | 4 | | |
| Clinical outcome | Disease free | With Metastasis | With local recur. | Died of breast Cancer | Died of unrelated diseases | |
| n= | 81 | 7 | 5 | 20 | 7 | |

Methods

Materials

RNA extraction kit and RT kit were obtained from AbGene Ltd, Surrey, England, UK. PCR primers were designed using Beacon Designer (Palo Alto, California, USA) and synthesised by Invitrogen Ltd (Pasley, Scotland, UK). Molecular biology grade agarose and DNA ladder were from Invitrogen (Pasley, Scotland, UK. Master mix for routine PCR and quantitative PCR was from AbGene (Surrey, England, UK). Rabbit anti-human VE-cadherin and anti-factor-8 (von Willebrand Factor), goat anti-rabbit and an universal staining kit were from Santa Cruz Biotechnology Ltd (Santa Cruz, CA, USA), and Vector Laboratories (Nottingham, England, UK), respectively.

Sample collection

Breast cancer tissues (n = 120) and normal background tissues (n = 32) were collected immediately after surgery and stored in the deep freezer until use. Patients were routinely followed clinically after surgery. The median follow-up period was 120 months (June 2004). The presence of tumour cells in the collected tissues was verified by a consultant pathologist, who examined H&E stained frozen sections. Details of histology were obtained from Pathology reports and re-verified by a consultant pathologist (ADJ) (table 1). Patients were routinely followed up on a regular basis and details stored in a database.

Tissue processing, RNA extraction and cDNA synthesis

Frozen sections of tissues were cut at a thickness of 5–10 μm and were kept for immunohistochemistry and routine histology. Additional 15–20 sections were mixed and homogenised using a hand-held homogeniser, in ice cold RNA extraction solution. The concentration of RNA was determined using a UV spectrophotometer. Reverse transcription was carried using a RT kit with an anchored oligo-dt primer supplied by AbGene, using 1 μg total RNA in 96-well plate. The quality of cDNA was verified using β -actin primers.

Quantitative analysis of the transcripts of angiomin, angiomin-related molecule, and endothelial markers

The level of angiomin, angiomin-like-1 and angiomin-like-2, PECAM-1, VE-cadherin and factor-8 transcripts from the above-prepared cDNA was determined using a real-time quantitative PCR, based on the Amplifluor™ technology, modified from a method previously reported [17,18]. Briefly, pairs of PCR primers were designed using the Beacon Designer software (version 2, California, USA). To one of the primers, an additional sequence, known as the Z sequence (5'actgaacctgacctgaca'3) which is complementary to the universal Z probe (Intergen Inc., Oxford, England, UK), was added. Sequences of the respective primers were: Angiomin (5'atagctgtagggagaacag'3 and 5'ctgaagaactgcgactgtg'3),

Angiomin-like-1 (5'catgagagctgaccaga'3, and 5'cctcattctactgtccatct'3), angiomin-like-2 (5'caccatcaccaccacct'3 and 5'agaaacagcagcagcagtag'3), PECAM-1 (CD31)(5'ccatcatggaggatgag'3 and 5' actgaacctgacctgacctgacctgacctgttct'3), VE-cadherin (5' gggagaccagcctctgtc'3 and 5' actgaacctgacctgacctgacctgacctgacctgtc'3), vWF (5' ggagagatgggacactaaca'3 and 5' actgaacctgacctgacctgacctgacctgacctgaggt'3) and β -actin: 5'atgatatcgccgctcg'3 and 5'cgctcgtgtaggatttca'3 A Taqman detection kit for β -actin was purchased from Perkin-Elmer. The reaction was carried out using the following: Hot-start Q-master mix (Abgene), 10 pmol of specific forward primer, 1 pmol reverse primer which has the Z sequence, 10 pmol of FAM-tagged probe (Intergen Inc., Oxford, England, UK), and cDNA from approximate 50 ng RNA. The reaction was carried out using IcylerIQ™ (Bio-Rad) which is equipped with an optic unit that allows real time detection of 96 reactions, using the following condition: 94°C for 12 minutes, 50 cycles of 94°C for 15 seconds, 55°C for 40 seconds and 72°C for 20 seconds. The levels of the transcripts were generated from a standard that was simultaneously amplified with the samples. Cytokeratin-19 (CK19) was used for comparison of cellularity during the analysis and primers for CK19 were 5'-caggctccagggtactgac-3' and 5'-actgaacctgacctgacctgacctgacctgacctgtcttc-3' respectively [19,20]. The levels of the angiomin and the angiomin like transcripts are shown here as the number of the respective copies and the ratio of angiomin (or its like proteins) vs cytokeratin 19 (CK19).

Assessment of micro-vessel density (MVD) using anti-von Willebrand Factor (vWF) and VE-cadherin

The assessment of MVD was as previously reported [21,22]. Frozen sections of breast tumour and background tissue were cut at a thickness of 6 μm using a cryostat. The sections were mounted on super frost plus microscope slides, air dried and then fixed in a mixture of 50% Acetone and 50% methanol. The sections were then placed in "Optimax" wash buffer for 5–10 minutes to rehydrate. Sections were incubated for 20 mins in a 0.6% BSA blocking solution and probed with the primary antibody (1:50 dilution) for 2 hours at room temperature. Following extensive washings, sections were incubated for 30 mins in the secondary biotinylated antibody (Multilink Swine anti- goat/mouse/rabbit immunoglobulin, Dako Inc.). Following washings, Avidin Biotin Complex (Vector Laboratories) was then applied to the sections followed by extensive washings. Diamino benzidine chromogen (Vector Labs) was then added to the sections which were incubated in the dark for 5 mins. Sections were then counter stained in Gill's Haematoxylin and dehydrated in ascending grades of methanol before clearing in xylene and mounting under a cover slip. Microvessels were counted using X20 objectives by 3 independent researchers, as we have recently reported [21].




Figure 1

Expression of angiomin and angiomin like transcript in normal and tumour tissues using conventional RT-PCR (A) and quantitative real-time PCR (B). A: paired normal (N) and tumour (T) tissues from selected patients. B: levels of the respective transcript and the transcript:CK19 ratio (inserts) from all the samples (normal n = 32 and tumour n = 120). * p < 0.05 vs tumours.




Figure 2

Levels of expression of angiomotins and their relationship with grade (Top panel, A-C) and histological types (bottom panel, D-F). A and D: angiomin, B and E: angiomin-like-1, C and F: angiomin-like-2. *** $p = 0.01$, * $p = 0.049$, vs grade 1 tumours. Inserts: respective angiomin transcript:CK19 transcript ratio.

Statistical analysis was carried out using Mann-Whitney U test and the Kruskal-Wallis test. Survival analysis was carried out using the Kaplan-Meier's and Cox Proportional tests with SPSS12 package.

Results

Expression of angiomotins in mammary tissues

The presence of angiomin and the angiomin-like transcripts were detected in both normal and tumour mammary tissues (figure 1A). In these selective paired samples, it was seen that most tumour samples had a stronger angiomin signal, however, signals for angiomin-like-1 and -like-2 were not different between normal and tumour tissues.

A quantitative analysis of the molecules indicated that there was a significantly higher level of angiomin in

breast tumour tissues, than in normal tissues (figure 1B-a). The same trend was reflected when the transcripts were normalised by CK19, as measure of controlling the cellularity (figure 1B-a insert). Angiomin like-1 and like-2 transcript were also high in tumour tissues compared with normal tissues (figure 1B-b and 1B-c, respectively), however, the difference was not statistically significant. Moreover, after being normalised by CK19, as shown in figure 1B inserts, tumour tissues exhibited a lower ratio, compared with normal tissue, although this was not statistically significant. No significant difference between normal and tumour was seen for the angiomin-like-2:CK19 ratio.

Angiomin and histological type and tumour grade

Grade-2 and grade-3 tumours had significantly higher levels of angiomin transcript than grade 1 tumours ($p =$




Figure 3

Angiominins in nodal status (top panel, A-C) and predicted prognosis (bottom panel, E-F). A and D: angiomin, B and E: angiomin-like-1, C and F: angiomin-like-2. A, B AND C : difference between node negative and node positive tumours. * $p < 0.01$ vs node negative tumours. D, E & F: angiominins and predicted clinical outcome based on Nottingham Prognostic Index (NPI = $(0.2 \times \text{size}) + \text{grade} + \text{Nodal status}$). NPI < 3.4, 3.4–5.4 and > 5.4 represented good (15 year survival rate 80%) (NPI1), moderate (15 year survival 42%) (NPI2) and poor prognosis (15 year survival 15%). Inserts: respective angiomin transcript:CK19 transcript ratio.




Figure 4
 Angiomin and clinical outcomes. Patients were assessed based on the outcome after 10 years from initial surgery, and were divided into those who remained disease free, with metastasis, with local recurrence and who died of breast cancer related causes. Inserts: respective angiomin transcript:CK19 transcript ratio. * p = 0.03 vs disease free group.

0.01 Grade 1 vs grade 2, p < 0.05 grade 1 vs grade 3) (figure 2A). The angiomin:CK19 ratio was also significantly higher in grade 2 and grade 3 tumours compared with grade 1 tumours (figure 2A insert). No significant difference between different grades was seen with angiomin-Like-1 or its ratio to CK19 (figure 2B and insert). A marginally high level of angiomin-like-2 was seen in grade 3 tumours, however neither the transcript nor the transcript:CK19 reached a statistical difference (figure 2C and insert).

Ductal tumours had higher levels of angiomin (91.4 ± 13.7) than lobular tumours (48.1 ± 16, p = 0.026) (figure 2D). A similar trend was seen with angiomin-like-1 (298.8 ± 51.4 for ductal and 78.1 ± 25.7 for lobular, p = 0.045). Levels of angiomin like-2 transcript were marginally higher in ductal tumours (139 ± 92), compared with lobular tumours (29.2 ± 7.3) p = 0.24) (figure 2E and 2F).

Angiomin is associated with nodal involvement and predicted clinical outcome

There was a significantly higher level of angiomin and angiomin:CK19 ratio in tumour with positive axillary nodes (p = 0.0018) (figure 3A and insert). Although there was a trend of higher levels of angiomin-like-1 and

like-2 in node positive tumours, this was not significant (p = 0.08 and p = 0.6, respectively, figure 3B and 3C).

We have used Nottingham Prognostic Index (NPI) as an indicator for predicted clinical outcome. As shown in figure 3D, although there was a trend of high level of angiomin transcript in moderate and poor prognostic tumours, statistical difference was only seen with angiomin:CK19 ratio (figure 3D and its insert). No significant difference was otherwise seen with angiomin-like-1 and angiomin-like-2 (figure 3E and 3F).

Correlation of levels of angiomin with angiogenesis and other angiogenic markers

We also quantified the levels of VE-cadherin, PECAM-1 and vWF as an indicator of degree of angiogenesis in breast cancer, as we previously reported [21]. There was an increased level of VE-cadherin in breast tumour tissues (2.4 ± 0.64), compared with normal background tissues (1.9 ± 0.61), p = 0.7. Similarly, a high level of PECAM1 (CD31) in tumour tissues (275.0 ± 73.7), compared with normal background tissues (145.7 ± 30.0), p = 0.8. A Spearman correlation test between PECAM1, VE-cadherin and angiomin revealed a significant correlation between angiomin transcript and VE-cadherin as well as PECAM1 (r = 0.35 and r = 0.338, p < 0.05 respectively).




Figure 5
 The correlation between Angiotensin II receptor antagonist (A and D), angiotensin-like-1 (B and E) and angiotensin-like-2 (C and F) and Overall Survival (A-C) and disease free survival (D-F), using Kaplan-Meier survival analysis. Levels of angiotensin transcript were associated with the overall survival (A).

Angiotensin transcript levels were also significantly correlated with microvessel count (MVD) using anti-vWF ($r = 0.400$, $p < 0.05$). There was no significant correlation between levels of these factors and angiotensin-like-1 and angiotensin-like-2.

Implications of angiotensin with clinical outcome

Following a median 120 month followup, patients were divided into the following groups: those who remained disease free, who developed metastasis or local recurrence, and those who died of breast cancer related disease (excluding non-cancer related deaths). Angiotensin was seen at significantly higher levels in patients who developed metastatic disease compared with those who remained disease free ($p = 0.03$) (figure 4 left). Although tumours from patients with local recurrence and who died of breast cancer also had a higher level of angiotensin, the

difference was nonetheless not significant. Levels of both angiotensin like-1 and like-2 transcripts were lower in patients who had a poor clinical outcome, although the difference was not statistically significant (figure 4 middle and right).

We have divided patients into two groups, those with high levels of angiotensin and those with low levels of angiotensin, by using the prognostic index, NPI, as a general reference. Where a tumour had angiotensin level higher than the mean level of NPI-2 group (NPI 3.4–5.4, with moderate prognosis), it was assigned as high. Kaplan-Meier survival analysis has shown that patients bearing tumours with high levels of angiotensin were associated with a shorter overall survival (119.9 (105.2–134.7 95% CI) months vs 136.5 (124.5–148.4, 95%CI) months for patients with low angiotensin transcript, $p < 0.05$) (figure

Table 2: Levels of angiotenin transcript and their relationship with ER status. Shown are number of the respective transcript.

| | ER(-) | ER(+) | P value |
|---------------|--------------|--------------|-----------|
| Angiotenin | 87.2 ± 16.8 | 101.7 ± 29.7 | P = 0.67 |
| Angiotenin-L1 | 383.7 ± 86.0 | 167.6 ± 39.1 | P = 0.024 |
| Angiotenin-L2 | 174.0 ± 120 | 31.7 ± 8.6 | P = 0.24 |

5A). No significant correlation was seen between overall survival and Angiotenin-L1 and L2 (figure 5B and 5C). Similarly, patients bearing tumours with high levels of angiotenin were associated with a shorter disease free survival (108.6 (92.3–125.0 95% CI) months vs 134.3 (122.0–146.7, 95%CI) months for patients with low angiotenin transcript, however, the difference was not statistically significant, $p = 0.0634$) (figure 5D). No significant correlation was seen between disease free survival and Angiotenin-L1 and L2 (figures 5E and 5F). Using multivariate analysis of the following factors, nodal status, tumour grade, angiotenin, angiotenin-like-1 and angiotenin-like-2, we have found that nodal status ($p = 0.0185$) and angiotenin transcript ($p = 0.031$) were independent survival factors.

Furthermore, we have found that angiotenin-L1 was expressed at a significantly lower level in ER positive tumours compared with ER negative tumours (table 2). No significant difference was seen with angiotenin, angiotenin-L2 and ER status.

Discussion

Angiogenesis is the essential process in the development and spread of breast cancer, by providing blood supply to tumours and escape route for tumour cells to other part of the body. Here we report that angiotenin, a protein that regulate the motility and morphology of endothelial cells is highly expressed in human breast tissues and that its levels are associated with other angiogenic markers and with the clinical outcome in patients with breast cancer.

Angiotenin is a motility regulator of vascular endothelial cells and may be a molecule that links to breast cancer growth and spread by way of stimulating angiogenesis. The current study provides lines of evidence to support this possibility. First, angiotenin is highly expressed in aggressive tumours (grade 2 and 3 and tumours with nodal involvement) than in less aggressive tumours. Second, levels of angiotenin are correlated with levels of angiogenic markers. Third, significantly higher levels of angiotenin transcript are seen in patients with metastatic disease. These data, together with the report that angiotenin directly enhances angiogenesis *in vitro* and *in vivo*, suggest that angiotenin is linked to the angiogenic and aggressive nature of breast cancer. However, it would require additional work to verify if angiotenin and angio-

motin like proteins can be reliable surrogate markers for angiogenesis.

Although angiotenin-like-1 and -like-2 proteins are also expressed at higher levels in breast tumour tissues, the difference is not significant. There are no consistent patterns to suggest these two analogues of angiotenin are also linked to the aggressiveness of breast tumours, although both of the angiotenin related proteins are expressed in endothelial cells [23]. In fact, the levels of both analogues decrease in tumours that are associated with metastasis and mortality, which is in clear contrast with that of angiotenin. Currently there is no clear explanation to the seemingly different role between angiotenin and angiotenin like proteins. However, angiotenin-like-1 protein is known to be a tight junction related molecule and is highly located in tight junctions [10]. Recent years have seen significant advances in the understanding of the biology and role of tight junction in endothelium as defence mechanism in preventing blood borne cancer cells to escape. Tight junctions in endothelial cells may act as a 'sealing' structure to separate the blood circulation from tissue space [24]. Endothelial tight junctions act as a natural barrier for the vascular spread of cancer cell, by keeping the circulating cancer cells 'at bay-in the circulation'. In epithelial cells, as well as being a permeability barrier, tight junctions act as a strong cell adhesion mechanism and have a potential tumour suppressor role [25]. It has been reported that loss of certain tight junctional molecules, such as ZO-1, ZO-2, and occludin are frequently seen in clinical tumours and the loss of these TJ molecules is associated with the aggressiveness of tumours [26-29]. Thus, it is anticipated that angiotenin-like-1 protein, may act very different from angiotenin, potentially through their participation in tight junctions. Clearly, this is a fertile area to investigate in the future. The study has also indicated that ER negative tumours had a significantly higher level of angiotenin-L1 transcripts than in ER positive tumours. Although angiotenin-L2 transcripts were higher in ER negative tumours than in ER positive tumours, the difference was not statistically significant. In contrast, no difference was seen with angiotenin transcript between the two groups. This presents an interesting link between ER status and the angiotenin-L1 transcript. It has been established that ER negative tumours are associated with a poorer prognosis than ER positive tumours. It is possible that high levels of angiotenin-L1 in the ER

negative tumours contribute to this clinical link. Another potential link is that an intrinsic relationship between the expression of angiomin-1 and oestrogen in breast cancer may exist. Clearly, more experimental work is required here.

Conclusion

Although it is at early stage in the investigation of angiomin and its family into breast cancer, the current study and recent reports have clearly shown the important role of angiomin in angiogenesis and in the aggressive nature of breast tumours. These data suggest that angiomin is not only a highly useful prognostic indicator in breast cancer, it may also be a valid therapeutic target in cancer.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

WGJ was responsible for study design, data analysis, and preparation of MS, ADJ for pathological information and verification, GW for immunohistochemical work and image analysis, LH for study design and MS preparation, REM for clinical follow-up.

Acknowledgements

The authors would like to thank the Breast Cancer Campaign (BCC), Cancer Research Wales (CRW) and Cancer Research UK for their support.

References

- Folkman J: **Angiogenesis-retrospect and outlook.** *EXS* 1992, **61**:4-13.
- Bicknell R, Harris AL: **Mechanisms and therapeutic implications of angiogenesis.** *Curr Opin Oncol* 1996, **8**:60-5.
- Fidler IJ: **Critical determinants of cancer metastasis: rationale for therapy.** *Cancer Chemother Pharmacol* 1999, **43(Suppl)**:S3-S10.
- Cao G, O'Brien CD, Zhou Z, Sanders SM, Greenbaum JN, Makrigianakis A, DeLisser HM: **Involvement of human PECAM-1 in angiogenesis and in vitro endothelial cell migration.** *Am J Physiol Cell Physiol* 2002, **282**:C1181-90.
- Lampugnani MG, Resnati M, Raiteri M, Pigott R, Pisacane A, Houen G, Ruco LP, Dejana E: **A novel endothelial-specific membrane protein is a marker of cell-cell contacts.** *J Cell Biol* 1992, **118**:1511-22.
- Shih SC, Robinson GS, Perruzzi CA, Calvo A, Desai K, Green JE, Ali IU, Smith LE, Senger DR: **Molecular profiling of angiogenesis markers.** *Am J Pathol* 2002, **161**:35-41.
- Troyanovsky B, Levchenko T, Mansson G, Matvijenko O, Holmgren L: **Angiomin, an angiostatin binding protein that regulates endothelial cell migration and tube formation.** *J Cell Biol* 2001, **152**:1247-54.
- O'Reilly MS, Homgren L, Shing Y, Chen C, Rosenthal RA, Moses M, Lane WS, Cao Y, Sage EH, Folkman J: **Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma.** *Cell* 1994, **79**:315-28.
- Bratt A, Wilson WJ, Troyanovsky B, Aase K, Kessler R, Meir EGV, Holmgren L: **Angiomin belongs to a novel protein family with conserved coiled-coil and PDZ domains.** *Gene* 2002, **298**:69-77.
- Nishimura M, Kakizaki M, Ono Y, Morimoto K, Takeuchi M, Inoue Y, Imai T, Takai Y: **JEAP, a novel component of tight junctions in exocrine cells.** *J Biol Chem* 2002, **277**:5583-87.
- Shimono A, Behringer RR: **Angiomin regulates visceral endoderm movements during mouse embryogenesis.** *Curr Biol* 2003, **13**:613-7.
- Jiang WG, Puntis MCA, Hallett MB: **The molecular and cellular basis of cancer invasion and metastasis and its implications for treatment.** *Br J Surg* 1994, **81**:1576-90.
- Horak ER, Leek R, Klenk N, Lejeune S, Smith K, Stuart N, Greenall M, Stepniwska K, Harris AL: **Angiogenesis, assessed by platelet/endothelial cell adhesion molecule antibodies, as indicator of node metastases and survival in breast cancer.** *Lancet* 1992, **340**:1120-4.
- Aird WC, Jahroudi N, Weiler-Guettler H, Rayburn HB, Rosenberg RD: **Human von Willebrand factor gene sequences target expression to a subpopulation of endothelial cells in transgenic mice.** *Proc Natl Acad Sci USA* 1995, **92**:4567-71.
- Teo NB, Shoker BS, Jarvis C, Martin L, Sloane JP, Holcombe C: **Angiogenesis and invasive recurrence in ductal carcinoma in situ of the breast.** *Eur J Cancer* 2003, **39**:38-44.
- Salven P, Joensuu H, Heikkila P, Matikainen MT, Wasenius VM, Alanko A, Alitalo K: **Endothelial Tie growth factor receptor provides antigenic marker for assessment of breast cancer angiogenesis.** *Br J Cancer* 1996, **74**:69-72.
- Jiang WG, Douglas-Jones A, Mansel RE: **Level of expression of PPAR-gamma and its co-activator (PPAR-GCA) in human breast cancer.** *Int J Cancer* 2003, **106**:752-7.
- Nazarenko IA, Bhatnagar SK, Hohman RJ: **A closed tube format for amplification and detection of DNA based on energy transfer.** *Nucleic Acids Res* 1997, **25**:2516-21.
- Jiang WG, Watkins G, Fodstad O, Douglas-Jones A, Mokbel K, Mansel RE: **Differential expression of the CCN family members Cyr61 from CTGF and Nov in human breast cancer.** *Endocrine Rel Cancer* 2004, **11**:781-91.
- Jiang WG, Grimshaw D, Lane J, Martin TA, Parr C, Davies G, Laterra J, Mansel RE: **Retroviral hammerhead transgenes to cMET and HGF/SF inhibited growth of breast tumour, induced by fibroblasts.** *Clin Cancer Res* 2003, **9**:4274-81.
- Martin TA, Watkins G, Jiang WG: **Clinical value of VE-cadherin in assessing angiogenesis in human breast cancer.** *Histopathology* 2005, **46**:422-4.
- Martin TA, Parr C, Davies G, Matsumoto K, Nakamura T, Mansel RE, Jiang WG: **NK4 reduced HGF/SF and fibroblast induced breast tumour growth.** *Carcinogenesis* 2003, **24**:1317-25.
- Moreau J, Lord M, Boucher M, Belleau P, Fernandes MJ: **Protein diversity is generated within the motin family of proteins by alternative pre-mRNA splicing.** *Gene* 2005, **350**:137-48.
- Tsukita S, Furuse M, Iton M: **Multifunctional strands in tight junctions.** *Nat Rev Mol Cell Biol* 2001, **2**:285-293.
- Tobioka H, Sawada N, Zhong Y, Mori M: **Enhanced paracellular barrier function of rat mesothelial cells partially protects against cancer cell penetration.** *Br J Cancer* 1996, **74**:439-445.
- Satoh H, Zhong Y, Isomura H, Saitou M, Enomoto K, Sawada N, Mori M: **Localization of 7H6 tight junction-associated antigen along the cell border of vascular endothelial cells correlates with paracellular barrier function against ions, large molecules, and cancer cells.** *Exp Cell Res* 1996, **22**:269-74.
- Martin TA, Watkins G, Cunnick GH, Mansel RE, Jiang WG: **Reduced expression of tight junction molecules, ZO-1, ZO-2 and ZO-3, in breast cancer is associated with poor outcome.** *Eur J Cancer* 2004, **40**:2717-25.
- Hoover KB, Liao SY, Bryant P: **Loss of the tight junction MAGUK ZO-1 in breast cancer - Relationship to glandular differentiation and loss of heterozygosity.** *Am J Pathol* 1998, **153**:1767-1773.
- Kleeff J, Shi X, Bode HP, Hoover K, Shrikhande S, Bryant PJ, Korc M, Buchler MW, Friess H: **Altered expression and localization of the tight junction protein ZO-1 in primary and metastatic pancreatic cancer.** *Pancreas* 2001, **23**:259-65.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/6/16/prepub>