



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- 1 WHO. Coronavirus disease 2019 (COVID-19) situation report – 113. Geneva: World Health Organization, May 12, 2020. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200512-covid-19-sitrep-113.pdf?sfvrsn=feac3b6d\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200512-covid-19-sitrep-113.pdf?sfvrsn=feac3b6d_2) (accessed May 20, 2020).
- 2 Alho H, Dematteis M, Lembo D, Maremmanni I, Roncero C, Somaini L. Opioid-related deaths in Europe: Strategies for a comprehensive approach to address a major public health concern. *Int J Drug Policy* 2020; **76**: 102616.
- 3 Knopf A. NYC begins home methadone deliveries. Will it have lasting impact on the rules? *Filter*, April 30, 2020. <https://filtermag.org/new-york-home-methadone-delivery/> (accessed May 9, 2020).
- 4 FAQs: provision of methadone and buprenorphine for the treatment of opioid use disorder in the COVID-19 emergency. Rockyville: Substance Abuse and Mental Health Services, March 19, 2020. <https://www.samhsa.gov/sites/default/files/faqs-for-oud-prescribing-and-dispensing.pdf> (accessed May 20, 2020).
- 5 COVID-19: ACMD advice on proposed legislative changes to enable supply of controlled drugs during a pandemic. London: Advisory Council on the Misuse of Drugs, April 7, 2020. [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/878524/COVID-19\\_ACMD\\_advice\\_on\\_proposed\\_legislative\\_changes\\_to\\_enable\\_supply\\_of\\_controlled\\_drugs\\_during\\_a\\_pandemic1.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/878524/COVID-19_ACMD_advice_on_proposed_legislative_changes_to_enable_supply_of_controlled_drugs_during_a_pandemic1.pdf) (accessed May 20, 2020).
- 6 Ahamad K, Bach P, Brar R, et al. Risk mitigation in the context of dual public health emergencies. Vancouver, Canada: British Columbia Centre on Substance Use, 2020. <https://www.bccsu.ca/wp-content/uploads/2020/04/Risk-Mitigation-in-the-Context-of-Dual-Public-Health-Emergencies-v1.5.pdf> (accessed May 20, 2020).
- 7 Taha S, Maloney-Hall B, Buxton J. Lessons learned from the opioid crisis across the pillars of the Canadian drugs and substances strategy. *Subst Abuse Treat Prev Policy* 2019; **14**: 32.

## How might the NHS protect the mental health of health-care workers after the COVID-19 crisis?



Throughout the COVID-19 crisis, many health-care workers have worked long hours in high-pressured novel circumstances characterised by trauma and moral dilemmas.<sup>1</sup> Health-care workers have contended with the risk of infection, and by extension infecting their families, with outcomes seemingly worse for some, including black, Asian, and minority ethnic staff. Additionally, remote working is likely to have had its own challenges. Some staff will undoubtedly thrive in such circumstances, but we should now plan how to identify and support those who do not.

Post-trauma social support and stressors experienced during recovery are the risk factors most strongly predictive of longer-term mental health status.<sup>2</sup> Such stressors might be directly attributable to the crisis (eg, a colleague's death) or secondary (such as relationship or employment difficulties).<sup>3</sup>

Much evidence shows that supportive managers foster better mental health.<sup>4</sup> Furthermore, there are lessons from military practice<sup>5</sup> that can be applied to the post-COVID-19 health-care landscape. There are four key elements in an evidence-based staff National Health Service recovery plan. First, giving thanks, both written and verbally, which acknowledges the challenging work undertaken, can foster individual resilience.<sup>6</sup> This communication should include accurate up-to-date information about potential psychological difficulties and supports. Second, return-to-normal work interviews by supervisors who feel confident speaking about mental health. These interviews allow for a better understanding

of a staff member's experiences, while identifying secondary stressors in order to collaboratively design individualised recovery plans. Such discussions reduce sickness absence in other trauma-exposed occupations.<sup>7</sup> Third, active monitoring for anyone exposed to potentially traumatic events, particularly individuals considered to be at higher risk of developing mental health problems.<sup>8</sup> Although such monitoring is another function of good management, evidence supports proactive case finding, which proved successful after the London bombings.<sup>9</sup> An anonymous online self-check tool might encourage honest and meaningful responses while providing automated tailored feedback. Fourth, group discussions to help staff to develop a meaningful narrative that reduces risks of harm. Schwartz rounds, a structured forum for clinical and non-clinical staff to discuss emotional and social aspects of work, are one such evidence-based model.

Successful recovery planning<sup>10</sup> should minimise the onset of mental illness while maximising the opportunity for psychological growth.<sup>1</sup> Proactive managers should follow the evidence, which is both legally required and what staff deserve.

NG runs a psychological health consultancy that provides resilience training for a wide range of organisations, including a few UK National Health Service (NHS) teams. The work was supported by the UK National Institute for Health Research (NIHR) Health Protection Research Unit in Emergency Preparedness and Response at King's College London (London, UK), in partnership with Public Health England and in collaboration with the University of East Anglia (Norwich, UK) and Newcastle University (Newcastle upon Tyne, UK). The views expressed are those of the authors and not necessarily those of the NHS, NIHR, UK Department of Health and Social Care, or Public Health England. All other authors declare no competing interests.



Mark Thomas/Science Photo Library

Published Online  
May 28, 2020  
[https://doi.org/10.1016/S2215-0366\(20\)30224-8](https://doi.org/10.1016/S2215-0366(20)30224-8)

\*Neil Greenberg, Samantha K Brooks, Simon Wessely,  
Derek K Tracy  
neil.greenberg@kcl.ac.uk

The Health Protection Research Unit (NG, SKB, SW) and Cognition, Schizophrenia, and Imaging Laboratory, Department of Psychosis Studies (DK), Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London SE5 8AF, UK; and Oxleas NHS Foundation Trust, London, UK (DK)

- 1 Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ* 2020; **368**: m1211.
- 2 Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull* 2003; **129**: 52–73.
- 3 Tempest EL, English National Study on Flooding and Health Study Group, Carter B, Beck CR, Rubin GJ. Secondary stressors after flooding: a cross-sectional analysis. *Eur J Public Health* 2017; **27**: 1042–47.
- 4 Brooks SK, Dunn R, Amlot R, Rubin GJ, Greenberg N. A systematic, thematic review of social and occupational factors associated with psychological outcomes in healthcare employees during an infectious disease outbreak. *J Occ Environ Med* 2018; **60**: 248–57.
- 5 Carter N. Land post-operational stress management (POSM). April, 2014. [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/428920/LFSO\\_3209\\_Redacted.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/428920/LFSO_3209_Redacted.pdf) (accessed May 20, 2020).
- 6 McCanlies EC, Gu JK, Andrew ME, Violanti JM. The effect of social support, gratitude, resilience and satisfaction with life on depressive symptoms among police officers following Hurricane Katrina. *Int J Soc Psychiatry* 2018; **64**: 63–72.
- 7 Milligan-Saville JS, Tan L, Gayed A, et al. Workplace mental health training for managers and its effect on sick leave in employees: a cluster randomised controlled trial. *Lancet Psychiatry* 2017; **4**: 850–58.
- 8 National Institute for Health and Care Excellence. Post-traumatic stress disorder. NICE guideline [NG116]. Dec 5, 2018. <https://www.nice.org.uk/guidance/ng116> (accessed May 20, 2020).
- 9 Brewin CR, Fuchkan N, Huntley Z, et al. Outreach and screening following the 2005 London bombings: usage and outcomes. *Psychol Med* 2010; **40**: 2049–57.
- 10 Royal College of Psychiatrists. NHS staff recovery plan post COVID-19 (outbreak 1). Royal College of Psychiatrists. <https://www.rcpsych.ac.uk/about-us/responding-to-covid-19/responding-to-covid-19-guidance-for-clinicians/wellbeing-and-support/nhs-staff-recovery-plan> (accessed May 20, 2020).