

# Association between maternal pre-delivery body mass index and offspring overweight/obesity at 1 and 2 years of age among residents of a suburb in Taiwan

Hsien-Kuan Liu, Chien-Yi Wu, Yung-Ning Yang, Pei-Ling Wu, Zong-Rong He, San-Nan Yang and Shu-Leei Tey

Department of Pediatrics, E-Da Hospital, I-Shou University, Kaohsiung, Taiwan

## ABSTRACT

**Background:** Overweight and obesity among children can cause metabolic syndrome in adulthood and are a significant public health issue. Some studies suggest that maternal pre-pregnancy body mass index (BMI) and excessive gestational weight gain during pregnancy are associated with overweight and obesity in offspring. However, it is difficult to collect information on accurate pre-pregnancy BMI and pregnancy weight gain for women living in areas where medical resources are scarce. Maternal pre-delivery BMI might be predictive of the risk of overweight and obesity among offspring of pregnant mothers living in suburban areas.

**Methods:** We retrospectively collected data on term neonates with appropriate weights for their gestational age born between April 2013 and October 2015. We excluded neonates with major congenital anomalies or diseases and incomplete data. Mothers with systemic diseases or drug abuse were also excluded.

Offspring body weights and heights at 1- and 2-years-old were recorded. Maternal pre-delivery BMI was divided into following groups:  $<25$ ,  $25\text{--}29.9$ , and  $\geq 30$   $\text{kg/m}^2$ .

**Results:** We included 261 mother-child pairs in this study. The BMIs of the offspring differed significantly among the three maternal pre-delivery BMI groups at the age of 2 years ( $15.18 \pm 1.04$ ,  $15.83 \pm 1.28$ , and  $16.29 \pm 1.61$   $\text{kg/m}^2$ ,  $p < 0.001$ , respectively). After adjusting for potential cofounders possibly affecting weight using multivariate linear regression, the children's BMIs (adjusted 95% CI: 0.71 [0.31–1.11];  $p = 0.001$ ) and BMI percentiles (adjusted 95% CI 15.80 [7.32–24.28];  $p < 0.001$ ) at the age of 2 years were significantly higher in those born to mothers with pre-delivery BMIs of  $25\text{--}29.9$   $\text{kg/m}^2$  compared to mothers with pre-delivery BMIs  $<25$   $\text{kg/m}^2$ . Maternal pre-delivery BMI  $\geq 30$   $\text{kg/m}^2$  was significantly associated with increased BMIs (adjusted 95% CI: 1.17 [0.72–1.63];  $p < 0.001$ ) and BMI percentiles (adjusted 95% CI: 23.48 [13.87–33.09];  $p < 0.001$ ) in their children. A maternal pre-delivery BMI of  $27.16$   $\text{kg/m}^2$  was the optimal cut-off for predicting offspring overweight/obesity at the age of 2 years.

**Discussion:** Our results indicate that the maternal pre-delivery BMI was significantly associated with offspring BMI and weight gain at the age of 2 years. A maternal pre-delivery BMI of  $27.16$   $\text{kg/m}^2$  might be a useful predictor for estimating the risk of overweight or obesity in offspring at the age of 2 years.

Submitted 8 June 2018  
Accepted 18 January 2019  
Published 25 February 2019

Corresponding author  
Shu-Leei Tey,  
ed104496@edah.org.tw

Academic editor  
José Derraik

Additional Information and  
Declarations can be found on  
page 12

DOI 10.7717/peerj.6473

© Copyright  
2019 Liu et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Global Health, Gynecology and Obstetrics, Pediatrics, Public Health, Women's Health

**Keywords** Obesity, Maternal pre-delivery BMI, Overweight, Offspring, Body weight

## INTRODUCTION

Overweight and obesity among children is a public health issue that can result in adult metabolic syndrome (*De Onis, Blössner & Borghi, 2010; Kim, Lee & Lim, 2017*).

There are many factors associated with overweight/obesity in children, including early childhood lifestyle and sedentary behavior, dietary patterns, sleep patterns and durations, and parental body mass index (BMI) (*Reilly et al., 2005; Dev et al., 2013*).

The Developmental Origins of Health and Disease (DOHaD) hypothesis, which proposed by *Barker (2007)*, expounded the high correlation between maternal environment and their offspring. For the past few years, DOHaD hypothesis had been broadly adapted in subsequent studies concerning early life environment influences the later onset of obesity (*Wadhwa et al., 2009; Kappil, Wright & Sanders, 2016*). This concept has enticed an increasing number of researchers to study the associations between maternal obesity and their offspring's obesity (*Guo et al., 2015; Wrotniak et al., 2008*). During the gestational period, the maternal metabolic state can affect both maternal health and fetal growth, even predisposing the offspring to the development of metabolic disorders (*Catalano & Ehrenberg, 2006*). As there may be a connection between maternal BMI and offspring health, this is a potentially useful means of predicting and preventing child overweight and obesity.

The Institute of Medicine (IOM) of the National Academies proposed a recommendation of appropriate gestational weight gain (GWG) based on maternal pre-pregnancy BMI (*Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines, 2009*). Currently, studies have revealed that high pre-pregnancy BMI and excessive GWG were associated with overweight/obesity in their offspring (*Li et al., 2013; Schack-Nielsen et al., 2010*). In Taiwan, nearly all pregnant women deliver in a metropolitan hospital or medical provision clinic. Therefore, maternal pre-pregnancy BMI or GWG can be accessed easily during the pre-natal visits. However, for those pregnant women living in suburbs where medical resources are scarce and prenatal care was inadequate, it was difficult to collect accurate pre-pregnancy BMI and GWG (*Kisuule et al., 2013; Islam et al., 2017; Debessai et al., 2016; Fobelets et al., 2015*). For these women, and their future children, it became a priority to identify an alternate predictor of overweight/obesity in their offspring.

Pre-delivery BMI, which can be obtained just prior to delivery, is an easier measurement to obtain from pregnant women who lived in the suburbs. If a high pre-pregnancy BMI or excessive GWG could cause adverse pregnancy outcomes and increased offspring body weight later in life, then we considered that a high pre-delivery BMI might more or less have a similar effect. To answer this question, we designed a study to compare the association between maternal pre-delivery BMI and the BMI of the offspring at the ages of 1 and 2 years. Our goal for this study, therefore, was to find a new parameter to predict the risk of overweight and obesity in the offspring of the pregnant mothers in the suburbs,

and try to find out a cut-off point of maternal pre-delivery BMI for predicting overweight and obesity in their offspring.

## MATERIALS AND METHODS

### Study designs and subjects

E-Da Hospital is located in the Yanchao District (a suburb area in Kaohsiung), occupies 6,300 ha of land and has a population of about 29,000 people. Patients in the E-Da Hospital mostly reside in the Yanchao District and the adjacent regions. Owing to the vast territory with few residents, this district is a medical resource-limited area.

This was a retrospective cohort study. Both pregnant women and their children's health care records from E-Da Hospital were collected. We collected 1,753 records of both the delivering mothers and their infants between April 2013 and October 2015. The children's health records included the birth-related information (date of birth, sex, gestational weeks at time of birth, birth weight, birth height, Apgar score, among others) and the infancy health examinations during the first 24 months. Neonates who were preterm, post-term, small for GA (birth weight < 10th percentile), large for GA (birth weight > 90th percentile), with presence of major congenital anomalies, congenital disease, or incomplete data were excluded from this survey.

Pregnant women's health records included general information (age, occupation, education, number of pregnancies, smoking habits, among others), any history of diseases, all clinical measurements (height, weight, gynecological examinations, ultrasonography, gestational diabetes mellitus (GDM) screening test, and other lab tests), complications experienced during pregnancy, and pregnancy outcomes (delivery modes, labor complications, placenta weight). The pre-delivery BMI was calculated using the weight and height recorded upon admission for delivery. BMIs were calculated by dividing the weight in kilograms by the square of the height in meters. The pre-delivery BMI was categorized into three groups: <25, 25–29.9, and  $\geq 30$  kg/m<sup>2</sup>. Mothers who were diabetic, or had pregnancy-induced hypertension, a history of illegal substance abuse, human immunodeficiency virus infected, hyper- or hypothyroidism, or incomplete medical records were all excluded.

This study included the information and clinical measurements of 261 mother-child pairs, after excluding premature infants, post-term infants, congenital anomaly, small for gestational age + large for gestational age, pregnancy-induced hypertension, GDM, GDM + pregnancy-induced hypertension, maternal substance abuse, human immunodeficiency virus infected mother, maternal hyper/hypothyroidism, and incomplete data of neonate and mother.

We obtained approval from the E-Da Hospital's Institutional Review Board (EMRP02107N). All patient information was de-identified before analysis.

### Assessment of infant body weight

All infants had a preventative health visit for vaccinations at 1 and 2 years of age according to the Taiwan vaccination programs. Body weight, body height and health status were recorded at every visit. No infants had acute illnesses at the vaccination health visits.

The body weights were measured with light clothing using a digital scale, and the body heights were measured with bare feet using a recumbent length stadiometer. Weight gain was calculated as the difference between the weight at birth and weight at 1 or 2 year of age. BMIs were calculated by dividing the weight in kilograms by the square of the height in meters. The definition of children obesity is according to new growth charts for Taiwanese children (*Chen & Chang, 2010*). Those with missing data on body weight at 1 or 2 year of age were excluded.


### Statistical analysis

Data were analyzed using SPSS statistical software version 20 (IBM Corp., Armonk, NY, USA). Proportions are presented for categorical variables and means  $\pm$  standard deviations are presented for continuous variables. The Chi-square test (for categorical variables) and Kruskal–Wallis test were used for comparisons among three groups of maternal pre-delivery BMIs. We conducted a multivariate linear regression to adjust the potential variables associated with outcomes including sex, mode of delivery, and maternal pre-delivery BMI. We report 95% confidence intervals (CI) with *p*-values. Covariate variables were selected from among the significant predictors ( $p < 0.05$ ) between the three groups according to maternal pre-delivery BMI, as well as by the use of clinical judgment. Statistical significance was defined as a *p*-value  $< 0.05$ .

Receiver operating characteristic (ROC) curves were drawn to determine the optimal threshold for predicting overweight and obesity in children at 2 years of age depending on maternal pre-delivery BMI. The optimal threshold value was used as the cut-off point to determine the sensitivity, specificity, and negative predictive values (NPVs) of maternal pre-delivery BMI to detect overweight and obesity at 2 years of age.

## RESULTS

This study included the information and clinical measurements of 261 mother-child pairs, after excluding premature infants ( $n = 314$ ), post-term infants ( $n = 7$ ), congenital anomaly ( $n = 9$ ), small for gestational age + large for gestational age ( $n = 274$ ), loss to follow up after discharge ( $n = 84$ ), pregnancy-induced hypertension ( $n = 119$ ), GDM ( $n = 85$ ), GDM + pregnancy-induced hypertension ( $n = 61$ ), maternal substance abuse ( $n = 3$ ), human immunodeficiency virus infected mother ( $n = 3$ ), maternal hyper/hypothyroidism ( $n = 18$ ), and maternal incomplete data ( $n = 7$ ) (*Fig. 1*). Among the 261 children, 137 (52.5%) were boys. The mean gestational age was  $38.97 \pm 0.99$  weeks, and the mean birth weight was  $3.18 \pm 0.28$  kg. The participants were stratified into three groups according to the maternal pre-delivery BMI ( $<25$ ,  $25\text{--}29.9$ , and  $\geq 30$  kg/m<sup>2</sup>). The maternal and offspring demographic characteristics are presented in *Table 1*. The overall overweight and obesity rates in children aged 1 and 2 years were 10.0% (26/261) and 9.2% (24/261), respectively. The rate of overweight/obesity showed no significant difference at 1 year of age in each maternal pre-delivery BMI category (5/85 (2.5%), 11/96 (11.5%), and 10/63 (15.9%), respectively;  $p = 0.142$ ). However, the rate differed



**Figure 1** Participants' selection after excluding neonatal and maternal factors from 1,753 neonates. Full-size [DOI: 10.7717/peerj.6473/fig-1](https://doi.org/10.7717/peerj.6473/fig-1)

significantly at 2 years of age (2/79 (2.5%), 10/85 (11.8%), and 12/57 (21.1%), respectively;  $p = 0.003$ ). The infants birth weights ( $3.08 \pm 0.25$ ,  $3.24 \pm 0.28$ , and  $3.22 \pm 0.27$  kg, respectively;  $p = 0.001$ ) and birth BMIs ( $12.29 \pm 0.90$ ,  $12.62 \pm 1.02$ , and  $12.76 \pm 1.00$  kg/m<sup>2</sup>, respectively;  $p = 0.002$ ) were significantly higher in children whose maternal pre-delivery BMIs were  $\geq 25$  kg/m<sup>2</sup> than in those whose maternal pre-delivery BMIs were  $< 25$  kg/m<sup>2</sup>. Our results also revealed that 56 infants were delivered by Cesarean section (21.5%). The rate of Cesarean section differed significantly in each maternal pre-delivery BMI category (10/90 (11.1%), 27/105 (25.7%), and 19/66 (28.8%), respectively;  $p = 0.011$ ). The overweight and obesity rates at 1 year showed no significant association among the three groups ( $p = 0.142$ ), whereas it differed significantly at 2 years ( $p = 0.003$ ).

**Table 1** Baseline maternal and offspring demographic characteristics according to pre-delivery maternal body mass index (BMI).

	BMI			<i>p</i> *
	25 kg/m <sup>2</sup> ( <i>n</i> = 90)	25–29.9 kg/m <sup>2</sup> ( <i>n</i> = 105)	≥ 30 kg/m <sup>2</sup> ( <i>n</i> = 66)	
<b>Maternal demographics</b>				
Maternal age				0.257
<35 years old	74 (82.2%)	77 (73.3%)	48 (72.7%)	
≥ 35 years old	16 (17.8%)	28 (26.7%)	18 (27.3%)	
Placenta weight	653.70 ± 151.95	663.23 ± 135.42	654.67 ± 116.97	0.549
Parity				0.402
Primipara	48 (53.3%)	66 (62.9%)	38 (57.6%)	
Multipara	42 (46.7%)	39 (37.1%)	28 (42.4%)	
<b>Offspring demographics</b>				
Gestational age (days)	39.02 ± 0.92	39.01 ± 0.99	38.81 ± 1.07	0.461
Birth weight (kg)	3.08 ± 0.25	3.24 ± 0.28	3.22 ± 0.27	0.001
Birth BMI (kg/m <sup>2</sup> )	12.29 ± 0.90	12.62 ± 1.02	12.76 ± 1.00	0.002
Sex				0.293
Boy	46 (51.1%)	51 (48.6%)	40 (60.6%)	
Girl	44 (48.9%)	54 (51.4%)	26 (39.4%)	
Mode of delivery				0.011
Vaginal	80 (88.9%)	78 (74.3%)	47 (71.2%)	
Cesarean section	10 (11.1%)	27 (25.7%)	19 (28.8%)	
Apgar score				
1 min	7.97 ± 0.18	7.92 ± 0.49	7.82 ± 0.89	0.891
5 min	8.98 ± 0.15	8.95 ± 0.35	8.81 ± 0.58	0.824
Data collected				
1-year-old (days) <sup>a</sup>	380.10 ± 31.43	377.51 ± 23.08	377.93 ± 15.67	0.222
2-year-old (days) <sup>b</sup>	820.32 ± 48.43	816.68 ± 51.11	813.59 ± 45.99	0.319
1-year-old body type <sup>a</sup>				
Normal	80 (94.1%)	85 (88.5%)	53 (84.1%)	0.142
Overweight + Obesity	5 (5.9%)	11 (11.5%)	10 (15.9%)	
2-year-old body type <sup>b</sup>				
Normal	77 (97.5%)	75 (88.2%)	45 (78.9%)	0.003
Overweight + Obesity	2 (2.5%)	10 (11.8%)	12 (21.1%)	

**Notes:**

Data are presented as means ± standard deviations or as numbers (proportion).

\* *p*-values were analyzed using the Kruskal–Wallis (for continuous variables) and Chi-square tests (for categorical variables).

<sup>a</sup> Total 244 participants were enrolled in the analysis at 1 year of age.

<sup>b</sup> Total 221 participants were enrolled in the analysis at 2 years of age.

## Offspring weight index among the three maternal pre-delivery BMI groups

The anthropometric measurements of the offspring were assessed at ages 1 and 2 years. The children's BMIs at age 1 year were significantly different among the three maternal pre-delivery BMI groups ( $16.37 \pm 1.15$ ,  $16.67 \pm 1.11$ , and  $17.19 \pm 1.61$  kg/m<sup>2</sup>,

**Table 2** Offspring weight index at 1 and 2 years of age according to maternal pre-delivery body mass index (BMI).

	Maternal pre-delivery BMI			<i>p</i> *
	<25 kg/m <sup>2</sup> ( <i>n</i> = 85)	25–29.9 kg/m <sup>2</sup> ( <i>n</i> = 96)	≥ 30 kg/m <sup>2</sup> ( <i>n</i> = 63)	
<b>1 year old<sup>a</sup></b>				
BW (kg)	9.31 ± 0.91	9.55 ± 0.87	9.83 ± 1.03	0.007
BH (cm)	75.36 ± 2.74	75.66 ± 2.58	75.60 ± 2.35	0.861
BMI (kg/m <sup>2</sup> )	16.37 ± 1.15	16.67 ± 1.11	17.19 ± 1.61	0.005
BMI percentile	45.94 ± 25.68	52.00 ± 25.54	60.07 ± 25.76	0.006
Weight gain <sup>b</sup> (kg)	6.21 ± 0.87	6.32 ± 0.78	6.60 ± 0.99	0.044
<b>2 years old<sup>c</sup></b>				
BW (kg)	12.04 ± 1.19	12.31 ± 1.39	12.93 ± 1.62	0.004
BH (cm)	88.80 ± 3.12	88.25 ± 3.08	89.12 ± 3.16	0.461
BMI (kg/m <sup>2</sup> )	15.18 ± 1.04	15.83 ± 1.28	16.29 ± 1.61	<0.001
BMI percentile	35.24 ± 24.74	49.86 ± 27.72	56.68 ± 29.97	<0.001
Weight gain <sup>d</sup> (kg)	8.90 ± 1.13	9.12 ± 1.29	9.72 ± 1.57	0.004

**Notes:**

BW, body weight; BH, body height; BMI, body mass index.

\* *p*-value was analyzed by Kruskal–Wallis test.

<sup>a</sup> Total 244 participants were enrolled in the analysis at 1 year of age.

<sup>b</sup> The weight difference between birth and 1 year of age.

<sup>c</sup> Total 221 participants were enrolled in the analysis at 2 years of age.

<sup>d</sup> The weight difference between birth and 2 years of age.

respectively;  $p = 0.005$ ). The children BMI percentile ( $45.94 \pm 25.68$ ,  $52.00 \pm 25.54$ , and  $60.07 \pm 25.76$ , respectively) and weight gain ( $6.21 \pm 0.87$ ,  $6.32 \pm 0.78$ , and  $6.60 \pm 0.99$ , respectively) at age 1 years showed a significant association among the three groups ( $p = 0.006$  and  $p = 0.044$ ). Similarly, the children's BMIs at age 2 years were significantly different among the three maternal pre-delivery BMI groups ( $15.18 \pm 1.04$ ,  $15.83 \pm 1.28$ , and  $16.29 \pm 1.61$  kg/m<sup>2</sup>, respectively;  $p < 0.001$ ). Additionally, the children BMI percentile ( $35.24 \pm 24.74$ ,  $49.86 \pm 27.72$ , and  $56.68 \pm 29.97$ , respectively) and weight gain ( $8.80 \pm 1.13$ ,  $9.12 \pm 1.29$ , and  $9.72 \pm 1.57$ , respectively) at age 2 years showed a significant association among the three groups ( $p < 0.001$  and  $p = 0.004$ ) (Table 2).

### Multivariable linear regression of factors associated with offspring body weight parameters at ages 1 and 2 years of age

We adjusted for the potential confounding factors that could have affected the children's body weights at 1 and 2 years of age (sex, mode of delivery, and maternal pre-delivery BMI). The resulting multivariate linear regression analyses are presented in Table 3. At 1 year of age, BMIs ( $p < 0.001$ ), BMI percentiles ( $p = 0.001$ ), and weight gain ( $p = 0.015$ ) were significantly higher in children born to mothers with pre-delivery BMIs of  $\geq 30$  kg/m<sup>2</sup> compared to those born to mothers with pre-delivery BMIs of  $<25$  kg/m<sup>2</sup>. At 2 years of age, BMIs ( $p = 0.001$ ) and BMI percentiles ( $p < 0.001$ ) were significantly higher in children born to mothers with pre-delivery BMIs of 25–29.9 kg/m<sup>2</sup>

**Table 3** Multivariable linear regression for factors associated with offspring body weight parameters at 1 and 2 years of age.

<b>1-year-old (n = 244)</b>				
<b>BMI</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	0.32 [−0.01 to 0.65]	0.053	0.28 [−0.05 to 0.60]	0.092
Mode of delivery	0.11 [−0.29 to 0.51]	0.578	−0.00 [−0.40 to 0.39]	0.987
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	0.17 [−0.39 to 0.29]	0.769	0.31 [−0.07 to 0.68]	0.109
≥ 30 kg/m <sup>2</sup>	0.19 [0.29–1.03]	<0.001	0.80 [0.37–1.23]	<0.001
<b>BMI percentile</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	−2.55 [−9.14 to 4.05]	0.448	−3.34 [−9.88 to 3.17]	0.312
Mode of delivery	3.50 [−4.49 to 11.49]	0.389	1.32 [−6.72 to 9.35]	0.747
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	3.43 [−6.71 to 6.80]	0.989	5.80 [−1.81 to 13.42]	0.134
≥ 30 kg/m <sup>2</sup>	3.76 [3.50–18.32]	0.004	14.35 [5.72–22.97]	0.001
<b>Weight gain</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	0.36 [0.14–0.58]	0.001	0.34 [0.12–0.55]	0.003
Mode of delivery	0.06 [−0.21 to 0.33]	0.675	0.01 [−0.26 to 0.28]	0.943
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	0.12 [−0.29 to 0.17]	0.608	0.11 [−0.14 to 0.37]	0.379
≥ 30 kg/m <sup>2</sup>	0.13 [0.08–0.59]	0.010	0.36 [0.07–0.65]	0.015
<b>2-year-old (n = 221)</b>				
<b>BMI</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	0.20 [−0.16 to 0.56]	0.283	0.14 [−0.20 to 0.49]	0.413
Mode of delivery	−0.12 [−0.56 to 0.33]	0.606	−0.32 [−0.74 to 0.11]	0.145
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	0.19 [−0.19 to 0.55]	0.337	0.71 [0.31–1.11]	0.001
≥ 30 kg/m <sup>2</sup>	0.21 [0.37–1.18]	<0.001	1.17 [0.72–1.63]	<0.001
<b>BMI percentile</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	−2.21 [−9.81 to 5.39]	0.567	−3.19 [−10.45 to 4.08]	0.388
Mode of delivery	−2.58 [−11.82 to 6.66]	0.583	−6.80 [−15.75 to 2.15]	0.136
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	5.59 [−2.12 to 13.30]	0.154	15.80 [7.32–24.28]	<0.001
≥ 30 kg/m <sup>2</sup>	13.96 [5.42–22.49]	0.001	23.48 [13.87–33.09]	<0.001
<b>Weight gain</b>	<b>Crude</b>		<b>Adjusted*</b>	
Variables	B (95% CI)	P	B (95% CI)	p
Sex	0.27 [−0.09 to 0.62]	0.143	0.21 [−0.14 to 0.55]	0.239



**Table 3 (continued).**

Weight gain	Crude		Adjusted*	
Mode of delivery	-0.42 [-0.85 to 0.02]	0.060	-0.55 [-0.97 to -0.12]	0.012
Maternal pre-delivery BMI <sup>†</sup>				
25–29.9 kg/m <sup>2</sup>	-0.12 [-0.49 to 0.24]	0.503	0.32 [-0.09 to 0.72]	0.124
$\geq 30$ kg/m <sup>2</sup>	0.72 [0.32–1.12]	0.001	0.92 [0.46–1.38]	<0.001

**Notes:**

B, unstandardized regression coefficient; CI, confidence interval; BMI, body mass index.

\* Co-variables included in the linear regression analysis: sex, mode of delivery, and maternal pre-delivery body mass index

<sup>†</sup> Maternal pre-delivery BMI was grouped as follows: <25, 25–29.9, and  $\geq 30$  kg/m<sup>2</sup>; the reference category is <25 kg/m<sup>2</sup>.

**Table 4 Risk factors associated to children overweight/obesity at 2 years of age.**

Variable	OR	95% CI	p-value
Maternal pre-delivery BMI	1.206	[1.10–1.33]	<0.001
Maternal pre-delivery BMI (>27.16/<27.16 kg/m <sup>2</sup> ) <sup>a</sup>	7.312	[2.41–22.20]	<0.001
Children sex (male/female)	1.304	[0.55–3.08]	0.545
Mode of delivery (Cesarean section/Vaginal delivery)	0.738	[0.24–2.28]	0.597
Overweight/obesity at 1 year of age	2.680	[0.88–8.13]	0.082

**Notes:**

OR, Odd ratio; CI, confidence interval; BMI, body mass index.


<sup>a</sup> We transformed the continuous variables of maternal pre-delivery BMI into categorical variables by the cutoff point.

compared to those born to mothers with pre-delivery BMIs of <25 kg/m<sup>2</sup>.

Similarly, maternal pre-delivery BMI  $\geq 30$  kg/m<sup>2</sup> was also significantly associated with an increase in the children's BMIs, BMI percentiles, and weight gain (all  $p < 0.001$ ) at 2 years of age (Table 3).

### Diagnostic value of maternal pre-delivery BMI for predicting their offspring's overweight/obesity

The relative contributor of explanatory variables to offspring overweight/obesity at 2 year of age were analyzed (Table 4). Among these variables, pre-delivery BMI remained a significant risk factors that would contribute offspring overweight/obesity at 2 year of age (OR: 1.206, 95% CI [1.10–1.33];  $p < 0.001$ ). Children overweight/obesity at 1 year of age was another risk factor (OR: 2.680) but this factor was not significant ( $p = 0.082$ ). In addition, the ROC curves of maternal pre-delivery BMI for predicting offspring overweight/obesity at 2 years is shown in Fig. 2. The area under the ROC curve (AUROC) was 0.750% (95% CI [0.654–0.847]);  $p < 0.001$ ) for diagnosing the offspring with overweight/obesity at 2 years, according to new growth charts for Taiwanese children (Chen & Chang, 2010). The optimal cut-off point (by Youden index) for maternal pre-delivery BMI to detect their offspring's overweight/obesity at 2 years was 27.16 kg/m<sup>2</sup>, with a NPV of 96.69%, a sensitivity of 83.3%, and a specificity of 59.4%. We also transformed the continuous variable of maternal pre-delivery BMI into categorical variable by this cutoff point. The odd ratios of maternal pre-delivery BMI >27.16 kg/m<sup>2</sup> was 7.312% (95%CI [2.41–22.20];  $p < 0.001$ ) (Table 4).



**Figure 2** ROC curve of maternal pre-delivery body mass index for predicting offspring obesity at 2 years of age. [Full-size](#) DOI: 10.7717/peerj.6473/fig-2

## DISCUSSION

Many studies have shown that both higher maternal pre-pregnancy BMI and greater GWG are associated with increased BMIs in early childhood (*Li et al., 2013; Schack-Nielsen et al., 2010*). There are few studies that have evaluated the correlation between maternal pre-delivery BMI and offspring obesity. This study indicated that maternal pre-delivery BMI might also predict the risk of children developing overweight/obesity later in life. We found that the maternal pre-delivery BMI was linearly associated with the BMI of children aged 1–2 years even after adjustment for birth weight, sex, and mode of delivery. We also noticed that the risk of childhood overweight was highest among the mothers whose pre-delivery BMI was  $\geq 30$  kg/m<sup>2</sup>.

In recent years, some studies have reported that greater maternal pre-pregnancy BMIs may have an impact on the childhood body weights in their offspring (*Yu et al., 2013; Xiong et al., 2016*). Our results showed similar results to these studies. We found that women with higher pre-delivery BMIs were associated with higher risks for Cesarean section, higher birth weight babies, and an increased risk of offspring with overweight/obesity in the first 2 years of life. There are a couple of advantages with using maternal pre-delivery BMI. First, maternal pre-delivery BMI can be obtained easily before delivery and can be applied in all medical facilities, even in areas with limited medical resources. Second, recall errors and biases associated with pre-pregnancy BMI could be avoided by using the pre-delivery BMI.

There are a number of possible mechanisms responsible for the association between maternal pre-delivery BMI and overweight in their offspring. The DOHaD hypothesis, also called the “Barker hypothesis,” proposed by *Barker (2007)*,

*Armitage, Poston & Taylor (2008)* and *Zheng et al. (2014)*, could explain this relationship. According to this hypothesis, the energy excess in the maternal diet causes the accumulation of excess adipose tissue, which might modify DNA methylation and gene expression in their offspring (*Zheng et al., 2014; Morales et al., 2014*). This methylation process could result in offspring adiposity (*Godfrey et al., 2011*). This theory indicates that high maternal pre-delivery BMI might play an important role in offspring overweight and might contribute to the overweight epidemic among infants and children.

Historically, there has been an emphasis on promoting sufficient weight gain during pregnancy in an effort to reduce low-birth weight deliveries and adverse perinatal outcomes. In 2009, the IOM published new recommendations for weight gain during pregnancy (*Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines, 2009*). However, obesity rather than insufficient weight gain during pregnancy became a more common problem nowadays. Between 1997 and 2007, approximately 46% of pregnant women in the US gained more weight than the IOM recommended. Additional studies showed that maternal pre-pregnancy obesity and excessive GWG were associated with greater risks of future offspring obesity. Nevertheless, the recommended GWG might not be suitable for reproductive-aged women in different countries due to variations in body physique or body composition, which differed from race and ethnicity (*Wagner & Heyward, 2000*). *Yang et al. (2015)* investigated the recommended GWG for Chinese women as recommended by the IOM, and they found that the GWG suggested by the IOM might not be helpful for Chinese women. In our study, we found that maternal pre-delivery BMI  $27.16 \text{ kg/m}^2$  was a reliable cut-off value for predicting offspring overweight/obesity at 2 years of age in Chinese reproductive-aged women with an odd ratio of 7.312. High NPV of this cut-off point can offer us a guide to educate the mother to control their BMI before delivery. If the mother can control their pre-delivery BMI less than  $27.16 \text{ kg/m}^2$ , there will be a 96.69% chance that their offspring will not have overweight/obesity at 2 years of age. In addition, maternal pre-delivery BMI is easier to assess than GWG for pregnant women residing in the suburbs of Taiwan.

There are several strengths in our study. The pre-delivery BMI can be measured easily and accurately despite the different scales of medical facilities and can even be easily measured in medical resource-limited hospitals. Furthermore, this measurement would not be affected by irregular or delayed prenatal care. Thus, potential misreporting and participation bias with respect to the pre-pregnant BMI can be avoided. There are also some limitations in our study. First, this was a retrospective study, so it is possible that selection bias could have influenced our results. However, this issue was minimized by stratifying the groups according to maternal pre-delivery BMIs and adjusting the possible confounding factors in this study. Second, although we created multivariable models to adjust for the potential confounders, the sample size of this study was still small and the participants were from a single hospital located in southern Taiwan. Thus, the study result may not be generalizable to other countries or other ethnicities. Third, there are still a few possible confounders that we were not able to adjust for, including childhood nutrition and physical activity. Our sample population was mainly from a suburb where

the lifestyle factors are similar, so these effects are expected to be small. Last, a large number of cases were excluded due to incomplete data. However, the basal characteristics of maternal and offspring we excluded was not significantly different than those enrolling participants, suggesting that this limitation may not have introduced a significant selection bias (Supplementary Table).

Our hospital is located in a suburb area in Taiwan and, also, is the largest hospital nearby. Some infants would receive their regular health examination and vaccine administration in the local clinic instead of our hospital after birth. Therefore, anthropometric measurement and healthy status of participants could not be obtained due to loss following up. With these incomplete data, the result might not be generalized. A further large-scale study may minimize this bias.

## CONCLUSION

In conclusion, this study result indicates that maternal pre-delivery BMI might be a new parameter to predict the risk of overweight and obesity among the offspring of pregnant mothers in a suburb of Taiwan. Furthermore, the maternal pre-delivery BMI value of 27.16 kg/m<sup>2</sup> may suggest a useful predictor when estimating the offspring's risk of overweight or obesity at age 2 years. These findings indicate that more attention needs to be paid to infants of mothers with higher pre-delivery BMIs, especially those  $\geq 27.16$  kg/m<sup>2</sup>.

## ACKNOWLEDGEMENTS

The authors are grateful to Chun-Hua Yang and Tzu-Shan Chen, who were consultants on statistical methods in this study.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The authors received no funding for this work.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Hsien-Kuan Liu conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Chien-Yi Wu performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Yung-Ning Yang performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Pei-Ling Wu analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

- Zong-Rong He analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- San-Nan Yang conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Shu-Leei Tey conceived and designed the experiments, performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

## Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

We obtained approval from the E-Da Hospital's Institutional Review Board (EMRP02107N). All patient information was de-identified before analysis.

## Data Availability

The following information was supplied regarding data availability:

The raw data are available in the [Supplemental Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6473#supplemental-information>.

## REFERENCES

- Armitage JA, Poston L, Taylor PD. 2008.** Developmental origins of obesity and the metabolic syndrome: the role of maternal obesity. *Frontiers of Hormone Research* **36**:73–84  
DOI [10.1159/000115355](https://doi.org/10.1159/000115355).
- Barker DJ. 2007.** The origins of the developmental origins theory. *Journal of Internal Medicine* **261**:412–417.
- Catalano PM, Ehrenberg HM. 2006.** The short- and long-term implications of maternal obesity on the mother and her offspring. *BJOG* **113**(10):1126–1133 DOI [10.1111/j.1471-0528.2006.00989.x](https://doi.org/10.1111/j.1471-0528.2006.00989.x).
- Chen W, Chang M-H. 2010.** New growth charts for Taiwanese children and adolescents based on World Health Organization standards and health-related physical fitness. *Pediatrics & Neonatology* **51**(2):69–79 DOI [10.1016/S1875-9572\(10\)60014-9](https://doi.org/10.1016/S1875-9572(10)60014-9).
- Debessai Y, Costanian C, Roy M, El-Sayed M, Tamim H. 2016.** Inadequate prenatal care use among Canadian mothers: findings from the maternity experiences survey. *Journal of Perinatology* **36**(6):420–426 DOI [10.1038/jp.2015.218](https://doi.org/10.1038/jp.2015.218).
- De Onis M, Blössner M, Borghi E. 2010.** Global prevalence and trends of overweight and obesity among preschool children. *American Journal of Clinical Nutrition* **92**(5):1257–1264  
DOI [10.3945/ajcn.2010.29786](https://doi.org/10.3945/ajcn.2010.29786).
- Dev DA, McBride BA, Fiese BH, Jones BL, Cho H, Behalf of the Strong Kids R Team. 2013.** Risk factors for overweight/obesity in preschool children: an ecological approach. *Childhood Obesity* **9**(5):399–408 DOI [10.1089/chi.2012.0150](https://doi.org/10.1089/chi.2012.0150).
- Fobelets M, Beeckman K, Hoogewys A, Embo M, Buyl R, Putman K. 2015.** Predictors of late initiation for prenatal care in a metropolitan region in Belgium. A cohort study. *Public Health* **129**(6):648–654 DOI [10.1016/j.puhe.2015.03.008](https://doi.org/10.1016/j.puhe.2015.03.008).

- Godfrey KM, Sheppard A, Gluckman PD, Lillycrop KA, Burdge GC, McLean C, Rodford J, Slater-Jefferies JL, Garratt E, Crozier SR, Emerald BS, Gale CR, Inskip HM, Cooper C, Hanson MA. 2011. Epigenetic gene promoter methylation at birth is associated with child's later adiposity. *Diabetes* 60(5):1528–1534 DOI 10.2337/db10-0979.
- Guo L, Liu J, Ye R, Liu J, Zhuang Z, Ren A. 2015. Gestational weight gain and overweight in children aged 3–6 years. *Journal of Epidemiology* 25(8):536–543 DOI 10.2188/jea.JE20140149.
- Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. 2009. In: Rasmussen KM, Yaktine AL, eds. *Weight Gain During Pregnancy: Reexamining the Guidelines*. Washington D.C.: National Academies Press, 254–256.
- Islam MJ, Broidy L, Baird K, Mazerolle P. 2017. Exploring the associations between intimate partner violence victimization during pregnancy and delayed entry into prenatal care: evidence from a population-based study in Bangladesh. *Midwifery* 47:43–52 DOI 10.1016/j.midw.2017.02.002.
- Kappil M, Wright RO, Sanders AP. 2016. Developmental origins of common disease: epigenetic contributions to obesity. *Annual Review of Genomics and Human Genetics* 17(1):177–192 DOI 10.1146/annurev-genom-090314-050057.
- Kim J, Lee I, Lim S. 2017. Overweight or obesity in children aged 0 to 6 and the risk of adult metabolic syndrome: a systematic review and meta-analysis. *Journal of Clinical Nursing* 26(23–24):3869–3880 DOI 10.1111/jocn.13802.
- Kisuule I, Kaye DK, Najjuka F, Ssematimba SK, Arinda A, Nakitende G, Otim L. 2013. Timing and reasons for coming late for the first antenatal care visit by pregnant women at Mulago hospital, Kampala Uganda. *BMC Pregnancy and Childbirth* 13(1):121 DOI 10.1186/1471-2393-13-121.
- Li N, Liu E, Guo J, Pan L, Li B, Wang P, Liu J, Wang Y, Liu G, Hu G. 2013. Maternal prepregnancy body mass index and gestational weight gain on offspring overweight in early infancy. *PLOS ONE* 8(10):e77809 DOI 10.1371/journal.pone.0077809.
- Morales E, Groom A, Lawlor DA, Relton CL. 2014. DNA methylation signatures in cord blood associated with maternal gestational weight gain: results from the ALSPAC cohort. *BMC Research Notes* 7(1):278 DOI 10.1186/1756-0500-7-278.
- Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, Steer C, Sherriff A, Avon Longitudinal Study of Parents and Children Study Team. 2005. Early life risk factors for obesity in childhood: cohort study. *BMJ* 330:1357 DOI 10.1136/bmj.38470.670903.E0.
- Schack-Nielsen L, Michaelsen KF, Gamborg M, Mortensen EL, Sørensen TI. 2010. Gestational weight gain in relation to offspring body mass index and obesity from infancy through adulthood. *International Journal of Obesity* 34(1):67–74 DOI 10.1038/ijo.2009.206.
- Wadhwa PD, Buss C, Entringer S, Swanson JM. 2009. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Seminars in Reproductive Medicine* 27(5):358–368 DOI 10.1055/s-0029-1237424.
- Wagner DR, Heyward VH. 2000. Measures of body composition in blacks and whites: a comparative review. *American Journal of Clinical Nutrition* 71(6):1392–1402 DOI 10.1093/ajcn/71.6.1392.
- Wrotniak BH, Shults J, Butts S, Stettler N. 2008. Gestational weight gain and risk of overweight in the offspring at age 7 y in a multicenter, multiethnic cohort study. *American Journal of Clinical Nutrition* 87(6):1818–1824 DOI 10.1093/ajcn/87.6.1818.
- Xiong C, Zhou A, Cao Z, Zhang Y, Qiu L, Yao C, Wang Y, Zhang B. 2016. Association of pre-pregnancy body mass index, gestational weight gain with cesarean section in term deliveries of China. *Scientific Reports* 6(1):37168 DOI 10.1038/srep37168.

- Yang S, Peng A, Wei S, Wu J, Zhao J, Zhang Y, Wang J, Lu Y, Yu Y, Zhang B. 2015.** Pre-pregnancy body mass index, gestational weight gain, and birth weight: a cohort study in China. *PLOS ONE* **10(6)**:e0130101 DOI [10.1371/journal.pone.0130101](https://doi.org/10.1371/journal.pone.0130101).
- Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. 2013.** Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis. *PLOS ONE* **8(4)**:e61627 DOI [10.1371/journal.pone.0061627](https://doi.org/10.1371/journal.pone.0061627).
- Zheng J, Xiao X, Zhang Q, Yu M. 2014.** DNA methylation: the pivotal interaction between early-life nutrition and glucose metabolism in later life. *British Journal of Nutrition* **112(11)**:1850–1857 DOI [10.1017/S0007114514002827](https://doi.org/10.1017/S0007114514002827).