



Review

Purinergic Signalling in the Cochlea

Srdjan M. Vlajkovic ^{1,*}  and Peter R. Thorne ^{1,2}

¹ Department of Physiology and The Eisdell Moore Centre, Faculty of Medical and Health Sciences, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

² Section of Audiology, School of Population Health, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

* Correspondence: s.vlajkovic@auckland.ac.nz; Tel.: +64-9-9239782

Abstract: The mammalian cochlea is the sensory organ of hearing with a delicate, highly organised structure that supports unique operating mechanisms. ATP release from the secretory tissues of the cochlear lateral wall (stria vascularis) triggers numerous physiological responses by activating P2 receptors in sensory, supporting and neural tissues. Two families of P2 receptors, ATP-gated ion channels (P2X receptors) and G protein-coupled P2Y receptors, activate intracellular signalling pathways that regulate cochlear development, homeostasis, sensory transduction, auditory neurotransmission and response to stress. Of particular interest is a purinergic hearing adaptation, which reflects the critical role of the P2X₂ receptor in adaptive cochlear response to elevated sound levels. Other P2 receptors are involved in the maturation of neural processes and frequency selectivity refinement in the developing cochlea. Extracellular ATP signalling is regulated by a family of surface-located enzymes collectively known as “ectonucleotidases” that hydrolyse ATP to adenosine. Adenosine is a constitutive cell metabolite with an established role in tissue protection and regeneration. The differential activation of A₁ and A_{2A} adenosine receptors defines the cochlear response to injury caused by oxidative stress, inflammation, and activation of apoptotic pathways. A₁ receptor agonism, A_{2A} receptor antagonism, and increasing adenosine levels in cochlear fluids all represent promising therapeutic tools for cochlear rescue from injury and prevention of hearing loss.



Citation: Vlajkovic, S.M.; Thorne, P.R. Purinergic Signalling in the Cochlea.

Int. J. Mol. Sci. **2022**, *23*, 14874.

<https://doi.org/10.3390/ijms232314874>

ijms232314874

Academic Editor: Rashid Giniatullin

Received: 27 October 2022

Accepted: 25 November 2022

Published: 28 November 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: ATP; P2X receptors; P2Y receptors; ectonucleotidases; adenosine; adenosine receptors; cochlea; hearing loss

1. Introduction

The cochlea of the inner ear provides exquisite sensitivity and frequency selectivity to sound detection in the mammalian auditory system [1]. It contains three fluid-filled compartments: scala media filled with endolymph (high K⁺ and low Na⁺ concentration), scala vestibuli and scala tympani filled with perilymph (high Na⁺ and low K⁺). In the mammalian cochlea, the sensory epithelium of the organ of Corti contains sensory hair cells surrounded by various supporting cells, including Deiters’, Hensen’s and pillar cells [2] (Figure 1). Inner hair cells (IHC) are responsible for sound transduction (conversion of acoustic stimuli to electrical signals to the brain), whilst outer hair cells (OHC) are known as “cochlear amplifiers”, providing enhanced frequency selectivity and sensitivity at low sound levels [3]. Primary auditory neurons located in the spiral ganglion provide afferent innervation mostly to the IHC, and the efferent innervation from the brainstem structures (superior olivary complex) modifies the activity of the OHC [2]. Cochlear supporting cells are essential constituents of an epithelial network that regulates the unique ionic composition of cochlear extracellular fluids [4]. The spiral ligament and stria vascularis form the lateral wall of the cochlea and contribute to the maintenance of ion homeostasis, generation of the endocochlear potential (a positive electrical potential confined to scala media), and regulation of cochlear metabolism [2].




Figure 1. Inner ear anatomy. The mammalian inner ear contains two sensory organs, the organ of hearing (cochlea) and the organ of balance (vestibular organ). The cochlea comprises three fluid-filled compartments. Scala vestibuli and scala tympani contain Na^+ -rich perilymph, whereas scala media contains K^+ -rich endolymph. The lateral wall of the cochlea comprises the spiral ligament (SL) made up of fibrocytes, providing structural support to the secretory tissues of the stria vascularis (SV). The organ of Corti is the central structure for sensory transduction in the cochlea. It is covered by the tectorial membrane (TM) and contains two types of sensory cells, inner hair cells (IHC) and outer hair cells (OHC). These sensory cells are surrounded by supporting cells, such as Deiters' cells (DC), Hensen's cells (HC) and pillar cells (PC). The inner sulcus cells (ISC) are epithelial cells that form the medial border of the organ of Corti. Created with [BioRender.com](https://www.biorender.com).

Amongst the multiple signaling pathways that regulate cochlear function, significant attention has been focused on purinergic regulation, particularly its protective adaptation to acoustic stimuli [5,6]. Adenosine 5'-triphosphate (ATP) release from marginal cells in the stria vascularis [7] exerts a homeostatic regulatory mechanism promoting K^+ efflux from scala media and reduction of the endocochlear potential, thus reducing cochlear sensitivity to loud sound [8,9]. Purinergic signalling in the cochlea can affect several essential functions, from ion homeostasis to active mechanical amplification by the outer hair cells [10].

The release of ATP and activation of purinergic receptors in cochlear sensory, supporting, secretory and neural tissues play a critical role in cochlear physiology and pathophysiology [11,12]. The ionotropic P2X and the metabotropic P2Y receptors are widely distributed in cochlear tissues [6,13]. To date, seven subtypes of the P2X family (P2X₁₋₇)

and eight subtypes of the P2Y family (P2Y_{1,2,4,6,11–14}) have been cloned and functionally characterised [12]. P2X receptors provide direct routes for Ca²⁺ entry into the cell, whereas P2Y receptors either activate phospholipase C (PLC) and release intracellular Ca²⁺ or alter cAMP levels [12,14]. Intracellular Ca²⁺ seems to be the principal second messenger in ATP-mediated signaling [15].

P2 receptors have been implicated in the sensory cell sensitivity adjustment, regulation of cochlear electrochemical homeostasis, blood flow, and cochlear development [6]. P2 receptor signalling is regulated by ectonucleotidases, which hydrolyse extracellular ATP to adenosine [16,17]. Adenosine activates protective mechanisms in the cochlea by acting on adenosine (P1) receptors expressed on cochlear sensory and non-sensory cells [13,18].

A substantial body of data suggests that extracellular ATP is a critical player in processes that guide the activation of the mammalian auditory system in response to sound during ear development. Spontaneous activity in afferent auditory pathways appears before the onset of acoustically evoked signal processing [19,20]. During this critical developmental period, IHC fire bursts of action potentials that stimulate synaptic maturation and refinement of auditory circuits to establish a precise tonotopic organization in the developing cochlea and receptive fields in central auditory nuclei [19,20]. Endogenous ATP release in the organ of Corti triggers this activity by activating diverse P2 receptors in supporting cells and the IHC. ATP initiates Ca²⁺ waves spreading through the developing cochlea, causing a K⁺ efflux from these cells through Ca²⁺-activated K⁺ channels and the leak K⁺ channels [19].

P2 receptor signalling contributes to neuronal firing by increasing the excitability of auditory nerve fibers, spiral ganglion and auditory brainstem neurons [19,20]. ATP release during cochlear development is central to this process. The neonatal cochlea has a transient epithelium (the Kölliker's organ) which spontaneously secretes ATP in rhythmical bursts that activate P2 receptors on the IHC [21]. This, in turn, synchronises the output of adjacent IHC, which may refine tonotopic maps in the brain auditory nuclei.

Several lines of evidence also support the notion that ATP-dependent mechanisms safeguard against the harmful effects of acoustic overstimulation in the adult cochlea [8]. Purinergic-mediated adaptation of the cochlear sensory structures to loud sound decreases the activity of the sensory cells and protects them and their afferent synapses from sustained noise exposure and acoustic injury [22].

This review highlights the role of purinergic P2 (ATP) and P1 (adenosine) receptors in regulating cochlear function in health and disease, focusing on the recent developments in the field.

2. Methodology

A literature search of relevant publications from 2000 to 2022 was performed using PubMed and Google Scholar. We have also used reference lists from the relevant papers. Following Boolean search logic, we used the following keywords: (cochlea) and (ATP OR P2 receptors OR adenosine receptors OR purinergic signalling OR ATP release OR ectonucleotidases OR adenosine transport); (ATP OR P2 receptors) and (cochlear development); (vestibular system) and (P2 receptors OR purinergic signalling). The search results were then examined according to their relevance to this review. Only English language publications were included in this review.

3. ATP Release in the Cochlea

Various mechanisms are involved in ATP release from cells, including exocytosis, transporter proteins, and ion channel-mediated release [23]. Tissue injury can also lead to lytic release from damaged cells, rapidly raising the extracellular ATP concentrations [23].

The principal conduits for ATP release in the cochlea during development and adulthood are integral membrane proteins from the gap junction family known as pannexin and connexin hemichannels [24]. The opening of these channels allows the efflux of ATP down its concentration gradient and release into the endolymph and perilymph [24–26].

Connexin and pannexin hemichannels are essential for the cellular release of ATP and regulate purinergic receptor activation [25]. In turn, P2 receptor activation by ATP can amplify purinergic signaling through a positive feedback loop via inositol 1,4,5-trisphosphate (IP₃), giving rise to the concept of ATP-induced ATP release [27]. IP₃, activated by P2Y receptors, mobilises intracellular Ca²⁺ which opens pannexin hemichannels, enabling the tide of calcium waves across epithelial cells [28].

Several P2X receptor subtypes (P2X₂, P2X₄, P2X₇) can also activate the pannexin 1 (Panx1) channel to carry through large molecules involved in initiating inflammatory responses and apoptotic cell death [28,29].

Pannexins have different electrophysiological and pharmacological properties than connexins [30]. Three isoforms of the pannexin hemichannel (Panx1, 2 and 3) have been identified in the cochlea, mostly in the supporting cells, the spiral limbus, and the lateral wall [31]. Panx1 is a dominant pannexin isoform in the cochlea [24]. *Panx1* deletion abolishes cochlear ATP release and ATP-mediated K⁺ cycling essential for maintaining the endocochlear potential (EP) in the mammalian cochlea [32,33]. EP is a driving force for hair cell transduction and is essential for normal hearing. *Panx1* deficiency causes moderate-to-severe progressive hearing loss and the progressive loss of sensory hair cells by activating the caspase-3 apoptotic pathway [33]. Interestingly, the deletion of predominant connexin isoforms in the cochlea, connexin 26 (Cx26) and connexin 30 (Cx30), does not reduce ATP release under physiological conditions, suggesting that Panx1 channels dominate ATP release in the cochlea [24] (Figure 2).




Figure 2. The elements of the purinergic signalling system in the cochlea. Pannexin hemichannels (e.g., Panx1) are the principal conduits of ATP release in the cochlea. Extracellular ATP activates ATP-gated ion channels (P2X receptors) and G protein-coupled receptors (P2Y receptors). ATP is hydrolysed to adenosine 5'-diphosphate (ADP) and adenosine 5'-monophosphate (AMP) by surface-located ecto-nucleoside triphosphate diphosphohydrolases (NTPDases), whereas AMP is dephosphorylated to adenosine by ecto-5'-nucleotidase (CD73). Adenosine activates four types of adenosine receptors (AR: A₁, A_{2A}, A_{2B} and A₃). Bi-directional adenosine transporters mediate cellular uptake of adenosine. In the intracellular space, adenosine is hydrolysed to inosine by adenosine deaminase (Ada) or phosphorylated to AMP by adenosine kinase (Adk) and ATP by ATP synthase (AS). Created with [BioRender.com](https://www.biorender.com).

4. P2 Receptors in the Cochlea

4.1. P2X Receptors in the Cochlea

All P2X receptors are transiently expressed in the developing mammalian cochlea, but their immunoexpression is limited in the adult cochlea [6,13].

P2X₁ receptor (P2X₁R) is transiently expressed in the otic capsule, spiral limbus, epithelial cells of the Reissner's membrane and spiral ganglion neurons (SGN) during early postnatal development in rats but is absent after hearing onset [34].

P2X₂ receptor (P2X₂R) is the predominant P2X subtype expressed in the epithelial cells lining the cochlear partition of the rat and mouse cochlea, including the inner and outer sensory hair cells, supporting Deiters' cells in the organ of Corti and Reissner's membrane that separates endolymph in scala media from perilymph in scala vestibuli [35,36].

P2X₃ receptor (P2X₃R) is also transiently expressed in the developing cochlear tissues. Its expression is detected in perinatal and juvenile C57BL/6 mice from embryonic day 18 (E18) to postnatal day 6 (P6) in the SGN, sensory hair cells, and peripheral neurites projecting towards the sensory hair cells [37]. The P2X₃R immunoexpression in the peripheral neurites and the hair cells diminishes by P6 and is absent after the onset of hearing (P11 to P17) [37]. The transient expression of these receptors, particularly P2X₃R, follows a precise spatio-temporal profile suggesting the role of this receptor in synaptic pruning [37]. The synaptic pruning likely involves transiently expressed heterodimeric P2X_{2/3} receptors inhibiting neurotrophic support for SGN during synaptic reorganisation [38].

In the adult guinea pig cochlea, P2X₄R is expressed in the outer hair cells [39] and spiral ligament capillaries [40], the latter suggesting that extracellular ATP regulates blood flow in the cochlear lateral wall by activating P2X₄R in endothelial cells.

There is little evidence for the immunoexpression of P2X₅ and P2X₆ receptors in the mammalian cochlea [41].

All P2X receptors are expressed in the developing rat spiral ganglion, but only P2X₂ and P2X₇ are sustained into adulthood [6]. P2X₂ was immunolocalised to the postsynaptic membranes of Type I and Type II SGN [35,42]. Strong P2X₇ expression was also observed in the olivocochlear efferent fibres innervating the sensory hair cells from E18 through to adulthood, suggesting a role for these receptors in auditory neurotransmission [43]. Recent evidence, however, indicates that P2X₇R is immunolocalised to peripheral glial cells rather than afferent neurons in the auditory nerve of small rodents [44]. Physiological responses in the peripheral glia are characterised by classical features of P2X₇R activation, including the formation of ion- and macromolecule-permeable pores. These properties suggest that P2X₇R could contribute to glial-mediated inflammatory processes under pathologic conditions, potentially contributing to auditory neuropathy and hearing loss [44].

With sustained elevated sound levels, ATP is released into the endolymph and ATP-gated ion channels (P2X receptors) on the epithelial cells lining the endolymphatic compartment shunt K⁺ outside scala media, reducing the driving force for sensory transduction and contributing to protective hearing adaptation [8]. The P2X₂ receptor subunit on Reissner's membrane and other epithelial tissues lining the cochlear endolymphatic compartment is essential for this shunt conductance evoked by noise exposure [45].

A significant component of temporary hearing loss that develops with sustained exposure to moderate noise has been attributed to the release of ATP in the cochlea, activating the P2X₂ receptor in scala media [8]. This purinergic hearing adaptation enables the cochlea to detect sounds in background noise and may also protect the cochlea from permanent damage and hearing loss (Figure 3). In the study that established this hearing adaptation mechanism [8], the role of the P2X₂ receptor was determined in *P2rx2* knockout and age-matched wildtype mice using auditory brainstem responses (ABR) measured during sustained noise exposure. The knockout mice failed to exhibit the temporary threshold shift (TTS) observed in wildtype mice after exposure to sustained moderate noise levels (85 dB SPL). This finding was a paradigm shift in understanding the mechanism of TTS, as the study demonstrated that the P2X₂R almost exclusively mediated TTS under moderate noise conditions. In the absence of TTS, the *P2rx2* KO mice exhibited normal hearing

sensitivity at a young age but developed accelerated age-related hearing loss compared to wildtype mice. In addition, *P2rx2* KO mice demonstrated increased vulnerability to sustained loud sound at higher noise levels (95 dB SPL), developing significantly higher permanent threshold shifts (PTS) than wildtype mice [8,22,46].




Figure 3. Purinergic hearing adaptation. Exposure to moderate noise levels induces ATP release from the stria vascularis (SV). ATP activates $P2X_2$ receptors in the tissues lining the endolymphatic compartment (e.g., Reissner's membrane, sensory and supporting cells). The activation of $P2X_2$ receptors mediates K^+ shunt conductance from scala media, thus reducing the endocochlear potential (EP), the driving force for sensory transduction. Reduced EP results in decreased sensitivity of the cochlea to sound stimulation, protecting the ear from permanent damage and hearing loss. The role of $P2X_2$ receptors in hearing adaptation was demonstrated using *P2rx2* knockout (KO) mice. Exposure to moderate noise levels caused temporary threshold shift (TTS) in wildtype mice but not in *P2rx2* KO mice. Without purinergic hearing adaptation, *P2rx2* KO mice showed accelerated age-related hearing loss (ARHL) and elevated permanent threshold shift (PTS) in response to higher noise levels. Created with [BioRender.com](https://www.biorender.com).

The study by Housley et al. [8] complements a report by Yan et al. [47], which demonstrated that the absence of cochlear $P2X_2$ R signaling in two Chinese families due to a dominant-negative mutation (conversion of 178G>T (p.V60L)) at chromosome 12, removed intrinsic purinergic otoprotection and induced the autosomal-dominant progressive hearing loss designated as DFNA41. Members of the family with DFNA41 with a history of noise exposure demonstrated enhanced high-frequency hearing loss, previously modelled in *P2rx2*-null mice [8]. Another knock-in mouse model based on human p.V60L mutation exhibited hearing loss at 21 days of age and progressed to deafness by six months [48]. Abnormal morphology of the inner hair cells and ribbon synapses was observed in those mice [48]. Other studies demonstrated that mutations in human $P2X_2$ R could cause hearing loss without completely disrupting channel function [49], highlighting the important role of these receptors in hearing protection.

Hearing protection is also regulated by ATP-evoked Ca^{2+} signaling in the supporting cells of the organ of Corti [50]. Extracellular ATP controls the intercellular Ca^{2+} waves,

which travel through supporting cells regulating the repair mechanisms following acoustic trauma [15]. Lahne and Gale [51] showed that two distinct Ca^{2+} waves are triggered during cochlear damage in organotypic tissue cultures, both elicited by extracellular ATP. A slower Ca^{2+} wave in Deiters' cells was mediated by P2Y receptors and Ca^{2+} release from IP_3 -sensitive stores. The faster Ca^{2+} wave propagated through sensory hair cells and was likely mediated by the P2X₄ receptor [51]. Periodic Ca^{2+} waves have been linked to gene regulation and likely play a crucial role in developing the organ of Corti and the acquisition of hearing [52].

Liu et al. [53] have shown that type II unmyelinated cochlear afferents that innervate OHC are activated when OHC are damaged. This response depends on both P2X and P2Y receptors and is activated by ATP released from nearby supporting cells in response to hair cell damage. Type II afferents may thus represent cochlear nociceptors, and their activation may reflect evasion of further injury to the inner ear after irreversible damage to OHC [53].

4.2. P2Y Receptors in the Cochlea

The immunoexpression of the P2Y receptors (P2Y₁, P2Y₂, P2Y₄, P2Y₆, and P2Y₁₂) was demonstrated in the developing (E16–P28) and the adult rat cochlea (P49–P56) by Huang et al. [54]. In the sensory epithelium, the earliest expression of P2Y receptors (P2Y₂ and P2Y₄) was observed in the greater epithelial ridge at embryonic day 18 (E18), and this expression pattern was retained at birth (P0). At P0, the P2Y₆ receptor was immunolocalised to the immature IHC and OHC [54], whereas at the early postnatal age (P6–P12), the P2Y₆ receptor localisation resembled the immunoexpression in adults. In adult rats, P2Y₆ becomes the predominant subtype in the IHC, and both P2Y₁ and P2Y₄ receptors are immunolocalised to the OHC [54]. The predominant P2Y receptor in the supporting cells is the P2Y₂ receptor immunolocalised to Deiters' cells, Hensen's cells, pillar cells and Claudius' cells, supporting cells that have a role in intercellular communication [51,54].

P2Y receptors (P2Y₂, P2Y₄, P2Y₆, and P2Y₁₂) were detected in the SGN at birth except for P2Y₁, which was expressed later in the postnatal age, and this expression pattern was retained until adulthood [54]. In the lateral wall tissues, P2Y receptor expression was first observed at early postnatal age (P0–P6), with P2Y₁ and P2Y₂ as the predominant subtypes. Following the onset of hearing, the P2Y expression in the lateral wall shifted to P2Y₂ and P2Y₄ [54]. This distribution of purinergic P2Y receptors suggests their multiple roles in cochlear development, maintaining cochlear homeostasis, and regulating sound transduction and neurotransmission [6].

Interestingly, pharmacological inhibition of the P2Y₁ receptor dramatically reduces spontaneous activity in the developing cochlea [55]. Spontaneous bursts of electrical activity in the developing auditory system arise within the cochlea before hearing onset to promote the maturation of auditory neurons. ATP release from supporting cells and activation of P2Y₁ receptors invokes coordinated excitation of neurons that will process similar sound frequencies [55]. The role of P2Y receptors in cochlear development is discussed further in the next section.

5. ATP and Cochlear Development

From birth to hearing onset, the auditory system relies on intrinsic mechanisms that elicit the coordinated firing of neurons processing similar sound frequencies in the adult cochlea [20,21]. ATP is released from the supporting cells in the greater epithelial ridge (GER) of the neonatal rat cochlea, a transient non-sensory cell population that disappears during postnatal cochlear maturation. Lysosomes are the organelles involved in ATP storage and release from GER cells [56]. ATP is also stored in the stria vascularis, and ATP-containing vesicles in marginal cells have also been identified as lysosomes [57]. ATP release from marginal cells and GER involves Ca^{2+} -dependent lysosomal exocytosis [56,57]. Lysosomal exocytosis of ATP is coupled to the P2Y₂ receptor in marginal cells via the P2Y₂R-phospholipase C- IP_3 pathway [58]. Connexin hemichannels mediate the release of ATP responsible for Ca^{2+} wave propagation in the developing mouse cochlea [58].

Before the onset of hearing, immature IHC and primary auditory neurons in the spiral ganglion experience sound-independent activity, which is believed to be important in retaining and refining neural connections in the absence of sound [21]. This activity originates in a group of transient epithelial supporting cells forming Kölliker's organ (as part of GER), which is only present during cochlear development. ATP released through connexin hemichannels may activate P2 receptors in both Kölliker's organ and the adjacent IHC, leading to the generation of electrical activity in the auditory system [59,60]. It was proposed that inner border cells have a major role in generating spontaneous morphological activity within Kölliker's organ [60,61].

More recently, it was proposed that Ca^{2+} waves in the supporting GER cells cause increased and synchronized Ca^{2+} activity in the OHC via ATP-induced activation of P2X_3 receptors [62]. This synchronization is required for the refinement of their immature afferent innervation. Ceriani et al. proposed that the correct maturation of the afferent connectivity of OHCs requires sound-independent Ca^{2+} signalling from sensory and non-sensory cells, which was P2X_3 receptor-dependent [62].

At all developmental stages, pharmacological inhibition of the P2Y_1 receptor dramatically reduces spontaneous activity in sensory and non-sensory cells [55]. The frequency of the spontaneous activity increases progressively during the postnatal prehearing period but remains dependent on the $\text{P2Y}_1\text{R}$ located on the cochlear supporting cells. When $\text{P2Y}_1\text{R}$ is activated, it triggers the release of Ca^{2+} in supporting cells and the activation of Ca^{2+} -dependent potassium channels. The efflux of K^+ to the extracellular space activates the sensory hair cells, but it also causes supporting cells to shrink due to water egress [63]. Conversely, when $\text{P2Y}_1\text{R}$ is inhibited, this causes the supporting cells to swell, entrapping potassium ions near the sensory cells. Bursts of electrical activity are thus controlled by the rhythmic swelling and shrinking of supporting cells mediated by the $\text{P2Y}_1\text{R}$ [63].

Other P2 receptors may also be involved in cochlear maturation. Outer sulcus cells (OSC) adjacent to the lateral wall of the cochlea may have a role in maintaining an adequate K^+ concentration in the cochlear endolymph in response to variable intensities of auditory stimulation [64]. Temporal changes in $\text{P2Y}_4\text{R}$ expression during OSC development likely contribute to the endolymphatic ion composition required to generate the endocochlear potential through the activation of K^+ channels [64].

In addition, P2X_3 receptors may be required for the differentiation of Type I SGN and their branch refinement [65]. Synaptic refinement and strengthening are activity-dependent processes aiding the orderly arrangement of cochleotopic maps in the central auditory system. The maturation of auditory brainstem circuits is guided by the electrical activity of the IHC in the developing cochlea, and modulated by paracrine ATP signalling [66]. Using slice recordings before hearing onset and in vivo recordings after hearing onset, Jovanovic et al. showed that cell-specific purinergic modulation follows a precise tonotopic pattern in the ventral cochlear nucleus in gerbils, which was mediated by the heterologous $\text{P2X}_{2/3}$ receptor [66].

6. Ectonucleotidases in the Cochlea

Ectonucleotidases are a large family of surface-located enzymes that hydrolyze extracellular nucleotides (ATP, UTP) to their respective nucleosides and thus regulate complex extracellular P2 receptor signalling pathways in mammalian tissues [16,67]. The best-characterised ectonucleotidase family in the mammalian cochlea is the ecto-nucleoside triphosphate diphosphohydrolase (NTPDase) family [18]. All enzymes from this family (NTPDase1-8) are expressed in the adult rat cochlea (Figure 2). The spatial and temporal expression of NTPDases in various cell types in the vasculature, sensory and neural tissues in the cochlea impacts multiple physiological and pathophysiological processes, including cochlear response to noise [68].

Vlajkovic et al. provided a detailed description of NTPDase1 and NTPDase2 distribution in mouse and rat cochlear tissues using immunocytochemistry [17,69]. These two cell surface-located enzymes have different hydrolytic profiles: NTPDase1 hydrolyses

nucleoside 5'-triphosphates (NTPs) and nucleoside 5'-diphosphates (NDPs) to a similar extent, whilst NTPDase2 has a high preference for NTPs [67]. NTPDase1 immunoexpression was most prominent in the cochlear vasculature and cell bodies of the spiral ganglion neurons, whereas considerable NTPDase2 immunoreactivity was detected in the stria vascularis [17,69]. Both NTPDases were localised in the cuticular plates of the sensory hair cells, and auditory nerve fibres projecting from the synaptic area underneath the inner and outer hair cells. Their localisation corresponds to the reported distribution of the P2X₂ receptor in sensory, supporting and neural cells and P2Y receptor distribution in the cochlear vasculature and secretory tissues of the lateral wall. The putative role of NTPDase1 and 2 in the cochlea is to regulate extracellular ATP signalling involved in cochlear blood flow, electrochemical regulation of sound transduction and neurotransmission in the cochlea [17,69].

NTPDase3 (NTPase activity > NDPase) immunoreactivity was observed in the primary afferent neurons of the spiral ganglion and their neurites extending to the synapses beneath the inner and outer hair cells, suggesting a role for NTPDase3 in regulating ATP signaling associated with auditory neurotransmission [70]. Semi-quantitative immunohistochemistry revealed increased NTPDase3 immunolabeling in the synaptic regions of the inner and outer hair cells at elevated sound levels. NTPDase3 upregulation in the noise-exposed cochlea can prevent the activation of the cytotoxic P2X₇ receptor, suggesting the potential neuroprotective nature of this ectonucleotidase [70].

O'Keefe et al. reported the dynamic changes in the expression of NTPDase5 and 6 in the developing and adult rat cochlea [71,72]. These two intracellular members of the NTPDase family can be released in a soluble form and show a preference for nucleoside 5'-diphosphates, such as uridine 5'-diphosphate (UDP) and guanosine 5'-diphosphate (GDP). NTPDase6 immunolocalisation in the developing cochlea underpins its putative role in hair cell bundle development, while NTPDase5 may have an extracellular role in the development of sensory and neural tissues [72]. In the adult rat cochlea, upregulation of NTPDase5 after exposure to loud sound indicates a possible role for NTPDase5 in cochlear response to stress [71]. In addition, NTPDase6 immunolocalisation in the vestibular end organ could be linked to the maintenance of vestibular hair bundles [73].

P2 receptors in the cochlea initiate various signaling pathways that could be involved in the noise-induced cochlear injury. Stimuli such as noise or hypoxia could induce the excessive release of ATP into the cochlear fluid spaces [7], which may exert a cytotoxic effect mainly acting on the P2X₇ receptor. Membrane-bound NTPDases appear essential for regulating extracellular nucleotide concentrations and P2 receptor signaling in the cochlea in physiological and pathophysiological conditions. In the rat cochlea exposed to traumatic noise (110 dB SPL), we observed increased expression of NTPDase1 and NTPDase2 mRNA transcripts, while mild noise (90 dB SPL) altered only NTPDase1 mRNA expression levels [68]. Functional studies revealed increased ATPase activities in the cochlea after exposure to traumatic noise, consistent with the up-regulation of NTPDases. The changes in NTPDase expression may reflect the adaptive response of cochlear tissues to limit ATP signaling during noise exposure and thus protect the cochlea [68].

7. Adenosine Receptor Signalling in the Cochlea

Adenosine is a naturally occurring purine nucleoside that mediates its physiological actions by interacting with four cell surface-located adenosine receptors (A₁, A_{2A}, A_{2B}, A₃) distributed throughout the body [74]. Adenosine can be released from cells via specific bidirectional adenosine transporters and is also the end-product of ATP hydrolysis (Figure 2). Adenosine receptors are G protein-coupled receptors activating diverse cellular signaling pathways that define their tissue-specific roles [75]. The specific tissue distribution of adenosine receptors in the mammalian cochlea implicates the role of adenosine signalling in cochlear blood flow, sensory transduction and auditory neurotransmission [18]. The ability of adenosine A₁ receptors to reduce oxidative stress and inflammation in the cochlea and thus prevent cochlear injury caused by acoustic trauma or ototoxic drugs has opened a new chapter in the preventative treatment of sensorineural hearing loss [18,76]. The

balance between A₁ and A_{2A} receptors appears to be a critical factor for cochlear response to oxidative stress, which has been established as an underlying mechanism of several inner ear pathologies (e.g., noise-induced, age-related and drug-related hearing loss) [77]. Preclinical studies have demonstrated the extraordinary potential of adenosine receptor ligands (agonists and antagonists) in regulating the cochlear response to stress and injury and opened new avenues for the pharmacological management of hearing loss [18].

The distribution of A₁, A_{2A}, A₃ receptors was identified by immunohistochemistry [78,79]. Adenosine receptors were differentially expressed in the organ of Corti sensory and supporting cells, spiral ganglion neurons, lateral wall tissues and cochlear blood vessels. The distribution of adenosine receptors in sensory and neural tissues and the vasculature coincided with other elements of purinergic signalling (P2 receptors, ectonucleotidases), supporting the role of extracellular nucleotides and nucleosides in the regulation of cochlear function [78].

Studies on mice with global deletion of A₁ or A_{2A} receptors demonstrated the distinct roles of these receptors in cochlear physiology and response to injury [77]. Genetic deletion of the A₁R resulted in early-onset high-frequency hearing loss at ambient sound levels; this hearing loss was aggravated by noise exposure [77]. In contrast, the A_{2A}R deletion did not affect auditory thresholds but improved the survival of sensorineural tissues in the cochlea after exposure to traumatic noise. The A_{2A}R-null mice demonstrated better preservation of OHC and afferent synapses and minimal loss of SGN after noise exposure. This study suggests that the loss of A₁R results in increased susceptibility to cochlear neural injury and hearing loss, whilst the absence of A_{2A}R increases cochlear resistance to acoustic trauma [77].

Extracellular adenosine concentrations in the mammalian cochlea are regulated by selective adenosine uptake with adenosine transporters and intracellular enzymes such as adenosine deaminase (Ada) and adenosine kinase (Adk) [18] (Figure 2). Two equilibrative (ENT1 and ENT2) and two concentrative (CNT1 and CNT2) nucleoside transporters are expressed in the rat cochlea [80]. Exogenous adenosine perfused through the cochlear perilymphatic compartment was effectively taken up by cells lining this fluid compartment, and the uptake of adenosine was inhibited by an adenosine uptake blocker nitrobenzylthioinosine. The study demonstrates the bi-directional nucleoside transport in the cochlea, supporting adenosine recycling and regulating adenosine concentrations in cochlear fluid spaces [80].

Intracellularly, adenosine is hydrolysed by Ada to inosine, whilst Adk phosphorylates adenosine to adenosine 5'-monophosphate (AMP) [18]. It was reported that Ada-deficient patients have bilateral SNHL and severe immune deficiency, which were faithfully reproduced in *Ada*-null mice [81]. Ada deficiency in those mice was associated with hearing deficits and damage to cochlear hair cells, but early initiation of enzyme replacement therapy improved hearing and immune abnormalities [82]. Interestingly, elevated cochlear adenosine levels in untreated mice were associated with enhanced expression of *Adora2b* gene encoding A_{2B}R. Treatment with an A_{2B}R antagonist significantly improved hearing loss in *Ada*-null mice, nerve fiber and myelin density, suggesting that the activation of A_{2B}R aggravates sensorineural hearing loss (SNHL) [81].

8. Adenosine A₁ Receptors and Sensorineural Hearing Loss

Hearing loss is a global health issue. The World Health Organization estimates that by 2050, over 700 million people, or one in every ten people, will experience disabling hearing loss [83]. Noise-induced hearing loss (NIHL) has become a leading occupational health risk in developed countries and may result from unsafe recreational, social, and residential noise exposures. However, pharmacological treatments for NIHL are still lacking.

Adenosine is a constitutive cell metabolite with an established role in tissue protection and regeneration. The adenosine A₁ receptors are the primary mediators of cytoprotection in the cochlea [18]. The activation of A₁R protects from hearing loss by inhibiting oxidative stress, inflammation and apoptotic pathways in the cochlea [76]. The overproduction

of reactive oxygen species (ROS) induces expression of the A₁R via activation of the nuclear factor kappa B (NF-κB) [84]. Mice with genetic deletion of the NF-κB p50 subunit demonstrate altered expression of A₁R and A_{2A}R and distinctive behavioral phenotypes, suggesting a critical role of NF-κB in expression levels of adenosine receptors [85]. However, exogenously administered adenosine receptor agonists are required to boost the protective capacity of these receptors under oxidative stress conditions.

Previous studies have shown that A₁R agonists can prophylactically reduce noise-induced cochlear injury. For example, intratympanic administration of A₁R agonist R-phenylisopropyladenosine (R-PIA) before acoustic overexposure significantly improved auditory thresholds and hair cell survival in the chinchilla cochlea [86].

However, the post-exposure treatment of NIHL is more appealing from a clinical perspective. Wong et al. demonstrated, for the first time, that A₁R agonists (adenosine, 2-chloro-N-cyclopentyladenosine or CCPA), applied to the round window membrane of the cochlea 6 hours after noise exposure, effectively reduced auditory brainstem threshold shifts in rats by reducing oxidative stress and noise-induced hair cell loss [87]. In contrast, the selective A_{2A}R agonist CGS-21680 and A₃R agonist CI-IB-MECA did not protect the cochlea from injury and hearing loss [87].

More recently, a selective A₁ adenosine receptor agonist, adenosine amine congener (ADAC), emerged as a potentially effective treatment for noise-induced cochlear injury and hearing loss [88–90]. The post-exposure treatment with ADAC led to a significantly greater recovery of hearing thresholds and improved survival of sensory hair cells in rats compared with non-treated controls [89]. We have also demonstrated the dose-dependent rescue effect of ADAC on noise-induced cochlear injury and established the time window for treatment [88]. ADAC was most effective in the first 24 hours after noise exposure (8–16 kHz, 110 dB sound pressure level for 2 hours), providing up to 21 dB protection averaged across the frequencies (8–28 kHz). The drug was effective at doses 50–200 µg/kg administered as five consecutive daily intraperitoneal injections. Even delayed treatment 48 hours after noise exposure provided clinically significant improvement of auditory brainstem thresholds (>10 dB) at some frequencies [88]. These data show that ADAC mitigates noise-induced hearing loss in a dose- and time-dependent manner and support ADAC development as a potential clinical otological treatment for acute sensorineural hearing loss caused by exposure to traumatic noise.

The chemotherapeutic agent cisplatin can also cause the upregulation of adenosine receptors in the cochlea, which likely represents a compensatory mechanism to counter the toxic effects of cisplatin-induced ROS overproduction [91]. More recent studies implicate ROS-induced inflammatory and apoptotic processes in the cochlea by activating signal transducer and activator of transcription (STAT1) [79]. A₁R activation protects against cisplatin ototoxicity by suppressing inflammatory and oxidative stress responses initiated by ROS generation [79]. Intratympanic or parenteral administration of the A₁R agonists R-PIA, CCPA and ADAC significantly reduced cisplatin-induced threshold shifts and protected against cisplatin-induced hair cell damage [92,93]. These studies suggest that the A₁R contributes significantly to cochlear protection from ototoxic drugs and mitigates drug-induced hearing loss.

In contrast, inhibition of the A₁R by a broad spectrum adenosine receptor antagonist caffeine potentiated cisplatin-induced hearing loss in a rat model of cisplatin ototoxicity [94]. A single-dose oral administration of caffeine exacerbated cisplatin-induced hearing loss by increasing synaptopathy and inflammation in the cochlea, whereas multiple doses of caffeine were associated with enhanced damage to OHC. This study suggests that caffeine consumption should be limited in cancer patients treated with cisplatin [94].

Aminoglycoside antibiotics can also cause sensorineural hearing loss [95]. Using an established organotypic tissue culture model of the neonatal mouse cochlea, Lin et al. investigated the effect of P1 (adenosine) and P2 (ATP) receptor activation on the sensory hair cell survival after exposure to the ototoxic aminoglycoside neomycin [96]. Neomycin-induced ototoxicity was aggravated by the addition of slowly hydrolyzable ATP analog

ATP γ S, whilst the activation of adenosine receptors by ADAC or adenosine conferred partial protection from neomycin ototoxicity. It was inferred that adenosine A₁ receptors are critical for maintaining cochlear homeostasis and survival following ototoxic injury [96].

9. Adenosine Kinase and Age-Related Hearing Loss

Age-related hearing loss (ARHL) is the most common sensory disability, characterized by a decline in hearing sensitivity and speech discrimination, impaired sound localization, and delayed auditory information processing [97]. Currently, there are no treatments that can prevent or delay ARHL. A preclinical study [98] demonstrated that enhanced adenosine receptor signalling in the cochlea could provide partial protection from ARHL in C57BL/6J mice characterized by accelerated hearing loss [99]. That study [98] targeted adenosine kinase (Adk), the key enzyme in adenosine metabolism that regulates intracellular and extracellular adenosine concentrations, using a selective Adk inhibitor ABT-702 administered intraperitoneally twice a week, from three to nine months of age. At nine months, mice treated with ABT-702 showed lower auditory brainstem threshold shifts and improved survival of sensory hair cells compared to control non-treated mice. The study provides the first evidence that ARHL can be partially prevented or delayed by enhancing endogenous adenosine receptor signalling in the cochlea [98].

10. Regulators of G Protein Signalling and Hearing Loss

Previous studies have shown that the “post-exposure” treatment with adenosine A₁R agonists confers partial protection against acoustic trauma and other forms of SNHL [87,88]. Another study attempted to rescue noise-induced cochlear injury by post-exposure treatment targeting A₁R without using receptor agonists [100]. This was achieved by pharmacological inhibition of the protein that regulates G protein signalling by the A₁R (Regulator of G protein Signalling 4, RGS4). A molecular complex between RGS4 and neurabin, an intracellular scaffolding protein expressed in neural and cochlear tissues, is the key negative regulator of A₁R activity [101]. In that study, Wistar rats (6–8 weeks) were exposed to traumatic noise (110 dB SPL, 8–16 kHz for 2 h), and a small molecule RGS4 inhibitor CCG-4986 was delivered by intratympanic injection 24 or 48 h after noise exposure. Intratympanic administration of CCG-4986 48 h after noise exposure attenuated noise-induced auditory threshold shifts by up to 19 dB, whilst the earlier drug administration (24 h) led to even better preservation of auditory thresholds (up to 32 dB). Functional improvement was linked to improved survival of sensorineural tissues and afferent synapses in the cochlea. RGS inhibitors thus represent a novel paradigm for treating noise-induced cochlear injury and possibly other forms of SNHL [100].

Subsequent studies have shown that other members of the RGS family of proteins may be involved in cisplatin-induced ototoxicity. Cisplatin upregulates RGS17 expression in the cochlea, leading to increased oxidative stress, upregulation of inflammatory genes and enhanced apoptosis in the cochlea [102]. The silencing of *RGS17* suppressed cisplatin-induced hearing loss in rats, while overexpression of *RGS17* enhanced cisplatin-induced hearing loss. RGS17 represents a novel mediator of cisplatin ototoxicity and a potential therapeutic target for cisplatin-induced hearing loss [102].

11. Adenosine A_{2A} Receptors and Hearing Loss

The role of A_{2A} receptors in cochlear injury development appears to be opposite to the A₁R. Previous studies in mice with genetic deletion of the adenosine A_{2A} receptor have demonstrated better preservation of cochlear afferent synapses and spiral ganglion neurons after acoustic overexposure compared to control wildtype mice [77]. This has informed our alternative approaches to cochlear neuroprotection based on pharmacological inhibition of the A_{2A}R. In a rat organotypic tissue culture model of excitotoxic injury (combined exposure to NMDA and kainic acid), the co-administration of istradefylline (a clinically approved A_{2A}R antagonist) reduced deafferentation of the inner hair cells and improved the survival of afferent synapses after excitotoxic injury [103]. This study may

have implications for the treatment of cochlear neuropathy and the prevention of hidden hearing loss as its clinical manifestation.

The A_{2A} receptor targeting may also be relevant for the preventative treatment of age-related hearing loss. Middle-aged C57BL/6J mice, prone to early onset ARHL, were given weekly istradefylline injections (1 mg/kg) from 6 to 12 months of age [104]. Auditory function was assessed using ABR to tone pips (4–32 kHz) at 6, 9, and 12 months of age. Weekly injections of istradefylline attenuated ABR threshold shifts by approximately 20 dB at mid to high frequencies (16–32 kHz) and improved hair cell survival in a turn-dependent manner. This study presents the first evidence for the rescue potential of istradefylline in ARHL [104]. This and other options for the therapeutic targeting of adenosine receptors in the cochlea are shown in Figure 4.




Figure 4. Cochlear protection by adenosine receptors. The preclinical studies established four distinct strategies targeting adenosine receptors to rescue the cochlea from injury and prevent sensorineural hearing loss. These pharmacological interventions include the administration of A_1 receptor agonists, A_{2A} receptor antagonists, adenosine kinase (Adk) inhibitors and RGS4 (Regulator of G protein signalling 4) inhibitors. The most common route for drug delivery to the inner ear is the intratympanic injection onto the round window membrane of the cochlea, which precludes off-target effects associated with systemic administration. Created with [BioRender.com](https://www.biorender.com).

12. Conclusions

Purinergic signalling is an intricate system of extracellular receptors, enzymes and transporters that regulates multiple physiological and pathophysiological processes in the mammalian inner ear. P2 receptors have an essential role in cochlear development, regulation of electrochemical homeostasis, auditory neurotransmission, and adaptation to elevated sound levels. Surface-located ectonucleotidases (NTPDases and CD73) hydrolyse extracellular ATP to adenosine and thus convert the P2 receptor environment into the P1 (adenosine) receptor environment. The two main types of adenosine receptors, A_1 and A_{2A} , have differential roles in cochlear injury response, highlighting their clinical significance as prospective therapeutic targets. Pharmacological manipulation of P2 receptor signalling is also a developing strategy for the therapeutic management of hearing loss.

Author Contributions: Conceptualisation, methodology and writing—original draft preparation, S.M.V.; review and editing, P.R.T.; illustrations, S.M.V.; funding acquisition, S.M.V. and P.R.T. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by Auckland Medical Research Foundation (Auckland, New Zealand), grant number 3720285.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Elliott, S.J.; A Shera, C. The cochlea as a smart structure. *Smart Mater. Struct.* **2012**, *21*, 064001. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Raphael, Y.; A Altschuler, R. Structure and innervation of the cochlea. *Brain Res. Bull.* **2003**, *60*, 397–422. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Liberman, M.C.; Gao, J.; He, D.Z.Z.; Wu, X.; Jia, S.; Zuo, J. Prestin is required for electromotility of the outer hair cell and for the cochlear amplifier. *Nature* **2002**, *419*, 300–304. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Cohen-Salmon, M.; Ott, T.; Michel, V.; Hardelin, J.P.; Perfettini, I.; Eybalin, M.; Wu, T.; Marcus, D.C.; Wangemann, P.; Willecke, K.; et al. Targeted ablation of connexin26 in the inner ear epithelial gap junction network causes hearing impairment and cell death. *Curr. Biol.* **2002**, *12*, 1106–1111. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Thorne, P.R.; Munoz, D.J.; Nikolic, P.; Mander, L.; Jagger, D.J.; Greenwood, D.; Vlajkovic, S.; Housley, G.D. Potential Role of Purinergic Signalling in Cochlear Pathology. *Audiol. Neurotol.* **2002**, *7*, 180–184. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Housley, G.D.; Bringmann, A.; Reichenbach, A. Purinergic signaling in special senses. *Trends Neurosci.* **2009**, *32*, 128–141. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Muñoz, D.J.; Kendrick, I.S.; Rassam, M.; Thorne, P.R. Vesicular storage of adenosine triphosphate in the guinea-pig cochlear lateral wall and concentrations of ATP in the endolymph during sound exposure and hypoxia. *Acta Oto-Laryngol.* **2001**, *121*, 10–15.
8. Housley, G.D.; Morton-Jones, R.; Vlajkovic, S.M.; Telang, R.S.; Paramanathasivam, V.; Tadros, S.F.; Wong, A.C.Y.; Froud, K.E.; Cederholm, J.M.; Sivakumaran, Y.; et al. ATP-gated ion channels mediate adaptation to elevated sound levels. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 7494–7499. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Thorne, P.R.; Muñoz, D.J.B.B.; Housley, G.D. Purinergic Modulation of Cochlear Partition Resistance and Its Effect on the Endocochlear Potential in the Guinea Pig. *JARO-J. Assoc. Res. Otolaryngol.* **2004**, *5*, 58–65. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Housley, G.D.; Jagger, D.J.; Greenwood, D.; Raybould, N.P.; Salih, S.G.; Järleback, L.E.; Vlajkovic, S.M.; Kanjhan, R.; Nikolic, P.; Muñoz, D.J.M.M.; et al. Purinergic regulation of sound transduction and auditory neurotransmission. *Audiol. Neuro-Otol.* **2002**, *7*, 55–61. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Berekméri, E.; Szepeszy, J.; Köles, L.; Zelles, T. Purinergic signaling in the organ of Corti: Potential therapeutic targets of sensorineural hearing losses. *Brain Res. Bull.* **2019**, *151*, 109–118. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Burnstock, G.; Verkhratsky, A. Special Senses. In *Purinergic Signalling and the Nervous System*; Springer: Berlin/Heidelberg, Germany, 2012; pp. 627–679.
13. Köles, L.; Szepeszy, J.; Berekméri, E.; Zelles, T. Purinergic signalling and cochlear injury-targeting the immune system? *Int. J. Mol. Sci.* **2019**, *20*, 2979. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Abbracchio, M.P.; Burnstock, G.; Verkhratsky, A.; Zimmermann, H. Purinergic signalling in the nervous system: An overview. *Trends Neurosci.* **2009**, *32*, 19–29. [\[CrossRef\]](#)
15. Piazza, V.; Ciubotaru, C.D.; Gale, J.E.; Mammano, F. Purinergic signalling and intercellular Ca²⁺ wave propagation in the organ of Corti. *Cell Calcium* **2007**, *41*, 77–86. [\[CrossRef\]](#) [\[PubMed\]](#)
16. Zimmermann, H. History of ectonucleotidases and their role in purinergic signaling. *Biochem. Pharmacol.* **2021**, *187*, 114322. [\[CrossRef\]](#)
17. Vlajkovic, S.M.; Thorne, P.R.; Sévigny, J.; Robson, S.C.; Housley, G.D. NTPDase1 and NTPDase2 immunolocalization in mouse cochlea: Implications for regulation of p2 receptor signaling. *J. Histochem. Cytochem.* **2002**, *50*, 1435–1441. [\[CrossRef\]](#)
18. Vlajkovic, S.M.; Housley, G.D.; Thorne, P.R. Adenosine and the auditory system. *Curr. Neuropharmacol.* **2009**, *7*, 246–256. [\[CrossRef\]](#)
19. Jovanovic, S.; Milenkovic, I. Purinergic Modulation of Activity in the Developing Auditory Pathway. *Neurosci. Bull.* **2020**, *36*, 1285–1298. [\[CrossRef\]](#)
20. Mammano, F. ATP-dependent intercellular Ca²⁺ signaling in the developing cochlea: Facts, fantasies and perspectives. *Semin. Cell Dev. Biol.* **2013**, *24*, 31–39. [\[CrossRef\]](#)
21. Tritsch, N.X.; Yi, E.; Gale, J.E.; Glowatzki, E.; Bergles, D.E. The origin of spontaneous activity in the developing auditory system. *Nature* **2007**, *450*, 50–55. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Housley, G.D.; von Jonquieres, G.; Pinyon, J.L.; Matheson, J.-A.T.; Pearson, L.J.; Salthouse, T.P.; Cederholm, J.M. Cochlear homeostasis: A molecular physiological perspective on maintenance of sound transduction and auditory neurotransmission with noise and ageing. *Curr. Opin. Physiol.* **2020**, *18*, 106–115. [\[CrossRef\]](#)
23. Dahl, G. ATP release through pannexon channels. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* **2015**, *370*, 20140191. [\[CrossRef\]](#)

24. Chen, J.; Zhu, Y.; Liang, C.; Chen, J.; Zhao, H.B. Pannexin1 channels dominate ATP release in the cochlea ensuring endocochlear potential and auditory receptor potential generation and hearing. *Sci. Rep.* **2015**, *5*, 10762. [[CrossRef](#)]
25. Belousov, A.B.; Fontes, J.D.; Freitas-Andrade, M.; Naus, C.C. Gap junctions and hemichannels: Communicating cell death in neurodevelopment and disease. *BMC Cell Biol.* **2017**, *18* (Suppl. S1), 4. [[CrossRef](#)]
26. Zhu, Y.; Zhao, H.B. ATP activates P2X receptors to mediate gap junctional coupling in the cochlea. *Biochem. Biophys. Res. Commun.* **2012**, *426*, 528–532. [[CrossRef](#)]
27. Ceriani, F.; Pozzan, T.; Mammano, F. Critical role of ATP-induced ATP release for Ca²⁺ signaling in nonsensory cell networks of the developing cochlea. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E7194–E7201. [[CrossRef](#)]
28. Locovei, S.; Wang, J.; Dahl, G. Activation of pannexin 1 channels by ATP through P2Y receptors and by cytoplasmic calcium. *FEBS Lett.* **2006**, *580*, 239–244. [[CrossRef](#)]
29. Baroja-Mazo, A.; Barberà-Cremades, M.; Pelegrín, P. The participation of plasma membrane hemichannels to purinergic signaling. *Biochim. Et Biophys. Acta-Biomembr.* **2013**, *1828*, 79–93. [[CrossRef](#)]
30. Sahu, G.; Sukumaran, S.; Bera, A.K. Pannexins form gap junctions with electrophysiological and pharmacological properties distinct from connexins. *Sci. Rep.* **2014**, *4*, 4955. [[CrossRef](#)]
31. Zhao, H.B. Expression and function of pannexins in the inner ear and hearing. *BMC Cell Biol.* **2016**, *17* (Suppl. S1), 16. [[CrossRef](#)]
32. Zhu, Y.; Zhao, H.B. ATP-mediated potassium recycling in the cochlear supporting cells. *Purinergic Signal.* **2010**, *6*, 221–229. [[CrossRef](#)] [[PubMed](#)]
33. Zhao, H.-B.; Zhu, Y.; Liang, C.; Chen, J. Pannexin 1 deficiency can induce hearing loss. *Biochem. Biophys. Res. Commun.* **2015**, *463*, 143–147. [[CrossRef](#)] [[PubMed](#)]
34. Nikolic, P.; Housley, G.D.; Luo, L.; Ryan, A.F.; Thorne, P.R. Transient expression of P2X₁ receptor subunits of ATP-gated ion channels in the developing rat cochlea. *Dev. Brain Res.* **2001**, *126*, 173–182. [[CrossRef](#)] [[PubMed](#)]
35. Järlebark, L.E.; Housley, G.D.; Raybould, N.P.; Vlajkovic, S.; Thorne, P.R. ATP-gated ion channels assembled from P2X₂ receptor subunits in the mouse cochlea. *Neuroreport* **2002**, *13*, 1979–1984. [[CrossRef](#)]
36. Järlebark, L.E.; Housley, G.D.; Thorne, P.R. Immunohistochemical localization of adenosine 5'-triphosphate-gated ion channel P2X₂ receptor subunits in adult and developing rat cochlea. *J. Comp. Neurol.* **2000**, *421*, 289–301. [[CrossRef](#)]
37. Huang, L.C.; Ryan, A.F.; Cockayne, D.A.; Housley, G.D. Developmentally regulated expression of the P2X₃ receptor in the mouse cochlea. *Histochem. Cell Biol.* **2006**, *125*, 681–692. [[CrossRef](#)]
38. Greenwood, D.; Jagger, D.J.; Huang, L.C.; Hoya, N.; Thorne, P.R.; Wildman, S.S.; King, B.F.; Pak, K.; Ryan, A.F.; Housley, G.D. P2X receptor signaling inhibits BDNF-mediated spiral ganglion neuron development in the neonatal rat cochlea. *Development* **2007**, *134*, 1407–1417. [[CrossRef](#)]
39. Szűcs, A.; Szappanos, H.; Tóth, A.; Farkas, Z.; Panyi, G.; Csernoch, L.; Sziklai, I. Differential expression of purinergic receptor subtypes in the outer hair cells of the guinea pig. *Hear. Res.* **2004**, *196*, 2–7. [[CrossRef](#)]
40. Wu, T.; Dai, M.; Shi, X.R.; Jiang, Z.G.; Nuttall, A.L. Functional expression of P2X₄ receptor in capillary endothelial cells of the cochlear spiral ligament and its role in regulating the capillary diameter. *Am. J. Physiol. Circ. Physiol.* **2011**, *301*, H69–H78. [[CrossRef](#)]
41. Xiang, Z.; Bo, X.; Burnstock, G. P2X receptor immunoreactivity in the rat cochlea, vestibular ganglion and cochlear nucleus. *Hear. Res.* **1999**, *128*, 190–196. [[CrossRef](#)]
42. Jagger, D.J.; Housley, G.D. Membrane properties of type II spiral ganglion neurones identified in a neonatal rat cochlear slice. *J. Physiol.* **2003**, *552*, 525–533. [[CrossRef](#)]
43. Nikolic, P.; Housley, G.D.; Thorne, P.R. Expression of the P2X₇ Receptor Subunit of the Adenosine 5'-Triphosphate-Gated Ion Channel in the Developing and Adult Rat Cochlea. *Audiol. Neurotol.* **2003**, *8*, 28–37. [[CrossRef](#)] [[PubMed](#)]
44. Prades, S.; Heard, G.; Gale, J.E.; Engel, T.; Kopp, R.; Nicke, A.; Smith, K.E.; Jagger, D.J. Functional P2X₇ Receptors in the Auditory Nerve of Hearing Rodents Localize Exclusively to Peripheral Glia. *J. Neurosci.* **2021**, *41*, 2615–2629. [[CrossRef](#)] [[PubMed](#)]
45. Morton-Jones, R.T.; Vlajkovic, S.M.; Thorne, P.R.; Cockayne, D.A.; Ryan, A.F.; Housley, G.D. Properties of ATP-gated ion channels assembled from P2X₂ subunits in mouse cochlear Reissner's membrane epithelial cells. *Purinergic Signal.* **2015**, *11*, 551–560. [[CrossRef](#)] [[PubMed](#)]
46. Cederholm, J.M.E.; Ryan, A.F.; Housley, G.D. Onset kinetics of noise-induced purinergic adaptation of the 'cochlear amplifier'. *Purinergic Signal.* **2019**, *15*, 343–355. [[CrossRef](#)] [[PubMed](#)]
47. Yan, D.; Zhu, Y.; Walsh, T.; Xie, D.; Yuan, H.; Sirmaci, A.; Fujikawa, T.; Wong, A.C.Y.; Loh, T.L.; Du, L.; et al. Mutation of the ATP-gated P2X₂ receptor leads to progressive hearing loss and increased susceptibility to noise. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 2228–2233. [[CrossRef](#)] [[PubMed](#)]
48. Chen, X.; Abad, C.; Chen, Z.-Y.; I Young, J.; Gurumurthy, C.B.; Walz, K.; Liu, X.Z. Generation and characterization of a P2rx2 V60L mouse model for DFNA41. *Hum. Mol. Genet.* **2021**, *30*, 985–995. [[CrossRef](#)]
49. George, B.; Swartz, K.J.; Li, M. Hearing loss mutations alter the functional properties of human P2X₂ receptor channels through distinct mechanisms. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 22862–22871. [[CrossRef](#)]
50. Horváth, T.; Polony, G.; Fekete, Á.; Aller, M.; Halmos, G.; Lendvai, B.; Heinrich, A.; Sperlág, B.; Vizi, E.S.; Zelles, T. ATP-Evoked Intracellular Ca²⁺ Signaling of Different Supporting Cells in the Hearing Mouse Hemicochlea. *Neurochem. Res.* **2016**, *41*, 364–375. [[CrossRef](#)]

51. Lahne, M.; Gale, J.E. Damage-induced cell–cell communication in different cochlear cell types via two distinct ATP-dependent Ca^{2+} waves. *Purinergic Signal.* **2010**, *6*, 189–200. [[CrossRef](#)]
52. Sirko, P.; E Gale, J.; Ashmore, J.F. Intercellular Ca^{2+} signalling in the adult mouse cochlea. *J. Physiol.* **2019**, *597*, 303–317. [[CrossRef](#)] [[PubMed](#)]
53. Liu, C.; Glowatzki, E.; Fuchs, P.A. Unmyelinated type II afferent neurons report cochlear damage. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 14723–14727. [[CrossRef](#)] [[PubMed](#)]
54. Huang, L.C.; Thorne, P.R.; Vlajkovic, S.M.; Housley, G.D. Differential expression of P2Y receptors in the rat cochlea during development. *Purinergic Signal.* **2010**, *6*, 231–248. [[CrossRef](#)] [[PubMed](#)]
55. Babola, T.A.; Li, S.; Wang, Z.; Kersbergen, C.J.; Elgoyhen, A.B.; Coate, T.M.; Bergles, D.E. Purinergic Signaling Controls Spontaneous Activity in the Auditory System throughout Early Development. *J. Neurosci.* **2021**, *41*, 594–612. [[CrossRef](#)] [[PubMed](#)]
56. Chen, J.; Hou, S.; Yang, J. ATP is stored in lysosomes of greater epithelial ridge supporting cells in newborn rat cochleae. *J. Cell. Biochem.* **2019**, *120*, 19469–19481. [[CrossRef](#)] [[PubMed](#)]
57. Liu, J.; Liu, W.; Yang, J. ATP-containing vesicles in stria vascular marginal cell cytoplasm in neonatal rat cochlea are lysosomes. *Sci. Rep.* **2016**, *6*, 20903. [[CrossRef](#)]
58. Liu, B.; Cao, W.; Li, J.; Liu, J. Lysosomal exocytosis of ATP is coupled to P2Y2 receptor in marginal cells in the stria vascular in neonatal rats. *Cell Calcium* **2018**, *76*, 62–71. [[CrossRef](#)]
59. Dayaratne, M.W.; Vlajkovic, S.M.; Lipski, J.; Thorne, P.R. Kölliker’s organ and the development of spontaneous activity in the auditory system: Implications for hearing dysfunction. *BioMed Res. Int.* **2014**, *2014*, 367939. [[CrossRef](#)]
60. Dayaratne, M.W.; Vlajkovic, S.M.; Lipski, J.; Thorne, P.R. Putative role of border cells in generating spontaneous morphological activity within Kölliker’s organ. *Hear. Res.* **2015**, *330*, 90–97. [[CrossRef](#)]
61. Rabbitt, R.D.; Holman, H.A. ATP and ACh Evoked Calcium Transients in the Neonatal Mouse Cochlear and Vestibular Sensory Epithelia. *Front. Neurosci.* **2021**, *15*, 710076. [[CrossRef](#)]
62. Ceriani, F.; Hendry, A.; Jeng, J.; Johnson, S.L.; Stephani, F.; Olt, J.; Holley, M.C.; Mammano, F.; Engel, J.; Kros, C.J.; et al. Coordinated calcium signalling in cochlear sensory and non-sensory cells refines afferent innervation of outer hair cells. *EMBO J.* **2019**, *38*, e99839. [[CrossRef](#)]
63. Babola, T.A.; Kersbergen, C.J.; Wang, H.C.; Bergles, D.E. Purinergic signaling in cochlear supporting cells reduces hair cell excitability by increasing the extracellular space. *eLife* **2020**, *9*, 52160. [[CrossRef](#)] [[PubMed](#)]
64. Lee, J.H.; Heo, J.H.; Kim, C.H.; Chang, S.O.; Kim, C.S.; Oh, S.H. Changes in P2Y₄ receptor expression in rat cochlear outer sulcus cells during development. *Hear. Res.* **2007**, *228*, 201–211. [[CrossRef](#)]
65. Wang, Z.; Jung, J.S.; Inbar, T.C.; Rangoussis, K.M.; Faaborg-Andersen, C.; Coate, T.M. The purinergic receptor P2rx3 is required for spiral ganglion neuron branch refinement during development. *eNeuro* **2020**, *7*, ENEURO.0179-20.2020. [[CrossRef](#)] [[PubMed](#)]
66. Jovanovic, S.; Radulovic, T.; Coddou, C.; Dietz, B.; Nerlich, J.; Stojilkovic, S.S.; Rübsamen, R.; Milenkovic, I. Tonotopic action potential tuning of maturing auditory neurons through endogenous ATP. *J. Physiol.* **2017**, *595*, 1315–1337. [[CrossRef](#)]
67. Robson, S.C.; Sévigny, J.; Zimmermann, H. The E-NTPDase family of ectonucleotidases: Structure function relationships and pathophysiological significance. *Purinergic Signal.* **2006**, *2*, 409–430. [[CrossRef](#)]
68. Vlajkovic, S.M.; Housley, G.D.; Muñoz, D.J.; Robson, S.C.; Sévigny, J.; Wang, C.J.; Thorne, P.R. Noise exposure induces up-regulation of ecto-nucleoside triphosphate diphosphohydrolases 1 and 2 in rat cochlea. *Neuroscience* **2004**, *126*, 763–773. [[CrossRef](#)]
69. Vlajkovic, S.M.; Thorne, P.R.; Sévigny, J.; Robson, S.C.; Housley, G.D. Distribution of ectonucleoside triphosphate diphosphohydrolases 1 and 2 in rat cochlea. *Hear. Res.* **2002**, *170*, 127–138. [[CrossRef](#)]
70. Vlajkovic, S.M.; Vinayagamoorthy, A.; Thorne, P.R.; Robson, S.C.; Wang, C.J.; Housley, G.D. Noise-induced up-regulation of NTPDase3 expression in the rat cochlea: Implications for auditory transmission and cochlear protection. *Brain Res.* **2006**, *1104*, 55–63. [[CrossRef](#)]
71. O’Keeffe, M.G.; Thorne, P.R.; Housley, G.D.; Robson, S.C.; Vlajkovic, S.M. Distribution of NTPDase5 and NTPDase6 and the regulation of P2Y receptor signalling in the rat cochlea. *Purinergic Signal.* **2010**, *6*, 249–261. [[CrossRef](#)]
72. O’Keeffe, M.G.; Thorne, P.R.; Housley, G.D.; Robson, S.C.; Vlajkovic, S.M. Developmentally regulated expression of ectonucleotidases NTPDase5 and NTPDase6 and UDP-responsive P2Y receptors in the rat cochlea. *Histochem. Cell Biol.* **2010**, *133*, 425–436. [[CrossRef](#)] [[PubMed](#)]
73. O’Keeffe, M.G.; Thorne, P.R.; Housley, G.D.; Robson, S.C.; Vlajkovic, S.M. Hair cell specific NTPDase6 immunolocalisation in vestibular end organs: Potential role of purinergic signaling in vestibular sensory transduction. *J. Vestib. Res.* **2012**, *22*, 213–219. [[CrossRef](#)] [[PubMed](#)]
74. Jacobson, K.; Gao, Z.-G. Adenosine receptors as therapeutic targets. *Nat. Rev. Drug Discov.* **2006**, *5*, 247–264. [[CrossRef](#)] [[PubMed](#)]
75. Ijzerman, A.P.; Jacobson, K.A.; Müller, C.E.; Cronstein, B.N.; Cunha, R.A. International Union of Basic and Clinical Pharmacology. CXII: Adenosine Receptors: A Further Update. *Pharmacol. Rev.* **2022**, *74*, 340–372. [[CrossRef](#)]
76. Sheth, S.; Mukherjee, D.; Rybak, L.P.; Ramkumar, V. The contribution of anti-oxidant and anti-inflammatory functions of adenosine A₁ Receptor in mediating otoprotection. In *Inflammatory Mechanisms in Mediating Hearing Loss*; Springer: Berlin/Heidelberg, Germany, 2018; pp. 149–164.
77. Vlajkovic, S.M.; Ambepitiya, K.; Barclay, M.; Boison, D.; Housley, G.D.; Thorne, P.R. Adenosine receptors regulate susceptibility to noise-induced neural injury in the mouse cochlea and hearing loss. *Hear. Res.* **2017**, *345*, 43–51. [[CrossRef](#)]

78. Vlajkovic, S.M.; Abi, S.; Wang, C.J.H.; Housley, G.D.; Thorne, P.R. Differential distribution of adenosine receptors in rat cochlea. *Cell Tissue Res.* **2007**, *328*, 461–471. [\[CrossRef\]](#)
79. Kaur, T.; Borse, V.; Sheth, S.; Sheehan, K.; Ghosh, S.; Tupal, S.; Jajoo, S.; Mukherjea, D.; Rybak, L.P.; Ramkumar, V. Adenosine A₁ receptor protects against cisplatin ototoxicity by suppressing the NOX3/STAT1 inflammatory pathway in the cochlea. *J. Neurosci.* **2016**, *36*, 3962–3977. [\[CrossRef\]](#)
80. Khan, A.F.; Thorne, P.R.; Muñoz, D.J.; Wang, C.J.; Housley, G.D.; Vlajkovic, S.M. Nucleoside transporter expression and adenosine uptake in the rat cochlea. *NeuroReport* **2007**, *18*, 235–239. [\[CrossRef\]](#)
81. Manalo, J.M.; Liu, H.; Ding, D.; Hicks, J.; Sun, H.; Salvi, R.; Kellems, R.E.; Pereira, F.A.; Xia, Y. Adenosine A_{2B} receptor: A pathogenic factor and a therapeutic target for sensorineural hearing loss. *FASEB J.* **2020**, *34*, 15771–15787. [\[CrossRef\]](#)
82. Xu, X.; Negandhi, J.; Min, W.; Tsui, M.; Post, M.; Harrison, R.V.; Grunebaum, E. Early enzyme replacement therapy improves hearing and immune defects in adenosine deaminase deficient-mice. *Front. Immunol.* **2019**, *10*, 416. [\[CrossRef\]](#)
83. World Health Organization. *World Report on Hearing—Executive Summary*; World Health Organization: Geneva, Switzerland, 2021.
84. Ramkumar, V.; Whitworth, C.A.; Pingle, S.C.; Hughes, L.F.; Rybak, L.P. Noise induces A₁ adenosine receptor expression in the chinchilla cochlea. *Hear. Res.* **2004**, *188*, 47–56. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Ramkumar, V.; AJhaveri, K.; Xie, X.; Jajoo, S.; A Toth, L. Nuclear factor κ B and adenosine receptors: Biochemical and behavioral profiling. *Curr. Neuropharmacol.* **2011**, *9*, 342–349. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Hu, B.H.; Zheng, X.Y.; McFadden, S.L.; Kopke, R.D.; Henderson, D. R-phenylisopropyladenosine attenuates noise-induced hearing loss in the chinchilla. *Hear. Res.* **1997**, *113*, 198–206. [\[CrossRef\]](#)
87. Wong, A.C.Y.; Guo, C.X.; Gupta, R.; Housley, G.D.; Thorne, P.R.; Vlajkovic, S.M. Post exposure administration of A₁ adenosine receptor agonists attenuates noise-induced hearing loss. *Hear. Res.* **2010**, *260*, 81–88. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Vlajkovic, S.M.; Chang, H.; Paek, S.Y.; Chi, H.H.-T.; Sreebhavan, S.; Telang, R.S.; Tingle, M.; Housley, G.D.; Thorne, P.R. Adenosine Amine Congener as a Cochlear Rescue Agent. *BioMed Res. Int.* **2014**, *2014*, 841489. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Vlajkovic, S.M.; Lee, K.-H.; Wong, A.C.Y.; Guo, C.X.; Gupta, R.; Housley, G.D.; Thorne, P.R. Adenosine amine congener mitigates noise-induced cochlear injury. *Purinergic Signal.* **2010**, *6*, 273–281. [\[CrossRef\]](#) [\[PubMed\]](#)
90. Chang, H.; Telang, R.S.; Sreebhavan, S.; Tingle, M.; Thorne, P.R.; Vlajkovic, S.M. Pharmacokinetic Properties of Adenosine Amine Congener in Cochlear Perilymph after Systemic Administration. *BioMed Res. Int.* **2017**, *2017*, 8091462. [\[CrossRef\]](#)
91. Ford, M.S.; Nie, Z.; Whitworth, C.; Rybak, L.P.; Ramkumar, V. Up-regulation of adenosine receptors in the cochlea by cisplatin. *Hear. Res.* **1997**, *111*, 143–152. [\[CrossRef\]](#)
92. Whitworth, C.A.; Ramkumar, V.; Jones, B.; Tsukasaki, N.; Rybak, L.P. Protection against cisplatin ototoxicity by adenosine agonists. *Biochem. Pharmacol.* **2004**, *67*, 1801–1807. [\[CrossRef\]](#)
93. Gunewardene, N.; Guo, C.X.; Wong, A.C.; Thorne, P.R.; Vlajkovic, S.M. Adenosine amine congener ameliorates cisplatin-induced hearing loss. *World J. Otorhinolaryngol.* **2013**, *3*, 100–107. [\[CrossRef\]](#)
94. Sheth, S.; Sheehan, K.; Dhukhwa, A.; Al Aameri, R.F.; Mamillapalli, C.; Mukherjea, D.; Rybak, L.P.; Ramkumar, V. Oral administration of caffeine exacerbates cisplatin-induced hearing loss. *Sci. Rep.* **2019**, *9*, 9571. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Schacht, J.; Talaska, A.E.; Rybak, L.P. Cisplatin and Aminoglycoside Antibiotics: Hearing Loss and Its Prevention. *Anat. Rec.* **2012**, *295*, 1837–1850. [\[CrossRef\]](#)
96. Lin, S.C.Y.; Thorne, P.R.; Housley, G.D.; Vlajkovic, S.M. Purinergic Signaling and Aminoglycoside Ototoxicity: The Opposing Roles of P₁ (Adenosine) and P₂ (ATP) Receptors on Cochlear Hair Cell Survival. *Front. Cell. Neurosci.* **2019**, *13*, 207. [\[CrossRef\]](#) [\[PubMed\]](#)
97. Gates, G.A.; Mills, J.H. Presbycusis. *Lancet* **2005**, *366*, 1111–1120. [\[CrossRef\]](#)
98. Vlajkovic, S.M.; Guo, C.X.; Telang, R.; Wong, A.C.Y.; Paramanathanasivam, V.; Boison, D.; Housley, G.D.; Thorne, P.R. Adenosine kinase inhibition in the cochlea delays the onset of age-related hearing loss. *Exp. Gerontol.* **2011**, *46*, 905–914. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Johnson, K.R.; Erway, L.C.; Cook, S.A.; Willott, J.F.; Zheng, Q.Y. A major gene affecting age-related hearing loss in C57BL/6J mice. *Hear. Res.* **1997**, *114*, 83–92. [\[CrossRef\]](#)
100. Fok, C.; Bogosanovic, M.; Pandya, M.; Telang, R.; Thorne, P.R.; Vlajkovic, S.M. Regulator of G Protein Signalling 4 (RGS4) as a Novel Target for the Treatment of Sensorineural Hearing Loss. *Int. J. Mol. Sci.* **2020**, *22*, 3. [\[CrossRef\]](#) [\[PubMed\]](#)
101. Chen, Y.; Liu, Y.; Cottingham, C.; McMahon, L.; Jiao, K.; Greengard, P.; Wang, Q. Neurabin Scaffolding of Adenosine Receptor and RGS4 Regulates Anti-Seizure Effect of Endogenous Adenosine. *J. Neurosci.* **2012**, *32*, 2683–2695. [\[CrossRef\]](#) [\[PubMed\]](#)
102. Dhukhwa, A.; Al Aameri, R.F.H.; Sheth, S.; Mukherjea, D.; Rybak, L.; Ramkumar, V. Regulator of G protein signaling 17 represents a novel target for treating cisplatin induced hearing loss. *Sci. Rep.* **2021**, *11*, 8116. [\[CrossRef\]](#) [\[PubMed\]](#)
103. Han, B.R.; Lin, S.C.; Espinosa, K.; Thorne, P.R.; Vlajkovic, S.M. Inhibition of the Adenosine A_{2A} Receptor Mitigates Excitotoxic Injury in Organotypic Tissue Cultures of the Rat Cochlea. *Cells* **2019**, *8*, 877. [\[CrossRef\]](#)
104. Shin, M.; Pandya, M.; Espinosa, K.; Telang, R.; Boix, J.; Thorne, P.; Vlajkovic, S. Istradefylline Mitigates Age-Related Hearing Loss in C57BL/6J Mice. *Int. J. Mol. Sci.* **2021**, *22*, 8000. [\[CrossRef\]](#) [\[PubMed\]](#)