

The Expanding Field of Autoimmune Sleep–Wake Disorders—Implications for the Movement Disorders Clinical Practice

Devine MF, Feemster JC, Lieske EA, et al. Objective sleep profile in LGI1/CASPR2 autoimmunity. *Sleep* 2022; 45(2): zsab297.

Rapid eye movement (REM) sleep behavior disorder (RBD), a parasomnia, has long been considered to be a herald of neurodegenerative disease, in particular α -synucleinopathies. Sleep–wake disorders are now increasingly recognized in autoimmune neurological disease and may even lead to the discovery of new entities, like in the case of anti-IgLON5 disease.^{1,2} This increasing interest in autoimmune-mediated sleep–wake disorders led Devine and colleagues³ to retrospectively assess clinical and polysomnography (PSG) features in a cohort of nine patients with leucine-rich, glioma inactivated 1 (LGI1) and two with contactin-associated protein-like 2 (Caspr2) antibodies.

The overall spectrum of clinical sleep-related disturbances in this cohort encompassed insomnia, dream enactment behavior (DEB) (both together in 63.6% of cases), excessive daytime sleepiness (hypersomnia, 36.4%) and snoring (72.7%). DEB was only observed in the LGI1 subgroup (56%).

On PSG, 73% of patients were diagnosed with obstructive sleep apnea, and when compared to age and sex-matched controls, patients with LGI1/CASPR2 antibodies showed altered sleep architecture with increased N1 and decreased REM sleep. Increased rates of REM sleep without atonia (RSWA), a correlate of RBD, were only found in patients with LGI1 antibodies. In fact, 88.9% of LGI1+ subjects fulfilled the RSWA diagnostic thresholds for RBD—when calculated according to RSWA diagnostic standards established previously from the same research group⁴—although only one without DEB fulfilled the American Academy of Sleep medicine (AASM) criteria for RSWA in RBD.

Therefore, RSWA may have been overlooked with the sole use of AASM criteria. The notion that RSWA seems to be a key quantitative polysomnographic feature also in patients with LGI1 antibodies, is one of the main and most interesting findings of this first attempt to systematically and quantitatively assess RSWA in autoimmune neurological disease. However, sleep disorders and PSG findings should

only be looked in the context of the whole clinical phenotype.

In addition, in LGI1+/CASPR2+ patients sleep–wake symptoms preceded the neurological manifestations in 27.3%, developed in parallel in 45.5%, and occurred after the neurological manifestations in 18.1%. This might offer a window of opportunity for earlier recognition and treatment. Immunotherapy in this cohort was installed only after full clinical manifestation, but led to an overall neurological improvement and in 50% to an additional improvement of sleep–wake symptoms.

More research in this emerging area is needed. For example, actigraphy recordings rather than PSG could provide better real-life data, which may prove a useful marker for insomnia detection and monitoring. Given the frequent occurrence of cognitive impairments in these patients, objective measurements like the multiple sleep latency tests might be preferable to the Epworth sleepiness scale.

From a pathophysiological point of view, future studies of autoimmune-mediated sleep–wake disorders, in which targets of autoimmunity are known in detail, may facilitate a better understanding of the pathophysiology of specific sleep–wake disorders. For clinical practice, it is important to be aware that RSWA and sleep–wake disorders are not only heralding synuclein-related neurodegeneration, but also autoimmune disease (Table 1).

Author Roles

(1) Research Project: A. Conception, B. Organization, C. Execution; (2) Statistical analysis: A. Design, B. Execution, C. Review and critique; (3) Manuscript: A. Writing of the First Draft, B. Review and Critique.

E.E.: 1A, 1 C, 3A

C.B.: 3B

B. B.: 1A, 3B

***Correspondence to:** Dr. Evdokia Efthymiou, Department of Neurology, University Hospital Zurich (USZ), Frauenklinikstrasse 26, Zurich 8091, Switzerland; E-mail: evdokia.efthymiou@usz.ch

Keywords: LGI1/CASPR2, polysomnography, sleep disorders, RSWA. Relevant disclosures and conflict of interest are listed at the end of this article.

This is an open access article under the terms of the [Creative Commons Attribution–NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Received 21 March 2023; revised 13 June 2023; accepted 1 August 2023.

Published online 7 September 2023 in Wiley Online Library ([wileyonlinelibrary.com](https://www.wileyonlinelibrary.com)). DOI: 10.1002/mdc3.13869

TABLE 1 Summary of sleep symptoms and V-PSG findings in autoimmune encephalitis and neurodegenerative disease

Summary	Autoimmune encephalitis ²	Neurodegenerative disease ⁵
Sleep symptoms		
Insomnia	NMDAR (GluN1), AMPAR (GluA1, GluA2) ^a , LGI1, Caspr2, IgLON5, DPPX, mGluR5 ^b	PD, DLB, MSA, PSP, AD, HD
Hypersomnia (EDS)	NMDAR (GluN1), AMPAR (GluA1, GluA2) ^a , LGI1 ^a , IgLON5, DPPX ^a , mGluR5 ^b	PD, DLB, MSA, HD, ALS
V-PSG findings		
RBD	LGI1, IgLON5, DPPX ^a	PD (50%), DLB (80%), MSA (80%–100%), SCA3 (50%); (PSP, AD: very rare)
NREM parasomnia	NMDAR (GluN1) ^c , IgLON5	DLB ^c , AD ^c
Quasi-purposeful behaviors	Caspr2, IgLON5, DPPX ^a	None
Narcolepsy	Ma2, AQP4	None
Stridor	IgLON5	MSA, SCA1 + 3 ^a , ALS
OSAS	IgLON5, DPPX ^a	MSA, AD, ALS
PMLS	Caspr2, IgLON5, DPPX	PD, SCA 1–3 and 6
Ambiguous sleep	DPPX ^a	DLB

Note: Ambiguous sleep = sleep stages cannot be differentiated.

Abbreviations: V-PSG, video-polysomnography; PD, Parkinson's disease; DLB, dementia with Lewy bodies; MSA, multiple system atrophy; PSP, progressive supranuclear palsy; AD, Alzheimer's disease; HD, Huntington's disease; EDS, excessive daytime sleepiness; ALS, amyotrophic lateral sclerosis; RBD, REM sleep behavior disorder; REM, rapid eye movement; SCA, spinocerebellar ataxia; NREM, non-REM sleep; OSAS, obstructive sleep apnea; PMLS, periodic limb movements during sleep.

^aDescribed in single case reports or in small case series.²

^bDescribed in small case series.⁶

^cConfusional arousals.

Disclosures

Ethical Compliance Statement: The authors confirm that the approval of an institutional review board and patient consent were not required for this work.

Funding Sources and Conflicts of Interest: No specific funding was received for this work. The authors declare that there are no conflicts of interest relevant to this work.

Financial Disclosures for the Previous 12 Months: E.E. and C.R.B. reports no financial disclosures. B.B. has received royalties from Oxford University Press.

Acknowledgment

Open access funding provided by Universitat Zurich. ■

Evdokia Efthymiou, MD Christian R. Baumann, MD 

Bettina Balint, MD 

Department of Neurology, University Hospital Zurich, University of Zurich, Zurich, Switzerland

References

- Sabater L, Gaig C, Gelpi E, et al. A novel non-rapid-eye movement and rapid-eye-movement parasomnia with sleep breathing disorder associated with antibodies to IgLON5: a case series, characterisation of the antigen, and post-mortem study. *Lancet Neurol* 2014;13(6): 575–586.
- Muñoz-Lopetegui A, Graus F, Dalmau J, Santamaria J. Sleep disorders in autoimmune encephalitis. *Lancet Neurol* 2020;19(12): 1010–1022.
- Devine MF, Feemster JC, Lieske EA, McCarter SJ, Sandness DJ, Steele T. Objective sleep profile in LGI1/CASPR2 autoimmunity. *Sleep* 2022;45(2):zsab297.
- McCarter SJ, St Louis EK, Sandness DJ, Duwell EJ, Timm PC, Boeve BF. Diagnostic REM sleep muscle activity thresholds in patients with idiopathic REM sleep behavior disorder with and without obstructive sleep apnea. *Sleep Med* 2017;33:23–29.
- Iranzo A. Sleep in Neurodegenerative Diseases. *Sleep Med Clin* 2016; 11(1):1–18.
- Spatola M, Sabater L, Planagumà J, et al. Encephalitis with mGluR5 antibodies: symptoms and antibody effects. *Neurol* 2018;90: e1964–e1972.