

Published in final edited form as:

J Dev Biol. ; 1(2): 112–125. doi:10.3390/jdb1020112.

Epicardium Formation as a Sensor in Toxicology

Peter Hofsteen, Jessica Plavicki, Richard E. Peterson, and Warren Heideman*

Department of Pharmaceutical Sciences, University of Wisconsin, 777 Highland Ave, Madison, WI 53705, USA

Peter Hofsteen: phofsteen@wisc.edu; Jessica Plavicki: jsberndt@students.wisc.edu; Richard E. Peterson: repeterson@pharmacy.wisc.edu

Abstract

Zebrafish (*Danio rerio*) are an excellent vertebrate model for studying heart development, regeneration and cardiotoxicity. Zebrafish embryos exposed during the temporal window of epicardium development to the aryl hydrocarbon receptor (AHR) agonist 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) exhibit severe heart malformations. TCDD exposure prevents both proepicardial organ (PE) and epicardium development. Exposure later in development, after the epicardium has formed, does not produce cardiac toxicity. It is not until the adult zebrafish heart is stimulated to regenerate does TCDD again cause detrimental effects. TCDD exposure prior to ventricular resection prevents cardiac regeneration. It is likely that TCDD-induced inhibition of epicardium development and cardiac regeneration occur via a common mechanism. Here, we describe experiments that focus on the epicardium as a target and sensor of zebrafish heart toxicity.

Keywords

epicardium; proepicardium; TCDD; AHR; heart regeneration

1. Introduction

Heart disease remains the leading cause of death worldwide. In the United States, one third of all deaths can be attributed to cardiovascular disease [1]. It is believed that the etiology of heart disease results from a combination of factors including individual genetic makeup and either intentional or unintentional exposure to environmental factors such as tobacco smoke or environmental contaminants at different stages of life. Here, we describe recent work identifying the epicardium as the target for chemicals that cause heart failure in developing embryos.

© 2013 by the authors; licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

*Author to whom correspondence should be addressed; wheidema@wisc.edu; Fax: +1-608-262-5345.

Conflict of Interest

The authors declare no conflict of interest.

Although literature searches retrieve hundreds of articles using the keyword epicardium along with toxicity or chemical, the majority of these identify articles that are either about the epicardium as a surface for electrophysiological contact, or articles with the pericardium but no mention of the epicardium. Of the few remaining articles, most examine the epicardium in adults, rather than studies examining epicardium formation or even expansion or maintenance.

It has long been known that drugs used in treating cancer have effects on humans that include damage to the epicardium and other heart structures [2,3]. While not directly related to using the epicardium, it should be noted that the pig pericardium is of intense interest as a biomaterial used in valve reconstruction. Chemical treatment and effects on this pericardial tissue is of interest both as it pertains to human responses to latent material in the xenograft as well as the effectiveness in treatment in prolonging the material effectiveness, primarily reducing calcification [4–6]. Finally among asbestos workers, mesotheliomas grow on various internal tissues. Although primarily associated with the lung, these sometimes arise on the pericardium [7].

Mammalian models have been used to study various aspects of chemical interactions with the epicardium. Iminashi and Arita used ouabain to alter the action potential duration on the epicardium of Japanese monkeys [8]. In another study, the effects of kallikrein was studied on the adult dog epicardium, following the effects of exogenous bradykins and proteases as well as bradykin receptor antagonists on blood pressure and heart rate [9]. Isolated epicardial cells have been used to test the effects of 4-Aminopyridine in ischemia response models [10]. Canine epicardial cells have also been used to test the effects of different adrenergic receptor agonists and antagonists in dogs in the acute stage of Chagas disease [11]. Other studies with dog epicardium examined the effects of captopril on the production of free radicals after electric shock [12].

Others have used the epicardium to study the toxic effects of doxorubicin on conduction properties of the ventricle in the rat [13]. In another study examining mouse epicardial cells, arsenite was found to inhibit the epithelial to mesenchymal transition (EMT) of the epicardial cells, but not differentiation into smooth muscle cells [14]. Using the chick embryo as a model, Olivey *et al* found that transforming growth factor-beta 1 and 2 increase EMT in PE explants [15]. Mouse hearts have also been used as whole explants to follow the effects of antiarrhythmic drugs, often focusing on the excitability of the epicardial layer [16].

There are an increasing number of articles showing the ability of different chemicals to cause pericardial edema in developing fish [17–23]. As will be seen below, the pericardial edema may be a secondary response to the effects of the chemical on epicardium formation. Thus, fish models may prove very sensitive in identifying epicardium-toxic compounds.

Due to the paucity of reports investigating epicardium formation following exposure to environmental chemicals, we describe experiments conducted in our laboratory using zebrafish and mouse models that have identified the epicardium as a target underlying cardiotoxicity. As an example, we use 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), a

cardiotoxic environmental contaminant and prototypical AHR ligand to show how a common form of cardiotoxicity in zebrafish is connected to impaired epicardium development. We also discuss another cardiotoxic compound and non-AHR agonist, valproic acid (VPA), and its role in zebrafish epicardium development. We speculate that formation of the epicardium may be sensitive to disruption by environmental exposures during vertebrate development.

2. Zebrafish Heart and Epicardium Development

The zebrafish heart is made up of two chambers, an atrium and a ventricle. In addition, the bulbous arteriosus lies directly after the ventricle, storing pulse pressure in a role analogous to that of the mammalian aorta. At the inflow tract, the sinus venosus collects venous blood for delivery to the atrium. Blood flows from the sinus venosus through the two contractile chambers and out the bulbous arteriosus to the ventral aorta, which delivers blood to the pharyngeal arch arteries [24].

The vertebrate heart is the first internal organ to form and function [25]. For detailed reviews about early zebrafish heart development see [26,27]. Briefly, in the earliest stages of zebrafish heart development, just prior to gastrulation, bilateral clusters of cardiac progenitor cells are located in the lateral marginal zone. During gastrulation, cardiac progenitors migrate to the anterior lateral plate mesoderm where they are positioned near the midbrain-hindbrain boundary. Myocardial and endocardial precursors converge at the embryonic midline forming a cardiac disc at 19 hours post fertilization (hpf). The cardiac disc undergoes morphogenetic movements to form a contractile linear heart tube (24–28 hpf), which consists of an inner endocardial lining continuous with the vascular endothelium and an overlying myocardial layer. Distinct ventricular and atrial chambers form (30 hpf), the heart tube loops (36 hpf) and the atrioventricular (AV) canal becomes visible. Between the chambers, the endocardium gives rise to endocardial cushions, which are later elaborated into cardiac valves. Proper valve development is essential for directional blood flow and heart function. Continued development of the zebrafish heart is highly dependent on the formation of the third and outer most layer of cardiac tissue, the epicardium.

The epicardium (*epi*: outer, *cardium*: heart) is a mesodermally derived squamous epithelium that is critical for heart development, function and regeneration. Unlike the cells that form the heart tube, the epicardium is derived from a transient pool of progenitors located outside of the heart field, termed the proepicardium (PE) [28–30]. PE cells migrate towards the looped heart and envelop the naked myocardium. In chicks, following initial epicardial coverage of the heart, a subset of epicardial cells undergoes an epithelial-to-mesenchymal transition (EMT) to become epicardial derived progenitor cells (EPDCs; reviewed in [31]). EPDCs invade the underlying myocardium and differentiate directly to become vascular smooth muscle cells and cardiac fibroblasts. Furthermore, EPDCs play important roles in heart looping, valve and coronary vasculature development, cardiac morphogenesis, cardiomyocyte alignment and proliferation, and maturation of the cardiac conduction system (reviewed in [31–33]). Furthermore, it is thought that epicardial cells retain plasticity as cardiac stem cells that can assist adult cardiac regeneration [33–35]. However, the formation or role EPDCs in zebrafish remains undetermined.

3. Zebrafish as a Model to Study Cardiotoxicity

Zebrafish are an excellent vertebrate model for studying human health and disease. Development of zebrafish internal organs can be monitored *in vivo* because embryos are translucent and develop externally. These attributes combined with high fecundity and potential for high throughput screening make the zebrafish an ideal model to study heart development and chemical toxicity [27,36]. Zebrafish were first developed as a genetic model, and are highly amenable to forward and reverse genetic screens. These screens have yielded much new information about heart development. Identifying genes required for heart development is greatly facilitated by the fact that zebrafish embryo-larvae are able to rely on passive oxygen diffusion and do not require heart function during the first week of life [37–39]. This is very important because it means that one can follow the effects of mutations interfering with heart function as the heart ceases to function. In mammals, these mutations often produce immediate death of the fetus at the onset of effect.

4. DLCs and Heart Defects

Halogenated aromatic hydrocarbons (HAH) are ubiquitous environmental contaminants produced in industrial manufacturing processes and combustion. Included in this class are polychlorinated biphenyls, polychlorinated dibenzofurans, and polychlorinated dibenzo-*p*-dioxins. A subset of these different chemicals shares a similar molecular shape, matching the binding site of the ligand-activated aryl hydrocarbon receptor (AHR). The prototype agonist for the AHR is 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD), commonly referred to as “dioxin”. Thus, members of this group of AHR agonists can be referred to as dioxin-like compounds (DLCs).

Embryonic exposure to TCDD disrupts cardiac development and function in fish, birds, and mammals [40–45]. TCDD is known to be highly toxic. In humans, epidemiological studies have linked TCDD exposure to congenital heart defects such as hypoplastic left heart syndrome [46,47] and ischemic heart disease [48,49].

The most sensitive organisms to the effects of DLCs are fish, especially fish in early life stages [42,50]. Using zebrafish as a model, it was quickly determined that TCDD exposure during early development causes decreased cardiomyocyte proliferation, a block and lack of erythrocyte development, reduced blood flow and cardiac output, and lack endocardial valve cushions. Ultimately, this leads to ventricular standstill and death [43,51–55]. Understanding why TCDD and related AHR agonists cause cardiotoxicity has been a question for quite some time.

5. TCDD and the Zebrafish Epicardium

In zebrafish, the temporal window of cardiotoxicity is peculiar. The first 48 hours of cardiac development proceeds normally when zebrafish are exposed to TCDD immediately following fertilization. The primitive heart tube forms, the heart loops and heart function is indistinguishable from controls. Circulation appears normal at 48 hpf in these animals. After 48 hpf, TCDD treated hearts diverge from experimental controls: the heart unloops and elongates, the ventricle becomes constricted, and pronounced pericardial edema develops

[53]. This unusual sensitivity to the cardiotoxic effects of TCDD corresponds to the developmental period in which the PE is specified and migrates over the myocardium to form the epicardium [28,56]. Furthermore, a similar phenotype was observed in the zebrafish *heartstrings* mutant [57]. These fish lack *tbx5a*, which was later shown to be required for PE specification [58].

We now know that TCDD-induced activation of AHR prevents the development of both the PE and epicardium. This results in severe heart malformations that culminate in death [56]. This explains the peculiar lag in toxicity until approximately 48 hpf. Furthermore, after the epicardium has formed it is no longer affected by TCDD; in adults a lethal TCDD exposure fails to produce cardiotoxicity [50,59]. The failure to form an epicardium explains most of the heart malformations caused by TCDD.

Because many other chemicals produce a syndrome including pericardial edema and heart malformations in zebrafish, the epicardium may be useful as a sensor of chemical cardiotoxicity. Furthermore, disruption of epicardial development may prove to be a common cause of cardiotoxicity in developing vertebrates.

6. TCDD and the Murine Epicardium

Consistent with the high sensitivity that developing fish have to DLCs, the mouse heart is less sensitive to embryonic TCDD exposure than the zebrafish heart. Unlike zebrafish, TCDD-exposed mice do not exhibit pronounced heart malformations; however, measurable defects such as bradycardia and cardiac hypertrophy have been observed [40]. Additionally, in the fetal murine heart, TCDD-exposure causes global transcriptomic changes in genes that regulate the cell cycle and extracellular matrix [60]. In contrast to the effect in fish, the changes in the mouse heart are not embryonically lethal. Whether they reduce fitness or affect adult health is not known.

We have recently found that TCDD exposure does affect the developing mouse epicardium. While TCDD did not prevent epicardium formation in the developing mouse, the epicardium appeared detached from the underlying myocardium (Figure 1). Epicardial detachment has also been observed in mice carrying mutations in RXR-alpha [61], Tcf21 [62], PDGF [63], and Flrt2 [64]. It is generally thought that epicardial detachment in these mutants is due to defects in extracellular matrix composition and EMT processes. Why the epicardium responses differ between mouse and zebrafish is an intriguing question for further investigation.

7. The Zebrafish Epicardium as a Sentinel for Other Cardiotoxic Compounds

Pericardial edema coupled with an elongated, unlooped heart is a phenotype that is not unique to fish exposed to DLCs. This syndrome has been observed in zebrafish carrying mutations that disrupt the cardiovascular and other systems, as well as following exposure to a variety of cardiotoxic chemicals [65–67]. Based on our results with TCDD, a possible link between these factors and the syndrome of heart malformation and pericardial edema could

be effects on epicardium formation. We have made a preliminary test of this idea using valproic acid (VPA), a compound that is not a DLC, yet produces the type of cardiac malformations described for TCDD. VPA is an anticonvulsant known to affect cardiovascular development in zebrafish [68]. Despite critical differences in the mechanism of action, both agents produced an elongated heart with a thin atrium, a compacted ventricle, and failure of epicardium development (Figure 2).

Of particular interest, VPA-exposed zebrafish appeared completely normal until the PE to epicardium transition, in which the cluster of cells forming the PE begins to cross over to the heart to form the epicardium. In VPA-treated zebrafish, there were no signs of pericardial edema during the period from 0–72 hpf. While the PE formed in these fish, there was no progression to forming the epicardium. In the ensuing period from ~72–120 hpf, during which the epicardial cells would normally envelop the heart, the hearts in the VPA-treated zebrafish became unlooped and elongated. This was accompanied by marked pericardial edema.

While the mechanism remains a mystery, epicardium formation was sensitive to blockade by two compounds with very distinct mechanisms of action. In both cases, the initial patterning of the heart was unaffected, and cardiotoxicity was only manifested at the point at which epicardial cells normally form. It may be that epicardium formation is more sensitive to disruption than other processes needed to produce a healthy functional heart. If so, the epicardium could be a sentinel for cardiotoxicity.

8. The Epicardium and Heart Regeneration

Following myocardial infarction, the human heart can lose billions of cardiomyocytes, resulting in the loss of a significant portion of myocardium [69]. The damaged myocardium is not replaced with new myocardial cells, but rather healed by formation of non-contractile scar tissue. This inability to replace lost heart tissue with contractile myocardium is a critical medical problem [70,71].

The epicardium has received great interest in the field of regenerative medicine due to the plasticity of epicardial progenitors (reviewed in [32–35,72]). Furthermore, many reports have identified the epicardium as a critical signaling center following myocardial injury.

Unlike mammals, zebrafish retain the ability to regenerate damaged myocardial tissue into adulthood [73]. This occurs mainly by differentiation and proliferation of spared resident cardiomyocytes [74,75]. Following surgical resection of the adult zebrafish ventricle at its apex, profuse bleeding occurs followed by the formation of a fibrin-rich blood clot. Subsequently, an epicardial-derived white epithelial sheath of tissue migrates and surrounds the newly formed blood clot [76]. The combination of sealing the wound with epithelial-like sheath of tissue, cardiomyocyte dedifferentiation and proliferation orchestrate regeneration of a new contractile myocardium over a period of 1–2 months.

During heart regeneration in zebrafish, the epicardium is thought to revert to an “embryonic-like” state and act as a signaling center by expressing embryonic genes such as the retinoic acid (RA) synthesizing enzyme, *raldh2* [76,77]. Expression of *raldh2* in

epicardial and endocardial cells during regeneration is thought to secrete trophic RA to assist in cardiomyocyte proliferation. Inhibition of RA signaling during heart regeneration reversed the regenerative capacity by halting cardiomyocyte proliferation [78]. While controversy remains regarding the role of the epicardium during heart regeneration, it appears the adult epicardium may play a supportive role by acting as a signaling center to direct differentiation, proliferation and recruitment of cardiac progenitors to the wound site. Thus, embryonic-like epicardial cells may be needed for wound healing in the adult heart.

Once the epicardium has formed, the heart appears to no longer be a target for DLCs in zebrafish, and even when exposed to a lethal concentration of TCDD the adult heart is unaffected [50,59]. However, as indicated above, embryonic-like epicardial progenitor cells are required for heart wound healing in the adult zebrafish. It is perhaps then not surprising that during wounding, the adult zebrafish regains sensitivity to TCDD. In experiments in which the apex of the adult ventricle were amputated, control hearts quickly regenerated the missing tissue, while TCDD-pretreated hearts did not progress past the initial stage of blood clot formation [59]. TCDD prevented the formation of the epicardially-derived sheath of tissue that normally envelops the wound, and reduced cardiomyocyte proliferation at the wound site. If TCDD was administered one day following surgical amputation, the regenerative capacity was not affected; the epithelial-like sheath of tissue formed and the blood clot was replaced with new contractile myocardium. This suggests that activated epicardial progenitors in the adult partially amputated zebrafish heart are targeted by TCDD.

9. Conclusions

The epicardium and epicardial progenitors are potentially vital for development, function and regeneration of the vertebrate heart. In zebrafish, the epicardium appears to be sensitive to disruption by xenobiotic chemicals during development, but not once formed. In humans, long-term exposure to TCDD is associated with ischemic heart disease [48,49]. It is interesting that other studies of ischemic heart disease have found a reduction in epicardial cell numbers in affected patients [48,49,79]. We speculate that epicardium development may play an important role in environmentally induced heart disease, worth further study. The zebrafish heart is now being used to screen drug candidates for Q-T prolongation and other forms of cardiotoxicity [80–82]. Due to high throughput screening capability, zebrafish have many advantages in screening for teratogenic effects over other vertebrate models. It may be that the tools available with the zebrafish—transgenic lines that mark the epicardium, translucent external development, and high fecundity—will make following epicardium development an important screen as well. Thus, we may see future use for epicardium formation as a marker of cardiotoxicity.

Acknowledgments

We would like to thank Dorothy Nesbit for technical services and fish husbandry, Andrew Schneider for assisting in procedures with mice, Henry Saint Clair Fredericks, and the entire zebrafish group for helpful discussions.

Supported by the National Institutes of Health (NIH) grant R01 ES012716 from the National Institute of Environmental Health Sciences (NIEHS) and the University of Wisconsin Sea Grant Institute, National Sea Grant College Program, National Oceanic and Atmospheric Administration, U.S. Department of Commerce grant number NA 16RG2257, Sea Grant Project R/BT-22 and 25.

References and Notes

1. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N, Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A, McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W, Sacco R, Sorlie P, Stafford R, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J. Executive summary: Heart disease and stroke statistics--2010 update: A report from the American Heart Association. *Circulation*. 2010; 121:948–954. [PubMed: 20177011]
2. Tsibiribi P, Bui-Xuan C, Bui-Xuan B, Lombard-Bohas C, Duperret S, Belkhiria M, Tabib A, Maujean G, Descotes J, Timour Q. Cardiac lesions induced by 5-fluorouracil in the rabbit. *Hum Exp Toxicol*. 2006; 25:305–309. [PubMed: 16866187]
3. Yeh ET, Tong AT, Lenihan DJ, Yusuf SW, Swafford J, Champion C, Durand JB, Gibbs H, Zafarmand AA, Ewer MS. Cardiovascular complications of cancer therapy: diagnosis, pathogenesis, and management. *Circulation*. 2004; 109:3122–3131. [PubMed: 15226229]
4. Guldner NW, Bastian F, Weigel G, Zimmermann H, Maleika M, Scharfschwerdt M, Rohde D, Sievers HH. Nanocoating with titanium reduces iC3b- and granulocyte-activating immune response against glutaraldehyde-fixed bovine pericardium: A new technique to improve biologic heart valve prosthesis durability? *J Thorac Cardiovasc Surg*. 2012; 143:1152–1159. [PubMed: 22153860]
5. Guldner NW, Jasmund I, Zimmermann H, Heinlein M, Girndt B, Meier V, Fluss F, Rohde D, Gebert A, Sievers HH. Detoxification and endothelialization of glutaraldehyde-fixed bovine pericardium with titanium coating: A new technology for cardiovascular tissue engineering. *Circulation*. 2009; 119:1653–1660. [PubMed: 19289635]
6. Neethling WM, Glancy R, Hodge AJ. Mitigation of calcification and cytotoxicity of a glutaraldehyde-preserved bovine pericardial matrix: Improved biocompatibility after extended implantation in the subcutaneous rat model. *J Heart Valve Dis*. 2010; 19:778–785. [PubMed: 21214104]
7. Abejie BA, Chung EH, Nesto RW, Kales SN. Grand rounds: asbestos-related pericarditis in a boiler operator. *Environ Health Perspect*. 2008; 116:86–89. [PubMed: 18197304]
8. Imanishi S, Arita M. Electrophysiologic properties differ in the ventricular endocardium and epicardium of the Japanese monkey. *J Electrocardiol*. 1987; 20:185–192. [PubMed: 3655590]
9. Staszewska-Woolley J, Woolley G. Participation of the kallikrein-kinin-receptor system in reflexes arising from neural afferents in the dog epicardium. *J Physiol*. 1989; 419:33–44. [PubMed: 2576069]
10. Lukas A, Antzelevitch C. Differences in the electrophysiological response of canine ventricular epicardium and endocardium to ischemia. Role of the transient outward current. *Circulation*. 1993; 88:2903–2915. [PubMed: 8252704]
11. Han W, Barr SC, Pacioretty LM, Gilmour RF Jr. Restoration of the transient outward potassium current by noradrenaline in chagasic canine epicardium. *J Physiol*. 1997; 500:75–83. [PubMed: 9097934]
12. Pagan-Carlo LA, Garcia LA, Hutchison JL, Buettner GR, Kerber RE. Captopril lowers coronary venous free radical concentration after direct current cardiac shocks. *Chest*. 1999; 116:484–487. [PubMed: 10453880]
13. Kharin SN, Krandycheva VV, Strelkova MV, Tsvetkova AS, Shmakov DN. Doxorubicin-induced changes of ventricular repolarization heterogeneity: Results of a chronic rat study. *Cardiovasc Toxicol*. 2012; 12:312–317. [PubMed: 22618330]
14. Allison P, Huang T, Broka D, Parker P, Barnett JV, Camenisch TD. Disruption of canonical TGFbeta-signaling in murine coronary progenitor cells by low level arsenic. *Toxicol Appl Pharmacol*. 2013;10.1016/j.taap.2013.04.035
15. Olivey HE, Mundell NA, Austin AF, Barnett JV. Transforming growth factor-beta stimulates epithelial-mesenchymal transformation in the proepicardium. *Dev Dyn*. 2006; 235:50–59. [PubMed: 16245329]
16. Sabir IN, Li LM, Jones VJ, Goddard CA, Grace AA, Huang CL. Criteria for arrhythmogenicity in genetically-modified Langendorff-perfused murine hearts modelling the congenital long QT

syndrome type 3 and the Brugada syndrome. *Pflugers Arch.* 2008; 455:637–651. [PubMed: 17805561]

17. Tu W, Niu L, Liu W, Xu C. Embryonic exposure to butachlor in zebrafish (*Danio rerio*): endocrine disruption, developmental toxicity and immunotoxicity. *Ecotoxicol Environ Saf.* 2013; 89:189–195. [PubMed: 23294635]
18. Li X, Ma Y, Li D, Gao X, Li P, Bai N, Luo M, Tan X, Lu C, Ma X. Arsenic impairs embryo development via down-regulating *Dvr1* expression in zebrafish. *Toxicol Lett.* 2012; 212:161–168. [PubMed: 22613031]
19. Incardona JP, Linbo TL, Scholz NL. Cardiac toxicity of 5-ring polycyclic aromatic hydrocarbons is differentially dependent on the aryl hydrocarbon receptor 2 isoform during zebrafish development. *Toxicol Appl Pharmacol.* 2011; 257:242–249. [PubMed: 21964300]
20. Valesio EG, Zhang H, Zhang C. Exposure to the JNK inhibitor SP600125 (anthracycline) during early zebrafish development results in morphological defects. *J Appl Toxicol.* 2013; 33:32–40. [PubMed: 21751222]
21. Lin CC, Hui MN, Cheng SH. Toxicity and cardiac effects of carbaryl in early developing zebrafish (*Danio rerio*) embryos. *Toxicol Appl Pharmacol.* 2007; 222:159–168. [PubMed: 17559897]
22. Reimers MJ, Flockton AR, Tanguay RL. Ethanol- and acetaldehyde-mediated developmental toxicity in zebrafish. *Neurotoxicol Teratol.* 2004; 26:769–781. [PubMed: 15451041]
23. Lefebvre KA, Trainer VL, Scholz NL. Morphological abnormalities and sensorimotor deficits in larval fish exposed to dissolved saxitoxin. *Aquat Toxicol (Amsterdam, Netherlands).* 2004; 66:159–170.
24. Hu N, Sedmera D, Yost HJ, Clark EB. Structure and function of the developing zebrafish heart. *Anat Rec.* 2000; 260:148–157. [PubMed: 10993952]
25. Lee RK, Stainier DY, Weinstein BM, Fishman MC. Cardiovascular development in the zebrafish. II. Endocardial progenitors are sequestered within the heart field. *Development.* 1994; 120:3361–3366. [PubMed: 7821208]
26. Stainier DY. Zebrafish genetics and vertebrate heart formation. *Nat Rev Genet.* 2001; 2:39–48. [PubMed: 11253067]
27. Bakkers J. Zebrafish as a model to study cardiac development and human cardiac disease. *Cardiovasc Res.* 2011; 91:279–288. [PubMed: 21602174]
28. Serluca FC. Development of the proepicardial organ in the zebrafish. *Dev Biol.* 2008; 315:18–27. [PubMed: 18206866]
29. Manner J, Perez-Pomares JM, Macias D, Munoz-Chapuli R. The origin, formation and developmental significance of the epicardium: a review. *Cells Tissues Organs.* 2001; 169:89–103. [PubMed: 11399849]
30. Manner J, Schlueter J, Brand T. Experimental analyses of the function of the proepicardium using a new microsurgical procedure to induce loss-of-proepicardial-function in chick embryos. *Dev Dyn.* 2005; 233:1454–1463. [PubMed: 15977171]
31. Lie-Venema H, van den Akker NM, Bax NA, Winter EM, Maas S, Kekarainen T, Hoeben RC, deRuiter MC, Poelmann RE, Gittenberger-de Groot AC. Origin, fate, and function of epicardium-derived cells (EPDCs) in normal and abnormal cardiac development. *Sci World J.* 2007; 7:1777–1798.
32. Gittenberger-de Groot AC, Winter EM, Bartelings MM, Jose Goumans M, Deruiter MC, Poelmann RE. The arterial and cardiac epicardium in development, disease and repair. *Differentiation.* 2012; 84:41–53. [PubMed: 22652098]
33. Riley PR. An epicardial floor plan for building and rebuilding the mammalian heart. *Curr Top Dev Biol.* 2012; 100:233–251. [PubMed: 22449846]
34. Smart N, Riley PR. The epicardium as a candidate for heart regeneration. *Future Cardiol.* 2012; 8:53–69. [PubMed: 22185446]
35. Smart N, Dube KN, Riley PR. Epicardial progenitor cells in cardiac regeneration and neovascularisation. *Vascul Pharmacol.* 2012
36. Hill AJ, Teraoka H, Heideman W, Peterson RE. Zebrafish as a model vertebrate for investigating chemical toxicity. *Toxicol Sci.* 2005; 86:6–19. [PubMed: 15703261]

37. Wells P, Pinder A. The respiratory development of Atlantic salmon. II. Partitioning of oxygen uptake among gills, yolk sac and body surfaces. *J Exp Biol.* 1996; 199:2737–2744. [PubMed: 9320645]
38. Kopp R, Schwerte T, Pelster B. Cardiac performance in the zebrafish breakdance mutant. *J Exp Biol.* 2005; 208:2123–2134. [PubMed: 15914656]
39. Stainier DY, Fouquet B, Chen JN, Warren KS, Weinstein BM, Meiler SE, Mohideen MA, Neuhauss SC, Solnica-Krezel L, Schier AF, Zwartkruis F, Stemple DL, Malicki J, Driever W, Fishman MC. Mutations affecting the formation and function of the cardiovascular system in the zebrafish embryo. *Development (Cambridge, England).* 1996; 123:285–292.
40. Thackaberry EA, Nunez BA, Ivnitski-Steele ID, Friggins M, Walker MK. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on murine heart development: Alteration in fetal and postnatal cardiac growth, and postnatal cardiac chronotropy. *Toxicol Sci.* 2005; 88:242–249. [PubMed: 16120746]
41. Cheung MO, Gilbert EF, Peterson RE. Cardiovascular teratogenicity of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin in the chick embryo. *Toxicol Appl Pharmacol.* 1981; 61:197–204. [PubMed: 7324065]
42. Peterson RE, Theobald HM, Kimmel GL. Developmental and reproductive toxicity of dioxins and related compounds: cross-species comparisons. *Crit Rev Toxicol.* 1993; 23:283–335. [PubMed: 8260069]
43. Carney SA, Chen J, Burns CG, Xiong KM, Peterson RE, Heideman W. Aryl hydrocarbon receptor activation produces heart-specific transcriptional and toxic responses in developing zebrafish. *Mol Pharmacol.* 2006; 70:549–561. [PubMed: 16714409]
44. King-Heiden TC, Mehta V, Xiong KM, Lanham KA, Antkiewicz DS, Ganser A, Heideman W, Peterson RE. Reproductive and developmental toxicity of dioxin in fish. *Mol Cell Endocrinol.* 2012; 354:121–138. [PubMed: 21958697]
45. Kopf PG, Walker MK. Overview of developmental heart defects by dioxins, PCBs, and pesticides. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev.* 2009; 27:276–285. [PubMed: 19953399]
46. Cronk CE, Pelech AN, Malloy ME, McCarver DG. Excess birth prevalence of Hypoplastic Left Heart syndrome in eastern Wisconsin for birth cohorts 1997–1999. *Birth Defects Res.* 2004; 70:114–120.
47. Kuehl KS, Loffredo CA. A cluster of hypoplastic left heart malformation in Baltimore, Maryland. *Pediatric Cardiol.* 2006; 27:25–31.
48. Flesch-Janys D, Berger J, Gurn P, Manz A, Nagel S, Waltsgott H, Dwyer JH. Exposure to polychlorinated dioxins and furans (PCDD/F) and mortality in a cohort of workers from a herbicide-producing plant in Hamburg, Federal Republic of Germany. *Am J Epidemiol.* 1995; 142:1165–1175. [PubMed: 7485063]
49. Bertazzi PA, Bernucci I, Brambilla G, Consonni D, Pesatori AC. The Seveso studies on early and long-term effects of dioxin exposure: a review. *Environ Health Perspect.* 1998; 106(Suppl 2):625–633. [PubMed: 9599710]
50. Lanham KA, Peterson RE, Heideman W. Sensitivity to dioxin decreases as zebrafish mature. *Toxicol Sci.* 2012; 127:360–370. [PubMed: 22403156]
51. Henry TR, Spitsbergen JM, Hornung MW, Abnet CC, Peterson RE. Early life stage toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in zebrafish (*Danio rerio*). *Toxicol Appl Pharmacol.* 1997; 142:56–68. [PubMed: 9007034]
52. Dong W, Teraoka H, Tsujimoto Y, Stegeman JJ, Hiraga T. Role of aryl hydrocarbon receptor in mesencephalic circulation failure and apoptosis in zebrafish embryos exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicol Sci.* 2004; 77:109–116. [PubMed: 14657521]
53. Antkiewicz DS, Burns CG, Carney SA, Peterson RE, Heideman W. Heart malformation is an early response to TCDD in embryonic zebrafish. *Toxicol Sci.* 2005; 84:368–377. [PubMed: 15635151]
54. Belair CD, Peterson RE, Heideman W. Disruption of erythropoiesis by dioxin in the zebrafish. *Dev Dyn.* 2001; 222:581–594. [PubMed: 11748828]
55. Mehta V, Peterson RE, Heideman W. 2,3,7,8-Tetrachlorodibenzo-p-dioxin Exposure Prevents Cardiac Valve Formation in Developing Zebrafish. *Toxicol Sci.* 2008; 104:303–311. [PubMed: 18477685]

56. Plavicki J, Hofsteen P, Peterson RE, Heideman W. Dioxin inhibits zebrafish epicardium and proepicardium development. *Toxicol Sci.* 2013; 131:558–567. [PubMed: 23135548]
57. Garrity DM, Childs S, Fishman MC. The heartstrings mutation in zebrafish causes heart/fin Tbx5 deficiency syndrome. *Development.* 2002; 129:4635–4645. [PubMed: 12223419]
58. Liu J, Stainier DY. Tbx5 and Bmp signaling are essential for proepicardium specification in zebrafish. *Circulation Res.* 2010; 106:1818–1828. [PubMed: 20413782]
59. Hofsteen P, Mehta V, Kim MS, Peterson RE, Heideman W. TCDD inhibits heart regeneration in adult zebrafish. *Toxicol Sci.* 2013; 132:211–221. [PubMed: 23204111]
60. Thackaberry EA, Jiang Z, Johnson CD, Ramos KS, Walker MK. Toxicogenomic profile of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the murine fetal heart: Modulation of cell cycle and extracellular matrix genes. *Toxicol Sci.* 2005; 88:231–241. [PubMed: 16120747]
61. Jenkins SJ, Hutson DR, Kubalak SW. Analysis of the proepicardium-epicardium transition during the malformation of the RXRalpha-/- epicardium. *Dev Dyn.* 2005; 233:1091–1101. [PubMed: 15861408]
62. Acharya A, Baek ST, Huang G, Eskioçak B, Goetsch S, Sung CY, Banfi S, Sauer MF, Olsen GS, Duffield JS, Olson EN, Tallquist MD. The bHLH transcription factor Tcf21 is required for lineage-specific EMT of cardiac fibroblast progenitors. *Development.* 2012; 139:2139–2149. [PubMed: 22573622]
63. Smith CL, Baek ST, Sung CY, Tallquist MD. Epicardial-derived cell epithelial-to-mesenchymal transition and fate specification require PDGF receptor signaling. *Circulation Res.* 2011; 108:e15–e26. [PubMed: 21512159]
64. Muller PS, Schulz R, Maretto S, Costello I, Srinivas S, Bikoff E, Robertson E. The fibronectin leucine-rich repeat transmembrane protein Flrt2 is required in the epicardium to promote heart morphogenesis. *Development.* 2011; 138:1297–1308. [PubMed: 21350012]
65. Muto A, Calof AL, Lander AD, Schilling TF. Multifactorial origins of heart and gut defects in nipbl-deficient zebrafish, a model of Cornelia de Lange Syndrome. *PLoS Biol.* 2011; 9:e1001181. [PubMed: 22039349]
66. Mitchell IC, Brown TS, Terada LS, Amatruda JF, Nwariaku FE. Effect of vascular cadherin knockdown on zebrafish vasculature during development. *PLoS One.* 2010; 5:e8807. [PubMed: 20098718]
67. Pruvot B, Quiroz Y, Voncken A, Jeanray N, Piot A, Martial JA, Muller M. A panel of biological tests reveals developmental effects of pharmaceutical pollutants on late stage zebrafish embryos. *Reprod Toxicol.* 2012; 34:568–583. [PubMed: 22982570]
68. Gurvich N, Berman MG, Wittner BS, Gentleman RC, Klein PS, Green JB. Association of valproate-induced teratogenesis with histone deacetylase inhibition *in vivo*. *Faseb J.* 2005; 19:1166–1168. [PubMed: 15901671]
69. Reinecke H, Minami E, Zhu WZ, Laflamme MA. Cardiogenic differentiation and transdifferentiation of progenitor cells. *Circ Res.* 2008; 103:1058–1071. [PubMed: 18988903]
70. Malki Q, Sharma ND, Afzal A, Ananthsubramaniam K, Abbas A, Jacobson G, Jafri S. Clinical presentation, hospital length of stay, and readmission rate in patients with heart failure with preserved and decreased left ventricular systolic function. *Clin Cardiol.* 2002; 25:149–152. [PubMed: 12000071]
71. Bolli P, Chaudhry HW. Molecular physiology of cardiac regeneration. *Ann N Y Acad Sci.* 2010; 1211:113–126. [PubMed: 21062300]
72. Smart N, Bollini S, Dube KN, Vieira JM, Zhou B, Riegler J, Price AN, Lythgoe MF, Davidson S, Yellon D, Pu WT, Riley PR. Myocardial regeneration: expanding the repertoire of thymosin beta4 in the ischemic heart. *Ann N Y Acad Sci.* 2012; 1269:92–101. [PubMed: 23045976]
73. Poss KD, Wilson LG, Keating MT. Heart regeneration in zebrafish. *Science.* 2002; 298:2188–2190. [PubMed: 12481136]
74. Kikuchi K, Holdway JE, Werdich AA, Anderson RM, Fang Y, Egnaczyk GF, Evans T, Macrae CA, Stainier DY, Poss KD. Primary contribution to zebrafish heart regeneration by gata4(+) cardiomyocytes. *Nature.* 2010; 464:601–605. [PubMed: 20336144]

75. Jopling C, Sleep E, Raya M, Marti M, Raya A, Belmonte JC. Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature*. 2010; 464:606–609. [PubMed: 20336145]
76. Lepilina A, Coon AN, Kikuchi K, Holdway JE, Roberts RW, Burns CG, Poss KD. A dynamic epicardial injury response supports progenitor cell activity during zebrafish heart regeneration. *Cell*. 2006; 127:607–619. [PubMed: 17081981]
77. Choi WY, Poss KD. Cardiac regeneration. *Curr Top Dev Biol*. 2012; 100:319–344. [PubMed: 22449849]
78. Kikuchi K, Holdway JE, Major RJ, Blum N, Dahn RD, Begemann G, Poss KD. Retinoic acid production by endocardium and epicardium is an injury response essential for zebrafish heart regeneration. *Dev Cell*. 2011; 20:397–404. [PubMed: 21397850]
79. Di Meglio F, Castaldo C, Nurzynska D, Romano V, Miraglia R, Montagnani S. Epicardial cells are missing from the surface of hearts with ischemic cardiomyopathy: A useful clue about the self-renewal potential of the adult human heart? *Int J Cardiol*. 2010; 145:e44–e46. [PubMed: 19176252]
80. Milan DJ, Jones IL, Ellinor PT, MacRae CA. In vivo recording of adult zebrafish electrocardiogram and assessment of drug-induced QT prolongation. *Am J Physiol Heart Circ Physiol*. 2006; 291:H269–H273. [PubMed: 16489111]
81. Park MJ, Lee KR, Shin DS, Chun HS, Kim CH, Ahn SH, Bae MA. Predicted drug-induced bradycardia related cardio toxicity using a zebrafish in vivo model is highly correlated with results from in vitro tests. *Toxicol Lett*. 2013; 216:9–15. [PubMed: 23137660]
82. Parg C, Seng WL, Semino C, McGrath P. Zebrafish: a preclinical model for drug screening. *Assay Drug Dev Technol*. 2002; 1:41–48. [PubMed: 15090155]
83. Kikuchi K, Gupta V, Wang J, Holdway JE, Wills AA, Fang Y, Poss KD. tcf21+ epicardial cells adopt non-myocardial fates during zebrafish heart development and regeneration. *Development* (Cambridge, England). 2011; 138:2895–2902.




Figure 1.

TCDD exposure causes epicardium detachment from the underlying myocardium in developing mice. Murine dams were exposed to vehicle control (corn oil) or TCDD (5 $\mu\text{g/g}$) at E7 and embryos were collected at E12 for sectional hematoxylin and eosin staining (n = 3/group). Shown are representative sections of control and TCDD hearts at the epicardium. Black arrows denote epicardial cells and their detachment from myocardium.




Figure 2.

VPA exposed zebrafish larvae lack epicardium. Embryos from a zebrafish epicardial reporter line (*tcf21:DsRed*; [83]) were exposed to VPA (0.06 μ M), or vehicle (EtOH) shortly following fertilization. The control is shown at left, while the VPA heart is shown at right. The presence or absence of epicardium was assessed with immunohistochemistry at 120 hours post fertilization using an antibody against *tcf21*, expressed in epicardial cells (Red). Cell nuclei are stained with DAPI shown as blue. Cytoplasmic *cmlc2:GFP* shown as green outlines the cardiomyocytes. Representative single plane confocal microscopy images (n = 6/group) are shown with anterior to the left (V = ventricle, A = atrium).