

# Malaria Endemicity in the Rural Communities of Ebonyi State, Nigeria

David Ekene Nwele<sup>1,\*</sup> , Ikechukwu Oliver Onyali<sup>2</sup>, Milliam Okwudili Iwueze<sup>2</sup>, Michael Okpara Elom<sup>3</sup>, Ogbonna Elom Sabastian Uguru<sup>4</sup>

<sup>1</sup>Department of Applied Biology, Faculty of Science, Ebonyi State University, Abakaliki, Nigeria; <sup>2</sup>Department of Parasitology and Entomology, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Nigeria; <sup>3</sup>Department of Medical Laboratory Science, Faculty of Health Sciences, Ebonyi State University, Abakaliki, Nigeria; <sup>4</sup>Department of Medical Microbiology, Faculty of Clinical Medicine, Ebonyi State University, Abakaliki, Nigeria

**Abstract:** Malaria remains a global health threat. Approximately 97% of the population is at risk in sub-Saharan countries, particularly Nigeria. This study compared the performance of 2 diagnostic methods in assessing malaria endemicity in the rural communities of Ebonyi State, Nigeria. A total of 1,140 study participants were screened for malaria parasite using Rapid Diagnostic Test kits (RDT) in the field, while thick and thin films for microscopy were examined in the laboratory. Our result showed that malaria prevalence was 56.8 by RDT and 38.6% by microscopic test. Age group under 10 years had the highest prevalence of 28.9% (RDT) and 23.6% (microscopy), respectively. The highest prevalence of 19.5% by RDT was recorded in Onicha Local Government Area, while the highest prevalence of 13.4% with microscopy was recorded in Ezza North Local Government Area. The sensitivity and specificity of microscopic examination were both 100%, while those of RDT were 95.5% and 75.9%, respectively.

**Key words:** Malaria, endemicity, diagnostic performance, rapid diagnostic test kit, microscopic examination, Nigeria

## INTRODUCTION

Malaria caused by infection with *Plasmodium* species is transmitted by mosquito-biting. The disease has a great impact on human health [1]. Malaria affects 3.3 billion people worldwide in at least 106 countries [2], but is both preventable and curable. In 2019 alone, 229 million malaria cases was detected globally with 409,000 estimated deaths (3). Children under 5 years are particularly susceptible to malaria [2], which accounted for 67% (274,000) of all malaria deaths worldwide [3].

Malaria is a serious threat to human health, repeated infections from childhood place a burden on individual households, health service, and economic development of communities and countries [4]. Malaria is one of the most important mosquito-borne diseases in Nigeria. It is more transmitted in rural communities, where favorable breeding habitats, espe-

cially during the wet season, and human activities are readily available. Malaria is a risk for 97% of Nigeria population [2] and contributes to about 11% of maternal mortality [5].

The impact of malaria is particularly pronounced in rural areas where the disease is common during periods of greatest need for agricultural labour [6]. Malaria transmission and severity of the disease vary greatly from region to region, village to village and even from person to person [7].

Factors contributing to the spread and transmission of malaria depend on the interaction between humans, *Anopheles* vectors, parasite, and environmental conditions. Natural transmission of malaria infection occurs by the bites of infected female *Anopheles* mosquitoes, but transmission can also occur by transfusion of infected blood. Infants may become infected perinatally by maternal blood.

## MATERIALS AND METHODS

### Ethical approval

Ethical approval was obtained from Ebonyi State Ministry of Health (Approval no. SMOH/03/017). The approval letter from the State Ministry of Health were taken to the traditional rulers as well as community leaders of the study communities

•Received 16 November 2021, revised 18 January 2022, accepted 22 February 2022.

\*Corresponding author (daveolink@gmail.com)

© 2022, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

for their approval of the study in their respective communities. Following approval by the community leaders, informed consent of the participants was sort through community mobilization and sensitization. The community members were invited to their town hall through the help of the community leaders, where they were explained the potential benefits of the research and allowed to consent freely.


**Study areas**

Ebonyi State is located in the Southeastern part of Nigeria. It shares boundaries with Benue State to the North, Enugu State to the Northeast, Abia State to Southeast and Cross River State to the East (Fig. 1). It is located between latitude 6o15’N and longitude 8o05’E. Ebonyi State is made up of 3 senatorial zones and 13 Local Government Areas (LGAs). The estimated total population in 2016 was 2,968,698 [8]. The climate of the area is tropical and the vegetation typical of the rain forest type. Average annual rainfall is 1,300 mm and average monthly temperature is of 30±5°C. There are 2 distinct seasons, the wet (between April and October) and the dry seasons (from

November to March). Most rural communities in Ebonyi State is mainly flat tableland with the capacity to hold water after rainfall. These topographic features usually create breeding grounds for mosquito vectors. A number of streams and rivers traverse most rural communities in Ebonyi State and constitutes the major sources of water supply to the inhabitants of the communities. The occupation of most rural residents in Ebonyi State is primarily farming, growing major crops such as yam, cassava, and rice.

**Study population**

A total of 1,140 study participants aged from 1-100 years, comprised of males (n=540) and females (n=600) from 3 communities drawn from 3 LGAs in the 3 senatorial zones that make up Ebonyi State (1 LGA per senatorial zone) were screened for malaria parasite. Three hundred and eighty (380) participants were screened from each community. Participants consented to the study.



**Fig. 1.** Map of Ebonyi State, which shows the study Areas.

### Sampling size technique

The sample size for this study was calculated using the formula below [9]

$$ME = \frac{z\sqrt{p(1-p)}}{n}$$

Where ME = margin of error, Z = z-score for confidence level, P = prior judgment or estimated prevalence, and n = sample size.

For this study, the confidence level was set at 95% and the z-score is 1.96: Margin of error = 0.025 (2.5%). P (initial estimate of prevalence) was the prevalence of 24.5% in the rural community of Ezza North LGA of Ebonyi State [10] which puts "P" at 0.245.

### Inclusion criteria

The inclusion criteria for the malaria endemicity study were individuals who have resided in the study community for the past one month before the commencement of the study. They are not seriously sick at the time of the study.

### Exclusion criteria

The exclusion criteria for the malaria endemicity study were visitors to the community, infants below the age of 6 months (due to protection by passive immunity), people who have taken anti-malaria drug within the last month.

### Sample collection

Malaria endemicity study was carried out in 3 communities, one each from Izzi, Onicha, and Ezza North LGAs of Ebonyi State. A total of 1,140 individuals (380 from each community) were examined for the presence of malaria parasite. All age groups ranging from 1-100 years comprised males and females were included. Sample collection was done on weekly basis. Rapid diagnostic test kits was used for the rapid diagnosis in the field while thick and thin smear slides for microscopy were prepared and transferred to the Medical Microbiology Laboratory of Ebonyi State University, Abakaliki where the slide were stained with 10% Giemsa and examined under a light microscope using x100 objective. All the samples collected each day were processed immediately after field collections. Slides that could not be examined the same day were continued the next day after proper staining. Prevalence rate was determined by calculating the percentage number of individuals infected with

malaria parasite in the study population.

### Rapid diagnosis

Rapid diagnosis was done using Rapid diagnostic test kit (RDT; SD Bioline malaria AG P.F/PAN kit, Standard Diagnostic. Seoul, Korea). The patient's finger was cleaned and air dried. The cleaned finger was pricked with a sterile lancet and a drop of blood collected with a capillary device. The blood was dropped to the lower well (hole) on the RDT test kit. Two drops of buffer were added to the blood on the lower well. Fifteen minutes later, results were interpreted according to the manufacturer's instruction and guidelines described by World Health Organization [11].

### Preparation of thick blood film

A drop of blood was stirred in a circular motion with a corner of slide. The film was allowed to dry without fixative, after which the smear was stained with 10% Giemsa for 20 min. The slides were washed by placing in a jar containing water for 3 min [12], and were later examined under a light microscope using x100 objective.

### Preparation of thin blood film

We smeared the blood with a swift and steady sweep along the surface of the slide. The film was air-dried and fixed with absolute methanol. The sample was stained with 10% Giemsa for 20 min [12] and were later examined under a light microscope using x100 objective.


### Data analysis

Data was analyzed using SPSS version 20.0 (IBM, Chicago, Illinois, USA). Statistical difference in parasite prevalence among different age groups and sex was determined using Chi square.


## RESULTS

### Diagnostic performance of microscopic examination and RDT


Of the 1,140 individuals examined for malaria prevalence, 648 (56.8%) showed positive with RDT while 440 (38.6%) were re-confirmed positive with microscopy (Fig. 2). There was no significant difference in parasite prevalence between the 2 diagnostic methods ( $\chi^2 = 0.601$ ,  $df = 2$ ,  $P = 0.741$  and  $\chi^2 = 0.718$ ,  $df = 2$ ,  $P = 0.698$ ), respectively. The test performance



**Fig. 2.** Prevalence of malaria in the study using 2 diagnostic methods.



**Fig. 3.** Comparison of test performance between microscopy and RDT.



**Fig. 4.** Age Prevalence of malaria in the study communities.

of microscopy and RDT is presented in Fig. 3. The sensitivity and specificity values for microscopic examination were both 100%. Those of RDT were 95.5% and 75.9%, respectively. All cases recorded in this study were infection with *Plasmodium falciparum*.

**Prevalence of malaria in relation to age**

The prevalence of malaria by age group with RDT examination was highest in 1-10 years (28.9%) and lowest in 91-100 years old (0.1%). Microscopic examination showed the highest prevalence (23.6%) among 1-10 years of age and lowest (0.1%) among age groups 51-60 years (Fig. 4). Statistically, there was no significant difference in the prevalence among the age groups with both diagnostic methods ( $\chi^2 = 16.371$ ,  $df =$

16,  $P = 0.427$  and  $\chi^2 = 11.380$ ,  $df = 12$ ,  $P = 0.497$ ), respectively.

**Gender related malaria prevalence**

The prevalence of malaria by gender is shown in Fig. 5. RDT and microscopic examination revealed the higher prevalence (31.2% and 20.4%, respectively) in males than females (25.6% and 18.2%, respectively); however, no significance difference was observed in the prevalence between males and females ( $\chi^2 = 0.170$ ,  $df = 2$ ,  $P = 0.919$  and  $\chi^2 = 1.865$ ,  $df = 2$ ,  $P = 0.394$ ).

**Malaria prevalence according to locality**

The highest prevalence of 19.5% with RDT was recorded in Onicha LGA while the lowest prevalence of 18.6% was shown in Izzi LGA. Microscopic examination showed the highest




Fig. 5. Prevalence of malaria in the communities by gender.




Fig. 6. Prevalence of malaria in the 3 study communities.

Table 1. Comparison of the diagnostic performance of microscopy with RDT

	Positive	Negative	True positive	False positive	True negative	False negative
Microscopy	440	700	440	0	700	0
RDT	648	492	420	228	720	20

prevalence of 13.4% in Ezza North LGA and the lowest prevalence of 12.5% in Izzi LGA (Fig. 6). There were no significant differences in the prevalence of malaria among LGAs for both diagnostic methods ( $\chi^2=0.601$ ,  $df=2$ ,  $P=0.741$  and  $\chi^2=0.718$ ,  $df=2$ ,  $P=0.698$ ), respectively.

Comparison of diagnostic performance of microscopic examination and RDT

Table 1 shows a comparison of microscopic and RDT results. For microscopy, 440 samples tested were positive and 700 were negative. A total of 648 samples were positive and 492 were negative by RDT. Among the 440 positive cases diagnosed by microscopic examination, RDT failed to detect 20 cases. Out of the 700 negative cases by microscopic examination, 228 showed a positive reaction by RDT yielding 228 false positive results.

DISCUSSION

Over the past few decades, there have been great success in control of malaria. However, malaria is still prevalent in Nigeria, including Ebonyi State, where suitable climatic conditions have resulted in the wide distribution of malaria vector mosquitoes. Our results showed malaria prevalence of 56.8% with

RDT and 38.6% with microscopy. This result strongly suggest that malaria is still a major public health concern in the study area. Our results are consistent with a previous report [13] indicating that malaria is endemic in Nigeria and maintains a consistently stable foci in these areas.

The malaria prevalence of 38.6% in this study by microscopic examination which is the gold standard is high. This rate is higher than previous reports, which reported that malaria prevalence was 13.7% in the West (Lagos state), 27.3% in the North (Sokoto State), 17% and 24.5% in Eastern Nigeria [10,14,15]. Other previous investigations have reported that malaria prevalence ranged from 41.7%-86.7% in Nigeria [16-18]. The malaria risk map estimated that prevalence in Nigeria varied from less than 20% to more than 70% [19].

This study showed the highest prevalence among young age group (1-10 years) by both microscopy and RDT. The result was similar to other previous reports [10,20,21], which reported a higher prevalence in young age group. There might be a slow acquisition of active immunity to malaria, since no significant difference in parasite prevalence was observed between different age groups.

Females showed a high prevalence rate. Earlier reports have also indicated high prevalence in females [10,20], while another previous study showed that male has a high prevalence rate [14], due possibly to the high risk of malaria infection by frequent exposure and cultural determinants [22]. On the other hand, other previous studies showed similar prevalence rate among the male and female in Nigeria, Kenya, and Mozambique [18,23,24]. These collective results suggest strongly that malaria risk factors are highly heterogeneous and complex [23,24].

More repeated exposure to malaria in males may lead to the development of partial immunity, which may lower risk of clinical malaria compared to females [25]. However, women's agricultural labour, household responsibilities, such as cooking dinner outdoors, waking up before sunrise to fetch water or preparing housework, may be at greater risk of contracting malaria than men. This may explain the higher prevalence among females, as seen in this study.

It is very hard to trace the reasons for the difference in prevalence between females and males. The difference observed in this study may have been coincidental since the differences in prevalence according to the gender was not statistically significant ( $P > 0.05$ ). With respect to study location, there was no significant difference in the prevalence of malaria among the 3 LGAs studied. This shows that the study areas share similar characteristic that encourages vector breeding and facilitates human-mosquito vector contact.

In this study, microscopy showed higher sensitivity (100%) than RDT (95.5%). However, the reduced sensitivity of RDT is higher than the 78.4% sensitivity reported in Zaria Nigeria [26]. It is slightly higher than the 93.3% recorded by Mohammed et al. [20]. However, Brown and Azike, [27] reported very poor sensitivity of 37.7% and specificity of 89.0% in a study conducted in Port Harcourt Rivers State Nigeria. HRP-2-based malaria RDT is known for detecting malaria antigen that continues to circulate in the blood almost 2 months after the treatment of a malaria [28]. This could possibly explain the reason for the false-positive results obtained in this study.

In conclusion, malaria is still a public health problem in Ebonyi State, especially in the rural communities. The transmission appears to last most of the year due to the availability of vector breeding habitats with the consequent adult production and distribution. Accurate diagnosis of *Plasmodium* species is important, not only for establishing the correct treatment regime, but also for applying effective malaria control strategies in the study area.

## ACKNOWLEDGEMENTS

We thank in a special way the traditional rulers of the study communities; HRH Eze Anthony Njoku (Okposi Umuoghara community), HRH Eze Ogonnaya Ukwa (Igbeagu community) and HRH Eze Patrick Chukwu Ogudu (Onicha Igboeze community), and their town union leaders for their unalloyed support during the study in their respective communities. We

equally thank Dr. Francis Onwe, the secretary Ethical Committee of the Ebonyi State Ministry of Health for his assistance in securing approval to carry out the study in the State.

## CONFLICT OF INTEREST

The authors hereby declare that there is no conflict of interest among them.

## REFERENCES

1. Caminade C, Kovats S, Rocklov J, Tompkins AM, Morse AP, Colón-González FJ, Stenlund H, Martens P, Lloyd SJ. Impact of climate change on global malaria distribution. *Proc Natl Acad Sci USA* 2014; 111: 3286-3291. <https://doi.org/10.1073/pnas.1302089111>
2. Andrew O. An assesment of the spatial pattern of malaria infection in Nigeria. *Int J Med Med Sci* 2014; 6: 80-86. <https://doi.org/10.5897/IJMMS2013.1006>
3. World Health Organization. World Malaria Report 2020. World Health Organization. Geneva, Switzerland. 2020, pp 12. Available from: <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2020>
4. Sachs J, Malaney P. The economic and social burden of malaria. *Nature* 2002; 415: 680-685. <https://doi.org/10.1038/415680a>
5. United States Embassy in Nigeria. Nigeria malaria fact sheet [Internet]. Available from: <https://nairametrics.com/wp-content/uploads/2013/03/Malaria-Fact-Sheet.pdf>
6. World Health Organization. Severe and complicated malaria. *Trans R Soc Trop Med Hyg* 2000; 94: 10-12. <http://helid.digicollection.org/pdf/s13418e/s13418e.pdf>
7. Beales PF, Gilles HM. Rationale and technique of malaria control. In Warrell DA, Gilles HM eds, *Essential Malariology*. 4th ed. CRC Press. London, UK. 2017, pp 107-190.
8. City Population. National Population Commission of Nigeria [Internet]. Available from: [www.citypopulation.de](http://www.citypopulation.de)
9. Naing L, Winn T, Rusli BN. Practical Issues in Calculating the Sample Size for Prevalence Studies. *Arc Orofacial Sci* 2006; 1: 9-14. <https://www.academia.edu/23980977/>
10. Nwele DE, Nwaorgu OC. Use of insecticide treated net (ITNs) among households and its impact on malaria prevalence in the communities in Ezza North Local Government Area of Ebonyi State, Nigeria. *Nigerian J Parasitol* 2017; 38: 85-90. <https://doi.org/10.4314/njpar.v38i1.16>
11. World Health Organization. How to use a rapid diagnostic test (Generic Pf): A guide for training at a village and clinic level. World Health Organization, Geneva, Switzerland. 2008, pp 48. <https://www.who.int/publications/m/item/how-to-use-a-rapid-diagnostic-test-generic-pf>
12. Tangpukdee N, Duangdee C, Wilairatana P, Krdsood S. Malaria diagnosis: a brief review. *Korean J Parasitol* 2009; 47: 93-102.

- <https://doi.org/10.3347/kjp.2009.47.2.93>
13. Okeke TA, Uzochukwu BS, Okafor HU. An in depth study of patient medicine seller's perspective on malaria in rural Nigerian community. *Malaria J* 2006; 5: 97. <https://doi.org/10.1186/1475-2875-5-97>
  14. Faruk HM, Adeoye GO. Malaria among out patients in Ikoyi area, Lagos State. Book of Abstract. Parasitology and Public Health Society of Nigeria. No 61. 2010.
  15. Anumudu CI, Adepoju A, Adeniran M, Adeoye O, Kassim A, Oyewole I, Nwuba RI. Malaria prevalence and treatment seeking behavior of young Nigerian adults. *Ann Afr Med* 2006; 15: 82-88. <https://www.ajol.info/index.php/aam/article/view/8380>
  16. Odikamnoroo OO, Ikeh IM, Okoh FN, Ebiriekwe SC, Nnadozie IA, Nkwuda JO, Asobie GC. Incidence of malaria/typhoid co-infection among adult population in Uwana community, Afikpo North Local Government Area, Ebonyi State, Southeastern Nigeria. *Afr J Infect Dis* 2017; 12: 33-38. <https://doi.org/10.21010/ajid.v12i1.6>
  17. Ali IC, Okeke OP, Unekwu AG, Effiong EJ. Prevalence and intensity of malaria among pregnant women in North Esatenr part of Ebonyi State Nigeria. *World J Biol Med Sciences* 2019; 6: 57-66.
  18. Dawaki S, Al-Mekhlafi HM, Ibrahim J, Atroosh WM, Abdulsalam AM, Sady H, Elyana FN, Adamu AU, Yelwa SI, Ahmed A, Al-Areeqi MA, Subramaniam LR, Nasr NA, Lau YL. Is Nigeria winning the battle against malaria? Prevalence, risk factors and KAP assessment among Hausa communities in Kano State. *Malaria J* 2016; 15: 351-356. <https://doi.org/10.1186/s12936-016-1394-3>
  19. Onyiri N. Estimating malaria burden in Nigeria; a geostatistical modeling approach. *Geospat Health* 2015; 10: 306-311. <https://doi.org/10.4081/gh.2015.306>
  20. Mohammed B, Emmanuel S, Bilyaminu M, Rufa'I A, Halima I, Kabir HS, Mohammed KA, Ibrahim I. Evaluation of the efficacy of rapid diagnostic tests compared to microscopy in the diagnosis of malaria infection. *Int J Mosquito Res* 2019; 6: 37-41. <https://www.dipterajournal.com/pdf/2019/vol6issue3/PartA/6-2-14-173.pdf>
  21. Perkins BA, Zucker JR, Otieno J, Jafari HS, Paxton L, Redd SC, Nahlen BL, Schwartz B, Oloo AJ, Olango C, Gove S, Campbell CC. Evaluation of an algorithm for intergrated management of childhood illness in an area of Kenya with high malaria transmission. *Bull World Health Organ* 1997; 75 (suppl): 33-42.
  22. Loha E, Lindtjorn B. Pridictors of *Plasmodium falciparum* malaria incidence in Chano mille, South Ethiopia: a longitudinal study. *Am J Trop Med Hyg* 2012; 87: 450-459. <https://doi.org/10.4269/ajtmh.2012.12-0155>
  23. Temu EA, Coleman M, Abilio AP, Kleinschmidt I. High prevalence of malaria in Zambezia, Mozambique: the protective effect of IRS versus increased risks due to pig keeping and house construction. *PLoS One* 2012; 7: e31409. <https://doi.org/10.1371/journal.pone.0031409>
  24. Noland GS, Graves PM, Sallau A, Eigege A, Emukah E, Patterson AE, Ajiji J, Okorofo I, Oji OU, Umar M, Alphonsus K, Damen J, Ngondi J, Ozaki M, Cromwell E, Obiezu J, Eneiramo S, Okoro C, McClintic-Doyle R, Oresanya O, Miri E, Emerson PM, Richards FO Jr. Malaria prevalence, anaemia and baseline intervention coverage prior to mass net distribution in Abia and Plateau States Nigeria. *BMC Infect Dis* 2014; 14: 168-74. <https://doi.org/10.1186/1471-2334-14-168>
  25. Hay SI, Okiro EA, Gething PW, Patil AP, Tatem AJ, Guerra CA, et al. Estimating the global clinical burden of *Plasmodium falciparum* in malaria in 2007. *PLoS Med* 2010; 7: e1000290. <https://doi.org/10.1371/journal.pmed.1000290>
  26. Sheyin Z, Bigwan IE. Comparison of CARE START HRP2 rapid malaria test with light microscopy for guiding patient's treatment of fever in Nigeria endemic areas. *J Med Sci* 2013; 4: 353-356. <http://dx.doi.org/10.14303/jmms.2013.109>
  27. Brown H, Azike CA. A comparative study on malaria rapid diagnostic kit and conventional giemsa stained blood film microscopy. *Int J Sci Res* 2012; 3: 1433-1435.
  28. Mahende C, Ngasala B, Lusingu J, Yong TS, Lushino P, Lemnge M, Mmbando B, Premji Z. Performance of rapid diagnostic test, blood-film microscopy and PCR for the diagnosis of malaria infection among febrile children from Korogwe District, Tanzania. *Malaria J* 2016; 15: 391. <https://doi.org/10.1186/s12936-016-1450-z>

