



Plant-Based Diets Induce Transcriptomic Changes in Muscle of Zebrafish and Atlantic Salmon

Anusha K. S. Dhanasiri^{1,2*}, Amritha Johny³, Xi Xue⁴, Gerd M. Berge⁵, Andre S. Bogevik⁶, Matthew L. Rise⁴, Christiane K. Fæste³ and Jorge M. O. Fernandes^{1*}

¹ Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway, ² Department of Paraclinical Sciences, Faculty of Veterinary Medicine, Norwegian University of Life Sciences (NMBU), Oslo, Norway, ³ Toxinology Research Group, Norwegian Veterinary Institute, Oslo, Norway, ⁴ Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, NL, Canada, ⁵ Norwegian Institute of Food, Fisheries and Aquaculture Research (Nofima), Sunndalsøra, Norway, ⁶ Norwegian Institute of Food, Fisheries and Aquaculture Research (Nofima), Fyllingsdalen, Norway

OPEN ACCESS

Edited by:

Katia Aquilano,
University of Rome Tor Vergata, Italy

Reviewed by:

Tamar Sztal,
Monash University, Australia
Steffen Just,
Ulm University Medical Center,
Germany

*Correspondence:

Anusha K. S. Dhanasiri
anusha.dhanasiri@nmbu.no
Jorge M. O. Fernandes
jorge.m.fernandes@nord.no

Specialty section:

This article was submitted to
Nutrigenomics,
a section of the journal
Frontiers in Genetics

Received: 22 June 2020

Accepted: 28 September 2020

Published: 22 October 2020

Citation:

Dhanasiri AKS, Johny A, Xue X, Berge GM, Bogevik AS, Rise ML, Fæste CK and Fernandes JMO (2020) Plant-Based Diets Induce Transcriptomic Changes in Muscle of Zebrafish and Atlantic Salmon. *Front. Genet.* 11:575237. doi: 10.3389/fgene.2020.575237

With the expansion of the aquaculture industry in the last two decades, there has been a large increase in the use of plant ingredients in aquafeeds, which has created new challenges in fish growth, health and welfare. Fish muscle growth is an important trait that is strongly affected by diet, but our knowledge on the effect of plant protein-based diets on global gene expression in muscle is still scant. The present study evaluated nutrigenomic effects of the inclusion of proteins from pea, soy and wheat into aquafeeds, compared to a control diet with fishmeal as the main protein source using the zebrafish model by RNA-seq; these results were extended to an important aquaculture species by analyzing selected differentially expressed genes identified in the zebrafish model on on-growing Atlantic salmon fed with equivalent plant protein-based diets. Expression of selected Atlantic salmon paralogues of the zebrafish homologs was analyzed using paralogue-specific qPCR assays. Global gene expression changes in muscle of zebrafish fed with plant-based diets were moderate, with the highest changes observed in the soy diet-fed fish, and no change for the pea diet-fed fish compared to the control diet. Among the differentially expressed genes were *mylpfb*, *hsp90aa1.1*, *col2a1a*, and *odc1*, which are important in regulating muscle growth, maintaining muscle structure and function, and muscle tissue homeostasis. Furthermore, those genes and their paralogues were differentially expressed in Atlantic salmon fed with the equivalent percentage of soy or wheat protein containing diets. Some of these genes were similarly regulated in both species while others showed species-specific regulation. The present study expands our understanding on the molecular effects of plant ingredients in fish muscle. Ultimately, the knowledge gained would be of importance for the improved formulation of sustainable plant-based diets for the aquaculture industry.

Keywords: plant-based proteins, fast muscle, pea protein concentrate, soy protein concentrate, wheat gluten, gene expression

INTRODUCTION

Exploration of alternative feed ingredients for fishmeal and fish oil has become a necessity for sustainable growth of the aquaculture sector (Carter and Hauler, 2000; Gatlin et al., 2007; Hardy, 2010). The industry has seen a steady increase in utilization of plant-based ingredients, mainly legumes, cereal grains and oil seeds in the fish feeds throughout the past years. For instance, the Norwegian salmon industry reported a large increase in use of plant ingredients, from 10% in 1990 to 60.5% in 2016 (Ytrestøyl et al., 2015; Aas et al., 2019). This huge transition to plant-based ingredients in the diet of farmed fish has led to new challenges in fish growth, health and welfare, and product quality (Hardy, 2010). Some of these challenges have been successfully addressed through advanced feed processing technologies and extensive research on the influence of dietary incorporation of plant ingredients on health, growth and nutrient utilization (Drew et al., 2007; Gatlin et al., 2007), but many challenges remain.

Plant-based proteins in fish diets can affect growth as a consequence of their nutritional and non-nutritional characteristics (Gatlin et al., 2007). Plant proteins also have some nutritional limitations such as the presence of high amounts of carbohydrates and imbalance in amino acid profiles compared with fishmeal. Generally, they are limited in some essential amino acids such as lysine and/or methionine, and low in available phosphorus and cationic minerals that mainly exist in bound form with phytic acid (Gatlin et al., 2007). With the inclusion of plant ingredients in the diets, fish could be exposed to various phytochemicals including anti-nutritional factors (ANFs) and phytoestrogens. ANFs (e.g., fibers, enzyme inhibitors, lectins, saponins, etc.) can interfere with nutrient digestibility, absorption and utilization, ultimately negatively affecting growth and health (Francis et al., 2001; Krogdahl et al., 2010). Phytoestrogens are compounds with estrogenic activity in animals, which can have an impact on the processes regulated by estrogens (Rietjens et al., 2013). Phytoestrogens have been reported to mostly exert a positive effect on fish growth (Chakraborty et al., 2014).

Among the widely used plant-based protein sources in aquafeeds, soy protein concentrate (SPC) produced from acid alcohol extraction has become the preferred choice compared to soy bean meal due to the absence of the ANF saponins as well as low contents of trypsin inhibitors, storage globulins with antigenic properties and oligosaccharides (Drew et al., 2007; Zhou et al., 2018). Nevertheless, it has been shown that diets with SPC caused intestinal enteritis in salmonids (Escaffre et al., 2007; Penn et al., 2011). Furthermore, inclusion of SPC in diets affects growth, depending on the percentage of replacement and fish species, e.g., partial replacement (50%) in diet of Atlantic salmon, *Salmo salar* (Storebakken et al., 2000b) and total replacement (100%) in diet of rainbow trout, *Oncorhynchus mykiss* (Kaushik et al., 1995) of fishmeal with SPC caused no changes in the growth of respective fish species. In contrast, a decrease in growth was observed in Japanese flounder, *Paralichthys olivaceus* (Deng et al., 2006), and gilthead sea bream, *Sparus aurata* (Kokou et al., 2015), with 25% of fishmeal replacement and 40% of inclusion with SPC in the diets, respectively.

The inclusion of pea protein concentrate (PPC) in aquafeeds also showed its potential as an alternative to fishmeal in farmed fish (Thiessen et al., 2003; Øverland et al., 2009; Zhang et al., 2012). Production of PPC by applying dehulling and air classification processes has enhanced the protein content, reduced the starch content and lowered certain ANF such as tannins compared to pea meal (Drew et al., 2007). However, it can still contain protease inhibitors, phytic acid, α -galactosidases and saponins in considerable amounts (Drew et al., 2007). PPC caused intestinal inflammation in the Atlantic salmon, similarly to soy-induced enteritis, and reduction in growth when included at a level of 35% in the diet (Penn et al., 2011). However, another study has reported that even inclusion levels of 50% of the dietary protein did not show any unfavorable effect on growth in rainbow trout (Zhang et al., 2012).

Wheat gluten (WG) has become an attractive alternative protein source for aquafeeds in several farmed species (Apper-Bossard et al., 2013), since it has a comparatively high crude protein content than fishmeal. Moreover, WG is used as pellet binder in feed manufacturing. It is high in sulfur amino acids and glutamate but generally low in lysine compared to the other commonly used plant protein sources (Storebakken et al., 2000a; Apper-Bossard et al., 2013). In a study on rainbow trout, it was found that WG was able to replace FM completely or to a large proportion without any unfavorable effect on growth, when lysine was supplemented in the diet (Pfeffer et al., 1992; Davies et al., 1997). Furthermore, moderate FM replacement with WG did not show any effect on growth in Atlantic salmon (35%) (Storebakken et al., 2000a) and Nile tilapia, *Oreochromis niloticus* L. (15%) (Schneider et al., 2004).

Muscle growth is an important trait affected by the nutritional status and diet (Valente et al., 2013). Some previous studies have examined the transcriptomic changes linked to muscle growth and dietary manipulation in fish (Alami-Durante et al., 2010; Bower and Johnston, 2010; Ulloa et al., 2013, 2015; Valente et al., 2016). However, knowledge on the effect of plant protein-based diets on global gene expression in muscle is still scant. In the present study, we have evaluated nutrigenomic effects of the inclusion of plant proteins SPC, PPC, and WG into aquafeeds, compared to a control diet with fishmeal as the main protein source, using the zebrafish as a model. Global transcriptome changes were analyzed in fast muscle using RNA-seq technology. The analysis was extended to a commercially important species by testing selected genes from the zebrafish model on on-growing Atlantic salmon fed with custom-made feeds containing the same plant ingredients. Considering the salmonid-specific genome duplication event (Lien et al., 2016), expression of selected paralogues of respective selected genes was analyzed using paralogue-specific qPCR assays.

MATERIALS AND METHODS

Feeding Experiments

Feeds

Custom-made diets for zebrafish and Atlantic salmon were produced by extrusion at the Nofima Feed Technology Centre, Fyllingsdalen, Norway as described by Johny et al. (2019). The

control diet included 79.4 and 63.4% fishmeal as protein source for, respectively, zebrafish and salmon (dietary composition is presented in **Supplementary Table 1**). Both control diets had 12% wheat as pellet-binding component, and 4.6% of additives including monosodium phosphate-24% P, vitamins and mineral mix. Considering the high fish oil requirement of salmon, 20% fish oil was added into the salmon diets as compared to 4% in the zebrafish diets. Plant protein-based diets were prepared by replacing 30% fishmeal in the control diet with PPC, SPC, or WG as protein source. The total protein content in all diets was 50 to 59 g/100 g feed for zebrafish and 44 to 48 g/100g feed for salmon, while the lipid content ranged from 9 to 12% and from 24 to 26% for, respectively, zebrafish and salmon. The plant-based diets were not supplemented with amino acids.

Zebrafish Feeding Trials and Sampling

Feeding trials were conducted using four-month-old zebrafish (AB strain, mean weight 0.214 g) for 46 days at Nord University, Bodø, Norway. Sixteen fish (1:1 sex ratio) were distributed into each of four randomly assigned replicate tanks (3.5 L volume) per dietary group in a flow-through system with 20% water exchange per h (ZebTEC stand-alone toxicology rack; Tecniplast, Buguggiate, Italy). Standard husbandry conditions were maintained with a stable temperature of $28 \pm \text{SD } 0.5^\circ\text{C}$, pH 7.5, water conductivity of 1500 $\mu\text{S/cm}$ and photoperiod of 12 h light/12 h dark. Feeding was performed twice a day with a total daily feed amount equal to 2.5% (w/w) of the body weight. Feeding behavior and health were regularly observed. The fish were fasted for 24 h prior to sampling. During the sampling, the fish in each tank were separated by gender and weighed. After euthanasia with a lethal dose of 200 mg/L tricaine methanesulfonate (MS222) (Sigma-Aldrich, St. Louis, United States), buffered with an equal amount of sodium bicarbonate, the fish were immediately frozen in liquid nitrogen and stored at -80°C until use. The skin was removed from the frozen specimens and fast muscle for RNA-seq analysis was carefully dissected under the microscope. Great care was taken to obtain solely fast muscle but it is plausible that slow muscle, muscle progenitor cells, neurons, immune cells and connective tissue were included. Therefore, the samples comprise a mixed cell population that is fast-muscle. Six females were randomly selected, including at least one fish from each of the four tanks per treatment. Statistical analysis of differences in specific growth rate among the dietary groups was performed with one-way ANOVA using R¹. Assumptions for statistical analysis, normality of the distribution and homogeneity of variances were assessed, respectively, using the Shapiro-Wilk test and Levene's test. Pairwise comparisons were analyzed using Tukey's honest significance test, and *p*-value after adjustment for the multiple comparisons <0.05 was considered as statistically significant.

Salmon Feeding Trials and Sampling

Feeding trials were conducted using one-year-old post-smolt Atlantic salmon (Salmobreed, Lønningdal, Norway) with a

starting mean weight of 223 g for 63 days at Nofima Research Station, Sunndalsøra, Norway as detailed by Johny et al. (2019). Each feeding group was allocated to three randomly distributed tanks (1 m³ and *n* = 15) and fish were fed to satiation with automatic disk feeders. They were reared at an average temperature of $10.6 \pm \text{SD } 0.6^\circ\text{C}$ and a seawater flow of 20 L/min. The oxygen levels of the tank outlets were above 80%. At the end of the feeding trials, the fish were anesthetized with 60–80 mg/L MS222, weighed and then euthanized with a double dose (120–160 mg/L) of MS222. After removing the skin, fast muscle was carefully dissected from the dorsal region to obtain fast-muscle enriched samples, which were frozen in liquid nitrogen and stored at -80°C until RNA isolation for qPCR analysis. Nine fish were randomly selected, including three fish from each of the three tanks per treatment. Their sex was undetermined, since they were not sexually mature at the time of sampling. Specific growth rates among the dietary groups were analyzed using one-way ANOVA, followed by Duncan's multiple range test using SAS software (as detailed in the related manuscript under preparation). All the pertinent assumptions, normality of the distribution and homogeneity of variances were checked before performing ANOVA.

RNA Sequencing

Total RNA was extracted from zebrafish fast muscle samples using QIAzol lysis reagent (Qiagen, Hilden, Germany), following the manufacturer's protocol. Tissues were homogenized twice at $5000 \times \text{g}$ for 15 s with zirconium oxide beads (1.4 mm; Precellys, Montigny-le-Bretonneux, France) using MagNALyser (Roche, San Francisco, United States). RNA integrity was checked using an Agilent 2200 TapeStation (Agilent Technologies, Santa Clara, United States), and RNA quantity was determined with a Qubit fluorometer (Invitrogen, Thermo Fisher Scientific, Waltham, United States). Libraries for RNA-seq were prepared using the NEBNext ultra II directional RNA library preparation kit with a poly(A) mRNA magnetic isolation module (NEB #E7490) in accordance to the manufacturer's protocol (New England BioLabs Inc., Ipswich, United Kingdom). Briefly, after poly(A) enrichment from 1 μg total RNA, mRNA was fragmented to about 100–200 nt lengths and used for 1st and 2nd strand cDNA synthesis. After A-tailing, end repair and adaptor ligation, the second strand was selectively removed using uracil-specific excision reagent (USER[®] II Enzyme; New England BioLabs Inc.). The resulting directional RNA-seq libraries were amplified with 8 PCR cycles and later purified using AMPure XP beads (Beckman Coulter, Inc., Brea, United States). Quality and quantity of the RNA-seq libraries were assessed using the Agilent 2200 TapeStation. Sequencing was performed on the Illumina NextSeq platform (Illumina, San Diego, United States) at Nord University using a single-end 75 bp high-throughput sequencing kit with 4% Phix control DNA (Illumina) as internal control.

RNA-Seq Data Analysis

Raw sequencing data were processed for quality and adapter trimming using Cutadapt (Martin, 2011) with $-\text{q } 25, 20$, quality-base = 33, trim-n -m 20 parameters, followed by a further

¹<https://www.r-project.org/>

TABLE 1 | Atlantic salmon homologs related to zebrafish genes that are differentially expressed by feeding with plant-based diets.

Gene name (symbol)	Zebrafish		Atlantic salmon	
	NCBI accession number	Differential expression	Paralogues*	NCBI accession number
<i>myosin light chain, phosphorylatable fast skeletal muscle b (mylpfb)</i>	NM_001004668.1	Soy group 2.7-fold decrease	<i>mylpfba</i>	XM_014203362.1
			<i>mylpfbb</i>	NM_001123716.1
<i>heat shock protein hsp90-alpha1 (hsp90aa1.1)</i>	NM_131328.1	Wheat group 3.9-fold decrease	<i>mylpfbc</i>	XM_014161797.1
			<i>hsp90aa1.1a</i>	XM_014205881.1
<i>activating molecule in beclin-1-regulated autophagy (ambra1a)</i>	NM_001281992.1	Soy group 2.1-fold increase	<i>hsp90aa1.1b</i>	XM_014144832.1
			<i>ambra1aa</i>	XM_014175106.1
<i>collagen type II, alpha 1 (col2a1a)</i>	NM_131292.1	Soy group 2.6-fold decrease	<i>ambra1ab</i>	XM_014126406.1
			<i>ambra1ac</i>	XM_014147648.1
<i>betacellulin (btcb)</i>	NM_001044764.3	Wheat group 2.6-fold increase	<i>ambra1ad</i>	XM_014123983.1
			<i>col2a1aa</i>	XM_014134054.1
<i>ryanodine receptor 1a (skeletal) (ryr1a)</i>	XM_021479405.1	Soy group 2.2-fold increase	<i>col2a1ab</i>	XM_014168236.1
			<i>col2a1ac</i>	XM_014135018.1
<i>ornithine decarboxylase (odc1)</i>	NM_131801.2	Soy group 2.6-fold decrease	<i>col2a1ad</i>	XM_014145553.1
			<i>btca</i>	XM_014129351.1
<i>mylpfbc (XM_014161797.1), ambra1ad (XM_014123983.1), col2a1ac (XM_014135018.1), and col2a1ad (XM_014145553.1)</i>			<i>btcb</i>	XM_014138426.1
			<i>ryr1aa</i>	XM_014196267.1
<i>mylpfbc (XM_014161797.1), ambra1ad (XM_014123983.1), col2a1ac (XM_014135018.1), and col2a1ad (XM_014145553.1)</i>			<i>ryr1ab</i>	XM_014129250.1
			<i>odc1a</i>	XM_014211026.1
<i>mylpfbc (XM_014161797.1), ambra1ad (XM_014123983.1), col2a1ac (XM_014135018.1), and col2a1ad (XM_014145553.1)</i>			<i>odc1b</i>	XM_014192087.1

*Genomic location and pairwise alignment details of paralogues are presented in **Supplementary Table 2**. Paralogues used for qPCR analysis are shown in bold letters. Paralogues, *btcb* (XM_014138426.1) and *ryr1ab* (XM_014129250.1) were not used for qPCR analysis, since the primers tested failed to show amplification. Paralogues *mylpfbc* (XM_014161797.1), *ambra1ad* (XM_014123983.1), *col2a1ac* (XM_014135018.1), and *col2a1ad* (XM_014145553.1) had very low expression levels in all dietary groups and therefore were not included in the differential expression analysis.

quality check with FastQC². Quality trimmed reads were mapped to the zebrafish genome and transcriptome downloaded from Ensembl (release 91) with TopHat2, version 2.1.0 (Kim et al., 2013). Indexing of the genome prior to mapping was done with Bowtie 2 (Langmead and Salzberg, 2012) and Tophat2. HTSeq (Anders et al., 2014) was used to compute gene expression values. Differentially expressed genes in fish fed with plant protein-based aquafeeds in comparison to their control diet counterparts were determined using DESeq2 (Anders and Huber, 2010) with the criteria adjusted *p*-value (*q*) with the Benjamini-Hochberg procedure ≤ 0.05 and absolute fold-change ≥ 1.5 .

qPCR Analysis

We used paralogue-specific qPCR assays to determine the expression of Atlantic salmon putative orthologs of several differentially expressed genes in zebrafish fed with soy or wheat diets as compared to the fishmeal control diet (Table 1). The Atlantic salmon had been fed with customized diets containing equal percentages of the same plant proteins as the zebrafish. Total RNA was extracted from Atlantic salmon

fast muscle ($n = 9$) using the above protocol, treated with DNase I (RNase-Free DNase Set from Qiagen) to digest any residual genomic DNA, and subjected to column-purification using RNeasy Mini Kit (Qiagen), following the manufacturer's instructions. RNA integrity was confirmed by 1% (w/v) agarose gel electrophoresis, and purity was assessed using a NanoDrop UV spectrophotometry (NanoDrop, Wilmington, United States). First-strand cDNA templates for qPCR were synthesized in 20 μ L reactions from 1 μ g of DNaseI-treated, column-purified total RNA using random primers (250 ng; Invitrogen/Life Technologies), dNTPs (0.5 mM final concentration; Invitrogen/Life Technologies), and M-MLV reverse transcriptase (200 U; Invitrogen/Life Technologies) with the manufacturer's first strand buffer (1 \times final concentration) and DTT (10 mM final concentration) at 37°C for 50 min.

Putative paralogous genes in Atlantic salmon corresponding to selected zebrafish genes were identified by BLASTn searches against the non-redundant nucleotide and expressed sequence tags (EST) databases at NCBI. Genomic location and pairwise alignment details of paralogues are presented in **Supplementary Table 2**. Additionally, an overview of their genomic neighborhood was retrieved from Ensembl (see text footnote 3) as shown in **Supplementary File 1**. All identified

²<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>

³<http://www.ensembl.org>

paralogues were aligned using ClustalW⁴, and paralogue-specific primers were designed to target regions where there were at least 2 bases different between paralogues to ensure specificity. Alignments of the paralogues with the primers highlighted are presented in the **Supplementary File 2**. Primers were designed either with primer-BLAST (NCBI) or manually and analyzed with Net Primer⁵ (**Table 2**).

Primer quality was assessed to ensure amplification of a single product without primer dimers by melt curve analysis using cDNA prepared with a pool of RNA with equal contribution from

all the samples used in the qPCR study. The size of the amplicons corresponding to each primer pair was verified by 2% (w/v) agarose gel electrophoresis using 1 kb plus ladder (Invitrogen/Life Technologies). Amplification efficiencies (Pfaffl, 2001) of primer pairs were calculated using a 5-point 1:3 dilution series starting with the pooled cDNA representing 10 ng of input total RNA. qPCR was performed in triplicate using Power SYBR Green I dye chemistry in 384-well format on a ViiA 7 real time PCR system (Applied Biosystems, Foster City, CA, United States) for the normalizer and target genes. Each 13 μ L reaction mixture contained 1 \times Power SYBR Green PCR Master Mix (Applied Biosystems), 50 nM of both the forward and reverse primers, and 4 μ L diluted cDNA (corresponding to 5 ng of input total


⁴<https://www.ebi.ac.uk/Tools/msa/clustalw2/>

⁵<http://www.premierbiosoft.com/netprimer/>

TABLE 2 | Primers employed for paralogue-specific qPCR study in Atlantic salmon fast muscle.

Gene	Accession number	Sequence 5'-3' §	Efficiency (%)	Size (bp)
<i>mylpfba</i>	XM_014203362.1	F-CTGTTCTCTCATACTTCCTTCTCTT R-CACCCGCTCTCACACAAATATC	110.5	113
<i>mylpfbb</i>	NM_001123716.1	F-CTATTCTCTTTATGTGTACTCGTGTGC R-GCCCGCTCTCACACAAATGT	81.3	96
<i>hsp90aa1.1a</i>	XM_014205881.1	F-GAAGTGATAGAGCTTTTCACAGAACTC R-CAGGGAAACCATTTTCATCACC	87.7	178
<i>hsp90aa1.1b</i>	XM_014144832.1	F-AAAGTGATGGATCTTTTCATCGAG R-TCAGGGAGACCATTTCGTCAG	93.3	179
<i>ambra1aa</i>	XM_014175106.1	F-CGTAGCAGTGGGAAAAGCTAAC R-GCCATTGCATGGTGAAATCCA	86.1	78
<i>ambra1ab</i>	XM_014126406.1	F-GGCAACATTGTCATCAGCTCC R-CAGAACCATGTTTGTAACACTAGCTA	103.5	103
<i>ambra1ac</i>	XM_014147648.1	F-AACAACAGCAGCAGCAGAGATAG R-AGCATCGTCACTAGCATCAGTC	107.7	136
<i>col2a1aa</i>	XM_014134054.1	F-CATGGTGCCTCAGCCCTATC R-CATTGGCTGCCTCTTTCTCC	86.6	125
<i>col2a1ab</i>	XM_014168236.1	F-CAGCGGTCTGGGTCTGGC R-CAGTTGGCGGACGGAGG	100.0	112
<i>btca</i>	XM_014129351.1	F-CAGGGTATCCTACTAAACCGAGGA R-TAGGGCCAAAGCTGTGGCTAT	105.1	168
<i>ryr1aa</i>	XM_014196267.1	F-CTGCTGAACCTGCTCCTGCT R-TTTTCAGTGTCTGCCTTCTCGG	97.4	76
<i>odc1a</i>	XM_014211026.1	F-CTTGTCCAATCTTTTGTGACAATCT R-CAGGAAGACAAAATCAGGAGCA	90.5	78
<i>odc1b</i>	XM_014192087.1	F-CCAGTTTTCTTTTGTGACCAATCTA R-TCATCCGACATGGACATCTCA	87.9	143
<i>igf1*</i>	NM_001123623.1	F-CCTGTTCGCTAAATCTCACTTC R-TACAGCACATCGCACTCTTGA	91.5	227
<i>igfII*</i>	NM_001123647.1	F-GAAAAACAAGAATGAAGGTCAA R-CCACCAGCTCTCCTCCACATA	124.1	127
<i>igfbp-1a2*</i>	NM_001279137	F-CTAAACCCCAAAACCCAGAT R-GTTGTGGCCAGGAAGGTAGA	134.8	104
<i>ef1a2α</i>	BG933853	F-GCACAGTAACACCGAAACGA R-ATGCCTCCGCACTTGATAGAT	88.0	132
<i>actbα</i>	BG933897	F-CCAAGCCAAACAGGAGAAG R-AGGGACAACACTGCCTGGAT	99.5	91

§Alignments of the paralogues with the primers highlighted are presented in **Supplementary File 2**. *Genes were tested in addition to the selected genes that were differentially regulated in the zebrafish feeding experiment: *igfI*-insulin-like growth factor I, *igfII*-insulin-like growth factor II, *igfbp-1a2*-insulin-like growth factor binding protein 1 paralogue A2. α Genes selected as reference genes for qPCR analysis.



RNA). The thermocycling profile was as follows: 1 cycle at 50°C for 2 min and 1 cycle at 95°C for 10 min, followed by 40 cycles at 95°C for 15 s and at 60°C for 1 min. The fluorescence signal data were collected after each 60°C step. Minus reverse transcriptase and no-template reactions were used as negative controls to confirm that there were no genomic and reagent contaminations, respectively.

Reference genes for normalization were selected after analyzing six candidate reference genes, including *actb* (β -actin), two paralogues of *elongation factor 1 alpha* [*ef1a1* (Olsvik et al., 2005) and *ef1a2* (Xue et al., 2015)], *pabpc1* (*polyadenylate-binding protein 1*), *rpl32* (*60S ribosomal protein L32*) (Xue et al., 2015), and *eif3d* (*eukaryotic translation initiation factor 3 subunit D*) (Caballero-Solares et al., 2017). Evaluation of normalizer genes was carried out using all the replicate samples from each of the control and plant protein diet-fed groups using *geNorm* (Vandesompele et al., 2002). *actb* and *ef1a2* were selected as normalizer genes as they were found to be the most stable of the tested six reference genes (i.e., lowest M-value, a gene stability measure). The relative quantity (RQ) of each transcript was determined using a qBase relative quantification framework (Hellemans et al., 2007; Xue et al., 2019), with normalization to the expression levels of *actb* and *ef1a2*. The sample with the lowest normalized expression level was used as calibrator (i.e., assigned RQ 1.0) when determining the RQ of each gene.

Statistical analysis was performed using R (see text footnote 1). Changes in gene expression between the control and each of the plant-based diets were determined by either parametric unpaired *t*-test or the non-parametric Wilcoxon rank sum test with continuity correction, based on the fulfillment of all pertinent assumptions for statistical analysis. Normality of the distribution was analyzed using the Shapiro-Wilk test, and homogeneity of variances was assessed by *F*-test. Gene expression changes were considered statistically significant at $p \leq 0.05$.

RESULTS


Plant-Based Diets Induced Moderate Transcriptomic Changes in Zebrafish Muscle

We used RNA-seq to determine the global transcriptomic changes in the fast-muscle enriched samples of zebrafish in response to plant protein-based diets that included 30% PPC (pea diet), SPC (soy diet), or WG (wheat diet), as well as a control diet containing fishmeal as the main protein source (Supplementary Table 1). The number of raw sequences ranged from 18 to 35 million per library. We obtained 18 to 31 and 16 to 30 million trimmed and mapped reads, respectively, per library (Figure 1). Each library had less than 7% of multiple mapping.

The analysis of global transcriptomic changes in the fast muscle of fish fed with the plant-based diets in comparison to the control group revealed 137 and 29 significantly differentially expressed genes (DEGs, fold change ≥ 1.5 , Benjamini-Hochberg adjusted *p*-value < 0.05) in soy and wheat diet groups, respectively, with a higher proportion of downregulated genes (Figures 2A,B). However, the pea diet group did not show any significant changes compared to the control group within the applied statistical criteria.

Feeding Soy and Wheat Diets Resulted in Regulation of Genes Involved in Growth, Structure and Function in Zebrafish Muscle

GO enrichment analysis of DEGs in the soy or wheat diet groups failed to show any enriched biological processes, most likely due to the relatively low number of DEGs. However,



functional annotation revealed that several DEGs are involved in regulating fast muscle growth and its structure and function in zebrafish. The list of annotated DEGs in the soy and wheat diet groups compared to the control group and associated GO terms is presented in **Supplementary Tables 3, 4**, respectively. An important gene for fast and slow skeletal muscle development, *myosin light chain, phosphorylatable fast skeletal muscle b (mylpfb*, 2.7-fold), and a component of the myosin complex, *myosin heavy polypeptide 1.1 (myhz1.1*, 2.7-fold), were downregulated in zebrafish fed with the soy diet, but not with the wheat diet. Whereas *popeye domain-containing (popdc3)*, needed for skeletal muscle development, was upregulated in the same group of fish. Another gene involved in muscle development, *heat shock protein hsp90-alpha1 (hsp90aa1.1*, 3.9-fold), was downregulated in zebrafish fed with the wheat diet, but its expression did not significantly change in the zebrafish fed with the soy diet. Two of the upregulated genes in zebrafish fed with the soy diet were *activating molecule in beclin-1-regulated autophagy (ambra1a*, 2.1-fold), a positive regulator of autophagy during skeletal muscle development, and *ryanodine receptor 1a (ryr1a*, 2.2-fold), an essential component of all skeletal muscle fiber calcium-release channels. On the other hand, a gene coding for a vital component of the skeletal muscle extracellular matrix, *collagen type II, alpha 1 (col2a1a*, 2.6-fold), was downregulated in the same group of fish. Zebrafish fed with soy diet also showed downregulation of *ornithine decarboxylase (odc1*, 2.6-fold), which is important for polyamine biosynthetic processes and myoblast proliferation. *betacellulin (btc*, 2.6-fold), a member of the epidermal growth factor family that mediates diverse processes including proliferation and differentiation, was upregulated in fast muscle of wheat diet-fed zebrafish.

Furthermore, several genes involved in metabolic processes including proteolysis [i.e., *dipeptidyl-peptidase 6b, dpp6b* (3.3-fold increase)], cholesterol biosynthetic process [i.e., apolipoproteins, *apoa1a, apoa1b* and *apobb.1* (2.7-, 2.8-, 2.9-fold decrease, respectively)] and glucose metabolism [i.e., *glyceraldehyde-3-phosphate dehydrogenase, gapdh* (1.6-fold decrease)] were differentially regulated in soy diet-fed zebrafish.

Among the genes differentially regulated in both soy and wheat diets-fed fish were *PDZ and LIM domain 1 (pdlim1*, downregulation), involving in actin cytoskeleton organization; serotransferrin (*tfa*, downregulation), important for ion transport as well as response to bacterium; *mutS homolog 3 (msh3*, upregulation), vital for DNA repair; and *activating transcription factor 3 (atf3*, downregulation), needed for regulation of transcription and fish immune responses (Feng and Rise, 2011).

A considerable number of *crystallin* genes (*crystallin, gamma* and *crystallin, beta*), mainly known to be important for visual perception, were downregulated in the soy diet fed group. Even though chaperone-like activity was described for mammalian *crystallin, gamma* (Andley et al., 1996), to our knowledge, the importance of this gene for fast muscle growth and homeostasis in mammals or fish has not been reported.

Differential Expression of Atlantic Salmon Homologs Related to Zebrafish Genes That Are Differentially Expressed With Feeding Plant-Based Diets


The expression of selected genes that were differentially regulated with the diet in zebrafish fast muscle was subsequently examined in the fast-muscle enriched samples of Atlantic salmon fed with equivalent diets. Considering the salmonid genome duplication event, we used paralogue-specific qPCR assays to quantify the transcript levels of Atlantic salmon homologs of the selected zebrafish genes (**Table 1**).

Expression profiles of the selected paralogous genes in fast muscle of Atlantic salmon are presented in **Figures 3, 4**. The paralogues *mylpfba* and *mylpfbb* were upregulated in the soy diet group (**Figure 3A1**) whereas the zebrafish putative ortholog was downregulated in those fed with the same plant-based diet. There was no significant regulation of *mylpfb* observed for wheat diet fed zebrafish or Atlantic salmon. Both the *hsp90aa1.1a* and *hsp90aa1.1b* paralogues (**Figure 3B2**) were upregulated (2.5- and 2.2-fold, respectively) in wheat diet-fed Atlantic salmon, contrary to the downregulation observed in wheat diet-fed zebrafish. The expression levels of all three *ambra1a* paralogues did not change significantly in soy diet-fed Atlantic salmon (**Figure 3C1**) even though the putative ortholog was upregulated in the zebrafish counterparts. They were also not significantly changed in Atlantic salmon fed with the wheat diet (**Figure 3C2**). Similarly, to the findings in zebrafish, *col2a1aa* and *col2a1ab* were downregulated (1.7-fold each) in Atlantic salmon fed with the soy diet (**Figure 4A1**). The expression of *btca* and *ryr1aa* paralogues did not change significantly in soy or wheat diet-fed Atlantic salmon (**Figures 4B1,2**). Both *odc1* paralogues were downregulated (1.2-fold each) in soy diet-fed Atlantic salmon (**Figure 4C1**), comparable to their zebrafish counterparts. Further, *odc1* paralogues were also downregulated in Atlantic salmon fed with wheat diet (**Figure 4C2**) while they were not significantly changed in the zebrafish fed with the similar plant-based diet.

We also analyzed the expression of genes involved in insulin-like growth factor (IGF) signaling in the fast muscle of Atlantic salmon fed with soy and wheat diets. The transcript expression of *insulin-like growth factor I (igf1)* did not show any significant changes with respect to the diet (**Figures 5A,B**). In contrast, *insulin-like growth factor II (igfII*, 1.5- and 1.9-fold, respectively, for soy and wheat diet groups) and *insulin-like growth factor binding protein 1 paralog A2 (ifbp-1a2*, 2.1- and 31.8-fold, respectively, for soy and wheat diet groups) were upregulated with both soy and wheat diet groups (**Figures 5A,B**).

Growth Performance of Zebrafish and Atlantic Salmon During the Feeding Studies

The specific growth rate of zebrafish fed with diets including 30% PPC, SPC, or WG did not significantly change after 46 days of the feeding trial as compared to their counterparts fed with a diet containing fishmeal as the main protein source (**Figure 6**).








FIGURE 4 | Expression profiles of *col2a1a*, *btca*, *ryr1a*, and *odc1* paralogues in the fast muscle of Atlantic salmon fed with the soy (A1,B1,C1) and wheat (A2,B2,C2) diets compared to controls fed with a diet containing fishmeal as the main protein source. Values are means \pm S.E.M ($n = 7-9$). Asterisks above the bars indicate significant changes ($p < 0.05$) in transcript levels.



Moreover, inclusion of 30% SPC also did not show any significant changes in the specific growth rate of Atlantic salmon compared to counterparts fed with a diet containing fishmeal as the main protein source; on the other hand, inclusion of 30% WG had a significant reduction in specific growth rate in salmon compared to the control group (growth data are included in the related manuscript under preparation).

DISCUSSION

The present study reports the nutrigenomic effects of fish diets containing partial inclusion of plant-based proteins, e.g., pea, soy, and wheat, in the fast muscle of zebrafish. Global gene expression changes in the muscle tissues of zebrafish fed with plant-based diet were moderate yet included several genes important for the regulation of muscle growth, maintenance, function and homeostasis.

The specific growth rate of zebrafish was not significantly affected by any of the plant protein-based diets used in the present study after a 46-day feeding trial. Inclusion of plant-based proteins in aquafeeds can affect fish growth, depending on the percentage of replacement and species. For instance, 50% replacement of fishmeal with SPC did not affect the growth rate in Atlantic salmon (Storebakken et al., 2000b), whereas 25% replacement of fishmeal with SPC significantly reduced the growth rate of Japanese flounder (Deng et al., 2006). The total replacement of marine protein sources with plant-based proteins from 35 to 98 days post-fertilization reportedly decreased the specific growth rate of zebrafish (Ulloa et al., 2013). The plant-based diets used in our study (Johny et al., 2019) contained still a considerable amount of fishmeal (49.4%) in addition to the 30% plant protein, resulting in a crude protein content of 50 g pea, 55 g soy, and 59 g wheat per 100 g feed, which was similar to the control diet with 56 g crude protein/100 g feed. The relatively high ratio of fishmeal in the plant-based diets has probably helped the zebrafish to maintain a similar growth rate as their control counterparts fed with 79.4% fishmeal. Moreover, we found that Atlantic salmon fed with plant-based diets containing 33.4% fishmeal in addition to 30% SPC also did not show any significant changes in the specific growth rate; however, salmon fed with the diet containing 30% WG had a growth rate decreased by 11.1% compared to the controls that were fed with 63.4% fishmeal.

The analysis of DEGs in the muscle of zebrafish fed with the plant protein-containing diets showed that the DEG numbers varied with the plant ingredient. They were highest in fish fed with the soy diet followed by the fish fed with the wheat diet. Surprisingly, the pea diet did not induce any significant gene expression changes as compared to the fishmeal control group. Among the DEGs changed in the soy and wheat diet-fed groups, there were genes associated with muscle growth, function, metabolism and homeostasis, some of which were also differentially expressed in Atlantic salmon fed with the same plant-based protein diets, while others showed species

specific regulation. Even though those gene expression changes have not significantly influenced their growth, we believe that these molecular adjustments occurred in the muscle to maintain muscle homeostasis following imbalances caused by dietary plant ingredients. Moreover, our recently published study (Dhanasiri et al., 2020) from the same feeding trial also indicated plant-based diets induced epigenetic changes in the intestine with mild inflammation without significantly affecting their growth.

All teleosts share three rounds of whole-genome duplication (WGD), 1R, 2R, and 3R where the last was a teleost-specific WGD (Glasauer and Neuhauss, 2014; Lien et al., 2016). Additionally, a fourth (4R) WGD occurred in the common ancestor of salmonids (Glasauer and Neuhauss, 2014). WGDs are important for evolutionary adaptations and innovations; complexity and diversifications. The duplicated genes after WGDs can undergo different fates including: non-functionalization (deleterious mutations occur in one of the duplicates leading to loss of expression); subfunctionalization (mutations in paralogous genes leading to preservation of both duplicates); neofunctionalization (one of the genes acquires a novel function) (Glasauer and Neuhauss, 2014). Differential expression patterns of Atlantic salmon homologs (putative orthologs and paralogues) related to zebrafish genes regulated by dietary inclusion of plant-based diets observed in our study detailed in the following sections, provide possible evidence for gene duplication and divergence. However, functional studies are needed to clarify the fate of the Atlantic salmon paralogous genes to determine whether they were subfunctionalized or neofunctionalized.

Skeletal muscle is a highly adaptable tissue, which changes in size and cell composition in response to environmental and physiological conditions. Its homeostasis can be challenged by both short and long term nutritional changes (Matsakas and Patel, 2009). Skeletal muscle fibers consist of myofibrils that are made up of sarcomeres containing organized arrays of actin thin filaments and myosin thick filaments along with accessory proteins (Johnston et al., 2011). Several muscle subtypes are described based on the expression of numerous isoforms of myosin light and heavy chains. Myosin heavy chain is mainly responsible for the functional and phenotypic diversity of muscles (Pette and Staron, 2000). The regulatory light chain of myosin in fast skeletal muscle is encoded by the *mylpf* gene in mammals (Wang et al., 2007), and ortholog, *mylpfb*, exists in zebrafish. In mouse, *mylpf* knockdown resulted in the complete lack of skeletal muscle during development, indicating the importance of *mylpf* for the growth of both fast and slow skeletal muscle (Wang et al., 2007). *mylpfb* was downregulated 2.7-fold in muscle of zebrafish fed with the soy diet and there were no other differentially expressed myosin-related genes that could compensate for this change but its regulation did not have a significant influence on growth. On the other hand, its homologs *mylpfba* and *mylpfbb* were upregulated 1.5- and 1.4-fold in Atlantic salmon fed with the same diet and these gene expression changes also did not affect growth significantly. These puzzling species-specific observations reflect the complexity of myogenesis and may be due to diet-related changes at regulation levels that were not addressed in the present study, such as post-translational modifications or expression of non-coding RNAs.

Several studies in zebrafish and Atlantic salmon demonstrated that heat-shock proteins 90 α (Hsp90 α), especially Hsp90 α 1, play an important role in myosin folding, myofibril assembly and myosin thick filament organization (Du et al., 2008; Garcia de la serrana and Johnston, 2013; He et al., 2015). Loss of Hsp90 α 1 function resulted in increased myosin protein degradation (Du et al., 2008) and disruption of all sarcomeric structures, including both thick and thin filaments in skeletal muscles in zebrafish (Codina et al., 2010). We observed a downregulation of *hsp90aa1.1* in zebrafish fed with the wheat diet, however, we did not observe a significant change in the specific growth rate of these fish. On the other hand, both the *hsp90aa1.1a* and *hsp90aa1.1b* paralogues were upregulated in the same group of salmon, concomitantly with a significant reduction in growth. Increased expression of those paralogues was also previously reported under the rapid enhancement of nutritional status, such as the initiation of feeding after starvation in juvenile salmon and supplementation of complete amino acids into the starved primary myogenic cultures (Garcia de la serrana and Johnston, 2013). Wheat diet-fed fish showed a significant reduction in specific growth rate indicating a possible lower nutritional status compared to the control group. Further studies are required to understand the regulation of *hsp90aa1* paralogues in Atlantic salmon under different nutritional conditions as well as their species-specific regulation.

In addition to genes coding for fast muscle components or directly involved in muscle growth and function, some genes indirectly involved in muscle functions, such as *col2a1a*, were downregulated in zebrafish and salmon fed with the soy diet. The skeletal muscle extracellular matrix (ECM) is vital for muscle fiber force transmission, maintenance and repair (Gillies and Lieber, 2011). Collagen is a major structural protein in muscle ECM even though it is a minor constituent (1–10%) of the muscle mass dry weight. *collagen type II (col2a1)* is expressed during skeletal muscle development in mammals (Gillies and Lieber, 2011) and zebrafish embryogenesis (Dale and Topczewski, 2011). Therefore, the downregulation of *col2a1a* caused by the soy diet in adult zebrafish and Atlantic salmon warrants further investigation. Moreover, both zebrafish and Atlantic salmon fed with the soy diet showed downregulation of *odc1*, a gene coding for ornithine decarboxylase, which is important for polyamine biosynthetic processes. Studies in mouse myoblasts and human skeletal muscle cell lines showed that *odc1* gene promotes myoblast proliferation (Lee et al., 2011). The downregulation of *odc1* in both zebrafish and salmon fed with the soy diet could be linked to high polyamine levels present in SPC, since ornithine decarboxylase levels are modulated by the available polyamine concentration (Pegg, 2006). This could have ultimately delayed myoblast proliferation and affected myogenesis.

Muscle growth is regulated by growth hormones acting directly through sarcolemmal receptors and indirectly *via* the insulin-like growth factor (IGF) pathway. The IGF system comprises IGF-I, IGF-II, several receptors and six binding proteins (IGFBPs) (Johnston et al., 2011). IGFs stimulate muscle growth in fish by promoting the proliferation of myogenic cells, protein synthesis and hypertrophy, as well as inhibiting protein degradation and muscle atrophy (Fuentes et al., 2013).

The availability of IGFs is regulated by IGFs, and generally IGFBP-1 and -2 inhibit the growth-promoting functions of IGFs in both mammals and fish (Fuentes et al., 2013). Zebrafish selected for large body size showed higher *igf1* and lower *igfbp1a* mRNA levels compared to those selected for small body size in a study investigating responses to changes in the nutritional input (Amaral and Johnston, 2012). Interestingly, a study in Atlantic salmon showed that switching to fast growth involves the local upregulation of *igf1*, *igfbp-5.2* and *igfbp-4* accompanied by downregulation of *igfbp-2.1* and *igf1* in the skeletal muscle (Bower et al., 2008). We did not observe differential expression of genes from the GH-IGF signaling system in zebrafish fed with any of the three-plant protein-based diets as compared to the fishmeal diet, which is in agreement with the lack of growth rate differences between the feeding groups. On the other hand, *igf1* and *igfbp-1a2* were upregulated in Atlantic salmon fed with the soy and wheat diets with normal and reduced growth, respectively, compared to the control group. Further studies are needed to explain the involvement of specific components of the GH-IGF system in regulating the growth changes observed in Atlantic salmon fed with different plant-based diets.

Odc1 is the only salmon gene in our study whose paralogues show different expression patterns with the wheat diet, which suggests that they have undergone subfunctionalization following duplication. The identified paralogues of *mylpfb*, *hsp90aa1.1*, *ambra1*, and *col2a1* showed similar patterns of regulation with soy- and wheat diets, indicating functional redundancy. Nevertheless, cellular localization and functional studies are required to ascertain conclusively if these paralogues are indeed functionally redundant or undergoing subfunctionalization or neofunctionalization.

CONCLUSION

The partial inclusion of proteins from soy and wheat into zebrafish feed resulted in the regulation of, respectively, 137 and 29 genes in muscle. In contrast, the inclusion of pea protein concentrate did not induce changes in the gene expression as compared to a control group receiving a fishmeal-based diet. Among the differentially expressed genes, several were important for muscle growth, structure, function and homeostasis. Some of these genes and their paralogues were similarly regulated in Atlantic salmon fed with equivalent diets while others showed species-specific regulation. Our results revealed that molecular adjustments occurred in the muscle of zebrafish and Atlantic salmon, probably to maintain muscle homeostasis following imbalances caused by dietary plant ingredients, even if there were no significant differences observed in the growth. Ultimately, this knowledge may be applied for the improved formulation of sustainable plant-based diets for the aquaculture industry.

REFERENCES

Aas, T. S., Ytrestøyl, T., and Åsgård, T. (2019). *Resource utilization of Norwegian Salmon Farming in 2016: Professional Final Report*. Tromsø: Nofima.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://www.ncbi.nlm.nih.gov/>, PRJNA577226; <https://www.ncbi.nlm.nih.gov/>, SUB6971208.

ETHICS STATEMENT

The animal study was reviewed and approved by Norwegian Animal Research Authority, Norway; Nord University, Norway.

AUTHOR CONTRIBUTIONS

AD, JF, MR, and CF conceived and designed the study. CF, JF, and MR contributed reagents and materials for the experiments. AD, XX, and AJ performed laboratory work. AD and GB conducted the zebrafish and Atlantic salmon feeding trials, respectively. AB prepared the feeds. AD, JF, and XX analyzed the data. AD, JF, XX, MR, and CF wrote the manuscript. All authors read and approved the manuscript.

FUNDING

This work was supported by the Norwegian Research Council through a post-doctoral fellowship to AD (NFR 267944) and a research grant to CF (NFR 254822).

ACKNOWLEDGMENTS

We are thankful to Dr. Qirui Zhang at Lund University (Lund, Sweden) and Dr. Prabhugouda Siriyappagouder at Nord University (Bodø, Norway) for their guidance and recommendations during the bioinformatics analysis. We are also grateful to Dr. Albert Caballero-Solares (Memorial University of Newfoundland, St. John's, Canada) for his support during the paralogue-specific qPCR analysis.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2020.575237/full#supplementary-material>

Alami-Durante, H., Médale, F., Cluzeaud, M., and Kaushik, S. J. (2010). Skeletal muscle growth dynamics and expression of related genes in white and red muscles of rainbow trout fed diets with graded levels of a mixture of plant protein sources as substitutes for

- fishmeal. *Aquaculture* 303, 50–58. doi: 10.1016/j.aquaculture.2010.03.012
- Amaral, I. P. G., and Johnston, I. A. (2012). Experimental selection for body size at age modifies early life-history traits and muscle gene expression in adult zebrafish. *iJ. Exp. Biol.* 215, 3895–3904. doi: 10.1242/jeb.068908
- Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biol.* 11:R106. doi: 10.1186/gb-2010-11-10-r106
- Anders, S., Pyl, P. T., and Huber, W. (2014). HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31, 166–169. doi: 10.1093/bioinformatics/btu638
- Andley, U. P., Mathur, S., Griest, T. A., and Petrash, J. M. (1996). Cloning, expression, and chaperone-like activity of human α A-crystallin. *J. Biol. Chem.* 271, 31973–31980. doi: 10.1074/jbc.271.50.31973
- Apper-Bossard, E., Feneuil, A., Wagner, A., and Respondek, F. (2013). Use of vital wheat gluten in aquaculture feeds. *Aquat. Biosyst.* 9:21. doi: 10.1186/2046-9063-9-21
- Bower, N. I., and Johnston, I. A. (2010). Discovery and characterization of nutritionally regulated genes associated with muscle growth in Atlantic salmon. *Physiol. Genomics* 42A, 114–130. doi: 10.1152/physiolgenomics.00065.2010
- Bower, N. I., Li, X., Taylor, R., and Johnston, I. A. (2008). Switching to fast growth: the insulin-like growth factor (IGF) system in skeletal muscle of Atlantic salmon. *J. Exp. Biol.* 211, 3859–3870. doi: 10.1242/jeb.024117
- Caballero-Solares, A., Hall, J. R., Xue, X., Eslamloo, K., Taylor, R. G., Parrish, C. C., et al. (2017). The dietary replacement of marine ingredients by terrestrial animal and plant alternatives modulates the antiviral immune response of Atlantic salmon (*Salmo salar*). *Fish Shellfish Immunol.* 64, 24–38. doi: 10.1016/j.fsi.2017.02.040
- Carter, C. G., and Hauler, R. C. (2000). Fish meal replacement by plant meals in extruded feeds for Atlantic salmon, *Salmo salar* L. *Aquaculture* 185, 299–311. doi: 10.1016/S0044-8486(99)00353-1
- Chakraborty, S. B., Horn, P., and Hancz, C. (2014). Application of phytochemicals as growth-promoters and endocrine modulators in fish culture. *Rev. Aquac.* 6, 1–19. doi: 10.1111/raq.12021
- Codina, M., Li, J., Gutiérrez, J., Kao, J. P. Y., and Du, S. J. (2010). Loss of Smyhc1 or Hsp90 α 1 function results in different effects on myofibril organization in skeletal muscles of zebrafish embryos. *PLoS One* 5:e8416–e8416. doi: 10.1371/journal.pone.0008416
- Dale, R. M., and Topczewski, J. (2011). Identification of an evolutionarily conserved regulatory element of the zebrafish *col2a1a* gene. *Dev. Biol.* 357, 518–531. doi: 10.1016/j.ydbio.2011.06.020
- Davies, S. J., Morris, P. C., and Baker, R. T. M. (1997). Partial substitution of fish meal and full-fat soya bean meal with wheat gluten and influence of lysine supplementation in diets for rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquac. Res.* 28, 317–328. doi: 10.1111/j.1365-2109.1997.tb01048.x
- Deng, J., Mai, K., Ai, Q., Zhang, W., Wang, X., Xu, W., et al. (2006). Effects of replacing fish meal with soy protein concentrate on feed intake and growth of juvenile Japanese flounder, *Paralichthys olivaceus*. *Aquaculture* 258, 503–513. doi: 10.1016/j.aquaculture.2006.04.004
- Dhanasiri, A. K. S., Chen, X., Dahle, D., Siriyappagounder, P., Fæste, C. K., and Fernandes, J. M. O. (2020). Dietary inclusion of plant ingredients induces epigenetic changes in the intestine of zebrafish. *Epigenetics* 15, 1035–1051. doi: 10.1080/15592294.2020.1747777
- Drew, M. D., Borgeson, T. L., and Thiessen, D. L. (2007). A review of processing of feed ingredients to enhance diet digestibility in finfish. *Anim. Feed Sci. Technol.* 138, 118–136. doi: 10.1016/j.anifeedsci.2007.06.019
- Du, S. J., Li, H., Bian, Y., and Zhong, Y. (2008). Heat-shock protein 90 α 1 is required for organized myofibril assembly in skeletal muscles of zebrafish embryos. *Proc. Natl. Acad. Sci. U.S.A.* 105, 554–559. doi: 10.1073/pnas.0707330105
- Escaffre, A. M., Kaushik, S., and Mambrini, M. (2007). Morphometric evaluation of changes in the digestive tract of rainbow trout (*Oncorhynchus mykiss*) due to fish meal replacement with soy protein concentrate. *Aquaculture* 273, 127–138. doi: 10.1016/j.aquaculture.2007.09.028
- Feng, C. Y., and Rise, M. L. (2011). Identification and molecular cloning of Atlantic cod (*Gadus morhua*) activating transcription factor 3 (ATF3) transcript and its induction in spleen following intraperitoneal polyriboinosinic polyribocytidylic acid injection. *Fish Shellfish Immunol.* 31, 475–481. doi: 10.1016/j.fsi.2011.06.002
- Francis, G., Makkar, H. P. S., and Becker, K. (2001). Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish. *Aquaculture* 199, 197–227. doi: 10.1016/s0044-8486(01)00526-9
- Fuentes, E. N., Valdés, J. A., Molina, A., and Björnsson, B. T. (2013). Regulation of skeletal muscle growth in fish by the growth hormone – Insulin-like growth factor system. *Gen. Comp. Endocrinol.* 192, 136–148. doi: 10.1016/j.ygcen.2013.06.009
- García de la Serrana, D., and Johnston, I. A. (2013). Expression of heat shock protein (Hsp90) paralogs is regulated by amino acids in skeletal muscle of Atlantic salmon. *PLoS One* 8:e74295. doi: 10.1371/journal.pone.0074295
- Gatlin, D. M., Barrows, F. T., Brown, P., Dabrowski, K., Gaylord, T. G., Hardy, R. W., et al. (2007). Expanding the utilization of sustainable plant products in aquafeeds: a review. *Aquac. Res.* 38, 551–579. doi: 10.1111/j.1365-2109.2007.01704.x
- Gillies, A. R., and Lieber, R. L. (2011). Structure and function of the skeletal muscle extracellular matrix. *Muscle Nerve* 44, 318–331. doi: 10.1002/mus.22094
- Glaser, S. M. K., and Neuhauss, S. C. F. (2014). Whole-genome duplication in teleost fishes and its evolutionary consequences. *Mol. Genet. Genomics* 289, 1045–1060. doi: 10.1007/s00438-014-0889-2
- Hardy, R. W. (2010). Utilization of plant proteins in fish diets: effects of global demand and supplies of fishmeal. *Aquac. Res.* 41, 770–776. doi: 10.1111/j.1365-2109.2009.02349.x
- He, Q., Liu, K., Tian, Z., and Du, S. J. (2015). The Effects of Hsp90 α 1 mutations on myosin thick filament organization. *PLoS One* 10:e0142573. doi: 10.1371/journal.pone.0142573
- Hellemans, J., Mortier, G., De Paep, A., Speleman, F., and Vandesompele, J. (2007). qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol.* 8:R19. doi: 10.1186/gb-2007-8-2-r19
- Johnston, I. A., Bower, N. I., and Macqueen, D. J. (2011). Growth and the regulation of myotomal muscle mass in teleost fish. *iJ. Exp. Biol.* 214, 1617–1628. doi: 10.1242/jeb.038620
- Johny, A., Fæste, C. K., Bøgevik, A. S., Berge, G. M., Fernandes, J. M. O., and Ivanova, L. (2019). Development and validation of a liquid chromatography high-resolution mass spectrometry method for the simultaneous determination of mycotoxins and phytoestrogens in plant-based fish feed and exposed fish. *Toxins* 11:222. doi: 10.3390/toxins11040222
- Kaushik, S. J., Cravedi, J. P., Lalles, J. P., Sumpter, J., Fauconneau, B., and Laroche, M. (1995). Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture* 133, 257–274. doi: 10.1016/0044-8486(94)00403-b
- Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S. L. (2013). TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* 14:R36. doi: 10.1186/gb-2013-14-4-r36
- Kokou, F. J., Sarropoulou, E., Cotou, E., Rigos, G., Henry, M., Alexis, M., et al. (2015). Effects of fish meal replacement by a soybean protein on growth, histology, selected immune and oxidative status markers of gilthead sea bream, *Sparus aurata*. *J. World Aquac. Soc.* 46, 115–128. doi: 10.1111/jwas.12181
- Krogdahl, Å., Penn, M., Thorsen, J., Refstie, S., and Bakke, A. M. (2010). Important antinutrients in plant feedstuffs for aquaculture: an update on recent findings regarding responses in salmonids. *Aquac. Res.* 41, 333–344. doi: 10.1111/j.1365-2109.2009.02426.x
- Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9, 357–359. doi: 10.1038/nmeth.1923
- Lee, N. K., Skinner, J. P., Zajac, J. D., and MacLean, H. E. (2011). Ornithine decarboxylase is upregulated by the androgen receptor in skeletal muscle and regulates myoblast proliferation. *Am. J. Physiol. Endocrinol. Metab.* 301, E172–E179. doi: 10.1152/ajpendo.00094.2011
- Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., et al. (2016). The Atlantic salmon genome provides insights into rediploidization. *Nature* 533, 200–205. doi: 10.1038/nature17164
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 17, 10–12. doi: 10.14806/ej.17.1.200
- Matsakas, A., and Patel, K. (2009). Skeletal muscle fibre plasticity in response to selected environmental and physiological stimuli. *Histol. Histopathol.* 24, 611–629.

- Olsvik, P. A., Lie, K. K., Jordal, A.-E. O., Nilsen, T. O., and Hordvik, I. (2005). Evaluation of potential reference genes in real-time RT-PCR studies of Atlantic salmon. *BMC Mol. Biol.* 6:21. doi: 10.1186/1471-2199-6-21
- Øverland, M., Sørensen, M., Storebakken, T., Penn, M., Krogdahl, Å., and Skrede, A. (2009). Pea protein concentrate substituting fish meal or soybean meal in diets for Atlantic salmon (*Salmo salar*)—Effect on growth performance, nutrient digestibility, carcass composition, gut health, and physical feed quality. *Aquaculture* 288, 305–311. doi: 10.1016/j.aquaculture.2008.12.012
- Pegg, A. E. (2006). Regulation of ornithine decarboxylase. *J. Biol. Chem.* 281, 14529–14532. doi: 10.1074/jbc.R500031200
- Penn, M. H., Bendiksen, E. Å., Campbell, P., and Krogdahl, Å. (2011). High level of dietary pea protein concentrate induces enteropathy in Atlantic salmon (*Salmo salar* L.). *Aquaculture* 310, 267–273. doi: 10.1016/j.aquaculture.2010.10.040
- Pette, D., and Staron, R. S. (2000). Myosin isoforms, muscle fiber types, and transitions. *Microsc. Res. Tech.* 50, 500–509. doi: 10.1002/1097-0029(20000915)50:6<500::aid-jemt7>3.0.co;2-7
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29:e45. doi: 10.1093/nar/29.9.e45
- Pfeffer, E., Al-Sabty, H., and Haverkamp, R. (1992). Studies on lysine requirements of rainbow trout (*Oncorhynchus mykiss*) fed wheat gluten as only source of dietary protein. *J. Anim. Physiol. Anim. Nutr.* 67, 74–82. doi: 10.1111/j.1439-0396.1992.tb00585.x
- Rietjens, I. M. C. M., Sotoca, A. M., Vervoort, J., and Louisse, J. (2013). Mechanisms underlying the dualistic mode of action of major soy isoflavones in relation to cell proliferation and cancer risks. *Mol. Nutr. Food Res.* 57, 100–113. doi: 10.1002/mnfr.201200439
- Schneider, O., Amirkolaie, A. K., Vera-Cartas, J., Eding, E. H., Schrama, J. W., and Verreth, J. A. J. (2004). Digestibility, faeces recovery, and related carbon, nitrogen and phosphorus balances of five feed ingredients evaluated as fishmeal alternatives in Nile tilapia, *Oreochromis niloticus* L. *Aquac. Res.* 35, 1370–1379. doi: 10.1111/j.1365-2109.2004.01179.x
- Storebakken, T., Shearer, K. D., Baeverfjord, G., Nielsen, B. G., Åsgård, T., Scott, T., et al. (2000a). Digestibility of macronutrients, energy and amino acids, absorption of elements and absence of intestinal enteritis in Atlantic salmon, *Salmo salar*, fed diets with wheat gluten. *Aquaculture* 184, 115–132. doi: 10.1016/S0044-8486(99)00316-6
- Storebakken, T., Shearer, K. D., and Roem, A. J. (2000b). Growth, uptake and retention of nitrogen and phosphorus, and absorption of other minerals in Atlantic salmon *Salmo salar* fed diets with fish meal and soy-protein concentrate as the main sources of protein. *Aquac. Nutr.* 6, 103–108. doi: 10.1046/j.1365-2095.2000.00135.x
- Thiessen, D. L., Campbell, G. L., and Adelizi, P. D. (2003). Digestibility and growth performance of juvenile rainbow trout (*Oncorhynchus mykiss*) fed with pea and canola products. *Aquac. Nutr.* 9, 67–75. doi: 10.1046/j.1365-2095.2003.00203.x
- Ulloa, P. E., Peña, A. A., Lizama, C. D., Aranedo, C., Iturra, P., Neira, R., et al. (2013). Growth response and expression of muscle growth-related candidate genes in adult zebrafish fed plant and fishmeal protein-based diets. *Zebrafish* 10, 99–109. doi: 10.1089/zeb.2012.0823
- Ulloa, P. E., Rincón, G., Islas-Trejo, A., Aranedo, C., Iturra, P., Neira, R., et al. (2015). RNA sequencing to study gene expression and SNP variations associated with growth in zebrafish fed a plant protein-based diet. *Mar. Biotechnol.* 17, 353–363. doi: 10.1007/s10126-015-9624-1
- Valente, L. M. P., Cabral, E. M., Sousa, V., Cunha, L. M., and Fernandes, J. M. O. (2016). Plant protein blends in diets for *Senegalese sole* affect skeletal muscle growth, flesh texture and the expression of related genes. *Aquaculture* 453, 77–85. doi: 10.1016/j.aquaculture.2015.11.034
- Valente, L. M. P., Moutou, K. A., Conceição, L. E. C., Engrola, S., Fernandes, J. M. O., and Johnston, I. A. (2013). What determines growth potential and juvenile quality of farmed fish species? *Rev. Aquac.* 5, S168–S193. doi: 10.1111/raq.12020
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., et al. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 3:research0034.1.
- Wang, Y., Szczesna-Cordary, D., Craig, R., Diaz-Perez, Z., Guzman, G., Miller, T., et al. (2007). Fast skeletal muscle regulatory light chain is required for fast and slow skeletal muscle development. *FASEB J.* 21, 2205–2214. doi: 10.1096/fj.06-7538com
- Xue, X., Hixson, S. M., Hori, T. S., Booman, M., Parrish, C. C., Anderson, D. M., et al. (2015). Atlantic salmon (*Salmo salar*) liver transcriptome response to diets containing *Camelina sativa* products. *Comp. Biochem. Physiol. D* 14, 1–15. doi: 10.1016/j.cbd.2015.01.005
- Xue, X., Woldemariam, N. T., Caballero-Solares, A., Umasuthan, N., Fast, M. D., Taylor, R. G., et al. (2019). Dietary immunostimulant CpG modulates microRNA biomarkers associated with immune responses in Atlantic salmon (*Salmo salar*). *Cells* 8:1592. doi: 10.3390/cells8121592
- Ytrestøyl, T., Aas, T. S., and Åsgård, T. (2015). Utilisation of feed resources in production of Atlantic salmon (*Salmo salar*) in Norway. *Aquaculture* 448, 365–374. doi: 10.1016/j.aquaculture.2015.06.023
- Zhang, Y., Øverland, M., Sørensen, M., Penn, M., Mydland, L. T., Shearer, K. D., et al. (2012). Optimal inclusion of lupin and pea protein concentrates in extruded diets for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 344–349, 100–113. doi: 10.1016/j.aquaculture.2012.03.012
- Zhou, Z., Ringø, E., Olsen, R. E., and Song, S. K. (2018). Dietary effects of soybean products on gut microbiota and immunity of aquatic animals: a review. *Aquac. Nutr.* 24, 644–665. doi: 10.1111/anu.12532

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Dhanasiri, Johny, Xue, Berge, Bogeveik, Rise, Fæste and Fernandes. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.