



COVID-19: Possible Cause of Induction of Relapse of *Plasmodium vivax* Infection

Rashmi Kishore¹ · Shivram Dhakad² · Nazneen Arif² · Lalit Dar² · B. R. Mirdha² · Richa Aggarwal³ · S. K. Kabra¹

Received: 19 June 2020 / Accepted: 25 June 2020 / Published online: 3 July 2020
© Dr. K C Chaudhuri Foundation 2020

To the Editor: In ongoing pandemic of novel corona virus disease (COVID-19), clinicians are observing atypical manifestations of the disease. We hereby report a case of COVID-19 co-infection with vivax malaria in a 10-y-old boy, who previously had received incomplete radical cure with primaquine for vivax infection, suggesting a possible role of COVID-19 in inducing current malarial relapse.

A 10-y-old boy, resident of Delhi, presented to the pediatric emergency department of a tertiary care hospital with history of high grade fever with chills and rigors, headache, cold, cough and pain abdomen. Past history revealed admission with a similar episode of febrile illness six months back. Medical records from previous admission revealed that he was diagnosed with *P. vivax* infection and had not completed primaquine therapy as per history. At presentation, child was febrile with fever of 104° F, maintained saturation of 97–98% on room air. Rest of systemic examination was normal. Investigations revealed Hb of 11.7/dl, total leucocyte count (TLC) of 4500/mm³ with 15% neutrophils, 53% lymphocytes and 28% eosinophils (absolute eosinophil count of 1260/mm³), and platelet count of 52,000/mm³. Rapid diagnostic test for malaria was faintly positive for pan antigen and smear examination showed all stages of *Plasmodium vivax* (*P. vivax*), confirming it to be malaria. Nasal and pharyngeal swabs RT-PCR for corona virus 2 (SARS-CoV-2) were positive. Other investigations were normal.

The child was admitted and treated for malaria and supportive care for COVID-19 infection [1]. Primaquine eradication therapy was given for 14 d after ruling out G6PD deficiency. He was discharged after complete recovery.

This boy had been diagnosed with *P. vivax* infection six months back and had now reactivation of malaria. Relapse rates after *P. vivax* infection vary geographically from 8 to 80%, with considerable proportion of population harboring dormant but activatable hypnozoites in endemic areas [2]. Exact mechanism causing this activation is though unclear, associated cytokine response with systemic illness has been postulated in *P. vivax* relapse [3]. COVID-19 leads to a cytokine storm, which is responsible for the more severe manifestations of the disease [4, 5]. Hence, we postulate that the COVID-19 infection, with its cytokine response was responsible for induction of *P. vivax* relapse in our patient. Our hypothesis is based on circumstantial evidence in form of documented malaria in past, a possibility of re-activation in natural course or re-infection cannot be ruled out.

Acknowledgments Director General, Indian Council of Medical Research, for providing funding for support of diagnostics.

Compliance with Ethical Standards

Conflict of Interest None.

✉ Rashmi Kishore
rashmi.k186@gmail.com

- ¹ Department of Pediatrics, All India Institute of Medical Sciences, New Delhi 110029, India
- ² Department of Microbiology, All India Institute of Medical Sciences, New Delhi, India
- ³ Department of Anesthesiology, Critical and Intensive Care, JPNA Trauma Centre, All India Institute of Medical Sciences, New Delhi, India

References

1. Sankar J, Dhochak N, Kabra SK, Lodha R. COVID-19 in children: clinical approach and management. *Indian J Pediatr.* 2020;87:433–42.
2. World Health Organization. Guidelines for the Treatment of Malaria. 2015. Available at: https://www.who.int/docs/default-source/documents/publications/gmp/guidelines-for-the-treatment-of-malaria-eng.pdf?sfvrsn=a0138b77_2. Accessed 19 May 2020.
3. White NJ. Determinants of relapse periodicity in plasmodium vivax malaria. *Malar J.* 2011;10:297.

4. Guo YR, Cao QD, Hong ZS, et al. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Mil Med Res.* 2020;7:11.
5. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med.* 2020;8:420–2.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.