

POSTER PRESENTATION

Open Access

Interplay of periglomerular and granule cell inhibitory synapses on mitral cell spiking

Denise Arruda*, Rodrigo Publio, Antonio C Roque

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Mitral cells in the olfactory bulb (OB) receive dendrodendritic inhibitory synaptic inputs at two different levels of OB: at the glomerular level, inhibitory input is provided by periglomerular (PG) cells and, at a deeper level (the external plexiform layer), inhibitory input is provided by granule (Gr) cells [1]. It is thought that both kinds of inhibition control mitral cell spiking in

specific ways: PG early inhibition would regulate spike initiation while Gr inhibition would coordinate spike synchronization among mitral cells [1,2]. However, recent computational modeling studies have suggested that, although acting at separate locations, PG and Gr inhibitory synapses may have integrated effects on mitral cell spiking which may be beneficial for odor




Figure 1 Firing rate of mitral cell as a function of maximal conductances of inhibitory synapses made by PG (horizontal axis) and Gr (vertical axis) cells. The color-code bar on the right indicates frequency in Hz.

* Correspondence: denisearruda84@gmail.com

Department de Física, FFCLRP, Universidade de São Paulo, Ribeirão Preto, SP, 14040-901, Brazil

representation and processing [2]. In this work we constructed in NEURON [3] a model of a triad composed of reduced compartmental models of a mitral cell, a PG cell and a Gr cell to study the combined effect of the inhibitory dendrodendritic synapses of the PG and Gr cells on the firing frequency of the mitral cell. The mitral and Gr cell models were taken from [4] with all parameters and characteristics kept as in the original. The PG cell model was based on the five-compartment model used in [5]. The cells were coupled by models of their reciprocal dendrodendritic synapses. The main parameters investigated in our simulations were the conductances of the dendrodendritic synapses, the amplitudes, durations and initiation times of the excitatory inputs applied to the mitral and PG cells representing stimuli coming from olfactory sensory neurons, and the amplitude, duration and initiation time of the input applied to the Gr cell representing afferents from other cells in the olfactory bulb. We performed simulations in which these parameters were varied and studied their effects on the mitral cell's spiking pattern. Our results show that the response pattern of the mitral cell is strongly dependent of the two inhibitory synapses, and that their interaction can regulate both the spike frequency and timing of the mitral cell. Figure 1 illustrates one of these results.

Acknowledgements

DA is supported by CAPES, RP is supported by FAPESP and ACR is supported by a CNPq research grant.

Published: 18 July 2011

References

1. Lledo PM, Gheusi G, Vincent JD: Information processing in the mammalian olfactory system. *Physiol Rev* 2005, 85:281-317.
2. David F, Linster C, Cleland TA: Lateral dendritic shunt inhibition can regularize mitral cell spike patterning. *J Comput Neurosci* 2008, 25:25-38.
3. Carnevale NT, Hines ML: The NEURON Book. Cambridge, UK: Cambridge University Press; 2006.
4. Davison AP, Brown JFD: Spike synchronization in a biophysically-detailed model of the olfactory bulb. *Neurocomput* 2001, 38-40:515-521.
5. Cleland TA, Sethupathy P: Non-topographical contrast enhancement in the olfactory bulb. *BMC Neurosci* 2006, 7:7.

doi:10.1186/1471-2202-12-S1-P269

Cite this article as: Arruda et al.: Interplay of periglomerular and granule cell inhibitory synapses on mitral cell spiking. *BMC Neuroscience* 2011 12 (Suppl 1):P269.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

