

POSTER PRESENTATION

Open Access

Transcription profiling in papillary thyroid carcinoma reveals potential diagnostic markers and drug targets

Mateus CB Filho^{1*}, Fabio A Marchi², Silvia R Rogatto^{1,3}, Luiz P Kowalski¹

From São Paulo Advanced School of Comparative Oncology
Águas de São Pedro, Brazil. 30 September - 6 October 2012

Background

Papillary thyroid carcinoma (PTC) is the most frequent malignant endocrine neoplasia with an increasing prevalence in the last decades. We aim to identify transcripts and pathways associated with PTC tumorigenesis.

Materials and methods

RNA from tumor and adjacent normal samples was evaluated using Sure Print G3 8x60K slides (Agilent Technologies). Sixty-five tumor (T) and four normal (N) tissues were labeled with Cy5. A pool composed by nine normal samples (without the corresponding tumor assayed) was labeled with Cy3 and used in the co-hybridization. Statistical analysis was performed using two approaches, a paired (4N vs 4T) and an independent analysis (9N vs. 61T).

Results

Overlapping paired (paired Significance Analysis of Microarray with 3% False Discovery Ratio) and independent analysis (mean log ratios <-1 or >1 with 99% Confidence Interval) resulted in a list of 546 deregulated genes. Networks and functional analysis were generated through IPA software (Ingenuity[®] Systems). The major molecular network identified was related to endocrine system development and function and down regulation of tyrosine metabolism was the main canonical pathway. A preliminary validation was carried out with RT-qPCR for *HMGA2*. A higher expression was confirmed ($P < 0.001$) in an independent sample set (11N vs. 47T). *HMGA2* expression had also diagnostic ability, correctly classifying 117/121 samples according to tumor status (sensitivity=97%, specificity=94% and area under the ROC curve=0.989).

Conclusion

This study unveils transcription modulations during PTC genesis and *HMGA2* may be a potential diagnostic marker. Functional studies are required to confirm *HMGA2* as an oncogenic driver in PTC and with a possible role as a drug target.

Financial support

FAPESP and CAPES.

Author details

¹International Research Center/AC Camargo Hospital, Sao Paulo, Brazil. ²Sao Paulo University, Sao Paulo, Brazil. ³Faculty of Medicine/UNESP, Botucatu, SP, Brazil.

Published: 4 April 2013

doi:10.1186/1753-6561-7-S2-P51

Cite this article as: Filho et al.: Transcription profiling in papillary thyroid carcinoma reveals potential diagnostic markers and drug targets. *BMC Proceedings* 2013 **7**(Suppl 2):P51.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: mateusbarros@gmail.com

¹International Research Center/AC Camargo Hospital, Sao Paulo, Brazil
Full list of author information is available at the end of the article