

## 9-O-Butylberberrubinium bromide

Zhu Chen,<sup>a</sup> Xue-Gang Li,<sup>a</sup> Yong-Sheng Xie<sup>b</sup> and Xiao-Li Ye<sup>c\*</sup>

<sup>a</sup>College of Pharmaceutical Sciences, Southwest University, Chongqing 400715, People's Republic of China, <sup>b</sup>School of Chemistry and Chemical Engineering, Shandong University, Jinan 250100, People's Republic of China, and <sup>c</sup>College of Life Sciences, Southwest University, Chongqing 400715, People's Republic of China  
Correspondence e-mail: yexiaoli@swu.edu.cn


Received 14 April 2009; accepted 15 May 2009

Key indicators: single-crystal X-ray study;  $T = 295\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$ ; disorder in main residue;  $R$  factor = 0.060; wR factor = 0.161; data-to-parameter ratio = 13.0.

In the title compound,  $\text{C}_{23}\text{H}_{24}\text{NO}_4^+\cdot\text{Br}^-$ , the butyl chain is disordered between two conformations; the occupancies refined to 0.735 (7) and 0.265 (7). The dihedral angle between the naphthalene ring system and the phenyl ring is  $11.6(2)^\circ$ . In the crystal structure, the cations are packed via  $\pi-\pi$  interactions into stacks propagating in the [010] direction. Weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{Br}$  hydrogen bonds contribute further to the crystal packing stability.

### Related literature

For the bioactivity of berberine, see: Jiang *et al.* (1998); Kupeli *et al.* (2002). For the bioactivity of 9-O-butyl-berberrubine bromide, see Ye & Li (2007).



### Experimental

#### Crystal data

$\text{C}_{23}\text{H}_{24}\text{NO}_4^+\cdot\text{Br}^-$   
 $M_r = 458.34$   
Monoclinic,  $P2_1/n$   
 $a = 9.716(4)\text{ \AA}$   
 $b = 7.623(3)\text{ \AA}$   
 $c = 27.443(11)\text{ \AA}$   
 $\beta = 92.983(8)^\circ$   
 $V = 2029.9(14)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 2.06\text{ mm}^{-1}$   
 $T = 295\text{ K}$   
 $0.12 \times 0.10 \times 0.06\text{ mm}$

#### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker 2005)  
 $T_{\min} = 0.791$ ,  $T_{\max} = 0.887$   
10394 measured reflections  
3592 independent reflections  
1965 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.069$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$   
 $wR(F^2) = 0.161$   
 $S = 1.08$   
3592 reflections  
276 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.58\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.55\text{ e \AA}^{-3}$

**Table 1**  
Centroid-to-centroid distances ( $\text{\AA}$ ).

$Cg1\cdots Cg3^i$	3.780 (4)	$Cg2\cdots Cg3^{ii}$	3.775 (4)
-------------------	-----------	----------------------	-----------

Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 1$ .  $Cg1$ ,  $Cg2$  and  $Cg3$  are the centroids of the N1/C13/C12/C9/C10/C14, C5–C10 and C15/C19–C22/C16 rings, respectively.

**Table 2**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C12–H12 $\cdots$ Br1	0.93	2.77	3.658 (5)	161
C2–H2A $\cdots$ O2 $^{iii}$	0.97	2.64	3.404 (14)	136

Symmetry code: (iii)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by the National Natural Science Foundation of China (grant No. 20673084), the Natural Science Foundation Project of CQ CSTC (grant No. 2008BB5257) and the Major Technologies R & D Program of CQ CSTC (grant No. 2008AA5021).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2550).

### References

- Bruker (2005). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jiang, J. Y., Geng, D. S., Tursonjan, T. & Liu, F. (1998). *Chin. Pharmacol. Bull.* **14**, 434–437.
- Kupeli, E., Kosar, M., Yesilada, E., Husnu, K. & Baser, C. (2002). *Life Sci.* **72**, 645–657.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Ye, X. L. & Li, X. G. (2007). Chinese Patent No. 200710078505.X.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o1370 [doi:10.1107/S1600536809018388]

## 9-O-Butylberberrubinium bromide

Z. Chen, X.-G. Li, Y.-S. Xie and X.-L. Ye

### Comment

Berberine is a main compound presented in the rhizome of *Coptis chinensis* Franch. Berberine and its derivatives are used for treating diarrhoea (Kupeli *et al.*, 2002) and anti-inflammatory (Jiang *et al.*, 1998). Herewith we report the crystal structure of the title compound (**I**), which exhibits an excellent antibacterial activity (Ye & Li, 2007).

In (**I**) (Fig. 1), the dihedral angles formed by the benzene rings C5–C10 and C15/C19/C20/C21/C22/C16 with the pyridine ring are 2.2 (3) and 12.8 (3)°, respectively. The six-membered heterocyclic ring (C13/C15/C16/C17/C18/N1) adopts screw-boat conformation. In the crystal structure, weak intermolecular C—H···O and C—H···Br hydrogen bonds (Table 2) link the molecules. The aromatic rings in the cations are involved in  $\pi$ – $\pi$  interactions (Table 1). The cations are packed *via*  $\pi$ – $\pi$  interactions into stacks propagated in direction [010].

### Experimental

Berberrubine was obtained from berberine by pyrolysis at 180 °C for 1 h. Then dried berberrubine (10 mmol) was dissolved in DMF (200 ml) and incubated with *n*-butyl bromide (11 mmol) for 5 h at 100 °C to give the corresponding crude 9-*O*-butylberberrubine bromide. The product was purified by recrystallization from methanol at -18 °C. Crystals suitable for X-ray diffraction were obtained by slow evaporation of a solution of the solid dissolved in methanol at room temperature for 15 d.

### Refinement

All H atoms were placed in calculated positions, with C—H = 0.93–0.97 Å, and included in the final cycles of refinement using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  for aryl and methylene H atoms or  $1.5U_{\text{eq}}(\text{C})$  for methyl H atoms.

### Figures

Fig. 1. View of the title compound showing the atom-labeling scheme and 30% probability displacement ellipsoids.

## 9-O-Butylberberrubinium bromide

### Crystal data

$\text{C}_{23}\text{H}_{24}\text{NO}_4^+\cdot\text{Br}^-$	$F_{000} = 944$
$M_r = 458.34$	$D_x = 1.500 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 9.716 (4) \text{ \AA}$	Cell parameters from 1163 reflections
	$\theta = 2.8\text{--}19.6^\circ$

# supplementary materials

---

$b = 7.623 (3) \text{ \AA}$	$\mu = 2.06 \text{ mm}^{-1}$
$c = 27.443 (11) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 92.983 (8)^\circ$	Block, yellow
$V = 2029.9 (14) \text{ \AA}^3$	$0.12 \times 0.10 \times 0.06 \text{ mm}$
$Z = 4$	

## Data collection

Bruker SMART CCD area-detector diffractometer	3592 independent reflections
Radiation source: fine-focus sealed tube	1965 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.069$
$T = 295 \text{ K}$	$\theta_{\max} = 25.1^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\min} = 1.5^\circ$
Absorption correction: multi-scan (SADABS; Bruker 2005)	$h = -11 \rightarrow 11$
$T_{\min} = 0.791, T_{\max} = 0.887$	$k = -7 \rightarrow 9$
10394 measured reflections	$l = -32 \rightarrow 32$

## Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.060$	$w = 1/[\sigma^2(F_o^2) + (0.0623P)^2 + 1.1441P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.161$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.08$	$\Delta\rho_{\max} = 0.58 \text{ e \AA}^{-3}$
3592 reflections	$\Delta\rho_{\min} = -0.55 \text{ e \AA}^{-3}$
276 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0020 (7)
Secondary atom site location: difference Fourier map	

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Br1	0.19113 (6)	0.34507 (10)	0.58534 (2)	0.0683 (3)	
O1	0.4605 (4)	0.4269 (6)	0.31978 (15)	0.0691 (13)	
O2	0.8340 (4)	-0.0249 (7)	0.66220 (16)	0.0802 (14)	
O3	0.5968 (4)	0.0055 (6)	0.66073 (16)	0.0708 (12)	
O4	0.1935 (4)	0.5291 (6)	0.31514 (15)	0.0741 (13)	
N1	0.6413 (4)	0.2391 (6)	0.44906 (17)	0.0469 (12)	
C1'	0.5253 (10)	0.5997 (11)	0.3125 (3)	0.073 (2)	0.735 (7)
H1'1	0.5814	0.6382	0.3407	0.088*	0.735 (7)
H1'2	0.4589	0.6897	0.3027	0.088*	0.735 (7)
C1	0.4509 (12)	0.5865 (18)	0.2933 (5)	0.073 (2)	0.265 (7)
H1A	0.4497	0.6807	0.3170	0.088*	0.265 (7)
H1B	0.3617	0.5872	0.2757	0.088*	0.265 (7)
C2	0.5550 (11)	0.6342 (14)	0.2577 (4)	0.107 (3)	0.735 (7)
H2A	0.5042	0.6720	0.2282	0.128*	0.735 (7)
H2B	0.6033	0.7367	0.2708	0.128*	0.735 (7)
C3	0.6599 (11)	0.5126 (15)	0.2427 (4)	0.117 (3)	0.735 (7)
H3A	0.6196	0.3962	0.2410	0.140*	0.735 (7)
H3B	0.7335	0.5103	0.2679	0.140*	0.735 (7)
C4	0.7234 (15)	0.5492 (18)	0.1943 (4)	0.120 (4)	0.735 (7)
H4A	0.7620	0.4430	0.1822	0.180*	0.735 (7)
H4B	0.7948	0.6357	0.1990	0.180*	0.735 (7)
H4C	0.6537	0.5922	0.1713	0.180*	0.735 (7)
C2'	0.611 (3)	0.540 (3)	0.2709 (9)	0.107 (3)	0.265 (7)
H2'1	0.6905	0.4764	0.2843	0.128*	0.265 (7)
H2'2	0.5562	0.4586	0.2506	0.128*	0.265 (7)
C3'	0.659 (3)	0.687 (3)	0.2392 (8)	0.117 (3)	0.265 (7)
H3'1	0.7435	0.7352	0.2543	0.140*	0.265 (7)
H3'2	0.5904	0.7792	0.2382	0.140*	0.265 (7)
C4'	0.686 (5)	0.633 (5)	0.1874 (8)	0.120 (4)	0.265 (7)
H4'1	0.7240	0.7304	0.1705	0.180*	0.265 (7)
H4'2	0.6009	0.5976	0.1709	0.180*	0.265 (7)
H4'3	0.7498	0.5369	0.1880	0.180*	0.265 (7)
C5	0.3838 (6)	0.4217 (8)	0.3608 (2)	0.0544 (14)	
C6	0.2479 (6)	0.4712 (8)	0.3590 (2)	0.0570 (15)	
C7	0.1715 (6)	0.4554 (8)	0.4011 (2)	0.0592 (16)	
H7	0.0792	0.4880	0.3995	0.071*	
C8	0.2291 (6)	0.3941 (8)	0.4439 (2)	0.0533 (15)	
H8	0.1758	0.3835	0.4710	0.064*	
C9	0.3696 (5)	0.3464 (8)	0.4475 (2)	0.0482 (13)	
C10	0.4459 (5)	0.3590 (8)	0.4044 (2)	0.0479 (13)	
C11	0.0506 (6)	0.5753 (10)	0.3118 (2)	0.080 (2)	
H11A	-0.0039	0.4744	0.3190	0.121*	
H11B	0.0260	0.6157	0.2794	0.121*	
H11C	0.0339	0.6668	0.3348	0.121*	
C12	0.4368 (5)	0.2854 (7)	0.4901 (2)	0.0482 (14)	

## supplementary materials

---

H12	0.3883	0.2809	0.5184	0.058*
C13	0.5723 (5)	0.2312 (7)	0.4923 (2)	0.0460 (13)
C14	0.5833 (5)	0.3019 (7)	0.4082 (2)	0.0488 (14)
H14	0.6356	0.3086	0.3808	0.059*
C15	0.6459 (5)	0.1642 (7)	0.5359 (2)	0.0476 (13)
C16	0.7894 (5)	0.1470 (8)	0.5374 (2)	0.0507 (14)
C17	0.8631 (6)	0.2081 (8)	0.4940 (2)	0.0573 (16)
H17A	0.8781	0.3337	0.4964	0.069*
H17B	0.9524	0.1514	0.4938	0.069*
C18	0.7835 (5)	0.1681 (8)	0.4478 (2)	0.0567 (15)
H18A	0.7797	0.0421	0.4429	0.068*
H18B	0.8292	0.2195	0.4206	0.068*
C19	0.5729 (6)	0.1186 (7)	0.5774 (2)	0.0508 (14)
H19	0.4775	0.1284	0.5772	0.061*
C20	0.6445 (6)	0.0617 (8)	0.6165 (2)	0.0532 (15)
C21	0.7863 (6)	0.0423 (8)	0.6182 (2)	0.0597 (15)
C22	0.8615 (6)	0.0853 (8)	0.5790 (2)	0.0580 (15)
H22	0.9569	0.0738	0.5803	0.070*
C23	0.7160 (7)	-0.0309 (10)	0.6915 (2)	0.0759 (19)
H23A	0.7082	-0.1461	0.7061	0.091*
H23B	0.7252	0.0554	0.7174	0.091*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0502 (4)	0.0849 (6)	0.0714 (5)	-0.0009 (4)	0.0186 (3)	0.0016 (4)
O1	0.066 (3)	0.088 (3)	0.055 (3)	0.020 (2)	0.017 (2)	-0.005 (2)
O2	0.069 (3)	0.111 (4)	0.060 (3)	0.010 (3)	-0.007 (3)	0.011 (3)
O3	0.067 (3)	0.088 (3)	0.058 (3)	0.000 (3)	0.006 (2)	0.010 (2)
O4	0.051 (3)	0.116 (4)	0.056 (3)	0.018 (3)	0.003 (2)	-0.001 (3)
N1	0.039 (2)	0.056 (3)	0.047 (3)	0.002 (2)	0.010 (2)	0.000 (2)
C1'	0.077 (4)	0.080 (4)	0.064 (4)	0.002 (4)	0.020 (4)	0.015 (4)
C1	0.077 (4)	0.080 (4)	0.064 (4)	0.002 (4)	0.020 (4)	0.015 (4)
C2	0.117 (6)	0.108 (6)	0.098 (5)	0.002 (5)	0.034 (5)	0.020 (5)
C3	0.133 (6)	0.114 (6)	0.107 (6)	0.002 (6)	0.039 (5)	0.015 (5)
C4	0.140 (9)	0.132 (10)	0.092 (7)	-0.003 (8)	0.061 (6)	0.002 (7)
C2'	0.117 (6)	0.108 (6)	0.098 (5)	0.002 (5)	0.034 (5)	0.020 (5)
C3'	0.133 (6)	0.114 (6)	0.107 (6)	0.002 (6)	0.039 (5)	0.015 (5)
C4'	0.140 (9)	0.132 (10)	0.092 (7)	-0.003 (8)	0.061 (6)	0.002 (7)
C5	0.052 (3)	0.059 (3)	0.054 (3)	0.005 (3)	0.016 (3)	-0.002 (3)
C6	0.051 (3)	0.071 (4)	0.049 (3)	0.004 (3)	0.004 (3)	-0.002 (3)
C7	0.044 (3)	0.074 (4)	0.060 (4)	0.007 (3)	0.005 (3)	-0.005 (3)
C8	0.041 (3)	0.066 (4)	0.054 (3)	0.003 (3)	0.010 (3)	-0.003 (3)
C9	0.044 (3)	0.052 (3)	0.049 (3)	-0.001 (3)	0.011 (3)	-0.008 (3)
C10	0.043 (3)	0.051 (3)	0.050 (3)	0.005 (3)	0.008 (3)	-0.002 (3)
C11	0.061 (4)	0.114 (6)	0.065 (5)	0.016 (4)	-0.003 (4)	0.005 (4)
C12	0.044 (3)	0.055 (3)	0.047 (3)	0.001 (3)	0.012 (3)	0.000 (3)
C13	0.040 (3)	0.049 (3)	0.049 (3)	-0.004 (2)	0.010 (3)	-0.005 (3)

C14	0.043 (3)	0.058 (3)	0.047 (3)	-0.001 (3)	0.014 (3)	-0.004 (3)
C15	0.043 (3)	0.046 (3)	0.055 (3)	-0.003 (3)	0.006 (3)	0.001 (3)
C16	0.044 (3)	0.053 (3)	0.056 (3)	0.001 (3)	0.007 (3)	-0.004 (3)
C17	0.041 (3)	0.064 (4)	0.068 (4)	0.007 (3)	0.011 (3)	-0.001 (3)
C18	0.043 (3)	0.066 (4)	0.062 (4)	0.011 (3)	0.014 (3)	-0.001 (3)
C19	0.041 (3)	0.055 (3)	0.057 (3)	-0.005 (3)	0.007 (3)	-0.006 (3)
C20	0.048 (3)	0.058 (4)	0.053 (4)	-0.003 (3)	0.004 (3)	-0.010 (3)
C21	0.053 (3)	0.066 (4)	0.059 (4)	0.001 (3)	-0.006 (3)	0.001 (3)
C22	0.044 (3)	0.068 (4)	0.063 (4)	0.003 (3)	0.003 (3)	-0.004 (3)
C23	0.074 (4)	0.094 (5)	0.060 (4)	0.005 (4)	0.005 (4)	-0.001 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C5	1.382 (6)	C4'—H4'3	0.9600
O1—C1	1.418 (12)	C5—C6	1.372 (8)
O1—C1'	1.477 (8)	C5—C10	1.397 (8)
O2—C21	1.370 (7)	C6—C7	1.412 (8)
O2—C23	1.434 (7)	C7—C8	1.358 (8)
O3—C20	1.389 (7)	C7—H7	0.9300
O3—C23	1.424 (7)	C8—C9	1.412 (7)
O4—C6	1.361 (7)	C8—H8	0.9300
O4—C11	1.430 (7)	C9—C12	1.389 (7)
N1—C14	1.319 (7)	C9—C10	1.431 (7)
N1—C13	1.393 (6)	C10—C14	1.403 (7)
N1—C18	1.486 (6)	C11—H11A	0.9600
C1'—C2'	1.517 (8)	C11—H11B	0.9600
C1'—H1'1	0.9700	C11—H11C	0.9600
C1'—H1'2	0.9700	C12—C13	1.378 (7)
C1—C2	1.487 (7)	C12—H12	0.9300
C1—H1A	0.9700	C13—C15	1.456 (8)
C1—H1B	0.9700	C14—H14	0.9300
C2—C3	1.454 (7)	C15—C16	1.399 (7)
C2—H2A	0.9700	C15—C19	1.416 (7)
C2—H2B	0.9700	C16—C22	1.391 (8)
C3—C4	1.517 (7)	C16—C17	1.497 (7)
C3—H3A	0.9700	C17—C18	1.483 (8)
C3—H3B	0.9700	C17—H17A	0.9700
C4—H4A	0.9600	C17—H17B	0.9700
C4—H4B	0.9600	C18—H18A	0.9700
C4—H4C	0.9600	C18—H18B	0.9700
C2'—C3'	1.510 (8)	C19—C20	1.321 (8)
C2'—H2'1	0.9700	C19—H19	0.9300
C2'—H2'2	0.9700	C20—C21	1.384 (8)
C3'—C4'	1.516 (8)	C21—C22	1.370 (8)
C3'—H3'1	0.9700	C22—H22	0.9300
C3'—H3'2	0.9700	C23—H23A	0.9700
C4'—H4'1	0.9600	C23—H23B	0.9700
C4'—H4'2	0.9600		
Cg1···Cg3 <sup>i</sup>	3.780 (4)	Cg2···Cg3 <sup>ii</sup>	3.775 (4)

## supplementary materials

---

C5—O1—C1	114.8 (6)	C8—C7—H7	119.0
C5—O1—C1'	112.8 (5)	C6—C7—H7	119.0
C21—O2—C23	105.2 (5)	C7—C8—C9	120.1 (5)
C20—O3—C23	106.2 (5)	C7—C8—H8	119.9
C6—O4—C11	117.9 (5)	C9—C8—H8	119.9
C14—N1—C13	122.5 (5)	C12—C9—C8	123.7 (5)
C14—N1—C18	117.9 (4)	C12—C9—C10	118.5 (5)
C13—N1—C18	119.6 (5)	C8—C9—C10	117.8 (5)
O1—C1'—C2'	95.0 (10)	C5—C10—C14	122.4 (5)
O1—C1'—H1'1	112.7	C5—C10—C9	120.8 (5)
C2'—C1'—H1'1	112.7	C14—C10—C9	116.7 (5)
O1—C1'—H1'2	112.7	O4—C11—H11A	109.5
C2'—C1'—H1'2	112.7	O4—C11—H11B	109.5
H1'1—C1'—H1'2	110.2	H11A—C11—H11B	109.5
O1—C1—C2	121.2 (9)	O4—C11—H11C	109.5
O1—C1—H1A	107.0	H11A—C11—H11C	109.5
C2—C1—H1A	107.0	H11B—C11—H11C	109.5
O1—C1—H1B	107.0	C13—C12—C9	122.8 (5)
C2—C1—H1B	107.0	C13—C12—H12	118.6
H1A—C1—H1B	106.8	C9—C12—H12	118.6
C3—C2—C1	122.8 (8)	C12—C13—N1	116.9 (5)
C3—C2—H2A	106.6	C12—C13—C15	124.6 (5)
C1—C2—H2A	106.6	N1—C13—C15	118.6 (5)
C3—C2—H2B	106.6	N1—C14—C10	122.5 (5)
C1—C2—H2B	106.6	N1—C14—H14	118.7
H2A—C2—H2B	106.6	C10—C14—H14	118.7
C2—C3—C4	117.0 (8)	C16—C15—C19	119.6 (5)
C2—C3—H3A	108.0	C16—C15—C13	120.2 (5)
C4—C3—H3A	108.0	C19—C15—C13	120.2 (5)
C2—C3—H3B	108.0	C22—C16—C15	121.0 (5)
C4—C3—H3B	108.0	C22—C16—C17	121.2 (5)
H3A—C3—H3B	107.3	C15—C16—C17	117.7 (5)
C3—C4—H4A	109.5	C18—C17—C16	111.6 (5)
C3—C4—H4B	109.5	C18—C17—H17A	109.3
H4A—C4—H4B	109.5	C16—C17—H17A	109.3
C3—C4—H4C	109.5	C18—C17—H17B	109.3
H4A—C4—H4C	109.5	C16—C17—H17B	109.3
H4B—C4—H4C	109.5	H17A—C17—H17B	108.0
C3'—C2'—C1'	114.1 (9)	C17—C18—N1	110.4 (5)
C3'—C2'—H2'1	108.7	C17—C18—H18A	109.6
C1'—C2'—H2'1	108.7	N1—C18—H18A	109.6
C3'—C2'—H2'2	108.7	C17—C18—H18B	109.6
C1'—C2'—H2'2	108.7	N1—C18—H18B	109.6
H2'1—C2'—H2'2	107.6	H18A—C18—H18B	108.1
C2'—C3'—C4'	114.3 (10)	C20—C19—C15	118.0 (5)
C2'—C3'—H3'1	108.7	C20—C19—H19	121.0
C4'—C3'—H3'1	108.7	C15—C19—H19	121.0
C2'—C3'—H3'2	108.7	C19—C20—C21	123.0 (6)
C4'—C3'—H3'2	108.7	C19—C20—O3	128.7 (5)

H3'1—C3'—H3'2	107.6	C21—C20—O3	108.3 (6)
C3'—C4'—H4'1	109.5	O2—C21—C22	127.7 (6)
C3'—C4'—H4'2	109.5	O2—C21—C20	111.1 (6)
H4'1—C4'—H4'2	109.5	C22—C21—C20	121.2 (6)
C3'—C4'—H4'3	109.5	C21—C22—C16	117.3 (5)
H4'1—C4'—H4'3	109.5	C21—C22—H22	121.3
H4'2—C4'—H4'3	109.5	C16—C22—H22	121.3
C6—C5—O1	121.4 (6)	O3—C23—O2	108.1 (5)
C6—C5—C10	119.8 (5)	O3—C23—H23A	110.1
O1—C5—C10	118.8 (5)	O2—C23—H23A	110.1
O4—C6—C5	116.8 (5)	O3—C23—H23B	110.1
O4—C6—C7	123.7 (5)	O2—C23—H23B	110.1
C5—C6—C7	119.5 (6)	H23A—C23—H23B	108.4
C8—C7—C6	121.9 (5)		

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+1, -y+1, -z+1$ .

*Hydrogen-bond geometry (Å, °)*


$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C12—H12 $\cdots$ Br1	0.93	2.77	3.658 (5)	161
C2—H2A $\cdots$ O2 <sup>iii</sup>	0.97	2.64	3.404 (14)	136

Symmetry codes: (iii)  $x-1/2, -y+1/2, z-1/2$ .

## supplementary materials

---

Fig. 1



Br1