

Heparin-free veno-venous extracorporeal membrane oxygenation in a multiple trauma patient

A case report

Youn Young Lee, MD^a, Hee Jung Baik, MD, PhD^{b,*}, Heeseung Lee, MD, PhD^b, Chi Hyo Kim, MD, PhD^b, Rack Kyung Chung, MD, PhD^b, Jong In Han, MD, PhD^b, Hyunyoung Joo, MD^a, Jae Hee Woo, MD, PhD^b

Abstract

Rationale: Extracorporeal membrane oxygenation (ECMO) in multiple trauma patients with post-traumatic respiratory failure can be quite challenging because of the need for systemic anticoagulation, which may lead to excessive bleeding. In the last decade, there is a growing body of evidence that veno-venous ECMO (VV-ECMO) is lifesaving in multiple trauma patients with acute respiratory distress syndrome, thanks to technical improvements in ECMO devices.

Patient concerns: We report a case of a 17-year-old multiple trauma patient who was drunken and had confused mentality.

Diagnoses: She was suffered from critical respiratory failure (life-threatening hypoxemia and severe hypercapnia/acidosis lasting for 70 minutes) accompanied by cardiac arrest and trauma-induced coagulopathy during general anesthesia.

Interventions: We decided to start heparin-free VV-ECMO after cardiac arrest considering risk of hemorrhage.

Outcomes: She survived with no neurologic sequelae after immediate treatment with heparin-free VV-ECMO.

Lessons: Heparin-free VV-ECMO can be used as a resuscitative therapy in multiple trauma patients with critical respiratory failure accompanied by coagulopathy. Even in cases in which life-threatening hypoxemia and severe hypercapnia/acidosis last for >1 hours during CPR for cardiac arrest, VV-ECMO could be considered a potential lifesaving treatment.

Abbreviations: ABGA = arterial blood gas analysis, ACT = activated coagulation time, aPTT = activated partial thromboplastin time, ARDS = acute respiratory distress syndrome, ASV = adaptive support ventilation, BIS = bispectral index, BP = blood pressure, CPR = cardiopulmonary resuscitation, CRRT = continuous renal replacement therapy, EC = effect-site concentration, ECMO = extracorporeal membrane oxygenation, ECPR = extracorporeal cardiopulmonary resuscitation, ER = emergency room, ET = endotracheal tube, FFP = fresh frozen plasma, F_IO₂ = inspired oxygen fraction, Hb = hemoglobin, Hct = hematocrit, HR = heart rate, ICU = intensive care unit, INR = international normalized ratio, ISS = injury severity score, OR = operating room, PEEP = positive end-expiratory pressure, PIP = peak inspiratory airway pressure, POD = postoperative day, pRBCs = packed red blood cells, PRVC = pressure-regulated volume-controlled, PT = prothrombin time, ROSC = return of spontaneous circulation, RR = respiratory rate, SpO₂ = saturation on pulse oximetry, TIC = trauma-induced coagulopathy, V_T = tidal volume, VV-ECMO = veno-venous extracorporeal membrane oxygenation.

Keywords: acute respiratory failure, extracorporeal membrane oxygenation, heparin, trauma

1. Introduction

The use of extracorporeal membrane oxygenation (ECMO) in multiple trauma patients with refractory respiratory failure has

been suggested as salvage therapy that can improve outcomes.^[1-3] However, the clinical benefit of ECMO support in critical trauma patients remains unclear. Careful consideration should be given to the risks and benefits before starting ECMO support because of the risk of organ bleeding, especially in multiple trauma patients. Heparin-free VV-ECMO could be an effective treatment strategy in trauma patients with contraindications to systemic anticoagulation, including multiple trauma patients with traumatic brain injury.^[4] We report a case of a 17-year-old multiple trauma patient who suffered from critical respiratory failure with life-threatening hypoxemia and severe hypercapnia/acidosis lasting for 70 minutes accompanied by cardiac arrest and trauma-induced coagulopathy during general anesthesia. She survived with no neurologic sequelae after immediate treatment with heparin-free VV-ECMO. The patient had provided informed consent for publication of this case report.

2. Case presentation

A 17-year-old girl, 160 cm in height and weighing 48 kg (BMI, 18.75 kg/m²) came to the emergency room (ER) due to a

Editor: N/A.

The authors have no funding and conflicts of interest to disclose.

^aDepartment of Anesthesiology and Pain Medicine, Ewha Womans University Hospital, ^bDepartment of Anesthesiology and Pain Medicine, College of Medicine, Ewha Womans University, Seoul, South Korea.

*Correspondence: Hee Jung Baik, 1071 Anyangcheon-ro, Yangcheon-gu, Seoul, Zip Code 07985, South Korea (e-mail: baikhj@ewha.ac.kr).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lee YY, Baik HJ, Lee H, Kim CH, Chung RK, Han JI, Joo H, Woo JH. Heparin-free veno-venous extracorporeal membrane oxygenation in a multiple trauma patient: a case report. *Medicine* 2020;99:5(e19070).

Received: 6 June 2019 / Received in final form: 4 December 2019 / Accepted: 7 January 2020

<http://dx.doi.org/10.1097/MD.00000000000019070>

motorcycle accident. She was drunk and had a confused mental state. She had multiple fractures in her left femur, right pelvis, both mandibles, multiple right ribs (1–5th), and tooth #21, and severe right lung contusion with several lacerations on the scalp. Her injury severity score (ISS) was 34 points, representing a severe grade of injury.

Her initial vital signs and laboratory findings were as follows: blood pressure (BP), 98/60 mm Hg; heart rate (HR), 118 beats/min; respiratory rate (RR), 38/minute; saturation on pulse oximetry (SpO₂), 98% in room air; hemoglobin (Hb), 10.1 g/dL; hematocrit (Hct), 30.6%; platelets, 385,000; prothrombin time (PT), 14.4 seconds; INR, 1.27; and activated partial thromboplastin time (aPTT), 29.1 seconds. Five minutes after she entered the ER, her SpO₂ suddenly dropped to 85%, and she was intubated immediately and mechanically ventilated under pressure-regulated volume-controlled (PRVC) mode {tidal volume (V_T), 350 mL; inspired oxygen fraction (F_IO₂), 0.4; RR, 14 breaths/min; positive end-expiratory pressure (PEEP), 7 cm H₂O}, and a right-sided chest tube was inserted.

Catheters were inserted into both her subclavian veins for active fluid resuscitation and blood transfusion. Even though she was given 2000 mL of crystalloid fluid and 5 units of packed red blood cells (pRBCs) for an hour in the ER, her BP suddenly dropped to 66/39 mm Hg and her HR increased to 130 beats/min; the ER physicians decided to administer 0.1 µg/kg/min norepinephrine. At the same time, her Hb, Hct, and platelet levels decreased to 6.8 g/dL, 20.5%, and 110,000, respectively. She was transfused with an additional 5000 mL of crystalloid fluid, 5 units of pRBCs, and 10 units of fresh frozen plasma (FFP), and was evaluated via CT of the brain (Fig. 1), abdomen and pelvis, facial bone, and chest PA (Fig. 2(A)). The ER physicians decided that she should undergo emergency surgical procedures on her left supracondylar femur fracture and both mandible fractures.

She arrived at the operating room (OR) in an intubated state with continuous infusion of 0.2 µg/kg/min norepinephrine; we immediately applied ventilation under volume-controlled (VC) mode {V_T, 550 mL; F_IO₂, 1.0; RR, 15 breaths/min; PEEP, 5 cm H₂O}. Her left radial artery was cannulated for arterial BP monitoring. Initial BP, HR, SpO₂, and bispectral index (BIS) were 126/70 mm Hg, 120 beats/min, 100%, and 45, respectively. Fifteen minutes after the induction of total intravenous anesthesia using target effect-site concentration (EC)-controlled infusion (EC of propofol, 1.5 µg/mL; EC of remifentanyl, 2 ng/mL), the surgery was started.

Twenty-five minutes after the start of surgery, her BP suddenly dropped from 130/100 to 78/40 despite a high dose of norepinephrine (3 µg/kg/min), her HR decreased from 151 to 90 beats/min, and her SpO₂ to 75%. Peak inspiratory airway



Figure 1. CT of the brain. There are hematoma and emphysema along the right frontal, parietal scalp and both occipital scalp area, but there is no evidence of intracranial hemorrhage. CT=computed tomography (CT).

pressure (PIP) suddenly increased from 28 to 35 mm Hg with a large amount of pink frothy watery secretion discharged from the endotracheal tube (ET). We ventilated her lungs manually with frequent endotracheal suction, and her SpO₂ was 52%. Consecutively severe bradycardia (40 beats/min) followed by cardiac arrest occurred; cardiopulmonary resuscitation (CPR) was started and a 1-mg epinephrine bolus was given five times during a 45-minute CPR. Severe hypoxemia, hypercapnia, and acidosis developed (Table 1). After the 45-minute CPR, return of spontaneous circulation (ROSC) was achieved and a chest X-ray was taken (Fig. 2(B)), which showed increased haziness in both lungs suggesting pulmonary edema, consistent with the large amount (2.4 L/1.5 hours) of pink frothy watery secretion from the ET even after ROSC.

We decided to start VV-ECMO therapy because of acute refractory lung failure and impending cardiac arrest even after ROSC (Table 1). We decided not to give bolus heparin considering the risk of hemorrhage. Her activated coagulation time (ACT) was 510 seconds and other coagulation tests showed prolonged times (Table 1). VV-ECMO (PLS-Set and Quadrox PLS; Maquet Cardiopulmonary AG, Rastatt, Germany) was applied on the right femoral vein, with a 19-Fr access cannula



Figure 2. Chest images. (A) Preoperative chest PA (7 h before anesthesia), (B) intraoperative chest AP (5 min before ECMO), and (C) postoperative chest AP (3 h after ECMO). ECMO=immediately after extracorporeal membrane oxygenation.

Table 1
Laboratory findings of the patient in operating room.

	AN	OP	CPR start	CPR 20 min	ROSC	ECMO
ABGA						
pH	7.25	7.02	6.99	6.85	6.81	6.97
PaCO ₂ (mm Hg)	30	47	76	112	115	60
PaO ₂ (mm Hg)	74	100	46	32	31	60
HCO ₃ ⁻ (mmol/L)	13.2	12.1	18.6	19.6	18.3	13.8
BE (mmol/L)	-14	-18.9	-13.1	-14.1	-16.1	-18
SaO ₂ (%)	92	94	55	23	19	71
Hb/Hct (g/dL, %)	10.4/29	11.2/31	15.5/43	14.8/41	13.3/37	13.0/36
Electrolytes						
Na ⁺ (mmol/L)	140	138	142	144	144	143
K ⁺ (mmol/L)	5.4	5.3	3.6	4.3	3.8	3.7
Ca ²⁺ (mg/dL)	3.0	4.3	2.5	1.3	2.1	2.1
Lactate (mg/dL)	61	99	114	325	278	207
Coagulation tests						
PT (s)/INR	18/1.59				27/2.57	
aPTT (s)	44.8				>180	

ABGA = arterial blood gas analysis, AN = start of anesthesia, aPTT = activated partial thromboplastin time, BE = base excess, CPR = cardiopulmonary resuscitation, ECMO = immediately after extracorporeal membrane oxygenation, Hb = hemoglobin, Hct = hematocrit, INR = international normalized ratio, OP = start of operation, PT = prothrombin time, ROSC = return of spontaneous circulation. Normal reference range: ABGA; pH 7.34–7.44, PaCO₂ 35–45 mm Hg, PaO₂ 75–100 mm Hg, HCO₃⁻ 22–26 mmol/L, BE -2 to +2 mmol/L, SaO₂ 93–97%; Hb 12.0–15.5 g/dL, hematocrit (Hct) 37–48%. Electrolytes: Na⁺ 135–145 mmol/L, K⁺ 3.5–5.0 mmol/L, Ca²⁺ 4.6–5.4 mg/dL; lactate 2.7–13.5 mg/dL. Coagulation tests: PT 10.7–13.8 s, INR 0.9–1.2, aPTT 20.6–33.0 s.

positioned 10 cm lower than the return cannula (21 Fr) by a cardiothoracic surgeon. As the ECMO flow started, her severe hypoxemia and hypercapnia began to improve, as shown in Table 1. Even though we administered 20 mg furosemide twice, her urine output was <0.5 mL/kg/hour and the estimated blood loss was <30 mL. The surgery was discontinued and the patient was transferred to the intensive care unit (ICU); ECMO flow was 8.95 L/min with FiO₂ of 1.0 and gas flow of 4 L/min; continuous renal replacement therapy (CRRT) was immediately started.

Three days after she entered the ICU, the ECMO flow rate gradually decreased to 3.19 L/min, with FiO₂ of 0.21 under adaptive support ventilation (ASV) mode {FiO₂ 0.3, V_T 250 mL, PEEP 16 cm H₂O}. Her vital signs were stable without any inotropics and arterial blood gas analysis (ABGA) values showed a pH 7.382, PCO₂ 44.1, PO₂ 94.9, and SaO₂ 97.1%. We decided to discontinue the VV-ECMO. Ventilator care was stopped on postoperative day (POD) 7, and 7L T-piece oxygen was started. On POD 10, we decided to stop CRRT and apply 2 L/min O₂ via nasal prong, and her mental status improved from drowsy to alert.

During the 33 days of ICU care, she underwent three additional surgeries under general anesthesia. No intracranial hemorrhage was seen in serial brain CT, and she was discharged with no neurologic complications after 128 days of hospitalization.

3. Discussion

Although the technique of ECMO is highly invasive, it has been shown to be a lifesaving therapy in non-traumatic and traumatic respiratory failure with refractory hypoxemia that persists despite conventional treatment.^[2,3,5] The use of ECMO in multiple trauma patients may be challenging considering the risk of hemorrhage, especially in cases of severe coagulopathy, contraindications to anticoagulant use, or traumatic brain injury. Nevertheless, recent growing evidence suggests that applying ECMO as rescue therapy in trauma patients with severe respiratory failure may provide potential survival benefits that outweigh the associated risk of bleeding.^[2,6,7] Recent advances

in ECMO devices and the application of heparin-free ECMO have dramatically increased safety profiles and expanded the application of ECMO to severe multiple trauma patients with coexisting bleeding shock or even traumatic brain injury.^[4,8]

Our multiple trauma patient, who suffered from severe hypoxemia (31 mm Hg PaO₂ at F_IO₂ 1.0) and hypercarbia (115 mm Hg) accompanied by cardiac arrest due to critical respiratory failure, survived due to application of heparin-free VV-ECMO.

Due to the risk of clot formation in patients on ECMO, anticoagulation is necessary for the prevention of thrombosis while avoiding excessive bleeding. Although anticoagulation in VV-ECMO is controlled by an ACT of 160 to 220 seconds and aPTT of 50 to 80 seconds in general,^[9] aPTT of 40 to 55 seconds may be sufficient to maintain modern VV-ECMO using heparin-bonded tubing.^[10] We decided not to give her anticoagulants during or after insertion of the ECMO circuit because coagulation tests before induction of anesthesia indicated prolonged times (PT, 18 seconds; INR, 1.59; aPTT, 44.8 seconds) and her platelet count was 84,000, indicating that trauma-induced coagulopathy (TIC) had developed. TIC, which is acute intrinsic coagulopathy arising in severely injured trauma patients, occurs in the presence of both tissue hypoperfusion due to hemorrhage and severe anatomical tissue injury.^[11] She was intubated and received a massive blood transfusion and fluid resuscitation with vasopressor therapy due to hemorrhagic shock and severe hypoxemia in the ER, which was the likely cause of TIC. Immediately after ECMO started, her coagulation tests indicated more prolonged times (ACT, 510 seconds; PT, 27 seconds; INR, 2.59; and aPTT > 180 seconds). Her prolonged PT, INR, and aPTT, which were monitored daily, were sustained during (94 hours) and 4 hours after ECMO (PT, 16 seconds; INR, 1.41; and aPTT, 51.9 seconds). Therefore, there was no need to give her anticoagulants during the ECMO.

Indications for VV-ECMO in acute respiratory failure include hypoxemic and/or hypercarbic respiratory failure, respiratory failure in lung transplantation, bronchopleural fistulas and pulmonary air leaks, and complex airway management.^[9] The

most common indication for ECMO in respiratory failure is severe acute respiratory distress syndrome (ARDS).^[12] It has been reported that 4.6% of trauma patients develop ARDS and independent predictors for ARDS include injury severity, thoracic injury, polytrauma, pneumonia, and receiving more than 5 units of FFP and 6 to 10 units of pRBC, especially during early (first 24 hours) transfusion of pRBC.^[13] Although the cause of acute severe respiratory failure in our patient has not been fully elucidated and may involve transfusion-related acute lung injury, severe pulmonary edema or hemorrhage, or trauma-induced ARDS, she had many risk factors for traumatic ARDS including high ISS (34 points), polytrauma, severe right lung contusion, receiving early transfusion of pRBC (10 units within 8 hours) and receiving 10 units of FFP in the ER due to severe shock. Various triage criteria for ECMO in hypoxic and hypercarbic respiratory failure have been reported as follows:

- (1) $\text{PaO}_2/\text{F}_1\text{O}_2$ (P/F ratio) < 100 with $\text{F}_1\text{O}_2 > 90\%$ despite optimal care for 6 hours or more,^[14]
- (2) P/F ratio of < 80 with PEEP 15–20 cm H_2O , hypercarbia with high plateau pressures > 30 cm H_2O and a pH of 7.15,^[15]
- (3) P/F ratio of < 50 with $\text{F}_1\text{O}_2 > 80\%$ for > 3 hours, or
- (4) P/F ratio < 80 for 6 hours with pH < 7.25 for 6 hours despite optimum mechanical ventilation.^[9]

However, these criteria are applicable to ARDS patients in the ICU. In our case, cardiac arrest occurred due to severe acute respiratory failure with massive pulmonary edema during general anesthesia. Despite ROSC after a 45-minute CPR, severe hypoxemia (31 mm Hg PaO_2), hypercapnia (115 mm Hg PaCO_2), and acidosis (pH 6.81) continued with a continuous large amount of pink frothy watery secretion from the ET, preventing us from ventilating her lungs due to the need for frequent suction. Therefore, we applied VV-ECMO as a lifesaving treatment to prevent recurrence of cardiac arrest. This is the last choice in such an impending cardiac arrest situation. Thus, we consider that VV-ECMO in this patient was similar to extracorporeal cardiopulmonary resuscitation (ECPR) in spite of its application after ROSC. ECPR is a method of CPR using ECMO to support circulation in refractory cardiac arrest patients.^[16,17] Despite inconclusive evidence of the role of ECPR in cardiac arrest,^[16] there is a report of patient survival with good neurologic outcomes due to ECPR applied after an 80-minute CPR.^[17]

Absolute contraindication to ECMO in respiratory failure is the presence of severe irreversible respiratory failure, and relative contraindications include prolonged use of high-pressure ventilation or high F_1O_2 , limited vascular access, contraindications to the use of anticoagulation, the presence of disease or organ dysfunction (e.g., severe irreversible brain injury or untreatable metastatic cancer), inability to receive blood products, high-body mass index > 45, and major immunosuppression.^[9,12]

Neurological complications including intracranial bleeding, brain death, and ischemic stroke in VV-ECMO could be critical and major causes of death.^[18] Improvements in the ECMO technique have reduced the incidence of neurological complications to approximately 7%.^[18,19] The pre-ECMO risk factors for neurological complications during VV-ECMO have been reported to be cardiac arrest, CRRT, and hyperbilirubinemia.^[18] Two different mechanisms of neurological injury may be involved in VV-ECMO. One is preexisting neurological injury before ECMO including anoxic brain injury or hypoperfusion-related brain injury. The other is anticoagulation.^[18] In the

decision to start ECMO in our patient, life-threatening hypoxemia (PaO_2 , 31–46 mm Hg) and severe hypercapnia (PaCO_2 , 76–115 mm Hg)/acidosis (pH 6.81–6.99) lasting for 70 minutes and the 45-minute CPR were thought to be risk factors for neurological complications during VV-ECMO. Our minimal knowledge of the patient's neurological status due to her drunken state despite absence of intracranial hemorrhage in her initial brain CT, the episode of hypoxemia with hemorrhagic shock in the ER, TIC, or CRRT in the ICU were also possible risk factors. Fortunately, she recovered with no neurological complications after three days of ECMO therapy and three surgeries under general anesthesia. Her recovery is attributed to immediate and successful CPR, young age, good cardiac function, and early ECMO therapy. It has been recently reported that early initiation of ECMO may lead to positive outcomes.^[20] Patient selection, monitoring for early detection of stroke and intracranial bleeding, and timing of ECMO are important factors influencing improvement in survival of trauma patients undergoing ECMO.^[19,20]

4. Conclusions

Heparin-free VV-ECMO can be used as a resuscitative therapy in multiple trauma patients with critical respiratory failure accompanied by coagulopathy. Even in cases in which life-threatening hypoxemia and severe hypercapnia/acidosis last for > 1 hours during CPR for cardiac arrest, VV-ECMO could be considered a potential lifesaving treatment.

Author contributions

Data curation: Hyunyoung Joo.

Supervision: Hee Jung Baik, Heeseung Lee, Chi Hyo Kim, Rack Kyung Chung, Jong In Han, Jae Hee Woo.

Writing – original draft: Youn Young Lee, Hee Jung Baik.

Writing – review & editing: Youn Young Lee, Hee Jung Baik.

References

- [1] Madershahian N, Wittwer T, Strauch J, et al. Application of ECMO in multitrauma patients with ARDS as rescue therapy. *J Card Surg* 2007;22:180–4.
- [2] Biderman P, Einav S, Fainblut M, et al. Extracorporeal life support in patients with multiple injuries and severe respiratory failure: a single-center experience? *J Trauma Acute Care Surg* 2013;75:907–12.
- [3] Burke CR, Crown A, Chan T, et al. Extracorporeal life support is safe in trauma patients. *Injury* 2017;48:121–6.
- [4] Muellenbach RM, Kredel M, Kunze E, et al. Prolonged heparin-free extracorporeal membrane oxygenation in multiple injured acute respiratory distress syndrome patients with traumatic brain injury. *J Trauma Acute Care Surg* 2012;72:1444–7.
- [5] Peek GJ, Mugford M, Tiruvoipati R, et al. Efficacy and economic assessment of conventional ventilatory support versus extracorporeal membrane oxygenation for severe adult respiratory failure (CESAR): a multicentre randomised controlled trial. *Lancet* 2009;374:1351–63.
- [6] Bedeir K, Seethala R, Kelly E. Extracorporeal life support in trauma: worth the risks? A systematic review of published series. *J Trauma Acute Care Surg* 2017;82:400–6.
- [7] Bonnell M, Cairl NS, Heidt DH. Extracorporeal membrane oxygenation (ECMO) as initial therapy for blunt thoracic trauma with associated intracranial hemorrhage: a case report and review of the literature. *J Trauma Crit Care* 2018;2:1–3.
- [8] Arlt M, Philipp A, Voelkel S, et al. Extracorporeal membrane oxygenation in severe trauma patients with bleeding shock. *Resuscitation* 2010;81:804–9.
- [9] Sen A, Callisen HE, Alwardt CM, et al. Adult venovenous extracorporeal membrane oxygenation for severe respiratory failure: current status and future perspectives. *Ann Card Anaesth* 2016;19:97–111.

- [10] Lamb KM, Cowan SW, Evans N, et al. Successful management of bleeding complications in patients supported with extracorporeal membrane oxygenation with primary respiratory failure. *Perfusion* 2013;28:125–31.
- [11] White NJ. Mechanisms of trauma-induced coagulopathy. *Hematology Am Soc Hematol Educ Program* 2013;2013:660–3.
- [12] Abrams D, Brodie D. Extracorporeal membrane oxygenation for adult respiratory failure: 2017 update. *Chest* 2017;152:639–49.
- [13] Chaiwat O, Lang JD, Vavilala MS, et al. Early packed red blood cell transfusion and acute respiratory distress syndrome after trauma. *Anesthesiology* 2009;110:351–60.
- [14] Extracorporeal Life Support Organization. Extracorporeal Life Support (ECLS) Registry Report 2017 . Available at: <https://www.else.org/Registry/Statistics/InternationalSummary.aspx> [accessed date October 15, 2018].
- [15] Brodie D, Bacchetta M. Extracorporeal membrane oxygenation for ARDS in adults. *N Engl J Med* 2011;365:1905–14.
- [16] Holmberg MJ, Geri G, Wiberg S, et al. Extracorporeal cardiopulmonary resuscitation for cardiac arrest: a systematic review. *Resuscitation* 2018;131:91–100.
- [17] Moon SH, Kim JW, Byun JH, et al. Case of a cardiac arrest patient who survived after extracorporeal cardiopulmonary resuscitation and 1.5 hours of resuscitation: a case report. *Medicine* 2017;96:e8646.
- [18] Hirose H, Cavarocchi NC. Neurological complications during venovenous extracorporeal membrane oxygenation. *J Thorac Dis* 2017; 9:1796–7.
- [19] Lorusso R, Gelsomino S, Parise O, et al. Neurologic injury in adults supported with veno-venous extracorporeal membrane oxygenation for respiratory failure: findings from the extracorporeal life support organization database. *Crit Care Med* 2017;45:1389–97.
- [20] Bosarge PL, Raff LA, McGwin GJr, et al. Early initiation of extracorporeal membrane oxygenation improves survival in adult trauma patients with severe adult respiratory distress syndrome. *J Trauma Acute Care Surg* 2016;81:236–43.