

6-Benzylsulfanyl-9*H*-purine

Ismat Fatima, Munawar Ali Munawar,* Misbahul Ain Khan, Sohail Nadeem and Rana Amjad

Institute of Chemistry, University of the Punjab, New Campus, Lahore, Pakistan
Correspondence e-mail: munawarlimunawar@yahoo.com


Received 26 October 2009; accepted 28 October 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$;
 R factor = 0.040; wR factor = 0.088; data-to-parameter ratio = 18.7.

The phenyl ring of the title compound, $\text{C}_{12}\text{H}_{10}\text{N}_4\text{S}$, a purine derivative, is oriented at a dihedral angle of $76.65(6)^\circ$ with respect to the purine ring system. An intermolecular $\text{N}-\text{H} \cdots \text{N}$ hydrogen bonds stabilizes the crystal structure.

Related literature

For the biological activity of purine derivatives, see: Lepage *et al.* (1964); Mitsuya & Border (1986); Ragazzi *et al.* (1989).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{10}\text{N}_4\text{S}$

$M_r = 242.30$

Orthorhombic, $P2_12_12_1$
 $a = 5.5717(3)\text{ \AA}$

$b = 9.4733(4)\text{ \AA}$

$c = 22.4656(14)\text{ \AA}$

$V = 1185.79(11)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.26\text{ mm}^{-1}$

$T = 296\text{ K}$
 $0.29 \times 0.12 \times 0.09\text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.930$, $T_{\max} = 0.978$

7941 measured reflections
2941 independent reflections
2102 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.088$
 $S = 0.98$
2941 reflections
157 parameters
H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.17\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1207 Friedel pairs
Flack parameter: $-0.09(8)$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N}3-\text{H3N} \cdots \text{N}4^i$	0.894 (19)	1.892 (19)	2.773 (2)	167.9 (18)

Symmetry code: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

IF acknowledges the Higher Education Commission, Pakistan for providing funding for this research.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5117).

References

- Bruker (2007). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
Lepage, G. A., Junga, J. G. & Bowman, B. (1964). *Cancer Res.* **24**, 835–840.
Mitsuya, H. & Border, S. (1986). *Proc. Natl. Acad. Sci. USA*, **83**, 1911–1915.
Ragazzi, E., Froldi, G., Santi-Sonein, E., Borea, P. A. & Fassina, G. (1989). *Pharmacol. Res.* **21**, 707–717.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o2994 [doi:10.1107/S1600536809045140]

6-Benzylsulfanyl-9H-purine

I. Fatima, M. A. Munawar, M. A. Khan, S. Nadeem and R. Amjad

Comment

The synthesis of purine derivatives has received considerable attention on account of their biological activity especially as antitumor (Lepage *et al.*, 1964), anti HTLV (Mitsuya & Border, 1986) and anti asthmatic (Ragazzi *et al.*, 1989) agents. During our search to find new synthetic antithyroid agents, certain purine derivatives were prepared. 6-(benzylthio)-7H-purine was synthesized during such an effort. The compound is now under study for possible antithyroid activity.

The phenyl ring is oriented at adihedral angle of 76.65 (6) ° with respect to purine ring system. An intermolecular N—H···N hydrogen bonds stabilizes the crystal structure.

Experimental

To a solution of 6-mercaptopurine (0.171 g) 1 mmol in 2 N NaOH (10 ml), benzyl bromide 1 mmol (0.171 g) was added and stirred at room temperature for 30 minutes. The pH of the mixture was adjusted at 5 with glacial acetic acid and the precipitates were collected, washed with water and diethyl ether. The crystals suitable for X-ray diffraction were grown in dichloromethane by slow evaporation at room temperature.

Refinement

The H-atoms bonded to C were refined geometrically and treated as riding atoms with C_{aromatic}—H = 0.93 Å and C_{methylene}—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The N—H atom was refined at calculated position with N—H = 0.894 (19) Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$ (parent N-atom)

Figures




Fig. 1. The crystal structure diagram of the title compound duly labeled with 50% probability level of drawn thermal ellipsoids.




Fig. 2. Unit cell diagram showing the intermolecular hydrogen bonding using dashed lines. The hydrogen atoms not involved in hydrogen bonding have been omitted for clarity.

supplementary materials

6-Benzylsulfanyl-9*H*-purine

Crystal data

C ₁₂ H ₁₀ N ₄ S	$F_{000} = 504$
$M_r = 242.30$	$D_x = 1.357 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 2230 reflections
$a = 5.5717 (3) \text{ \AA}$	$\theta = 2.3\text{--}24.8^\circ$
$b = 9.4733 (4) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$c = 22.4656 (14) \text{ \AA}$	$T = 296 \text{ K}$
$V = 1185.79 (11) \text{ \AA}^3$	Needle, red
$Z = 4$	$0.29 \times 0.12 \times 0.09 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer	2941 independent reflections
Radiation source: fine-focus sealed tube	2102 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.032$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -7 \rightarrow 6$
$T_{\text{min}} = 0.930$, $T_{\text{max}} = 0.978$	$k = -12 \rightarrow 12$
7941 measured reflections	$l = -28 \rightarrow 29$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.040$	$w = 1/[\sigma^2(F_o^2) + (0.0419P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.088$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 0.98$	$\Delta\rho_{\text{max}} = 0.17 \text{ e \AA}^{-3}$
2941 reflections	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$
157 parameters	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1207 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: -0.09 (8)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.47086 (11)	0.20744 (5)	0.15044 (2)	0.05534 (18)
N1	0.3943 (4)	-0.07206 (16)	0.13660 (7)	0.0553 (5)
N2	0.6348 (4)	-0.25776 (16)	0.18001 (8)	0.0577 (5)
N3	0.9619 (4)	-0.16476 (15)	0.23955 (8)	0.0519 (5)
H3N	1.020 (4)	-0.2456 (19)	0.2541 (8)	0.062*
N4	0.8929 (3)	0.06585 (15)	0.22833 (7)	0.0488 (4)
C1	0.7639 (4)	-0.15403 (16)	0.20432 (9)	0.0454 (5)
C2	0.7219 (4)	-0.01004 (17)	0.19749 (8)	0.0432 (5)
C3	0.5298 (4)	0.02806 (17)	0.16181 (8)	0.0447 (5)
C4	0.4558 (4)	-0.2074 (2)	0.14777 (10)	0.0618 (6)
H4	0.3578	-0.2750	0.1302	0.074*
C5	1.0294 (4)	-0.03046 (17)	0.25248 (10)	0.0530 (5)
H5	1.1608	-0.0090	0.2764	0.064*
C6	0.2345 (4)	0.2002 (2)	0.09542 (10)	0.0662 (6)
H6A	0.2763	0.1328	0.0646	0.079*
H6B	0.0862	0.1694	0.1140	0.079*
C7	0.2003 (4)	0.3442 (2)	0.06836 (9)	0.0500 (5)
C8	0.3547 (5)	0.3944 (2)	0.02616 (10)	0.0646 (6)
H8	0.4883	0.3409	0.0158	0.077*
C9	0.3174 (5)	0.5222 (2)	-0.00144 (11)	0.0741 (8)
H9	0.4228	0.5532	-0.0307	0.089*
C10	0.1270 (6)	0.6027 (2)	0.01415 (12)	0.0708 (7)
H10	0.1011	0.6889	-0.0046	0.085*
C11	-0.0261 (5)	0.5576 (3)	0.05709 (12)	0.0780 (7)
H11	-0.1550	0.6139	0.0684	0.094*
C12	0.0092 (5)	0.4280 (3)	0.08406 (10)	0.0690 (6)
H12	-0.0976	0.3972	0.1131	0.083*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0640 (4)	0.0405 (3)	0.0616 (3)	-0.0021 (3)	-0.0128 (3)	0.0078 (2)

supplementary materials

N1	0.0659 (12)	0.0470 (9)	0.0530 (11)	-0.0109 (9)	-0.0074 (9)	0.0019 (8)
N2	0.0708 (12)	0.0373 (8)	0.0650 (12)	-0.0098 (9)	-0.0048 (11)	-0.0040 (8)
N3	0.0602 (12)	0.0309 (7)	0.0647 (11)	0.0005 (8)	-0.0058 (10)	0.0064 (7)
N4	0.0571 (11)	0.0330 (7)	0.0562 (11)	-0.0034 (8)	-0.0085 (9)	0.0020 (7)
C1	0.0582 (14)	0.0303 (9)	0.0478 (11)	-0.0028 (9)	0.0053 (10)	0.0011 (8)
C2	0.0525 (13)	0.0308 (9)	0.0464 (11)	-0.0040 (9)	0.0029 (9)	0.0022 (8)
C3	0.0526 (12)	0.0386 (9)	0.0431 (11)	-0.0025 (9)	0.0044 (10)	0.0049 (8)
C4	0.0758 (16)	0.0467 (11)	0.0627 (13)	-0.0193 (12)	-0.0046 (14)	-0.0087 (11)
C5	0.0587 (14)	0.0378 (9)	0.0625 (13)	-0.0042 (10)	-0.0083 (12)	0.0024 (9)
C6	0.0704 (15)	0.0566 (12)	0.0715 (15)	-0.0128 (12)	-0.0217 (12)	0.0201 (11)
C7	0.0496 (13)	0.0480 (11)	0.0525 (13)	-0.0036 (10)	-0.0095 (10)	0.0048 (10)
C8	0.0681 (16)	0.0550 (12)	0.0705 (15)	0.0111 (11)	0.0186 (13)	0.0048 (12)
C9	0.094 (2)	0.0577 (13)	0.0710 (17)	-0.0028 (15)	0.0181 (15)	0.0168 (12)
C10	0.0819 (19)	0.0491 (12)	0.0814 (18)	0.0038 (13)	-0.0133 (16)	0.0103 (13)
C11	0.0651 (17)	0.0740 (15)	0.0948 (19)	0.0241 (15)	0.0010 (17)	-0.0048 (14)
C12	0.0568 (15)	0.0833 (15)	0.0668 (15)	-0.0018 (15)	0.0091 (13)	0.0148 (12)

Geometric parameters (\AA , $^\circ$)

S1—C3	1.7495 (17)	C6—C7	1.506 (3)
S1—C6	1.808 (2)	C6—H6A	0.9700
N1—C3	1.338 (2)	C6—H6B	0.9700
N1—C4	1.350 (3)	C7—C8	1.366 (3)
N2—C4	1.322 (3)	C7—C12	1.374 (3)
N2—C1	1.334 (2)	C8—C9	1.376 (3)
N3—C5	1.358 (2)	C8—H8	0.9300
N3—C1	1.362 (3)	C9—C10	1.352 (4)
N3—H3N	0.894 (19)	C9—H9	0.9300
N4—C5	1.306 (2)	C10—C11	1.357 (4)
N4—C2	1.380 (2)	C10—H10	0.9300
C1—C2	1.392 (2)	C11—C12	1.383 (3)
C2—C3	1.385 (3)	C11—H11	0.9300
C4—H4	0.9300	C12—H12	0.9300
C5—H5	0.9300		
C3—S1—C6	101.51 (10)	C7—C6—H6A	109.8
C3—N1—C4	116.80 (18)	S1—C6—H6A	109.8
C4—N2—C1	111.39 (16)	C7—C6—H6B	109.8
C5—N3—C1	106.18 (16)	S1—C6—H6B	109.8
C5—N3—H3N	128.6 (13)	H6A—C6—H6B	108.2
C1—N3—H3N	124.8 (14)	C8—C7—C12	117.73 (19)
C5—N4—C2	104.28 (15)	C8—C7—C6	121.1 (2)
N2—C1—N3	128.28 (17)	C12—C7—C6	121.2 (2)
N2—C1—C2	125.9 (2)	C7—C8—C9	121.6 (2)
N3—C1—C2	105.86 (16)	C7—C8—H8	119.2
N4—C2—C3	133.49 (16)	C9—C8—H8	119.2
N4—C2—C1	109.82 (18)	C10—C9—C8	119.9 (2)
C3—C2—C1	116.67 (18)	C10—C9—H9	120.1
N1—C3—C2	119.75 (16)	C8—C9—H9	120.1
N1—C3—S1	121.40 (16)	C9—C10—C11	120.0 (2)

C2—C3—S1	118.85 (14)	C9—C10—H10	120.0
N2—C4—N1	129.52 (19)	C11—C10—H10	120.0
N2—C4—H4	115.2	C10—C11—C12	120.1 (2)
N1—C4—H4	115.2	C10—C11—H11	120.0
N4—C5—N3	113.9 (2)	C12—C11—H11	120.0
N4—C5—H5	123.1	C7—C12—C11	120.7 (2)
N3—C5—H5	123.1	C7—C12—H12	119.7
C7—C6—S1	109.51 (15)	C11—C12—H12	119.7
C4—N2—C1—N3	−179.25 (19)	C6—S1—C3—C2	173.43 (16)
C4—N2—C1—C2	−0.2 (3)	C1—N2—C4—N1	1.0 (3)
C5—N3—C1—N2	179.4 (2)	C3—N1—C4—N2	−0.8 (3)
C5—N3—C1—C2	0.2 (2)	C2—N4—C5—N3	0.4 (2)
C5—N4—C2—C3	−178.5 (2)	C1—N3—C5—N4	−0.4 (2)
C5—N4—C2—C1	−0.2 (2)	C3—S1—C6—C7	−165.98 (17)
N2—C1—C2—N4	−179.20 (19)	S1—C6—C7—C8	77.9 (2)
N3—C1—C2—N4	0.0 (2)	S1—C6—C7—C12	−103.9 (2)
N2—C1—C2—C3	−0.6 (3)	C12—C7—C8—C9	−2.2 (4)
N3—C1—C2—C3	178.62 (17)	C6—C7—C8—C9	176.1 (2)
C4—N1—C3—C2	−0.2 (3)	C7—C8—C9—C10	1.6 (4)
C4—N1—C3—S1	179.49 (15)	C8—C9—C10—C11	0.3 (4)
N4—C2—C3—N1	179.0 (2)	C9—C10—C11—C12	−1.4 (4)
C1—C2—C3—N1	0.8 (3)	C8—C7—C12—C11	1.0 (4)
N4—C2—C3—S1	−0.7 (3)	C6—C7—C12—C11	−177.3 (2)
C1—C2—C3—S1	−178.90 (15)	C10—C11—C12—C7	0.8 (4)
C6—S1—C3—N1	−6.23 (19)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3N···N4 ⁱ	0.894 (19)	1.892 (19)	2.773 (2)	167.9 (18)

Symmetry codes: (i) $-x+2, y-1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

