

Signatures of Mucosal Microbiome in Oral Squamous Cell Carcinoma Identified Using a Random Forest Model

This article was published in the following Dove Press journal:
Cancer Management and Research

Jianhua Zhou^{1,*}
Lili Wang^{2,*}
Rongtao Yuan¹
Xinjuan Yu²
Zhenggang Chen¹
Fang Yang¹
Guirong Sun³
Quanjiang Dong²

¹Department of Stomatology, Qingdao Municipal Hospital, Qingdao University, Qingdao 266071, Shandong, People's Republic of China; ²Central Laboratories and Department of Gastroenterology, Qingdao Municipal Hospital, Qingdao University, Qingdao 266071, Shandong, People's Republic of China; ³Clinical Laboratory, The Affiliated Hospital, Qingdao University, Qingdao 266011, Shandong, People's Republic of China

*These authors contributed equally to this work

Objective: The aim of this study was to explore the signatures of oral microbiome associated with OSCC using a random forest (RF) model.

Patients and Methods: A total of 24 patients with OSCC were enrolled in the study. The oral microbiome was assessed in cancerous lesions and matched paracancerous tissues from each patient using 16S rRNA gene sequencing. Signatures of mucosal microbiome in OSCC were identified using a RF model.

Results: Significant differences were found between OSCC lesions and matched paracancerous tissues with respect to the microbial profile and composition. Linear discriminant analysis effect size analyses (LEfSe) identified 15 bacteria genera associated with cancerous lesions. *Fusobacterium*, *Treponema*, *Streptococcus*, *Peptostreptococcus*, *Carnobacterium*, *Tannerella*, *Parvimonas* and *Filifactor* were enriched. A classifier based on RF model identified a microbial signature comprising 12 bacteria, which was capable of distinguishing cancerous lesions and paracancerous tissues (AUC = 0.82). The network of the oral microbiome in cancerous lesions appeared to be simplified and fragmented. Functional analyses of oral microbiome showed altered functions in amino acid metabolism and increased capacity of glucose utilization in OSCC.

Conclusion: The identified microbial signatures may potentially be used as a biomarker for predicting OSCC or for clinical assessment of oral cancer risk.

Keywords: oral squamous cell carcinoma, microbiome, random forest machine learning, predicted functions

Introduction

Oral cancer is a fatal malignancy and the sixth most common cancer worldwide. Oral squamous cell carcinoma (OSCC) is the most frequently occurring oral cancer, accounting for over 90% of all oral cancers.¹ In recent years, the incidence of OSCC has shown an increasing trend, which constitutes a major global health problem. OSCC is associated with a high rate of relapse, unpredictable metastasis and oral maxillofacial damage.^{2,3} The 5-year survival rate of patients with OSCC is approximately 50–60%.³

Mounting evidence supports the association between oral microbiome and OSCC. It has been suggested that the oral microbiome plays a role in the development of oral cancer.^{4–6} There are three potential mechanisms by which the oral microbiome is involved in the pathogenesis.⁶ Firstly, oral bacteria can induce chronic inflammation. Inflammatory mediators produced from the oral mucosa

Correspondence: Quanjiang Dong
Central Laboratories and Department of
Gastroenterology, Qingdao Municipal
Hospital, Qingdao University, Qingdao
266071, Shandong, People's Republic of
China
Tel +86-532-88905289
Fax +86-532-85968434
Email jiangacer@126.com

Guirong Sun
Clinical Laboratory, The Affiliated Hospital,
Qingdao University, Qingdao 266011,
Shandong, People's Republic of China
Email sungr@qduhospital.cn

promote cell hyper-proliferation, mutagenesis, oncogene activation and angiogenesis, leading to carcinogenesis. Secondly, certain oral bacteria are directly involved in the development of OSCC through enhancing cell proliferation, causing cytoskeletal rearrangement and activation of NF- κ B pathways, and inhibiting cellular apoptosis. Finally, some metabolites from oral bacteria were shown to possess carcinogenic potentials.⁶

A number of bacteria from the oral microbiome have been shown to be associated with OSCC. The relative abundance of *Fusobacterium* sp., *Porphyromonas gingivalis*, *Streptococcus* sp., *Peptostreptococcus* sp., and *Prevotella* sp. are increased in OSCC.^{7,8} Zhao et al reported significant enrichment of *Fusobacterium*, *Dialister*, *Peptostreptococcus*, *Filifactor*, *Peptococcus*, *Catonella*, and *Parvimonas* in OSCC patients compared with matched controls using linear discriminant analysis (LDA) effect size (LEfSe) analysis.⁹ In a prospective study, involving examination of mouth swab microbiome, *Parvimonas micra* and *Neisseria sicca* were associated with reduced risk of oral cavity cancer.¹⁰ The identified OSCC associated oral bacteria, however, vary among studies. This is partially attributable to the variations between studies with respect to the sample types used for the analysis of oral microbiome. There are substantial variations in the composition of the oral microbiome between various sample types including tissue, saliva, and mouth swab.¹¹ Few studies have characterized the OSCC related oral microbiome from tissue, which is closely associated with oral mucosa.

Features of the oral microbiome associated with OSCC have been explored by comparing OSCC patients with healthy controls, or by comparing tumor sites with the surrounding normal tissue.^{9,10} It has been alerted microbial diversity and a distinctive profile of the oral microbiome in OSCC.⁸⁻¹⁰ Many bacteria of the oral microbiome are enriched or depleted in OSCC.^{8,9} Recent studies have also revealed that the collective behaviors of the microbial community are more closely related to the development of cancer in comparison with individual bacteria genera.¹² To further explore how the oral microbiome is involved in the carcinogenic process, it is of importance to identifying signatures discriminative for the OSCC associated oral microbiome. The random forest (RF) method is a machine learning program capable of identifying an optimal set of variables with high discriminative power from a large number of dependent or independent variables. It is known for its ability to render complex dependency patterns between the outcome and the covariates.¹³ The method has been widely used to identify microbial signature for colorectal carcinoma, gastric cancer, and pancreatic cancer.¹⁴⁻¹⁶ In this

study, we applied the RF model to explore the signatures of the OSCC associated oral microbiome and validated these in an independent cohort.

Patients and Methods

Patients and Sample Collection

A total of 24 patients at different stages of OSCC were enrolled in the study (Table 1). The OSCC stages were classified according to the American Joint Committee on Cancer (AJCC) staging manual (8th edition).¹⁷ Of these patients, 19 were male. The average age of patients was 61.1 ± 12.4 years. None of the enrolled subjects had history of betel chewing, or history of diabetes mellitus or other severe complications including heart, liver, or renal failures. These patients had not taken antibiotics for at least two weeks immediately prior to sampling. Tissues from the site of tumor and paracancerous tissue from 2 cm outside the edge of the tumor were sampled. The paracancerous tissues are histologically normal. All samples were stored at -80°C until further processing. This study was approved by the Research Ethics Committee of Qingdao Municipal Hospital, China. Prior to sample collection, written informed consent for the study was obtained from all participants. All experiments were performed in accordance with approved guidelines and regulations of the institution.

Analyses of the Oral Microbiome

To analyze the microbial communities of the oral mucosa, genomic DNA was extracted from the oral mucosa samples as previously reported.¹⁸ The variable V3-V4 region of the 16S rRNA gene was PCR amplified with primers 341F/806R to generate the amplicon libraries. Sequencing was performed on an Illumina PE250 platform (Illumina, Hayward, CA, USA). A total of 16,333,787 paired-ends reads were obtained. After quality control and filtration, 2,163,267 reads were produced with an average of 45,068 reads per sample. The reads were analyzed using UPARSE.¹⁹ Following global trimming at 250 nucleotides, reads were dereplicated, and singletons were discarded. Subsequently, reads were clustered into operational taxonomic units (OTUs) assuming 97% identity. Chimeric reads were then removed. Taxonomy assignment was performed using UClust.²⁰ Analyses of alpha and beta microbial diversity were conducted as described previously.¹⁸ Comparisons of the relative abundance of taxa between groups were performed using version 1.0 of LEfSe.²¹ A LDA value greater than 3 with a p value less than 0.05 was considered to be statistically significant. To analyze the correlation network, Spearman

Table I Clinical Characteristics of 24 Patients with Oral Squamous Cell Carcinoma

Patients	Sex	Age	Location of Tumor	cTNM	Clinical Stage	Smoking Status	Alcohol Consumption
1	M	53	Mouth-floor	T2N0M0	II	NS	N
2	F	80	Cheek	T2N1M0	III	NS	N
3	F	47	Tongue	T2N2M0	IV	NS	N
4	M	83	Tongue	T2N0M0	II	FS	N
5	M	60	Oropharynx	T2N0M0	II	CS	D
6	M	54	Gingiva	T2N0M0	II	FS	D
7	F	62	Tongue	T1N0M0	I	NS	N
8	M	66	Cheek	T2N0M0	II	FS	D
9	M	59	Tongue	T3N0M0	III	CS	D
10	M	68	Gingiva	T2N0M0	II	CS	N
11	M	68	Tongue	T2N0M0	II	NS	N
12	M	75	Tongue	T2N1M0	III	NS	N
13	M	42	Tongue	T2N1M0	III	FS	N
14	M	63	Tongue	T2N1M0	III	NS	D
15	M	59	Cheek	T2N0M0	II	FS	D
16	F	74	Tongue	T2N0M0	II	NS	N
17	F	54	Gingiva	T2N0M0	II	NS	N
18	M	81	Cheek	T2N0M0	II	FS	D
19	F	63	Cheek	T1N1M0	III	NS	N
20	F	56	Tongue	T1N0M0	I	NS	N
21	M	42	Tongue	T1N0M0	I	CS	N
22	M	74	Cheek	T1N0M0	I	NS	D
25	M	47	Tongue	T2N1M0	III	CS	N
26	M	58	Cheek	T2N0M0	II	FS	D

Abbreviations: M, male; F, female; NS, non-smoker; CS, current smoker; FS, former smoker; D, alcohol drinker; N, non-alcohol drinker.

correlation coefficients were computed between the genera. Correlations that had an absolute Spearman coefficient value greater than or equal to 0.6 with a *p* value less than 0.05 were transformed into links between two genera in the genus network. Cytoscape v3.7.1 was then used to construct network figures. PICRUSt (v1.1.1) was used to predict the functions of the microbial community according to the reference data set of GreenGenes 16S rRNA database (v13.5).²² The accuracy of the predicted functions was assessed by the nearest sequenced taxon index (NSTI). The predicted functions were categorized with KEGG orthology. STAMP (v2.1.3) was used to compare functional differences between cancerous and paracancerous groups.²³

Statistical Analyses

To identify microbial signatures capable of distinguishing cancerous lesions from paracancerous lesions, a RF model was built using the AUC-RF algorithm.²⁴ The input variables comprised the relative abundance of taxa. A taxon was included only if it was present in more than 20% samples and had a relative abundance of greater than 0.05%. A 20-times repeated 10-fold cross-validation of the RF model was

performed. The performance of the RF model was demonstrated by receiver operating characteristic (ROC) curve analysis.²⁵ Validation of OSCC associated bacteria was explored in an independent cohort in Shanghai (China) with 80 samples, of these, 40 samples were from OSCC surface lesion and 40 samples were from anatomically matched normal sites (BioProject accession number: PRJNA362794).⁹

The Mann–Whitney *U*-test was performed to detect significant differences in alpha diversity or relative abundance between disease groups. Significant *P* values were corrected for multiple testing using the Benjamini-Hochberg procedure, with a cut-off false discovery rate (FDR) – adjusted *P* value lower than 0.1.²⁶ Constrained PCoA is a flexible method which allows for constrained ordination on the basis of any distance or dissimilarity measure.²⁷ It was performed in R to explore the influence of clinical stage on the microbiome of OSCC.

Data Availability

The next-generation sequencing datasets have been submitted to NCBI (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA597251/>). The BioProject accession number is PRJNA597251.

Results

Differences in Microbial Community Between Cancerous and Paracancerous Tissues

The microbial community between the cancerous lesions (C) and matched paracancerous tissues (P) was compared in 24 patients. Both Chao1 and Shannon indices which estimate the biodiversity of microbiome, were not significantly different between group C and group P ($P > 0.05$ for both) (Figure 1A and B). On multiple linear regression analyses, the alpha diversity indices of the microbiome showed no association with age, sex, smoking, alcohol consumption, degree of differentiation, or clinical stage in either group C or group P. Principal coordination analyses (PCoA) showed that the microbial community structure of group C was distinct from that of group P ($P = 0.048$) (Figure 1C). Constrained PCoA analyses were performed to explore the influence of clinical stage on the OSCC microbiome. The results demonstrated an apparent separation between the clinical stage I, II, and III in group C, especially between stages II and III. This suggested the progression of OSCC was associated with alteration of oral mucosa microbiome (Figure 1D). In group P, however, the microbial community structure was not discernible in the plots between stages (Figure 1E).

Altered Composition of the Microbiome in OSCC

Compositional analyses showed the phyla of *Proteobacteria*, *Firmicutes*, *Bacteroidetes* and *Fusobacteria* were dominant in the microbiome of both groups (Figure 1F). The relative abundance of *Firmicutes*, *Bacteroidetes*, *Fusobacteria*, and *Spirochaetes* was increased in cancerous lesions. Differences with respect to the relative abundance of *Fusobacteria* (9.80% vs 1.70%) and *Spirochaetes* (3.63% vs 0.18%) were statistically significant using Mann–Whitney *U*-test ($P = 0.043$ and 0.032 , FDR-adjusted $P = 0.089$ and 0.083 , respectively). The abundance of *Proteobacteria*, *Actinobacteria*, and *Cyanobacteria* in cancerous lesions was lower compared to paracancerous tissues. The differences of relative abundance of *Actinobacteria* (0.77% vs 1.78%) and *Cyanobacteria* (0.23% vs 1.33%) were statistically significant ($P = 0.021$ and $P = 0.013$, FDR-adjusted $P = 0.067$ and 0.051 , respectively). At the genus level, LefSe analyses identified 15 genera which had a LDA score >3.0 (Figure 2A). The relative abundance of *Fusobacterium*, *Treponema*, *Streptococcus*, *Peptostreptococcus*, *Carno*

bacterium, *Tannerella*, *Parvimonas*, and *Filifactor* were increased in cancerous lesions (Figure 2B). Of these, *Fusobacterium* had the highest LDA score. In contrast, the relative abundance of *Streptophyta*, *Brevundimonas*, *Paenibacillus*, *Microbacterium*, *Desulfovibrio*, *Mucispirillum*, and *Arthrobacter* were decreased in cancerous lesions (Figure 2C).

Microbial Signatures Associated with OSCC

To determine microbial signatures capable of discriminating cancerous lesions from paracancerous tissues, an RF classifying model was built with the AUC-RF algorithm. The results showed a minimal set of 12 bacteria genera that maximally differentiated cancer sites from paracancerous sites. It contained *Microbacterium*, *Streptococcus*, *Brevundimonas*, *Peptostreptococcus*, *Streptophyta*, *Paenibacillus*, *Actinomyces*, *Parvimonas*, *Mucispirillum*, *Arthrobacter*, *Rhizobium* and *Fusobacterium* (Figure 3A and B). Most of these (10/12) have been identified to have a LDA value > 3.0 by the LefSe analyses, including four genera (*Fusobacterium*, *Parvimonas*, *Peptostreptococcus* and *Streptococcus*) enriched in cancerous lesions and six depleted genera (*Arthrobacter*, *Brevundimonas*, *Microbacterium*, *Mucispirillum*, *Paenibacillus*, and *Streptophyta*). RF models trained with this optimal set of features resulted in an out-of-bag error rate of 33.33%. To assess the model classifying accuracy, a 20-times repeated 10-fold cross-validation was performed. The AUC value from cross-validations was 0.82 [95% confidence interval (CI): 0.70–0.94] (Figure 3C). This set of 12 genera was further validated in an independent Chinese Shanghai cohort with AUC of 0.69 (95% CI: 0.54–0.80) (Figure 3D). The results suggested that these oral mucosal bacteria associated with OSCC may predict the risk of OSCC.

Co-Occurrence Network in OSCC

The co-occurrence network differed substantially between cancerous lesions and paracancerous tissues (Figure 4). The network was simplified in group C with a decreased number of nodes (Table 2). The network density and average degree were also lower in group C, which demonstrated a reduction in the network complexity. The clustering coefficient of the network was lower, leading to an increased number of components and more fragmented sub-networks in group C (Figure 4).




Figure 1 Profiles of the oral microbiome in OSCC. The alpha diversity was estimated using Chao1 (**A**) and Shannon (**B**) indices. PCoA plots for comparing community structure between paraneoplastic and cancerous tissues (**C**), between clinical stages in cancerous tissues (**D**) and paraneoplastic tissues (**E**). The composition of microbiome at the phylum level was compared between paraneoplastic and cancerous tissues (**F**). (P) paraneoplastic tissues; (C) cancerous tissues.

Predicted Functions of the OSCC Microbiome

Functional capacities of the OSCC microbiome were predicted using PICRUSt. Compared with paraneoplastic tissues, the average relative frequencies of pathways related to metabolism, especially amino acid metabolism, were decreased at KEGG levels 1 and 2 (**Figure 5A** and **B**).

Furthermore, reduced relative frequencies in OSCC were found for pathways of KEGG level 3 related to amino acid metabolism including phenylalanine metabolism, tryptophan metabolism, arginine and proline metabolism, valine, leucine and isoleucine degradation, lysine degradation and tyrosine metabolism (**Figure 5C**). Nevertheless, the relative frequencies of methane metabolism, glucose-related




Figure 2 LEfSe analyses of microbiome composition between paracancerous and cancerous tissues. Bacteria genera enriched in cancerous tissues had a positive LDA score, while those depleted had a negative score. Bacteria with LDA scores >3 are shown (A). Relative abundance of genera enriched in cancerous tissues (B). Relative abundance of genera enriched in paracancerous tissues (C). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. All FDR-adjusted P values were <0.05 .

metabolism, such as phosphotransferase system (PTS) and glycolysis, were significantly enriched in cancer lesions (Figure 5D). These alterations suggested functional disorders of amino acid metabolism and increased capacity of glucose utilization in the microbiome of OSCC. To validate these results, the predicted functions were compared between two randomized groups. The results showed no significant differences between two groups at KEGG levels 1–3 (Supplementary Table 1).

Discussion

Our results demonstrated significant differences in the β -diversity (including PCoA and constrained PCoA analysis) of the oral microbiome between cancerous and paracancerous tissues. These findings indicate altered community structure of

the oral microbiome in cancerous lesions. However, the α -diversity including Chao1 and Shannon indices showed no significant differences. Consistent with our results, previous studies have found no significant differences in α -diversity between cancerous lesions and paratumoral mucosa in patients with colorectal cancer.²⁸ Constrained PCoA analyses demonstrated an apparent separation among the different clinical stages in cancerous lesions of OSCC. It should be noted that the sample size in different stages was small. Thus, the observed association between variations of the oral microbiome and cancer progression needs to be demonstrated in studies with a large sample size. Nonetheless, in a previous study, the oral microbiome in patients with stage IV OSCC was different from that in healthy controls.²⁹ The microbiome communities showed a progressive shift with the progression




Figure 3 Identification of microbial signature associated with OSCC. The random forest model was constructed using AUC-RF algorithm based on bacteria which were present in more than 20% samples and had a relative abundance of over 0.05%. The model containing 12 genera was selected as the optimal model based on the highest OOB-AUC value obtained from the backward elimination process performed using the AUC-RF algorithm with the median decrease Gini (MDG) importance measure (**A**). MDG of selected genera in the optimal set (**B**). The ROC curve of the optimal model for classifying cancerous tissues from paracancerous tissues (**C**). The ROC curve of validation in a independent cohort in Shanghai (**D**).

of OSCC stage.²⁹ These findings are consistent with our results wherein the community structure of the oral microbiome was found to change with the stages of OSCC, indicating the potential involvement of the oral microbiome in cancer progression.

In recent studies, microbial signatures identified using RF machine learning algorithm were used to predict different diseases.¹⁴⁻¹⁶ The identified signatures can be viewed as a minimized representation of the microbial community for exploring mechanisms of microbial dysbiosis and targets for manipulating the dysbiotic microbiome. In a study by Zackular et al, the final numbers of colon tumor were quantitatively predicted on the basis of the initial composition of the gut microbiome using RF algorithm.³⁰ In another report, a minimal set of 22 viral genera were identified that

differentiated CRC patients from controls and predicted CRC outcomes.¹⁴ An RF model for predicting exacerbation of radiation-induced mucositis was generated using oral microbiome, it showed a high predictive accuracy (AUC = 0.89).³¹ Currently, few studies have been conducted to identify the microbial signature in OSCC. Our study revealed an RF model consisting of 12 relevant bacteria with differential abundance between the cancerous lesions and paracancerous tissues of OSCC. The trained model was capable of predicting OSCC with considerably high accuracy (AUC = 0.82). Moreover, the validation in an independent cohort showed a moderate accuracy. Monitoring the composition and abundance of these 12 bacteria may help predict the risk of OSCC, facilitate early diagnosis and treatment of OSCC and help improve the prognosis of OSCC.




Figure 4 Correlation network in paracancerous tissues (A) and cancerous tissues (B). The correlation coefficient was calculated with Spearman's rank correlation test ($|r| \geq 0.6$). Cytoscape was used for network construction.

In our study, the LDA scores of 15 genera were >3.0 when comparing the oral microbiome of cancerous lesions with that of paracancerous tissues. Furthermore, these bacteria showed differential abundance between the two groups, suggesting a close association between these bacteria and OSCC. The majority (10/12) of bacteria in the optimal set identified by RF analyses were contained in these 15 bacteria genera. The highly consistent results between these analyses demonstrate that variations of the optimal set of bacteria reflect the dysbiotic features of the oral microbiome associated with OSCC, accounting for remarkable capacity of the set to discriminating OSCC from normal tissues. Of the optimal set of bacteria, *Fusobacterium*, *Parvimonas*, *Peptostreptococcus* and *Streptococcus* were enriched in cancerous lesions. The association of these four bacteria with OSCC has been

Table 2 Features of Co-Occurrence Network of the Oral Microbiome

Parameters	P (n = 24)	C (n = 24)	P values
Clustering coefficient	0.694±0.273	0.474±0.345	3.55E-04
Connected components	2	2	–
Average degree	13.970±7.606	7.935±6.222	6.83E-06
Number of nodes	66	62	–
Average Shortest Path Length	1.764±0.486	3.255±0.779	1.34E-17
Degree Centrality	0.215±0.117	0.130±0.102	9.46E-05
Closeness Centrality	0.601±0.131	0.334±0.135	1.34E-17
Betweenness Centrality	0.024±0.030	0.039±0.060	0.785

Notes: –, not applicable; P, paracancerous tissues; C, cancerous tissues.

reported. It has been found an increase in the abundance of *Fusobacteria* in OSCC, especially in the late stage.^{11,32} A significant enrichment of *Fusobacterium*, *Peptostreptococcus*, *Peptococcus* and *Parvimonas* in OSCC surface lesions has been demonstrated.^{9,33} Variations in the abundance of *Peptostreptococcus* and *Parvimonas* in the salivary microbiome have been observed between normal individuals, oral precursor lesion and cancer patients.³⁴ Infection with *S. anginosus* occurs frequently in OSCC.³⁵ Species of these genera have been isolated from the oral tumor and metastatic lymphatic nodes.^{36,37} These results support a close association between these enriched bacteria and OSCC. In addition, these enriched bacteria may have carcinogenic potential. Previous studies have shown that *F. nucleatum* may induce tumorigenesis through the inflammatory NF- κ B signaling pathway.^{38,39} *F. nucleatum* expresses the adhesion molecule FadA which binds to E-cadherin on epithelial cells, inhibiting its tumor-suppressor activity.⁴⁰ *Streptococcus spp.* possesses alcohol dehydrogenase activity leading to production of the carcinogenic acetaldehyde, which is potentially involved in the occurrence of oral cancer.⁴¹ These findings suggest that our RF analyses identified a group of bacteria with carcinogenic potentials enriched in OSCC. In addition to these bacteria, we also found depletion of six bacteria genera in OSCC. Further studies are required to unravel the causes of variations in the abundance of these bacteria and their roles in carcinogenesis.

Bacterial interaction is a determinant of the homeostasis of the microbiome. Studies have shown reduced network complexity in peritumoral or tumoral tissues of the stomach in comparison with normal gastric mucosa.⁴² In our study, strongly interacted genera were picked up according to $|r| \geq 0.6$ to construct microbial




Figure 5 Differential functions predicted using PICRUSt between paracancerous and cancerous tissues. The function of metabolism at level 1 in KEGG pathway (**A**) and amino acid metabolism at level 2 (**B**). The decreased (**C**) or increased (**D**) relative frequency of functions of amino acid metabolism at level 3 in cancerous tissues. Differences between groups in the predicted functions were compared using STAMP. Statistical differences are considered when $P < 0.05$.

network. Our results demonstrated a simplified microbial network in cancerous lesions of OSCC, with the reduction in the number of nodes, network density, average degree, and clustering coefficient in cancer sites, suggesting fragmented and simplified network. These findings indicated the loss of certain bacterial interactions in OSCC, leading to disruption of the homeostasis of oral microbiome. However, a recent study found increased network complexity in OSCC.⁹ The contradictory results may be attributed to the difference in the cut-off value of correlation coefficient used bacterial species for the construction of microbial network.

The functional profile in the cancer-associated microbiome shows substantial changes. In a previous study, the functions

related to protein and amino acid metabolism (such as valine, leucine and isoleucine, phenylalanine, tyrosine and tryptophan biosynthesis) showed an inverse association with OSCC progression.²⁹ Similar results were obtained in a study of saliva metabolomics wherein phenylalanine metabolism was found down-regulated in OSCC.⁴³ Consistent with this, in our study, the functional prediction of oral bacterial communities showed a significant decrease in genes involved in amino acid metabolism (such as phenylalanine metabolism, tryptophan metabolism, arginine, and proline metabolism) in OSCC lesions. However, methane metabolism and glucose-related metabolism (such as PTS and glycolysis) were significantly enriched in cancer lesions. Tryptophan metabolism is associated with the evasion of immune surveillance by cancer cells.⁴⁴

Decreased levels of tryptophan were shown to promote the generation of immune-suppressive T cells promoting cancer development.⁴⁵ Cancer cells appear to utilize glycolytic metabolism for energy production and precursors for the increased rate of anabolic processes.⁴⁶ These suggest the gene functional disorders of amino acid metabolism and increased capacity of glucose utilization in the microbiome are associated with OSCC.

In summary, this study revealed dysbiosis in OSCC featured with functional and compositional changes. We identified a microbial signature comprising 12 bacteria associated with OSCC, which may be used as a biomarker for clinical prediction of the risk of OSCC. Further studies are required to explore the potential clinical application of the microbial signature.

Funding

This work was supported by grants from the National Natural Science Foundation of China (31870777, 81602144).

Disclosure

The authors report no conflicts of interest in this work.

References

- Kademani D. Oral cancer. *Mayo Clin Proc.* 2007;82(7):878–887. doi:10.4065/82.7.878
- Hooper SJ, Crean SJ, Fardy MJ, et al. A molecular analysis of the bacteria present within oral squamous cell carcinoma. *J Med Microbiol.* 2007;56(Pt12):1651–1659. doi:10.1099/jmm.0.46918-0
- Crozier E, Sumer BD. Head and neck cancer. *Med Clin North Am.* 2010;94(5):1031–1046. doi:10.1016/j.mena.2010.05.014
- Yost S, Stashenko P, Choi Y, et al. Increased virulence of the oral microbiome in oral squamous cell carcinoma revealed by metatranscriptome analyses. *Int J Oral Sci.* 2018;10(4):32. doi:10.1038/s41368-018-0037-7
- Karpiński TM. Role of oral microbiota in cancer development. *Microorganisms.* 2019;7(1):20. doi:10.3390/microorganisms7010020
- Zhang Y, Wang X, Li H, Ni C, Du Z, Yan F. Human oral microbiota and its modulation for oral health. *Biomed Pharmacother.* 2018;99:883–893. doi:10.1016/j.biopha.2018.01.146
- Pushalkar S, Ji X, Li Y, et al. Comparison of oral microbiota in tumor and non-tumor tissues of patients with oral squamous cell carcinoma. *BMC Microbiol.* 2012;12:144. doi:10.1186/1471-2180-12-144
- Mager DL, Haffajee AD, Devlin PM, Norris CM, Posner MR, Goodson JM. The salivary microbiota as a diagnostic indicator of oral cancer: a descriptive, non-randomized study of cancer-free and oral squamous cell carcinoma subjects. *J Transl Med.* 2005;3:27. doi:10.1186/1479-5876-3-27
- Zhao H, Chu M, Huang Z, et al. Variations in oral microbiota associated with oral cancer. *Sci Rep.* 2017;7(1):11773. doi:10.1038/s41598-017-11779-9
- Hayes RB, Ahn J, Fan X, et al. Association of oral microbiome with risk for incident head and neck squamous cell cancer. *JAMA Oncol.* 2018;4(3):358–365. doi:10.1001/jamaoncol.2017.4777
- Zhang Z, Yang J, Feng Q, et al. Compositional and functional analysis of the microbiome in tissue and saliva of oral squamous cell carcinoma. *Front Microbiol.* 2019;10:1439. doi:10.3389/fmicb.2019.01439
- Darveau RP. Periodontitis: a polymicrobial disruption of host homeostasis. *Nat Rev Microbiol.* 2010;8(7):481–490. doi:10.1038/nrmicro2337
- Hornung R, Wright MN, Wright MN. Block forests: random forests for blocks of clinical and omics covariate data. *BMC Bioinformatics.* 2019;20(1):358. doi:10.1186/s12859-019-2942-y
- Nakatsu G, Zhou H, Wu WKK, et al. Alterations in enteric virome are associated with colorectal cancer and survival outcomes. *Gastroenterology.* 2018;155(2):529–541. doi:10.1053/j.gastro.2018.04.018
- Yu G, Hu N, Wang L, et al. Gastric microbiota features associated with cancer risk factors and clinical outcomes: a pilot study in gastric cardia cancer patients from Shanxi, China. *Int J Cancer.* 2017;141(1):45–51. doi:10.1002/ijc.30700
- Half E, Keren N, Reshef L, et al. Fecal microbiome signatures of pancreatic cancer patients. *Sci Rep.* 2019;9(1):16801. doi:10.1038/s41598-019-53041-4
- Amin MB, Edge SB, Greene FL, et al. *AJCC Cancer Staging Manual.* 8th ed. New York: Springer; 2017.
- Wang L, Zhou J, Xin Y, et al. Bacterial overgrowth and diversification of microbiota in gastric cancer. *Eur J Gastroenterol Hepatol.* 2016;28(3):261–266. doi:10.1097/MEG.0000000000000542
- Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods.* 2013;10(10):996–998. doi:10.1038/nmeth.2604
- Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics.* 2010;26(19):2460–2461. doi:10.1093/bioinformatics/btq461
- Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. *Genome Biol.* 2011;12(6):R60. doi:10.1186/gb-2011-12-6-r60
- Langille MG, Zaneveld J, Caporaso JG, et al. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat Biotechnol.* 2013;31(9):814–821. doi:10.1038/nbt.2676
- Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics.* 2014;30(21):3123–3124. doi:10.1093/bioinformatics/btu494
- Calle ML, Urrea V, Boulesteix AL, Malats N. AUC-RF: a new strategy for genomic profiling with random forest. *Hum Hered.* 2011;72(2):121–132. doi:10.1159/000330778
- Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology.* 1982;143(1):29–36. doi:10.1148/radiology.143.1.7063747
- Benjamini Y, Hochberg Y. Controlling the false discovery rate – a practical and powerful approach to multiple testing. *J R Statist Soc B.* 1995;57(1):289–300.
- Anderson MJ, Willis TJ. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology.* 2003;84(2):511–525. doi:10.1890/0012-9658(2003)084[0511:CAOPCA]2.0.CO;2
- Allali I, Delgado S, Marron PI, et al. Gut microbiome compositional and functional differences between tumor and non-tumor adjacent tissues from cohorts from the US and Spain. *Gut Microbes.* 2015;6(3):161–172. doi:10.1080/19490976.2015.1039223
- Yang CY, Yeh YM, Yu HY, et al. Oral microbiota community dynamics associated with oral squamous cell carcinoma staging. *Front Microbiol.* 2018;9:862. doi:10.3389/fmicb.2018.00862
- Zackular JP, Baxter NT, Chen GY, Schloss PD. Manipulation of the gut microbiota reveals role in colon tumorigenesis. *mSphere.* 2015;1(1):pii:e00001–15.
- Zhu XX, Yang XJ, Chao YL, et al. The potential effect of oral microbiota in the prediction of mucositis during radiotherapy for nasopharyngeal carcinoma. *EBioMedicine.* 2017;18:23–31. doi:10.1016/j.ebiom.2017.02.002
- Schmidt BL, Kuczynski J, Bhattacharya A, et al. Changes in abundance of oral microbiota associated with oral cancer. *PLoS One.* 2014;9(6):e98741. doi:10.1371/journal.pone.0098741

33. Li Y, He J, He Z, et al. Phylogenetic and functional gene structure shifts of the oral microbiomes in periodontitis patients. *ISME J*. 2014;8(9):1879–1891. doi:10.1038/ismej.2014.28
34. Lee WH, Chen HM, Yang SF, et al. Bacterial alterations in salivary microbiota and their association in oral cancer. *Sci Rep*. 2017;7(1):16540. doi:10.1038/s41598-017-16418-x
35. Sasaki M, Yamaura C, Ohara-Nemoto Y, et al. *Streptococcus anginosus* infection in oral cancer and its infection route. *Oral Dis*. 2005;11(3):151–156. doi:10.1111/j.1601-0825.2005.01051.x
36. Sakamoto H, Naito H, Ohta Y, et al. Isolation of bacteria from cervical lymph nodes in patients with oral cancer. *Arch Oral Biol*. 1999;44(10):789–793. doi:10.1016/S0003-9969(99)00079-5
37. Sakamoto H, Sasaki J, Nord CE. Association between bacterial colonization on the tumor, bacterial translocation to the cervical lymph nodes and subsequent postoperative infection in patients with oral cancer. *Clin Microbiol Infect*. 1999;5(10):612–616. doi:10.1111/j.1469-0691.1999.tb00417.x
38. Kostic AD, Chun E, Robertson L, et al. *Fusobacterium nucleatum* potentiates intestinal tumorigenesis and modulates the tumor-immune microenvironment. *Cell Host Microbe*. 2019;20(1):207–215. doi:10.1016/j.chom.2013.07.007
39. DiDonato JA, Mercurio F, Karin M. NF- κ B and the link between inflammation and cancer. *Immunol Rev*. 2012;246(1):379–400. doi:10.1111/j.1600-065X.2012.01099.x
40. Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW. *Fusobacterium nucleatum* promotes colorectal carcinogenesis by modulating E-cadherin/ β -catenin signaling via its FadA adhesin. *Cell Host Microbe*. 2013;14(2):195–206. doi:10.1016/j.chom.2013.07.012
41. Meurman JH, Uittamo J. Oral micro-organisms in the etiology of cancer. *Acta Odontol Scand*. 2008;66(6):321–326. doi:10.1080/00016350802446527
42. Liu X, Shao L, Liu X, et al. Alterations of gastric mucosal microbiota across different stomach microhabitats in a cohort of 276 patients with gastric cancer. *EBioMedicine*. 2019;40:336–348. doi:10.1016/j.ebiom.2018.12.034
43. Wang Q, Gao P, Wang X, Duan Y. The early diagnosis and monitoring of squamous cell carcinoma via saliva metabolomics. *Sci Rep*. 2014;4:6802. doi:10.1038/srep06802
44. Hashim NAA, Ab-Rahim S, Suddin LS, Saman MSA, Mazlan M. Global serum metabolomics profiling of colorectal cancer. *Mol Clin Oncol*. 2019;11(1):3–14. doi:10.3892/mco.2019.1853
45. Grohmann U, Bronte V. Control of immune response by amino acid metabolism. *Immunol Rev*. 2010;236:243–264. doi:10.1111/j.1600-065X.2010.00915.x
46. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*. 2009;324(5930):1029–1033. doi:10.1126/science.1160809

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>