

## RESEARCH ARTICLE

# The shrimp IKK–NF- $\kappa$ B signaling pathway regulates antimicrobial peptide expression and may be subverted by white spot syndrome virus to facilitate viral gene expression

Pei-Hui Wang<sup>1</sup>, Zhi-Hua Gu<sup>1</sup>, Ding-Hui Wan<sup>1</sup>, Bo-Du Liu<sup>1</sup>, Xian-De Huang<sup>1</sup>, Shao-Ping Weng<sup>1</sup>, Xiao-Qiang Yu<sup>2</sup> and Jian-Guo He<sup>1,3</sup>

The  $\kappa$ B kinases IKK $\alpha$  and IKK $\beta$  and the IKK-related kinases TANK-binding kinase 1 (TBK1) and IKK $\epsilon$  are the master regulators of the NF- $\kappa$ B signaling pathway. Although this pathway has been extensively studied in mammals, less attention has been paid in crustaceans, which have significant economic value. Here, we report the cloning and functional studies of two IKK homologs, LvIKK $\beta$  and LvIKK $\epsilon$ , from Pacific white shrimp, *Litopenaeus vannamei*. LvIKK $\beta$  and LvIKK $\epsilon$  mRNAs are widely expressed in different tissues and are responsive to white spot syndrome virus (WSSV) infection. When overexpressed in *Drosophila S2* cells, LvIKK $\beta$  but not LvIKK $\epsilon$  activates the promoters of NF- $\kappa$ B pathway-controlled antimicrobial peptide genes (AMPs), such as the *Penaeidins* (PENs). In HEK 293T cells, both LvIKK $\beta$  and LvIKK $\epsilon$  activate an NF- $\kappa$ B reporter. The silencing of LvIKK $\beta$  or LvIKK $\epsilon$  using double-stranded RNA (dsRNA)-mediated RNA interference (RNAi) decreases the expression of *L. vannamei* AMPs, including PENs, lysozyme and crustins. Intriguingly, LvIKK $\beta$ - or LvIKK $\epsilon$ -silenced *L. vannamei* are resistant to WSSV infection. We hypothesized that successful infection with WSSV requires the activation of the IKK–NF- $\kappa$ B signaling pathway to modulate viral gene expression. We constructed luciferase reporters for 147 WSSV genes. By screening, we found that the WSSV051, WSSV059, WSSV069, WSSV083, WSSV090, WSSV107, WSSV244, WSSV303, WSSV371 and WSSV445 promoters can be activated by LvIKK $\beta$  or LvIKK $\epsilon$  in *Drosophila S2* cells. Taken together, our results reveal that LvIKK $\beta$  and LvIKK $\epsilon$  may participate in the regulation of shrimp AMPs and that WSSV may subvert the *L. vannamei* IKK–NF- $\kappa$ B signaling pathway to facilitate viral gene expression. *Cellular & Molecular Immunology* (2013) 10, 423–436; doi:10.1038/cmi.2013.30; published online 19 August 2013

**Keywords:** antimicrobial peptide; IKK–NF- $\kappa$ B; innate immunity; *Litopenaeus vannamei*; WSSV

## INTRODUCTION

Innate immunity is activated when pathogen signature molecules are recognized by pattern-recognition receptors, such as Toll-like receptors (TLRs), retinoic acid-inducible gene-I-like receptors and nucleotide-binding oligomerization domain-like receptors.<sup>1,2</sup> These signature molecules, which are derived from viruses, pathogenic bacteria, pathogenic fungi and parasitic protozoa, are known as pathogen-associated molecular patterns and include lipopolysaccharide, lipoproteins, single-stranded RNA, double-stranded RNA (dsRNA) and unmethylated CpG-containing

DNA.<sup>2–4</sup> In mammals, the recognition of pathogen-associated molecular patterns by TLRs and nucleotide-binding oligomerization domain-like receptors (nucleotide-binding oligomerization domain 1 and nucleotide-binding oligomerization domain 2) activates the NF- $\kappa$ B pathway, which plays a central role in coordinating the expression of pro-inflammatory cytokines and chemokines to eliminate microbial infection by triggering inflammation and recruiting innate and acquired immune cells.<sup>3,4</sup> The activation of NF- $\kappa$ B is achieved by the signal-induced phosphorylation and subsequent degradative polyubiquitylation of the

<sup>1</sup>MOE Key Laboratory of Aquatic Product Safety/State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, Guangzhou, China; <sup>2</sup>Division of Cell Biology and Biophysics, School of Biological Sciences, University of Missouri-Kansas City, Kansas City, MO, USA and <sup>3</sup>School of Marine Sciences, Sun Yat-Sen University, Guangzhou, China

Correspondence: PH Wang, MOE Key Laboratory of Aquatic Product Safety/State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, 135 Xingang Road West, Guangzhou 510275, China.

E-mail: wph326@gmail.com

Or JG He, MOE Key Laboratory of Aquatic Product Safety/State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, 135 Xingang Road West, Guangzhou 510275, China.

E-mail: lsshjg@mail.sysu.edu.cn

Received: 14 March 2013; Revised: 30 May 2013; Accepted: 4 June 2013

cytoplasmic inhibitor protein inhibitor of NF- $\kappa$ B alpha (I $\kappa$ B $\alpha$ ). I $\kappa$ B kinase alpha (IKK $\alpha$ ) and IKK $\beta$  mediate the phosphorylation of I $\kappa$ B $\alpha$  and represent a point of convergence for most of the signal transduction pathways that lead to NF- $\kappa$ B activation.<sup>2-4</sup> The degradation of I $\kappa$ B $\alpha$  results in the nuclear translocation of activated NF- $\kappa$ B proteins. In mammals, the recognition of dsRNA by TLR3 and the retinoic acid-inducible gene-I-like receptors (retinoic acid-inducible gene-I and MDA5) also activates the IRF3/7 signaling pathways at the same time as the NF- $\kappa$ B signaling pathway.<sup>1,4,5</sup> The IRF3/7 signaling pathways, which are the key regulators of type I IFN expression, are directly activated in the cytoplasm *via* C-terminal phosphorylation by the IKK-related kinases TANK-binding kinase 1 (TBK1) and IKK $\epsilon$ .<sup>5,6</sup> The C-terminal phosphorylation of IRF3 and IRF7 by TBK1 and IKK $\epsilon$  promotes their homodimerization and subsequent nuclear translocation, resulting in type I IFN expression.<sup>5,6</sup> IRF7 can also be phosphorylated by IRAK1/IKK $\alpha$  in the TLR7- and TLR9-mediated signaling pathways, resulting in IFN $\alpha$  expression.<sup>2,7</sup>

In *Drosophila*, the Toll and IMD pathways are the best-characterized pattern-recognition receptor pathways that recognize pathogen-associated molecular patterns on the surface of invading bacteria or fungi.<sup>8-10</sup> Although no components of the *Drosophila* Toll and IMD pathways have been identified as virus sensors, both pathways can be activated by viruses, and their activation contributes to the restriction of viral replication.<sup>5,11,12</sup> Upon infection with Gram-negative and certain Gram-positive bacteria, the p100-like NF- $\kappa$ B precursor protein Relish, which is the central component of the IMD pathway, is phosphorylated by IKK $\beta$ , activated by proteolytic cleavage, and translocated into the nucleus, where it promotes the expression of immune-related genes, such as antimicrobial peptide genes (AMPs).<sup>8-10,13</sup> Following stimulation by fungi and many Gram-positive bacteria, DIF and Dorsal, which are two p65-like NF- $\kappa$ B proteins of the Toll pathway, are activated by the signal-induced degradation of the I $\kappa$ B-related inhibitor Cactus (*Drosophila* I $\kappa$ B $\alpha$ ) and are translocated into the nucleus, where they promote the expression of immune-related genes such as AMPs.<sup>8-10</sup> Like mammalian I $\kappa$ B $\alpha$ , Cactus is phosphorylated upon stimulation; however, its degradation does not require the IKK family proteins, although the IKK complex does act on I $\kappa$ B $\alpha$  in mammals.<sup>14</sup> Currently, it is unknown which kinases can act directly on Cactus. *Drosophila melanogaster* IKK $\beta$  functions only in the IMD pathway and not the Toll pathway. Instead of a Cactus (I $\kappa$ B $\alpha$ ) kinase, *D. melanogaster* IKK $\epsilon$  regulates F-actin assembly by mediating the function of non-apoptotic caspases *via* the degradation of DIAP1 but does not participate in NF- $\kappa$ B activation.<sup>15,16</sup> Whether this phenomenon occurs in other invertebrates and the mechanism of Cactus phosphorylation are still unclear.

Crustaceans include lobsters, crabs, crayfish and shrimp, some of which are of great economic importance, such as cultured penaeid shrimp. Because the production of cultured shrimp is increasing dramatically around the world and increased stress is imposed on aquatic habitats where crustaceans play major ecological roles, crustacean immunology has

attracted significant attention in recent years. The Toll and IMD pathways are the major regulators of the immune response in *Drosophila*.<sup>17</sup> Recently, we cloned several members of these two pathways in Pacific white shrimp, *Litopenaeus vannamei*, including LvToll1-3, LvIMD, the p100-like NF- $\kappa$ B precursor protein LvRelish and the NF- $\kappa$ B family protein LvDorsal.<sup>18-21</sup> However, whether the shrimp Toll and IMD pathways also play an essential role in the immune response is still elusive. The rapid and transient expression of AMPs such as the *Penaeidins* (PENs), lysozyme and *crustins* are reported to be pivotal for shrimp defense against microbial infections, and AMP levels are associated with a successful defense against microbial infections by shrimp immune responses.<sup>22</sup> In this study, we cloned two *L. vannamei* IKKs (LvIKK $\beta$  and LvIKK $\epsilon$ ) and investigated their roles in AMP regulation and viral infection using luciferase assays and dsRNA-mediated gene silencing *in vivo*. We found that successful infection with WSSV may rely on the activation of the IKK-NF- $\kappa$ B signaling pathway, which can significantly induce the promoters of WSSV immediate-early genes, such as WSSV051, WSSV069 (*iel*) and WSSV083.

## MATERIALS AND METHODS

### Microorganisms

Inocula containing the Gram-negative bacterium *Vibrio alginolyticus* and white spot syndrome virus (WSSV) were prepared as described previously.<sup>23,24</sup> Bacteria were quantified by counting microbial colony-forming units (CFUs) per milliliter on Luria broth agar plates following incubation at 30 °C overnight.

### Experimental shrimp

Pacific white shrimp, *L. vannamei* (approximately 8–10 g each for gene expression analysis; approximately 1–2 g each for dsRNA-mediated RNA interference experiments), were purchased from a local shrimp farm in Zhuhai, Guangdong Province, China. The shrimp were cultured in a recirculating water tank system filled with air-pumped seawater (2.5% salinity) at 24–26 °C and fed a commercial diet at 5% of their body weight twice per day. The shrimp were cultured for at least 7 days for acclimation before experiments.

### Cloning the cDNA and genomic DNA of LvIKK $\beta$ and LvIKK $\epsilon$

In tissue distribution studies of LvIKK $\beta$  and LvIKK $\epsilon$ , total RNA (0.5  $\mu$ g) was isolated from various tissues of healthy shrimp using the RNeasy Mini Kit (Qiagen, Hilden, Germany) and then reverse transcribed into cDNA using a PrimeScript First Strand cDNA Synthesis Kit (TaKaRa, Dalian, China). An LvIKK $\epsilon$  cDNA fragment was obtained by PCR amplification using degenerate primers (dpIKK-F and dpIKK-R; Table 1) and a cDNA template prepared from shrimp gills. Based on the cDNA fragment and an LvIKK $\beta$  EST (accession no. CK572131), the full-length cDNAs of LvIKK $\epsilon$  and LvIKK $\beta$  were obtained using a RACE-PCR approach as described previously.<sup>18,21</sup> All conditions for RACE-PCR were identical except for the primers (listed in Table 1). Genomic

**Table 1 PCR primers used in this study**

Primer	Primer sequence (5'–3')
<b>cDNA cloning</b>	
DPLvIKKε-F	GTRCARATGCGIARTTTGAA
DPLvIKKε-R	ACWGCICYTCATACAT
LvIKKε-5' RACE1	GCAAGATGTGAAAGCACCAAG
LvIKKε-5' RACE 2	CCATCACAATCACCTTCCCAC
LvIKKε-3' RACE 1	GGCTCGCTCTTCAACATTCT
LvIKKε-3' RACE 2	GCAGGGATGAAACACTTGAGG
LvIKKβ-5' RACE1	TGTTTTAGGGCAGTGAGTGAGC
LvIKKβ-5' RACE 2	CCCAGTAAGGAAGAACACTG
LvIKKβ-3' RACE1	CTCACTCACTGCCCTAACACC
LvIKKβ-3' RACE2	GCAGCAAGAACCGCACAAC
<b>Genome walking</b>	
LvIKKε-gw1	TAACGCCCTGAAAGACGG
LvIKKε-gw2	GCTGATCCTCGCAGAAATGAC
LvIKKβ-gw1	AAGGTTTCTTACCCTGTTATT
LvIKKβ-gw2	TCCCTCAACTCAACCCTCC
<b>dsRNA preparation<sup>a</sup></b>	
dsGFP-F	AGTGCTTCAGCCGCTACCC
dsGFP-R	GCGCTTCTCGTTGGGGTC
dsGFP(T7)-F	<u>TAATACGACTCACTATAGGAG-</u> TGCTTCAGCCGCTACCC
dsGFP(T7)-R	<u>TAATACGACTCACTATAGGG-</u> CGCTTCTCGTTGGGGTC
dsLvIKKβ-F	GCTGCTGTCCGTTCTCTGC
dsLvIKKβ-R	TTTCTCCATTGCGACCTTCA
dsLvIKKβ-F(T7)	<u>TAATACGACTCACTATAGGGC-</u> TGCTGTCCGTTCTCTGC
dsLvIKKβ-R(T7)	<u>TAATACGACTCACTATAGGTTT-</u> CTCCATTGCGACCTTCA
dsLvIKKε-F	GAGGCTCGCTCTTCAACATTC
dsLvIKKε-R	TGTTACGCCGACCTCCATAC
dsLvIKKε-F(T7)	<u>TAATACGACTCACTATAGGGA-</u> GGCTCGCTCTTCAACATTC
dsLvIKKε-R(T7)	<u>TAATACGACTCACTATAGGTG-</u> TTACGCCGACCTCCATAC
<b>qPCR analysis</b>	
LvIKKβ-F	ACCACACTTCCACCTTTGG
LvIKKβ-R	TCCCGATGAAGGAAGAACAC
LvIKKε-F	TTGGCTTCTTCCAGGACAC
LvIKKε-R	TTTTATGGCTGCCAGGAGTC
LvEF-1α-F	GAAGTAGCCGCCCTGGTTG
LvEF-1α-R	CGGTTAGCCTTGGGGTTGAG
LvPEN2-F	GCATCAAGTTCGGAAGCTGT
LvPEN2-R	ACCCACATCCTTCCACAAG
LvPEN3-F	CTCTGGCTTGGAATGGAT
LvPEN3-R	GCATGGATTCACCTCCTCGT
LvPEN4-F	ATGCTACGGAATTCCCTCCT
LvPEN4-R	ATCCTTGCAACGCATAGACC
Lvlysozyme-F	AAGACACCGAACGATGGAAG
Lvlysozyme-R	TGGGGACTCGTTCTTTATG
Lvcrustin1-F	GTCGCAGTGCAGTACTGGT
Lvcrustin1-R	TAGTCGTTGGAGCACGTCTG
Lvcrustin2-F	ATCAGCAGGGGAACAAGAGA
Lvcrustin2-R	CGGACTCGCAGCAATAGACT
<b>Protein expression<sup>b</sup></b>	
pAc5.1-LvIKKβ-F	AAGAAAAAAGCGGCCGCAA-
	CATGGCAGCAGCAGAAGA
pAc5.1-LvIKKβ-R	GCTCTAGACAAGGAAGTTTCAACTGCCTTC

**Table 1 Continue**

Primer	Primer sequence (5'–3')
pAc5.1-LvIKKε-F	CGGGGTACCGAATGGCATTCTGCGAGGAT
pAc5.1-LvIKKε-R	GCTCTAGACGCAACCTCAGTTTGTAACTCTTG
pCMV-LvIKKβ-F	CCGCTCGAGATGGCAGCAGCAGAAGA
pCMV-LvIKKβ-R	GCTCTAGACAAGGAAGTTTCAACTGCCTTC
pCMV-LvIKKε-F	GGAATTCACCATGGCATTCTGCGAGGAT
pCMV-LvIKKε-R	GCTCTAGACGCAACCTCAGTTTGTAACTCTTG

Abbreviations: dsRNA, double-stranded RNA; GFP, green fluorescent protein; PEN, Penaeidins qPCR, quantitative PCR.

<sup>a</sup> T7 RNA polymerase-binding site is underlined.

<sup>b</sup> Primers used in the cellular localization and luciferase reporter assays were the same.

DNA from shrimp muscle was extracted using the Universal Genomic DNA Extraction Kit Ver. 3.0 (TaKaRa, Dalian, China) according to the manufacturer's instructions. The genomic DNA sequences of *LvIKKβ* and *LvIKKε* were obtained by PCR amplification using shrimp genomic DNA and gene-specific primers (Table 1). The genomic DNA sequences adjacent to the 5' ends of *LvIKKβ* and *LvIKKε* were obtained using a Genome Walker Universal Kit (Clontech, California, USA) as previously described.<sup>25</sup>

### Bioinformatic analysis

Nucleotide blast searches of the NCBI database were performed to retrieve IKK-like genes. Multiple sequence alignments were performed using ClustalX 2.0 (<http://www.ebi.ac.uk/tools/clustalw2>). The amino-acid sequences of *LvIKKβ* and *LvIKKε* were deduced using Simple Modular Architecture Research Tool (SMART, <http://smart.embl-heidelberg.de>). Neighbor joining phylogenetic trees were constructed using MEGA 4.0 (<http://www.megasoftware.net/index.html>) based on the deduced amino-acid sequences of related genes in typical species. Bootstrap sampling was reiterated 1000 times.

### Real-time quantitative PCR

For the tissue distribution analysis, the hemocyte, eyestalk, gill, heart, hepatopancreas, stomach, intestine, nerve, muscle, pyloric caecum and epithelium were collected from healthy shrimp. For immune challenge experiments, healthy shrimp were injected intramuscularly in the third abdominal segment with 100 μl of phosphate-buffered saline (PBS) (control, 137mM NaCl, 2.7mM KCl, 4.3mM Na<sub>2</sub>HPO<sub>4</sub>, and 1.47mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.3), 100 μl of *V. alginolyticus* inoculum (approximately 2.4 × 10<sup>6</sup> CFU) or 100 μl of WSSV inoculum (approximately 10<sup>7</sup> copies/shrimp). At 0, 3, 6, 12, 24, 36, 48 and 72 h post-injection (hpi), five shrimp from each group were randomly selected, and the hemocyte, gill, hepatopancreas and intestine were collected for quantitative PCR (qPCR) analysis. Shrimp total RNA was isolated, and qPCR cDNA templates were prepared as previously described.<sup>18,24,26</sup> The expression of *LvIKKβ* and *LvIKKε* in healthy and immune-challenged shrimp was detected using 1 μl of cDNA template, the Master SYBR Green I system and a Light Cycler (Roche, Diagnostics, Mannheim, Germany) with the following cycling parameters: 1 cycle of 95 °C for 30 s followed by 40 cycles of

95 °C for 5 s, 57 °C for 20 s and 78 °C for 1 s. qPCR was performed on three replicates per sample, and three shrimp were analyzed for each sample. The expression of *L. vannamei* elongation factor 1 $\alpha$  (*LvEF-1 $\alpha$* ) was used as an internal control. Standard curves for *LvIKK $\beta$* , *LvIKK $\epsilon$*  and *LvEF-1 $\alpha$*  were generated by running triplicate reactions of a 10-fold dilution series (10 different cDNA concentrations). The primer amplification efficiencies for *LvIKK $\beta$* , *LvIKK $\epsilon$*  and *LvEF-1 $\alpha$*  were 2.032, 2.031 and 2.023, respectively. The relative standard curve method was used for the calculation of fold changes in gene expression.<sup>27</sup>

### Plasmid construction

To examine the protein expression in *Drosophila* Schneider S2 cells, pAc5.1/V5-His A (Invitrogen, California, USA) and PCR products amplified using pAcLvIKK $\beta$ -F and pAcLvIKK $\beta$ -R were double digested, purified, ligated and transformed into competent DH5 $\alpha$  cells to select clones for sequencing. The pAc5.1-LvIKK $\epsilon$ 1 and pAc5.1-LvIKK $\epsilon$ 2 expression vectors were successfully constructed using the same procedure. We constructed an expression plasmid, pAc5.1-N-GFP, which expresses green fluorescent protein (GFP) in *Drosophila* S2 cells.<sup>23</sup> For protein localization studies in *Drosophila* Schneider S2 cells, the complete LvIKK $\beta$  open reading frame (ORF) was inserted into pAc5.1-N-GFP to construct pAc5.1-LvIKK $\beta$ -GFP, which expresses a fusion protein comprising full-length LvIKK $\beta$  and GFP. pAc5.1-LvIKK $\epsilon$ 1-GFP and pAc5.1-LvIKK $\epsilon$ 2-GFP were also constructed using the same methods. For protein expression in HEK 293T cells, the *LvIKK $\beta$* , *LvIKK $\epsilon$ 1* and *LvIKK $\epsilon$ 2* ORFs were inserted into pCMV-C-MYC (Beyotime, Shanghai, China) to construct pCMV-LvIKK $\beta$ , pCMV-LvIKK $\epsilon$ 1 and pCMV-LvIKK $\epsilon$ 2, respectively. The luciferase reporter vectors pGL3-PEN453, pGL3-PEN309, pGL3-PEN4, pGL3-Drs or pGL3-AttA were constructed in previous studies and demonstrated to be regulated predominantly through NF- $\kappa$ B activation.<sup>18,19,21,28–30</sup> Using the same method, 147 WSSV luciferase reporters were constructed as described previously.<sup>25,31</sup> An NF- $\kappa$ B luciferase reporter (Clontech, California, USA) was used to study NF- $\kappa$ B signaling in HEK 293T cells. The pRL-TK luciferase reporter vector (Promega, Madison, WI, USA) was used as an internal standard.

### Cell culture, transfection and luciferase assays

*Drosophila* S2 cells were maintained at 28 °C in Schneider's *Drosophila* medium supplemented with 10% fetal bovine serum (Invitrogen, California, USA) without CO<sub>2</sub>. When the culture density reached approximately 6 $\times$ 10<sup>6</sup>–20 $\times$ 10<sup>6</sup> viable cells/ml, *Drosophila* S2 cells were passaged onto a new plate at a density of approximately 5 $\times$ 10<sup>5</sup> viable cells/ml. HEK 293T cells were cultured in DMEM supplemented with 10% fetal bovine serum at 37 °C in a humidified 5% CO<sub>2</sub> incubator. For luciferase reporter assays, the protein expression plasmid, luciferase reporter plasmid, and pRL-TK *Renilla* luciferase plasmid were cotransfected into *Drosophila* S2 cells or HEK 293T cells that were seeded in 96-well plates 24 h before transfection. *Drosophila* S2 cells were transfected using Effectene Transfection Reagent (Qiagen, Hilden, Germany), and HEK

293T cells were transfected using Lipofectamine 2000 (Invitrogen, California, USA). Cells were harvested 36 h later and lysed for the assessment of protein expression and luciferase activity using the Dual Luciferase Reporter Assay System (Promega, Madison, WI, USA) as previously described.<sup>21,25</sup>

### Analysis of LvIKK $\beta$ and LvIKK $\epsilon$ by confocal microscopy

*Drosophila* S2 cells were seeded onto coverslips coated with poly-L-lysine in 24-well plates. Approximately 24 h later, the cells were transfected with pAc5.1-N-GFP, pAc5.1-LvIKK $\beta$ -GFP, pAc5.1-LvIKK $\epsilon$ 1-GFP or pAc5.1-LvIKK $\epsilon$ 2-GFP. At 36 h post-transfection, the cells on the coverslips were washed twice with PBS, fixed using Immunol Staining Fix Solution (Beyotime, Shanghai, China) and stained with Hoechst 33258 Solution (Beyotime, Shanghai, China). The treated cells were observed using a Leica laser scanning confocal microscope.

### dsRNA preparation and silencing of LvIKK $\beta$ and LvIKK $\epsilon$ *in vivo* by dsRNA-mediated RNA interference (RNAi)

dsRNA corresponding to *LvIKK $\beta$* , *LvIKK $\epsilon$*  and *GFP* (dsLvIKK $\beta$ , dsLvIKK $\epsilon$  and dsGFP, respectively) were prepared using the T7 RiboMAX Express Kit (Promega, Madison, WI, USA) according to the manufacturer's protocol, as previously described.<sup>32</sup> Briefly, DNA templates for the production of dsLvIKK $\beta$ , dsLvIKK $\epsilon$  and dsGFP were amplified by PCR using gene-specific primers (Table 1) with the T7 RNA polymerase-binding site at the 5' terminus to produce sense and antisense RNA strands separately. Then, the single-stranded RNA was annealed to generate dsRNA. After purification, the dsRNA was quantified and then stored at –80 °C. For dsRNA-mediated RNAi experiments, the experimental group (1–2 g per shrimp) was injected with dsLvIKK $\beta$  or dsLvIKK $\epsilon$  (1  $\mu$ g/g shrimp) by intramuscular injection, whereas the control groups were injected with dsGFP or PBS, respectively. To determine silencing effects, gill samples from at least three shrimp per treatment group were collected at 0, 24, 72, 120 and 144 h post-dsRNA injection (hpi), and total RNA was extracted. Total RNA from the gills of the dsRNA-injected *L. vannamei* was reverse transcribed into cDNA for the analysis of *LvIKK $\beta$* , *LvIKK $\epsilon$* , *PEN*, *lysozyme* and *crustin* expression using qPCR as described in the section on 'Real-time quantitative PCR'.

### WSSV infection experiments in dsRNA-injected *L. vannamei*

The expression of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  was significantly reduced in dsLvIKK $\beta$ - and dsLvIKK $\epsilon$ -injected *L. vannamei*, respectively, compared with the dsGFP-injected control groups (>80%) at most of the time points examined by qPCR analysis. To further investigate the roles of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in viral infection, we performed WSSV infection experiments. We infected *L. vannamei* intramuscularly with 100  $\mu$ l of WSSV inoculum (approximately 10<sup>7</sup> copies/shrimp) at 48 h post-dsRNA injection, and mortalities were recorded.

### Statistical analysis

Student's *t*-test was used to compare means between two samples using Microsoft Excel. In all cases, differences were considered significant at *P*<0.05 and highly significant at *P*<0.01.



The data are presented as the mean  $\pm$  standard error (standard error of the mean).


## RESULTS

### Cloning and sequence analysis of *LvIKK $\beta$* and *LvIKK $\epsilon$*


The full-length cDNA of *LvIKK $\beta$*  was 2479 bp, with a 2376-bp ORF, a 49-bp 5' untranslated region and a 54-bp 3' untranslated region (Figure 1a). The sequence of *LvIKK $\beta$*  has been deposited in NCBI GenBank under accession no. AEK86518. A sequence analysis indicated that *LvIKK $\beta$*  contains N-terminal protein kinase domains (KDs) and more C-terminally located leucine zipper (LZ) and helix-loop-helix (HLH) motifs but lacks the IKK $\gamma$ - (also known as NEMO) binding domain (Figure 1b).<sup>6</sup> The full-length *LvIKK $\beta$*  protein exhibits 24.7% and 34.7% identity with *Drosophila* IKK $\beta$  and human IKK $\beta$ , respectively (Supplementary Figure 2). We obtained two isoforms of *LvIKK $\epsilon$* . The *LvIKK $\epsilon$ 1* cDNA was 2513 bp, with a 2205-bp ORF, a 17-bp 5' untranslated region and a 291-bp 3' untranslated region (Figure 1a). Compared with *LvIKK $\epsilon$ 1*, *LvIKK $\epsilon$ 2* lacks a 90-bp sequence from 1596 to 1686 bp of

*LvIKK $\epsilon$ 1*, which encodes a 30-amino acid (aa) protein sequence (Figure 1a). The sequences of *LvIKK $\epsilon$ 1* and *LvIKK $\epsilon$ 2* have been deposited in NCBI GenBank under accession nos. AEK86519 and AEK86520. *LvIKK $\epsilon$*  contains N-terminal protein KD and C-terminal LZ domains but lacks HLH motifs and the IKK $\gamma$ -binding domain (Figure 1b).<sup>6</sup> *LvIKK $\epsilon$*  exhibits 36.4% and 35.1% identity with human TBK1 and IKK $\epsilon$ , respectively (Supplementary Figure 2). Moreover, *LvIKK $\epsilon$*  shares 23% identity with *LvIKK $\beta$* .

The genomic sequence of *LvIKK $\beta$*  is 7866 bp and harbors 11 exons and 10 introns (Figure 1a). The genomic sequence of *LvIKK $\epsilon$*  is 8955 bp (Figure 1a). The *LvIKK $\epsilon$ 1* cDNA is composed of 14 exons, whereas the *LvIKK $\epsilon$ 2* cDNA lacks the 11th exon, which encodes a 30-aa protein, compared with *LvIKK $\epsilon$ 1*, suggesting that *LvIKK $\epsilon$ 1* and *LvIKK $\epsilon$ 2* are two alternatively spliced isoforms of *LvIKK $\epsilon$*  (Figure 1a). Using Genome Walker, we amplified the genomic regions upstream of the 5' untranslated regions of *LvIKK $\beta$*  and *LvIKK $\epsilon$* . In the 1989-bp region upstream of the *LvIKK $\beta$*  5' end, NF- $\kappa$ B, SP1, AP-1, GATA and STAT motifs involved in



**Figure 1** Genomic organization (a), domain topology (b) and tissue distribution of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  (c). (a) Schematic diagrams of the genomic structures of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  and the splicing isoforms of *LvIKK $\epsilon$* . The numbers represent exons. Exons are depicted as boxes, and introns are depicted as lines. The eleventh exon is red, indicating that it is the eleventh exon of *LvIKK $\epsilon$ 1* and a partial intron of *LvIKK $\epsilon$ 2*. (b) Schematic representation of the domain topology of *LvIKK $\beta$*  and *LvIKK $\epsilon$* . *LvIKK $\beta$*  contains N-terminal protein KDs and more C-terminally located LZ and HLH motifs, whereas *LvIKK $\epsilon$*  contains N-terminal protein KDs and a C-terminal LZ domain. (c) Tissue distribution of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in healthy *L. vannamei*. The hemocyte, hepatopancreas, epithelium, intestine, eyestalk, stomach, gill, heart, pyloric cecum, nerve and muscle were collected from healthy *L. vannamei* to extract total RNA for tissue distribution analysis. The expression was normalized to that of *LvEF-1 $\alpha$*  using the relative standard curve method. The expression of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in the hepatopancreas was set to 1.0. The data are expressed as the mean fold change (means  $\pm$  s.e.,  $n=3$ ). HLH, helix-loop-helix; KD, kinase domain; LZ, leucine zipper.



**Figure 2** Temporal expression of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in the gill, hemocyte, hepatopancreas and intestine after challenge with *V. alginolyticus* or WSSV. Healthy *L. vannamei* were injected intramuscularly at the third abdominal segment with 100  $\mu$ l of PBS (control group), 100  $\mu$ l of *V. alginolyticus* ( $2.4 \times 10^6$  CFU) or 100  $\mu$ l of WSSV inocula ( $10^7$  copies). At 0, 3, 6, 12, 24, 36, 48 and 72 hpi, five shrimp were randomly selected from each group from which the gill, hemocyte, hepatopancreas and intestine were obtained for qPCR analysis. The expression of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in the untreated control group (0 hpi) was set to 1.0. The expression was normalized to that of *LvEF-1 $\alpha$*  using the relative standard curve method. qPCR was performed in triplicate for each sample. The data are expressed as the mean fold change (means  $\pm$  s.e.,  $n=3$ ) relative to the untreated group (0 hpi). Statistical significance was calculated using Student's *t*-test. Bars with \* indicate statistically significant differences ( $P < 0.05$ ); bars with \*\* indicate highly statistically significant ( $P < 0.01$ ). CFU, colony-forming unit; hpi, post-injection; PBS, phosphate-buffered saline; qPCR, quantitative PCR; WSSV, white spot syndrome virus.

the transcriptional regulation of immune system genes in arthropods were identified (Supplementary Figure 1a). In the 401-bp region upstream of the *LvIKK $\epsilon$*  5' untranslated region, GATA, SP1 and TATA box motifs were found (Supplementary Figure 1b).

### Phylogenetic tree construction

Phylogenetic analysis of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  showed that *LvIKK $\beta$*  is a member of the IKK $\alpha$  and IKK $\beta$  group, whereas *LvIKK $\epsilon$*  belongs to the IKK $\epsilon$  and TBK1 group (Supplementary Figure 3). The results also revealed that invertebrates may only possess one member of the IKK $\alpha$  and IKK $\beta$  group and one member of the IKK $\epsilon$  and TBK1 group.

### Tissue distribution of *LvIKK $\beta$* and *LvIKK $\epsilon$* in healthy shrimp

In healthy shrimp, *LvIKK $\beta$*  was highly expressed in the muscle (65.3-fold), nerve (8.1-fold), heart (7.0-fold), eyestalk (4.3-fold), hemocyte (3.0-fold), pyloric cecum (3.0-fold), gill (2.9-fold), epithelium (2.9-fold), stomach (2.3-fold) and intestine (1.7-fold) compared with mRNA expression in the hepatopancreas (1.0-fold) (Figure 1c). In healthy shrimp, *LvIKK $\epsilon$*  was highly expressed in the muscle (6.2-fold), gill (3.0-fold), nerve (2.7-fold), stomach (2.5-fold), heart (2.4-fold), pyloric cecum (2.2-fold), intestine (2.1-fold), hemocyte (1.2-fold), epithelium (1.2-fold) and eyestalk (1.1-fold) compared with mRNA expression in the hepatopancreas (1.0-fold) (Figure 1d).

### Expression profiles of *LvIKK $\beta$* and *LvIKK $\epsilon$* after microbial challenge

After WSSV infection, the expression of *LvIKK $\beta$*  increased in the gill compared with the PBS injection group, but no significant changes were observed in the hemocyte, hepatopancreas or intestine (Figure 2). *LvIKK $\epsilon$*  was slightly upregulated in the gill, hemocyte and hepatopancreas but not in the intestine after WSSV infection (Figure 2). After *V. alginolyticus* infection, *LvIKK $\beta$*  was upregulated in the gill, hepatopancreas and intestine compared with the PBS injection group, but downregulated in the hemocyte (Figure 2). After *V. alginolyticus* infection, *LvIKK $\epsilon$*  was upregulated in the gill, hepatopancreas and intestine and downregulated in the hemocyte (Figure 2).


### Cellular localization of *LvIKK $\beta$* and *LvIKK $\epsilon$* in *Drosophila* S2 cells

Using fluorescent imaging by confocal microscopy of the *LvIKK $\beta$* -GFP and *LvIKK $\epsilon$* -GFP fusion proteins, we observed that *LvIKK $\beta$* -GFP and *LvIKK $\epsilon$* -GFP were localized to the cytoplasm and nucleus (Figure 3). However, the green fluorescence of the *LvIKK $\epsilon$* -2-GFP fusion protein was confined to the cytoplasm, suggesting that the 30-aa protein sequence in *LvIKK $\epsilon$* 1, which *LvIKK $\epsilon$* 2 lacks, is a potential nuclear localization signal and is necessary for *LvIKK $\epsilon$* 1 nuclear localization (Figure 3c).


### 3.6. *LvIKK $\beta$* but not *LvIKK $\epsilon$* activates shrimp AMP promoters

In *Drosophila* S2 cells, the expression of *LvIKK $\beta$*  increased the promoter activities of *Drosophila Drosomycin* (*Drs*) and *Attacin A* (*Atta*) approximately 3.00- and 3.09-fold, respectively (Figure 4a). *LvIKK $\beta$*  also induced *Penaeus monodon Penaeidin* promoter activities approximately 2.00- and 3.67-fold for PEN453 and PEN309, respectively, and increased *L. vannamei Penaeidin4* promoter activity approximately 4.82-fold (Figure 4a). However, neither *LvIKK $\epsilon$* 1 nor *LvIKK $\epsilon$* 2 could activate shrimp and *Drosophila* AMP genes in *Drosophila* S2 cells (Figure 4a). These results support the possibility that *LvIKK $\beta$*  serves as a positive regulator of the shrimp NF- $\kappa$ B signaling pathway for AMP activation.

When transfected into HEK 293T cells, *LvIKK $\beta$* , *LvIKK $\epsilon$* 1 and *LvIKK $\epsilon$* 2 induced NF- $\kappa$ B activity approximately 13.2-, 13.4- and 5.5-fold, respectively, compared with the control group (Figure 4b). In addition, *LvIKK $\epsilon$* 1 induced NF- $\kappa$ B activity to 2.4 times that of *LvIKK $\epsilon$* 2 ( $P < 0.01$ ). Therefore, the additional 30-aa nuclear localization signal of *LvIKK $\epsilon$* 1 may be vital for the greater activation of NF- $\kappa$ B luciferase reporters in HEK 293T cells.



**Figure 3** Subcellular localizations of *LvIKK $\beta$*  (a), *LvIKK $\epsilon$* 1 (b) and *LvIKK $\epsilon$* 2 (c) in *Drosophila* S2 cells by confocal microscopy. *Drosophila* S2 cells were transfected with pAc5.1-*LvIKK $\beta$* -GFP plasmid, pAc5.1-*LvIKK $\epsilon$* 1-GFP plasmid or pAc5.1-*LvIKK $\epsilon$* 2-GFP plasmid. At 36 h post-transfection, the cover slips were washed, fixed, and stained with Hoechst 33258. Localization of recombinant proteins was examined under a Leica laser scanning confocal microscope. (a) *Drosophila* S2 cells transfected with pAc5.1-*LvIKK $\beta$* -GFP showed cytoplasmic and nuclear localization of the *LvIKK $\beta$* -GFP fusion protein. (b) *Drosophila* S2 cells transfected with pAc5.1-*LvIKK $\epsilon$* 1-GFP showed cytoplasmic and nuclear localization of the *LvIKK $\beta$* -GFP fusion protein. (c) *Drosophila* S2 cells transfected with pAc5.1-*LvIKK $\epsilon$* 2-GFP showed cytoplasmic localization of the *LvIKK $\epsilon$* 2-GFP fusion protein. *LvIKK $\epsilon$* 1 contains 30 aa encoded by the eleventh exon of *LvIKK $\epsilon$* 1, suggesting that the 30-aa sequence is necessary for nuclear localization of *LvIKK $\epsilon$* 1. aa, amino acid; GFP, green fluorescent protein.



**Figure 4** (a) The overexpression of LvIKK $\beta$  but not LvIKK $\epsilon$  activated *Drosophila* and shrimp AMP gene promoters in *Drosophila* S2 cells. *Drosophila* S2 cells were transfected with 0.05  $\mu$ g of protein expression vector (LvIKK $\beta$ , LvIKK $\epsilon$ 1 or LvIKK $\epsilon$ 2), 0.05  $\mu$ g luciferase reporter plasmid (pGL3-Basic, pGL3-PEN453, pGL3-PEN309, pGL3-PEN4, pGL3-Drs, or pGL3-AttA) and 0.005  $\mu$ g pRL-TK *Renilla* luciferase plasmid (as an internal control; Promega, Madison, WI, USA). (b) LvIKK $\beta$  and LvIKK $\epsilon$  activated NF- $\kappa$ B luciferase reporter in the HEK 293T cells. HEK 293T cells were cotransfected with the pCMV empty vector (70 ng), pCMV-LvIKK $\epsilon$ 2 (70 ng), pCMV-LvIKK $\epsilon$ 1 (70 ng) or pCMV-LvIKK $\beta$  (70 ng) together with the NF- $\kappa$ B luciferase reporter vector (30 ng) and the internal control *Renilla* expression vector (3 ng). pCMV empty vector and the pTAL luciferase reporter vector (Clontech, California, USA) were used as negative controls (blue bars). At 36 h after transfection, the cells were harvested and analyzed using the Dual Luciferase kit (Promega). The bars indicate the mean  $\pm$  s.d. of luciferase activity ( $n=3$ ). Statistical significance was calculated by the Student's *t*-test (not statistical significant # $P>0.05$ ; statistical significant \* $P<0.05$ ; highly statistical significant \*\* $P<0.01$ ). AMP, antimicrobial peptide; CMV, cytomegalovirus; PEN, Penaeidins.

### ***In vivo* knockdown of LvIKK $\beta$ and LvIKK $\epsilon$ by dsRNA-mediated RNAi**


To further confirm the roles of LvIKK $\beta$  and LvIKK $\epsilon$  in the regulation of shrimp AMPs, dsRNA-mediated RNAi experiments were performed as described previously.<sup>32</sup> The results from the qPCR analysis indicate that we successfully suppressed the expression of LvIKK $\beta$  and LvIKK $\epsilon$ . In the gills of dsLvIKK $\beta$ -injected *L. vannamei*, the expression of LvIKK $\beta$  was significantly reduced to 25.7%, 23.4%, 12.2% and 27.6% at 24, 72, 120 and 144 hpi, respectively, of the levels observed in the dsGFP control group (Figure 5a). In the gills of dsLvIKK $\epsilon$ -injected *L. vannamei*, the expression of LvIKK $\epsilon$  was significantly reduced to 16.1%, 23.4%, 12.5% and 10.3% at 24, 72, 120

and 144 hpi, respectively, of the levels observed in the dsGFP control group (Figure 5b).

### **Decreased AMP expression in LvIKK $\beta$ - and LvIKK $\epsilon$ -silenced shrimp**

Using luciferase assays, we demonstrated that overexpressed LvIKK $\beta$  and LvIKK $\epsilon$  increase the promoter activities of LvPEN4 and *P. monodon* Penaeidin (Figure 4a) in *Drosophila* S2 cells. Next, we investigated the effects of LvIKK $\beta$  and LvIKK $\epsilon$  on the expression of LvPENS, *Lvlysozyme* and *Lvcrustins in vivo*. In the gills of LvIKK $\beta$ -silenced *L. vannamei*, the levels of LvPEN2, LvPEN3, LvPEN4, *Lvlysozyme*, *Lvcrustin1* and *Lvcrustin2* were quantitated using qPCR. We observed that





**Figure 5** Silencing of *LvIKK $\beta$*  (a) and *LvIKK $\epsilon$*  (b) by dsRNA-mediated RNAi. Shrimps were injected with PBS, dsGFP (control), dsLvIKK $\beta$  or dsLvIKK $\epsilon$ , and the gill was collected at the indicated time points for isolation of total RNA and synthesis of cDNA. The expression levels of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  were determined by qPCR. qPCR was performed in triplicate for each sample. The data are expressed as the mean fold change (means $\pm$ s.e.,  $n=3$ ) relative to the untreated group (0 hpi). GFP, green fluorescent protein; hpi, post-injection; PBS, phosphate-buffered saline; RNAi, RNA interference; qPCR, quantitative PCR.

at most of the time points, all of the shrimp AMPs detected, including *LvPEN2*, *LvPEN3*, *LvPEN4*, *Lvlysozyme*, *Lvcrustin1* and *Lvcrustin2*, were significantly downregulated (Figure 6). In the gills of *LvIKK $\epsilon$* -silenced *L. vannamei*, the expression of these shrimp AMPs, including *LvPEN2*, *LvPEN3*, *LvPEN4*, *Lvlysozyme*, *Lvcrustin1* and *Lvcrustin2*, was also significantly decreased (Figure 6).

#### Knockdown of *LvIKK $\beta$* and *LvIKK $\epsilon$* increases survival rates after WSSV infection

WSSV is one of the most common and destructive pathogens in shrimp aquaculture, and shrimp mortality can reach 100% within 3–10 days after infection. To further evaluate the role of *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in shrimp immune responses, we performed WSSV infection experiments in dsRNA-injected *L. vannamei*. At 48 h after dsRNA injection, *L. vannamei* were infected with WSSV, and mortalities were recorded. We observed that injection of dsGFP, dsLvIKK $\beta$  or dsLvIKK $\epsilon$  could delay the initial outbreak of WSSV compared with the PBS injection group (Figure 7a). Compared with the dsGFP


injection group, dsLvIKK $\beta$ - or dsLvIKK $\epsilon$ -injected shrimp exhibited significant resistance to WSSV infection from 35 to 98 hpi (Figure 7a).

#### Screening for viral genes regulated by *LvIKK $\beta$* and *LvIKK $\epsilon$*

Because both the *LvIKK $\beta$* - and *LvIKK $\epsilon$* -silenced shrimp were resistant to WSSV infection, we hypothesized that WSSV infection requires the activation of the IKK–NF- $\kappa$ B signaling pathway to facilitate viral gene expression. We constructed luciferase reporters for 147 WSSV genes (Figure 7b). By screening *Drosophila* S2 cells co-transfected with *LvIKK $\beta$*  or *LvIKK $\epsilon$* , we determined that the *WSSV051*, *WSSV059*, *WSSV069* (*ie1*), *WSSV083*, *WSSV090*, *WSSV107*, *WSSV244*, *WSSV249*, *WSSV303*, *WSSV371* and *WSSV445* promoters can be activated by *LvIKK $\beta$*  or *LvIKK $\epsilon$*  (Figure 7c). Among these, *WSSV051*, *WSSV069* (*ie1*) and *WSSV083* are immediate early genes.

#### DISCUSSION

IKK $\alpha/\beta$  and TBK1/IKK $\epsilon$  are the central regulators of the NF- $\kappa$ B signaling pathway in mammals and represent a point of



**Figure 6** Silencing of *LvIKK $\beta$*  or *LvIKK $\epsilon$*  led to decrease in expression of shrimp AMPs, including *LvPEN2*, *LvPEN3*, *LvPEN4*, *Lvlysozyme*, *Lvcrustin1* and *Lvcrustin2* in the gill. Shrimps were injected with PBS, dsGFP (control), dsLvIKK $\beta$  or dsLvIKK $\epsilon$ , and the gill was collected at the indicated time points for isolation of total RNA and synthesis of first-strand cDNA. The expression levels of *LvPEN2*, *LvPEN3*, *LvPEN4*, *Lvlysozyme*, *Lvcrustin1* and *Lvcrustin2* in the gill of dsRNA-injected shrimps were determined using qPCR. qPCR was performed in triplicate for each sample. The data are expressed as the mean fold change (means  $\pm$  s.e.,  $n=3$ ) relative to the untreated group (0 hpi). AMP, antimicrobial peptide; dsRNA, double-stranded RNA; hpi, post-injection; PBS, phosphate-buffered saline; PEN, Penaeidins; qPCR, quantitative PCR.

convergence for most of the signal transduction pathways that lead to NF- $\kappa$ B activation.<sup>33</sup> The IKK–NF- $\kappa$ B cascades are also common targets of various viruses, such as the Tax transactivator oncoprotein of HTLV-1, which can directly target IKK $\alpha$ / $\beta$  and lead to NF- $\kappa$ B activation, resulting in the upregulation of NF- $\kappa$ B signaling pathway-controlled genes.<sup>34</sup> However, in invertebrates, most studies of IKK family proteins have been performed in *D. melanogaster*. In this study, we investigated the roles of LvIKK $\beta$  and LvIKK $\epsilon$  in shrimp AMP regulation and WSSV infection.


Here, we report two isoforms of IKK $\epsilon$  from *L. vannamei*: LvIKK $\epsilon$ 1 and LvIKK $\epsilon$ 2. LvIKK $\epsilon$ 1 possesses an additional 30-aa protein sequence that is absent in LvIKK $\epsilon$ 2 and other species. Confocal microscopy indicates that this 30-aa protein sequence may be responsible for the cellular localization of LvIKK $\epsilon$ 1 (Figure 3). Moreover, this 30-aa protein sequence may be responsible for the higher activity (2.4-fold higher) of an NF- $\kappa$ B reporter induced by LvIKK $\epsilon$ 1 than by LvIKK $\epsilon$ 2 (Figure 4b). LvIKK $\beta$  is a unique invertebrate homolog of mammalian IKK $\alpha$ /IKK $\beta$  in *L. vannamei* (Supplementary Figure 3). LvIKK $\beta$  also strongly induced NF- $\kappa$ B activity (8.1-fold) in human HEK 293T cells (Figure 4b). In *Drosophila* S2 cells, LvIKK $\beta$  but not LvIKK $\epsilon$  activated the AMP reporters of *Drosophila* and shrimp, consistent with their induction of NF- $\kappa$ B activity in HEK 293T cells (Figure 4a). The lack of activity of LvIKK $\epsilon$  on AMP promoters may be explained by the absence of elements or components of the NF- $\kappa$ B signaling pathway in *Drosophila* S2 cells that are essential for LvIKK $\epsilon$  activity.

The expression of *Drosophila* and shrimp AMPs is believed to be primarily controlled by the NF- $\kappa$ B signaling pathway.<sup>10,23,30,35,36</sup> In our previous studies, we showed that other components of the shrimp Toll and IMD pathways, such as LvToll1-3, LvIMD, and the NF- $\kappa$ B family proteins LvDorsal and LvRelish, can activate AMP luciferase reporters.<sup>18–21,25</sup> We also reported that the shrimp NF- $\kappa$ B family proteins LvRelish and LvDorsal can not only activate a shrimp AMP luciferase reporter in *Drosophila* S2 cells, but that they can also bind to putative NF- $\kappa$ B-binding sites in the AMP and WSSV069 (*ie1*) promoter regions.<sup>19,20,25,31</sup> Here, using dsRNA-mediated gene silencing, we further investigated the regulation of shrimp AMPs by the Toll and IMD pathways. In LvIKK $\beta$ - or LvIKK $\epsilon$ -silenced shrimp, LvPEN2, LvPEN3, LvPEN4, Lvlysozyme, Lvcrustin1 and Lvcrustin2 were significantly reduced (Figure 6). Combined with the activation of AMP luciferase reporters by LvIKK $\beta$  and LvIKK $\epsilon$ , we propose that the IKK–NF- $\kappa$ B signaling pathway regulates shrimp AMP expression. However, the detailed mechanism by which the NF- $\kappa$ B signaling pathway is activated by LvIKK $\beta$  and LvIKK $\epsilon$  is still unknown. In the *Drosophila* IMD pathway, IKK $\beta$  phosphorylates the p100-like NF- $\kappa$ B precursor protein Relish, and phosphorylated Relish is activated by proteolytic cleavage and translocated into the nucleus where it promotes AMP expression.<sup>37</sup> In future studies, we will investigate whether and how LvIKK $\beta$  phosphorylates and activate LvRelish. In the

*Drosophila* Toll pathway, the activation of the p65-like NF- $\kappa$ B proteins DIF and Dorsal does not require IKK $\beta$  or IKK $\epsilon$ .<sup>14</sup> To date, it is still unknown which kinase can act directly on Cactus to liberate DIF and Dorsal from the cytoplasm to the nucleus for AMP expression.<sup>14</sup> This study is the first report that an invertebrate IKK $\epsilon$  (LvIKK $\epsilon$ ) participates in the activation of the NF- $\kappa$ B signaling pathway. Which pathway (Toll or IMD) LvIKK $\epsilon$  participates in and how LvIKK $\epsilon$  activates this pathway is of great interest. We are also attempting to study whether LvIKK $\epsilon$  directly phosphorylates LvCactus and activates LvDorsal.

Many viruses such as HIV-1 hijack and stimulate the host NF- $\kappa$ B signaling pathway as part of their life cycles, diverting NF- $\kappa$ B immune regulatory functions to favor viral replication.<sup>38</sup> The WSSV genome encodes a protein, WSSV449, with similarity to LvPelle (LvIRAK4) of the shrimp Toll pathway.<sup>25</sup> WSSV449 could activate the NF- $\kappa$ B pathway at the same time as LvPelle.<sup>25</sup> Several studies have also reported that WSSV infection can activate the shrimp NF- $\kappa$ B signaling pathway.<sup>25,29,31,39–41</sup> In this study, we found that LvIKK $\beta$ - or LvIKK $\epsilon$ -silenced shrimp in which the activities of the NF- $\kappa$ B signaling pathway are attenuated are resistant to WSSV infection (Figure 7a). At the same time, we observed that the activities of the NF- $\kappa$ B signaling pathway in LvIKK $\beta$ - or LvIKK $\epsilon$ -silenced shrimp are significantly reduced by measuring the expression level of NF- $\kappa$ B signaling pathway-targeted genes, including LvPEN2, LvPEN3, LvPEN4, Lvlysozyme, Lvcrustin1 and Lvcrustin2 (Figure 6). Like HTLV-1, HIV-1, Xenotropic murine leukemia virus-related virus, cytomegalovirus (CMV), herpesvirus, hepatitis B virus and Epstein–Barr virus, a successful WSSV infection may also rely on the effective activation of the NF- $\kappa$ B signaling pathway.<sup>38,42</sup> To investigate the potential molecular mechanism for activation, we performed a large-scale screening of 147 WSSV gene promoters (Figure 7b). We found that the overexpression of both LvIKK $\beta$  and LvIKK $\epsilon$  activates the promoters of several WSSV genes, including WSSV051, WSSV059, WSSV069 (*ie1*), WSSV083, WSSV090, WSSV107, WSSV244, WSSV249, WSSV303, WSSV371 and WSSV445 (Figure 7c, Supplemental Figure 4). Bioinformatic analysis indicates that there are NF- $\kappa$ B binding sites in some, but not all, of these viral gene promoters, suggesting the direct or indirect activation of their promoters by the IKK–NF- $\kappa$ B signaling pathway. WSSV051, WSSV069 (*ie1*) and WSSV083 are WSSV immediate-early genes that may be important for activating the expression of other WSSV genes and may be advantageous to the virus infection cycle.<sup>43,44</sup> WSSV069 (*ie1*) can function as a transcriptional regulator, exhibits transactivation and DNA-binding activities and can take advantage of the shrimp STAT pathway to enhance its own expression.<sup>45,46</sup> The functions of WSSV090, WSSV107, WSSV244, WSSV303, WSSV371 and WSSV445 in viral infection, which are of great interest, are still unknown.

In this study, we cloned LvIKK $\beta$  and LvIKK $\epsilon$  and characterized their functions in shrimp AMP regulation and WSSV infection. Our *in vitro* and *in vivo* studies demonstrate that



**Figure 7** (a) Silencing of *LvIKK $\beta$*  or *LvIKK $\epsilon$*  could delay the outbreak time of WSSV infection. The efficiency of gene silencing in ds*LvIKK $\beta$* - and ds*LvIKK $\epsilon$* -injected *L. vannamei* was significant compared with the control groups (>80%) at all the examined time points by qPCR analysis. In the WSSV infection experiments, *L. vannamei* were injected intramuscularly with WSSV inocula (approximately  $10^7$  copies/shrimp) at 48 h after dsRNA injection. The mortalities of shrimps injected with PBS, dsGFP (control), ds*LvIKK $\beta$*  or ds*LvIKK $\epsilon$*  after WSSV infection were recorded. (b) Identification of viral genes activated by *LvIKK $\beta$*  or *LvIKK $\epsilon$*  through large-scale screening. Because both the *LvIKK $\beta$* - and *LvIKK $\epsilon$* -silenced shrimp showed resistant to WSSV infection, we speculated that WSSV infection required activation of the IKK-NF- $\kappa$ B signaling pathway to facilitate viral gene expression. We constructed luciferase reporters for 147 WSSV genes and investigate their activation by *LvIKK $\beta$*  or *LvIKK $\epsilon$* . (c) Promoter activities of *WSSV051*, *WSSV059*, *WSSV069*, *WSSV083*, *WSSV090*, *WSSV107*, *WSSV244*, *WSSV303*, *WSSV371* and *WSSV445* were induced by *LvIKK $\beta$*  and *LvIKK $\epsilon$*  in *Drosophila* S2 cell. dsRNA, double-stranded RNA; GFP, green fluorescent protein; PBS, phosphate-buffered saline; qPCR, quantitative PCR; WSSV, white spot syndrome virus.



both LvIKKβ and LvIKKε can regulate shrimp AMP expression by activating the NF-κB signaling pathway. We also observed that *LvIKKβ*- or *LvIKKε*-silenced shrimp are resistant to WSSV infection, which may be explained by the requirement of IKK–NF-κB signaling pathway activation for WSSV infection. Finally, we also identified the WSSV genes that may be regulated by the IKK–NF-κB signaling pathway through a large-scale viral promoter activity screen. This study extends our knowledge of shrimp AMP regulation and the interaction between the shrimp IKK–NF-κB signaling pathway and WSSV infection.

## ACKNOWLEDGEMENTS

This research was supported by National Natural Science Foundation of China under grant No. U1131002, the National High Technology Research and Development Program of China (973 Program) 2012CB114401, Technology Planning Project of Guangdong Province under number 2011A020102002, and China Agriculture Research System CARS-47.

Supplementary Information accompanies the paper on *Cellular & Molecular Immunology* website.

- Kawai T, Akira S. The roles of TLRs, RLRs and NLRs in pathogen recognition. *Int Immunol* 2009; **21**: 317–337.
- Kumar H, Kawai T, Akira S. Toll-like receptors and innate immunity. *Biochem Biophys Res Commun* 2009; **388**: 621–625.
- Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010; **140**: 805–820.
- Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 2006; **124**: 783–801.
- Takeuchi O, Akira S. Innate immunity to virus infection. *Immunol Rev* 2009; **227**: 75–86.
- Hacker H, Karin M. Regulation and function of IKK and IKK-related kinases. *Sci STKE* 2006; **2006**: re13.
- Kumar H, Kawai T, Akira S. Pathogen recognition in the innate immune response. *Biochem J* 2009; **420**: 1–16.
- Ferrandon D, Imler JL, Hetru C, Hoffmann JA. The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections. *Nat Rev Immunol* 2007; **7**: 862–874.
- Hoffmann JA. The immune response of *Drosophila*. *Nature* 2003; **426**: 33–38.
- Lemaitre B, Hoffmann J. The host defense of *Drosophila melanogaster*. *Annu Rev Immunol* 2007; **25**: 697–743.
- Sabin LR, Hanna SL, Cherry S. Innate antiviral immunity in *Drosophila*. *Curr Opin Immunol* 2010; **22**: 4–9.
- Kemp C, Imler JL. Antiviral immunity in *Drosophila*. *Curr Opin Immunol* 2009; **21**: 3–9.
- Erturk-Hasdemir D, Broemer M, Leulier F, Lane WS, Paquette N, Hwang D *et al*. Two roles for the *Drosophila* IKK complex in the activation of Relish and the induction of antimicrobial peptide genes. *Proc Natl Acad Sci USA* 2009; **106**: 9779–9784.
- Ganesan S, Aggarwal K, Paquette N, Silverman N. NF-κB/Rel proteins and the humoral immune responses of *Drosophila melanogaster*. *Curr Top Microbiol Immunol* 2011; **349**: 25–60.
- Dubin-Bar D, Bitan A, Bakhrat A, Kaiden-Hasson R, Etzion S, Shaanan B *et al*. The *Drosophila* IKK-related kinase (Ik2) and Spindle-F proteins are part of a complex that regulates cytoskeleton organization during oogenesis. *BMC Cell Biol* 2008; **9**: 51.
- Kuranaga E, Kanuka H, Tonoki A, Takemoto K, Tomioka T, Kobayashi M *et al*. *Drosophila* IKK-related kinase regulates nonapoptotic function of caspases via degradation of IAPs. *Cell* 2006; **126**: 583–596.
- de Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. The Toll and Imd pathways are the major regulators of the immune response in *Drosophila*. *EMBO J* 2002; **21**: 2568–2579.
- Wang PH, Liang JP, Gu ZH, Wan DH, Zhu WB, Qiu W *et al*. Molecular cloning, characterization and expression analysis of two novel Toll (LvToll2 and LvToll3) and three putative Spatzle-like Toll ligands (LvSpz1–3) from *Litopenaeus vannamei*. *Dev Comp Immunol* 2012; **36**: 359–371.
- Huang XD, Yin ZX, Jia XT, Liang JP, Ai HS, Yang LS *et al*. Identification and functional study of a shrimp Dorsal homologue. *Dev Comp Immunol* 2010; **34**: 107–113.
- Huang XD, Yin ZX, Liao JX, Wang PH, Yang LS, Ai HS *et al*. Identification and functional study of a shrimp Relish homologue. *Fish Shellfish Immunol* 2009; **27**: 230–238.
- Wang PH, Gu ZH, Huang XD, Liu BD, Deng XX, Ai HS *et al*. An immune deficiency homolog from the white shrimp, *Litopenaeus vannamei*, activates antimicrobial peptide genes. *Mol Immunol* 2009; **46**: 1897–1904.
- de Lorigeril J, Gueguen Y, Goarant C, Goyard E, Mugnier C, Fievet J *et al*. A relationship between antimicrobial peptide gene expression and capacity of a selected shrimp line to survive a *Vibrio* infection. *Mol Immunol* 2008; **45**: 3438–3445.
- Wang PH, Wan DH, Gu ZH, Deng XX, Weng SP, Yu XQ *et al*. *Litopenaeus vannamei* tumor necrosis factor receptor-associated factor 6 (TRAF6) responds to *Vibrio alginolyticus* and white spot syndrome virus (WSSV) infection and activates antimicrobial peptide genes. *Dev Comp Immunol* 2011; **35**: 105–114.
- Wang PH, Wan DH, Pang LR, Gu ZH, Qiu W, Weng SP *et al*. Molecular cloning, characterization and expression analysis of the tumor necrosis factor (TNF) superfamily gene, TNF receptor superfamily gene and lipopolysaccharide-induced TNF-α factor (LITAF) gene from *Litopenaeus vannamei*. *Dev Comp Immunol* 2012; **36**: 39–50.
- Wang PH, Gu ZH, Wan DH, Zhang MY, Weng SP, Yu XQ *et al*. The shrimp NF-κB pathway is activated by white spot syndrome virus (WSSV) 449 to facilitate the expression of WSSV069 (ie1), WSSV303 and WSSV371. *PLoS ONE* 2011; **6**: e24773.
- Wang PH, Wan DH, Gu ZH, Deng XX, Weng SP, Yu XQ *et al*. *Litopenaeus vannamei* tumor necrosis factor receptor-associated factor 6 (TRAF6) responds to *Vibrio alginolyticus* and white spot syndrome virus (WSSV) infection and activates antimicrobial peptide genes. *Dev Comp Immunol* 2011; **35**: 105–114.
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001; **29**: e45.
- Ho SH, Song YL. Cloning of penaeidin gene promoter in tiger shrimp (*Penaeus monodon*). *Fish Shellfish Immunol* 2009; **27**: 73–77.
- Li F, Wang D, Li S, Yan H, Zhang J, Wang B *et al*. A Dorsal homologue (FcDorsal) in the Chinese shrimp *Fenneropenaeus chinensis* is responsive to both bacteria and WSSV challenge. *Dev Comp Immunol* 2010; **34**: 874–883.
- O’Leary NA, Gross PS. Genomic structure and transcriptional regulation of the penaeidin gene family from *Litopenaeus vannamei*. *Gene* 2006; **371**: 75–83.
- Huang XD, Zhao L, Zhang HQ, Xu XP, Jia XT, Chen YH *et al*. Shrimp NF-κB binds to the immediate-early gene ie1 promoter of white spot syndrome virus and upregulates its activity. *Virology* 2010; **406**: 176–180.
- Wang PH, Gu ZH, Wan DH, Zhu WB, Qiu W, Weng SP *et al*. *Litopenaeus vannamei* sterile-α and armadillo motif containing protein (LvSARM) is involved in regulation of penaeidins and antilipopolysaccharide factors. *PLoS ONE* 2013; **8**: e52088.
- Vallabhapurapu S, Karin M. Regulation and function of NF-κB transcription factors in the immune system. *Annu Rev Immunol* 2009; **27**: 693–733.
- Zhi H, Yang L, Kuo YL, Ho YK, Shih HM, Giam CZ. NF-κB hyperactivation by HTLV-1 tax induces cellular senescence, but can be alleviated by the viral anti-sense protein HBZ. *PLoS Pathog* 2011; **7**: e1002025.
- Wang PH, Gu ZH, Wan DH, Zhang MY, Weng SP, Yu XQ *et al*. *Litopenaeus vannamei* Toll-interacting protein (LvTollip) is a

- potential negative regulator of the shrimp Toll pathway involved in the regulation of the shrimp antimicrobial peptide gene penaeidin-4 (PEN4). *Dev Comp Immunol* 2013; **40**: 266–277.
- 36 Kadalayil L, Petersen UM, Engstrom Y. Adjacent GATA and kappa B-like motifs regulate the expression of a *Drosophila* immune gene. *Nucleic Acids Res* 1997; **25**: 1233–1239.
- 37 Valanne S, Wang JH, Ramet M. The *Drosophila* Toll signaling pathway. *J Immunol* 2011; **186**: 649–656.
- 38 Hiscott J, Nguyen TL, Arguello M, Nakhaei P, Paz S. Manipulation of the nuclear factor-kappaB pathway and the innate immune response by viruses. *Oncogene* 2006; **25**: 6844–6867.
- 39 Li F, Yan H, Wang D, Priya TA, Li S, Wang B *et al*. Identification of a novel relish homolog in Chinese shrimp *Fenneropenaeus chinensis* and its function in regulating the transcription of antimicrobial peptides. *Dev Comp Immunol* 2009; **33**: 1093–1101.
- 40 Zhang S, Li CZ, Yan H, Qiu W, Chen YG, Wang PH *et al*. Identification and function of myeloid differentiation factor 88 (MyD88) in *Litopenaeus vannamei*. *PLoS ONE* 2012; **7**: e47038.
- 41 Li C, Chen YX, Zhang S, Lü L, Chen YH, Chai J *et al*. Identification, characterization, and function analysis of the Cactus gene from *Litopenaeus vannamei*. *PLoS ONE* 2012; **7**: e49711.
- 42 Santoro MG, Rossi A, Amici C. NF-kappaB and virus infection: who controls whom. *EMBO J* 2003; **22**: 2552–2560.
- 43 Liu WJ, Chang YS, Wang CH, Kou GH, Lo CF. Microarray and RT-PCR screening for white spot syndrome virus immediate-early genes in cycloheximide-treated shrimp. *Virology* 2005; **334**: 327–341.
- 44 Li F, Li M, Ke W, Ji Y, Bian X, Yan X. Identification of the immediate-early genes of white spot syndrome virus. *Virology* 2009; **385**: 267–274.
- 45 Liu WJ, Chang YS, Wang HC, Leu JH, Kou GH, Lo CF. Transactivation, dimerization, and DNA-binding activity of white spot syndrome virus immediate-early protein IE1. *J Virol* 2008; **82**: 11362–11373.
- 46 Liu WJ, Chang YS, Wang AH, Kou GH, Lo CF. White spot syndrome virus annexes a shrimp STAT to enhance expression of the immediate-early gene ie1. *J Virol* 2007; **81**: 1461–1471.



This work is licensed under a Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>