

RESEARCH ARTICLE

# Combined effect of genetic polymorphisms of *AURKA* and environmental factors on oral cancer development in Taiwan

Chia-Hsuan Chou<sup>1,2</sup>, Ying-Erh Chou<sup>2,3</sup>, Chun-Yi Chuang<sup>3,4</sup>, Shun-Fa Yang<sup>1,2</sup>, Chiao-Wen Lin<sup>5,6\*</sup>

**1** Institute of Medicine, Chung Shan Medical University, Taichung, Taiwan, **2** Department of Medical Research, Chung Shan Medical University Hospital, Taichung, Taiwan, **3** School of Medicine, Chung Shan Medical University, Taichung, Taiwan, **4** Department of Otolaryngology, Chung Shan Medical University Hospital, Taichung, Taiwan, **5** Institute of Oral Sciences, Chung Shan Medical University, Taichung, Taiwan, **6** Department of Dentistry, Chung Shan Medical University Hospital, Taichung, Taiwan

\* [cwlin@csmu.edu.tw](mailto:cwlin@csmu.edu.tw)



## Abstract

### OPEN ACCESS

**Citation:** Chou C-H, Chou Y-E, Chuang C-Y, Yang S-F, Lin C-W (2017) Combined effect of genetic polymorphisms of *AURKA* and environmental factors on oral cancer development in Taiwan. PLoS ONE 12(2): e0171583. doi:10.1371/journal.pone.0171583

**Editor:** Masaru Kato, National Cancer Center, JAPAN

**Received:** November 9, 2016

**Accepted:** January 23, 2017

**Published:** February 2, 2017

**Copyright:** © 2017 Chou et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** This study was supported by CSH-2016-E-002-Y2 from Chung Shan Medical University Hospital Taiwan, respectively. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

## Background

Oral squamous cell carcinoma (OSCC) is the sixth and fourth most common cause of cancer death in men worldwide and in Taiwan, respectively. *AURKA*, which encodes a centrosome-related serine/threonine kinase, is frequently amplified and overexpressed in many human cancers, particularly advanced OSCC. We conducted a hospital-based case-control study to estimate *AURKA* single-nucleotide polymorphisms (SNPs) and environmental risk factors to determine OSCC susceptibility and clinicopathological characteristics.

## Methodology/Principal findings

We enrolled a total of 876 OSCC patients and 1200 controls. Four SNPs of *AURKA*, namely rs1047972, rs2273535, rs2064863, and rs6024836, were analyzed using real-time polymerase chain reaction (PCR). Among the 1420 smokers, the *AURKA* polymorphism carriers with the betel nut chewing habit had a higher risk of oral cancer than *AURKA* wild-type (WT) carriers without the betel nut chewing habit. Patients with the *AURKA* rs2064863 gene had a 1.365-fold higher risk of stage III or IV OSCC (95% confidence interval [CI] 1.029–1.811) than those with the rs2064863 WT gene. Furthermore, carriers of the *AURKA* rs1047972/rs2273535/rs2064863 C-A-T haplotype had a 1.736-fold (95% CI 1.110–2.715) higher risk of OSCC than controls (C-T-T, the most common haplotype). Among patients with the betel quid chewing habit, carriers of other haplotypes (C-T-T, C-A-G, T-A-T, T-A-G, T-T-T, and C-T-G) had a 12.857-fold (95% CI 10.731–15.404) increased risk, and carriers of the C-A-T haplotype had the highest risk (AOR: 31.120; 95% CI 13.864–69.850) of OSCC, compared with those without the betel quid chewing who harbored other haplotypes.

## Conclusions

In conclusion, betel nut chewing combined with the *AURKA* C-A-T haplotypes lead to a high risk of OSCC. These findings reveal a novel genetic-environmental predisposition for oral tumorigenesis.

## Introduction

More than 90% of all head and neck malignant tumors occur in oral squamous cell carcinoma (OSCC) patients [1]. OSCC is the sixth and fourth most common cause of cancer death in men worldwide and in Taiwan, respectively [2]. Patients usually seek treatment only at the advanced stage of OSCC, resulting in a relatively low 5-year survival rate [3]. Both genetic factors and carcinogen-exposure behaviors (for example: betel nut chewing, alcohol consumption, and tobacco) regulate OSCC development [4, 5]. Moreover, our previous studies have demonstrated that genetic polymorphism combined with betel nut carcinogens may increase susceptibility to OSCC [6–12]. The results illustrate the importance of single-nucleotide polymorphisms (SNPs) for predicting risk or prognosis of OSCC.

*AURKA*, also known as *Aurora Kinase A*, encodes a centrosome-related serine/threonine kinase and is frequently amplified and overexpressed in many human cancers [13–15], particularly advanced OSCC [16]. Moreover, this gene has been identified as a definite low-penetrance tumor susceptibility gene [17]. The high expression of *AURKA* might be induced by centrosome amplification, aberrant chromosome segregation, aneuploidy, and malignant transformation [18–20], thus mediating the molecular mechanisms underlying carcinogenesis. The genetic associations of *AURKA* with several conditions have been documented. Lee et al demonstrated that the AA genotype of *AURKA* rs2273535 T>A was associated with an increased risk of oral cancer [21]. Dai et al reported that Caucasians harboring *AURKA* rs1047972 T>C had a reduced breast cancer risk [22]. However, few genetic variants of *AURKA* have been associated with OSCC.

In this case-control study, we investigated the relationship of four *AURKA* polymorphisms—namely rs1047972, rs2273535, rs2064863, and rs6024836—with OSCC susceptibility in Taiwanese male patients with OSCC.

## Results

### Patient characteristics and distribution of oral cancer

The distributions of the demographic characteristics of the study subjects are summarized in Table 1. A total of 876 male patients with oral cancer and 1200 male controls were included in this study. The mean age  $\pm$  SD in the controls and patients was  $53.90 \pm 10.02$  and  $54.80 \pm 11.03$  years, respectively. A significant difference was observed in the prevalence of betel nut chewing, cigarette smoking, and alcohol drinking between oral cancer patients and controls.

### Associations between *AURKA* SNP and oral cancer

In the control group, the genotypic frequencies of *AURKA* SNP rs1047972 C/T, rs2273535 A/T, rs2064863, and rs6024836 A/G were in Hardy-Weinberg equilibrium ( $p > 0.05$ ). The genotypic and allelic frequencies of *AURKA* SNPs in oral cancer patients and controls are shown in Table 2. After adjustment for age, betel quid chewing, cigarette smoking, and alcohol drinking, no significant difference was observed between oral cancer patients and controls.

**Table 1. The distributions of demographical characteristics in 1200 controls and 876 male patients with oral cancer.**

Variable	Controls (N = 1200)	Patients (N = 876)	p value
Age (yrs)	53.90 ± 10.02	54.80 ± 11.03	p = 0.06
Betel quid chewing			
No	1001 (83.4%)	175 (20.0%)	
Yes	199 (16.6%)	701 (80.0%)	p <0.01*
Cigarette smoking			
No	564 (47.0%)	92 (10.5%)	
Yes	636 (53.0%)	784 (89.5%)	p <0.01*
Alcohol drinking			
No	963 (80.3%)	388 (44.3%)	
Yes	237 (19.8%)	488 (55.7%)	p <0.01*
Stage			
I+II		435 (49.7%)	
III+IV		441 (50.3%)	
Tumor T status			
T1+T2		512 (58.4%)	
T3+T4		364 (41.6%)	
Lymph node status			
N0		597 (68.2%)	
N1+N2+N3		279 (31.8%)	
Metastasis			
M0		868 (99.1%)	
M1		8 (0.9%)	
Cell differentiation			
Well differentiated		141 (16.1%)	
Moderately or poorly differentiated		735 (83.9%)	

Mann-Whitney U test or Fisher's exact test was used between healthy controls and patients with oral cancer.

\* p value < 0.05 as statistically significant.

doi:10.1371/journal.pone.0171583.t001

In [Table 3](#), among the 1420 smokers, the subjects with *AURKA* rs1047972, rs2273535, rs2064863, and rs6024836 polymorphisms who exhibited the betel nut chewing habit respectively had 10.589-fold (95% confidence interval [CI] 6.994–16.032), 12.663-fold (95% CI 8.633–18.575), 17.912-fold (95% CI 6.596–48.643), and 13.912-fold (95% CI 9.392–20.607) significantly higher risks of OSCC than did smokers with wild-type genes without the betel nut chewing habit.

### Associations between *AURKA* SNPs and the clinicopathologic status of OSCC

We further clarified the role of *AURKA* polymorphisms in the clinicopathologic status of OSCC, such as the tumor clinical stage, tumor size, lymph node metastasis, and cell differentiation. Among the 876 oral cancer patients, only patients with the *AURKA* rs2064863 gene had a 1.365-fold higher risk of stage III or IV OSCC (95% CI 1.029–1.811) than did patients with the rs2064863 wild-type gene (p = 0.031). However, no significant difference was observed in the tumor size, lymph node metastasis clinical stage, lymph node metastasis, or cell differentiation ([Table 4](#)).

**Table 2. Genotyping and allele frequency of *AURKA* single nucleotide polymorphism in oral cancer and normal controls.**

Variable	Controls N = 1200(%)	Patients N = 876 (%)	AOR (95% CI)	p value
<b>rs1047972</b>				
CC	925 (77.1%)	668 (76.3%)	1.00	
TC	261 (21.8%)	190 (21.7%)	0.93 (0.70–1.23)	p = 0.60
TT	14 (1.1%)	18 (2.0%)	1.95 (0.74–5.13)	p = 0.18
TC+TT	275 (22.9%)	208 (23.7%)	0.97 (0.74–1.27)	p = 0.84
C allele	2111 (88.0%)	1526 (87.1%)	1.00	
T allele	289 (12.0%)	226 (12.9%)	1.02 (0.80–1.31)	p = 0.87
<b>rs2273535</b>				
TT	583 (48.6%)	404 (46.1%)	1.00	
AT	490 (40.8%)	368 (42.0%)	1.16 (0.91–1.48)	p = 0.25
AA	127 (10.6%)	104 (11.9%)	1.03 (0.70–1.51)	p = 0.88
AT+AA	617 (51.4%)	472 (53.9%)	1.13 (0.90–1.42)	p = 0.31
T allele	1656 (69.0%)	1176 (67.1%)	1.00	
A allele	774 (31.0%)	576 (32.9%)	1.06 (0.89–1.26)	p = 0.50
<b>rs2064863</b>				
TT	827 (68.9%)	590 (67.4%)	1.00	
GT	338 (28.2%)	252 (28.8%)	1.01 (0.79–1.31)	p = 0.92
GG	35 (2.9%)	34 (3.8%)	1.21 (0.64–2.30)	p = 0.55
GT+GG	373 (31.1%)	286 (32.6%)	1.03 (0.81–1.32)	p = 0.80
T allele	1992 (83.0%)	1432 (81.7%)	1.00	
G allele	408 (17.0%)	320 (18.3%)	1.05(0.85–1.29)	p = 0.67
<b>rs6024836</b>				
AA	525 (43.8%)	352 (40.2%)	1.00	
AG	530 (44.2%)	397 (45.3%)	1.15 (0.90–1.47)	p = 0.26
GG	145 (12.0%)	127 (14.5%)	1.19(0.83–1.70)	p = 0.36
AG+GG	675 (56.2%)	524 (59.8%)	0.86 (0.68–1.09)	p = 0.22
A allele	1580 (65.8%)	1101 (62.8%)	1.00	
G allele	820 (34.2%)	651 (37.2%)	1.11 (0.94–1.31)	p = 0.24

doi:10.1371/journal.pone.0171583.t002

### Haplotype analysis of the *AURKA* gene

We used Haploview software and the PHASE program to calculate pairwise linkage disequilibrium (LD) and analyzed the common haplotypes. As shown in Table 5, the p value for the global test of five haplotypes was 0.002 for OSCC development. The most common haplotype was C-T-T (68.4%) in the control group; thus, this haplotype was used as the haplotype reference. Compared with the reference group, carriers with C-A-T or other haplotypes had 1.736-fold (95% CI 1.110–2.715) and 2.788-fold (95% CI 1.387–5.605) significantly increased risks of OSCC (Table 5). Furthermore, we estimated the combined effect of betel quid chewing and *AURKA* haplotypes on OSCC development (Table 6). Patients with other haplotypes (C-T-T, C-A-G, T-A-T, T-A-G, T-T-T, and C-T-G) without the betel quid chewing habit were selected as a reference group. After adjustment for the effects of age, cigarette smoking, and alcohol drinking, patients with the betel quid chewing habit and other haplotypes had a 12.857-fold (95% CI 10.731–15.404) ( $p < 0.001$ ) increased risk. Moreover, those with the betel quid chewing habit and the C-A-T haplotype had the highest risk of OSCC (AOR 31.120; 95% CI 13.864–69.850;  $p < 0.001$ ).

**Table 3. Associations of the combined effect of *AURKA* gene polymorphisms and betel nut chewing with the susceptibility to oral cancer among 1420 smokers.**

Variable	Controls (n = 636) (%)	Patients (n = 784) (%)	OR (95% CI)	AOR (95% CI)
<b>rs1047972</b>				
CC genotype & non-betel nut chewing	348 (54.7%)	93 (11.9%)	1.00	1.00
TC or TT genotype or betel nut chewing	244 (38.4%)	535 (68.2%)	<b>8.21 (5.87–11.47)</b>	<b>6.80 (4.82–9.59)</b>
TC or TT genotype with betel nut chewing	44 (6.9%)	156 (19.9%)	<b>13.27 (8.09–21.77)</b>	<b>10.59 (6.38–17.57)</b>
<b>rs2273535</b>				
TT genotype & non-betel nut chewing	226 (35.5%)	53 (6.8%)	1.00	1.00
AT or AA genotype or betel nut chewing	313 (49.2%)	376 (48.0%)	<b>5.12 (3.40–7.71)</b>	<b>4.28 (2.81–6.51)</b>
AT or AA genotype with betel nut chewing	97 (15.3%)	355 (45.2%)	<b>15.61 (9.88–24.64)</b>	<b>12.66 (7.93–20.22)</b>
<b>rs2064863</b>				
TT genotype & non-betel nut chewing	318 (50.0%)	80 (10.2%)	1.00	1.00
GT or GG genotype or betel nut chewing	313 (49.2%)	676 (86.2%)	<b>8.59 (6.10–12.08)</b>	<b>7.25 (5.11–10.29)</b>
GT or GG genotype with betel nut chewing	5 (0.8%)	28 (3.6%)	<b>22.25 (6.70–73.89)</b>	<b>17.91 (5.29–60.69)</b>
<b>rs6024836</b>				
AA genotype & non-betel nut chewing	213 (33.5%)	46 (5.9%)	1.00	1.00
AG or GG genotype or betel nut chewing	314 (49.4%)	342 (43.6%)	<b>5.04 (3.27–7.77)</b>	<b>4.21 (2.70–6.58)</b>
AG or GG genotype with betel nut chewing	109 (17.1%)	396 (50.5%)	<b>16.82 (10.54–26.85)</b>	<b>13.91 (8.61–22.48)</b>

The adjusted odds ratios (AORs) with their 95% confidence intervals (CIs) were estimated by multiple logistic regression models after controlling for age and alcohol drinking.

doi:10.1371/journal.pone.0171583.t003

## Discussion

In this study, we observed that the combination of the *AURKA* gene polymorphisms with betel nut chewing increased OSCC susceptibility. Furthermore, we observed an interaction between the clinicopathological statuses and *AURKA* rs2064863 polymorphism. The results of pairwise allele analysis for rs1047972/rs2273535/rs2064863 revealed that the C-A-T haplotype is associated with the risk of OSCC.

**Table 4. Effect of *AURKA* rs2064863 polymorphism on clinical statuses in 786 male oral cancer.**

Variable	<i>AURKA</i> rs2064863		OR (95% CI)	AOR (95% CI) <sup>a</sup>
	TT (n = 590) n (%)	GT+GG (n = 286) n (%)		
<b>Clinical Stage</b>				
Stage I/II	278 (47.1%)	157 (54.9%)	1.00	1.00
Stage III/IV	312 (52.9%)	129 (45.1%)	<b>1.37 (1.04–1.82)<sup>b</sup></b>	<b>1.37 (1.03–1.81)<sup>c</sup></b>
<b>Tumor size</b>				
≤T2	333 (56.4%)	179 (62.6%)	1.00	1.00
> T2	257 (43.6%)	107 (37.4%)	0.78 (0.58–1.04)	0.78 (0.58–1.04)
<b>Lymph node metastasis</b>				
No	392 (66.4%)	205 (71.7%)	1.00	1.00
Yes	198 (33.6%)	81 (28.3%)	0.77 (0.57–1.05)	0.78 (0.57–1.06)
<b>Cell differentiation</b>				
well	97 (16.4%)	44 (15.4%)	1.00	1.00
Moderate/poor	493 (83.6%)	242 (84.6%)	2.70 (0.709–10.30)	2.68 (0.70–10.24)

<sup>a</sup> Adjusting for the effects of age, betel quid chewing, cigarette smoking and alcohol drinking

<sup>b</sup> p = 0.03

<sup>c</sup> p = 0.03

doi:10.1371/journal.pone.0171583.t004

**Table 5. Frequencies of *AURKA* haplotypes in OSCC patients and control subjects.**

Haplotype block			Controls n = 2400	Patients n = 1752	OR (95% CI)	AOR (95% CI) <sup>a</sup>
rs1047972 T/C	rs2273535 A/T	rs2064863 T/G				
Global haplotype test (p = 0.0018)						
C	T	T	1641 (68.4%)	1170 (66.8%)	1.00	1.00
C	A	G	399 (16.6%)	282 (16.1%)	0.99 (0.84–1.18)	0.95 (0.76–1.18)
T	A	T	269 (11.2%)	190 (10.8%)	0.99 (0.81–1.21)	0.92 (0.71–1.20)
C	A	T	69 (2.9%)	70 (4.0%)	1.42 (1.01–2.00) <sup>c</sup>	1.74 (1.11–2.72) <sup>e</sup>
Others <sup>b</sup>			22 (0.9%)	40 (2.2%)	2.55 (1.51–4.31) <sup>d</sup>	2.79 (1.39–5.61) <sup>f</sup>

<sup>a</sup> Adjusting for the effects of age, betel quid chewing, cigarette smoking and alcohol drinking

<sup>b</sup> Others: TAG (41), TTT (15), and CTG (6)

<sup>c</sup> p = 0.04

<sup>d</sup> p < 0.01

<sup>e</sup> p = 0.02

<sup>f</sup> p < 0.01

doi:10.1371/journal.pone.0171583.t005

*AURKA* overexpression has been demonstrated in various human cancers, including breast cancer, head and neck squamous cell carcinoma, colorectal cancer, ovarian cancer, and advanced OSCC [13–16, 23]. The abnormal expression of *AURKA* might lead to high chromosome instability in tumors and further increase susceptibility to malignant transformation [24]. These processes may result from the gaining of the chromosome 20q amplicon, which promotes the progression of adenoma to carcinoma [25]. Activation of the Wnt/ $\beta$ -catenin and PI3K/Akt signaling pathways plays a crucial role in cancer development [26]. Moreover, *AURKA* overexpression promotes cell proliferation and tumor progression and metastasis [27–29]. Furthermore, the blocking of *AURKA* induces autophagy and apoptosis in mice and significantly increases sensitivity to chemical treatment in OSCC [16, 30]. Taken together, these findings indicate that *AURKA* knockdown might be a valuable therapeutic strategy for OSCC.

A previous study demonstrated that the *AURKA* 91A (rs2273535) allele polymorphism is associated with a high risk of oral cancer [21]. Previous studies have also suggested that the *AURKA* rs2273535 polymorphism is associated with a high risk of breast cancer, particularly in Asians [31]. However, the *AURKA* rs1047972 polymorphism has been shown to reduce the incidence of breast cancer in Caucasians [22]. In this study, our data showed that *AURKA* polymorphisms were not associated with the risk of OSCC (Table 2). However, the significant interaction between the investigated *AURKA* polymorphisms and betel nut chewing was

**Table 6. Combined effect of betel quid chewing and *AURKA* haplotypes on OSCC development.**

Betel quid chewing	<i>AURKA</i> haplotype	Controls n = 2400	Patients n = 1752	AOR (95% CI) <sup>b</sup>
Yes	C-A-T	7 (0.3%)	55 (3.1%)	31.09 (13.86–69.79) <sup>c</sup>
Yes	Others <sup>a</sup>	391 (16.3%)	1347 (76.9%)	12.81 (10.70–15.35) <sup>c</sup>
No	C-A-T	62 (2.6%)	15 (0.9%)	1.44 (0.82–2.67)
No	Others <sup>a</sup>	1940 (80.8%)	335 (19.1%)	1.00

<sup>a</sup> Other haplotypes included C-T-T, C-A-G, T-A-T, T-A-G, T-T-T, and C-T-G

<sup>b</sup> Adjusting for the effects of age, cigarette smoking and alcohol drinking

<sup>c</sup> p < 0.01

doi:10.1371/journal.pone.0171583.t006

associated with a high incidence of OSCC in smokers. These results suggest that environmental risk factors such as betel nut chewing and tobacco smoking influence *AURKA* gene expression, subsequently leading to a high incidence of OSCC.

The present study demonstrated that the *AURKA* rs2064863 polymorphism was associated with a high risk of stage III/IV OSCC but not with tumor size, metastasis to the lymph node and distant organs, or cell differentiation (Table 4). Chuang et al reported that *AURKA* gene overexpression was strongly associated with the progression of colorectal adenoma to colorectal cancer. *AURKA* overexpression may cause high chromosome instability in tumors and further increase susceptibility to malignant transformation [24]. Sillard-Hardebol et al also observed the same results [25]. However, the possibility of an association among the advanced stage, *AURKA* expression level, and *AURKA* genotype, as well as the effects of the *AURKA* genotype on oral cancer risk, require further investigation.

A study found that the *AURKA* rs2273535 polymorphism increases the risk of breast cancer, and that the rs1047972 polymorphism is a protective factor for breast cancer [22]. A study conducted in Korea revealed that the *AURKA* rs2273535 polymorphism was in strong LD with the rs1047972 genotype, and that patients with the *AURKA* haplotype variants had high kinase activity and a high risk of progression to advanced stage gastric cancer [32]. In the present study, we observed that subjects with the rs1047972/rs2273535/rs2064863 C-A-T haplotype had a higher risk of OSCC than those with the common C-T-T haplotype. These results indicate that the *AURKA* gene variants might have different functional roles in different cancers.

The incidence of OSCC is particularly high in Taiwan because of the popularity of betel nut chewing. Approximately 2.5 million people in Taiwan have the betel nut chewing habit [33]. Several studies have indicated that betel quid may damage the oral mucosa, inducing genotoxic or nongenotoxic effects, which are associated with the initiation, promotion, and progression of OSCC [34–39]. In our study, betel nut chewing was related to OSCC development. Therefore, we estimated the combined effect of betel nut chewing and *AURKA* haplotypes on OSCC development. Among patients without the betel nut chewing habit, those with the *AURKA* rs1047972/rs2273535/rs2064863 C-A-T haplotype had a 1.443-fold (95% CI 0.801–2.601) higher risk of OSCC than those with other haplotypes, but the risk was nonsignificant. Moreover, we observed that betel nut users with other *AURKA* haplotypes had an increased risk of OSCC. Thus, the increased risk might result from betel nut chewing. Betel nut users with the *AURKA* rs1047972/rs2273535/rs2064863 C-A-T haplotype had a higher risk of OSCC. Previous studies have demonstrated that *AURKA* 91A (Ile31) is a low-penetrance tumor susceptibility allele, and individuals homozygous or heterozygous for this allele tend to exhibit an increased risk of several human cancers [40–45]. Matarasso et al confirmed that the expression level of *AURKA* mRNA was higher in Ile31/Ile31-homozygous (2.07-fold) and Phe31/Ile31-heterozygous (1.93-fold) samples from normal prostate tissues than in Phe31/Phe31-homozygous samples [46]. In addition, the *AURKA* rs1047972 polymorphism, which results in a valine-to-isoleucine substitution, has been investigated for its association with breast cancer, and this polymorphism has been found to be a protective factor for breast cancer in Caucasians but not in Asians [22, 47]. The effect of this SNP on OSCC has not been clarified. Betel nut chewing may induce OSCC, and betel quid components can stimulate the activation of Src and ERK for promoting the migration and motility of cancer cells [48]. Thus, betel nut chewing might alter *AURKA* expression, thereby impairing cell growth, differentiation, and inflammation.

Our study has some limitations. First, our data were collected from two medical centers and may have referral bias; future studies can therefore collect data from different regions in Taiwan to verify our present data. Second, the questionnaire on betel nut, tobacco, and alcohol use described users as “ever users” and “never users.” Data on the amount, length, and

history of betel nut, tobacco, and alcohol use could not be collected to conduct a comprehensive analysis.

In conclusion, betel nut users with the *AURKA* C-A-T haplotypes have a high risk of OSCC. In addition, the SNP rs2064863 is associated with a high risk of developing advanced-stage tumors. These findings reveal a novel genetic-environmental predisposition for oral tumorigenesis.

## Materials and methods

### Study subjects and specimen collection

In this study, we enrolled 876 male patients with OSCC from Chung Shan Medical University Hospital in Taichung, Taiwan from 2007 to 2015. In the 876 patients, the tumors were located in the buccal mucosa, tongue, gingiva, palate, floor of the mouth, and other sites. For the control group, we selected 1200 healthy male individuals with no self-reported history of cancer at any site from the Taiwan Biobank. Each subject completed a questionnaire on their demographic characteristics, betel nut chewing, tobacco use, and alcohol consumption, and medical history. Whole blood specimens collected from OSCC patients and controls were placed in tubes containing ethylenediamine tetra-acetic acid (EDTA), immediately centrifuged, and stored at  $-80^{\circ}\text{C}$ . All participants were provided with a written description of the study. The Institutional Review Board of Chung Shan Medical University Hospital approved this study (CSMUH No: CS13214-1), and informed written consent was obtained from each participant.

### Determination of genotypes

Genomic DNA was isolated from peripheral blood using QIAamp DNA blood mini kits (Qiagen, Valencia, CA, USA) following manufacturer instructions. DNA was dissolved in TE buffer (10 mM Tris and 1 mM EDTA acid; pH 7.8) and quantified by measuring the optical density at 260 nm. The final preparation was stored at  $-20^{\circ}\text{C}$  and used as templates for polymerase chain reaction (PCR). Allelic discrimination of the rs1047972, rs2273535, rs2064836, and rs6024836 *AURKA* polymorphisms was performed using TaqMan SNP genotyping assays with the ABI StepOne™ Real-Time PCR System (Applied Biosystems, Foster City, CA, USA), and these polymorphisms were further analyzed using SDS version 3.0 (Applied Biosystems).

### Statistical analysis

Hardy-Weinberg equilibrium was assessed using a chi-square goodness-of-fit test for bi-allelic markers. The Mann-Whitney U test and Fisher's exact test were used to compare differences in the distribution of age and demographic characteristics between the controls and OSCC patients. ORs with 95% CIs were estimated using logistic regression models. AORs with 95% CIs were used to assess association between genotype frequencies with OSCC risk and clinical factors. Moreover, Bonferroni correction is used to adjust for multiple comparisons. *p* values less than 0.05 were considered significant. The data were analysed with SPSS 12.0 statistical software (SPSS Inc., Chicago, IL, USA).

## Acknowledgments

This study was supported by CSH-2016-E-002-Y2 from Chung Shan Medical University Hospital Taiwan, respectively. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.



## Author contributions

**Conceptualization:** CWL.

**Data curation:** CHC.

**Formal analysis:** YEC.

**Funding acquisition:** SFY.

**Investigation:** CWL.

**Methodology:** CYC.

**Resources:** CYC.

**Validation:** CWL.

**Writing – original draft:** CHC CWL.

**Writing – review & editing:** CHC CWL.

## References

1. Liao KA, Tsay YG, Huang LC, Huang HY, Li CF, Wu TF. Search for the tumor-associated proteins of oral squamous cell carcinoma collected in Taiwan using proteomics strategy. *Journal of proteome research*. 2011; 10(5):2347–58. doi: [10.1021/pr101146w](https://doi.org/10.1021/pr101146w) PMID: [21322651](https://pubmed.ncbi.nlm.nih.gov/21322651/)
2. Chien MH, Liu YF, Hsin CH, Lin CH, Shih CH, Yang SF, et al. Impact of VEGF-C gene polymorphisms and environmental factors on oral cancer susceptibility in Taiwan. *PloS one*. 2013; 8(4):e60283. doi: [10.1371/journal.pone.0060283](https://doi.org/10.1371/journal.pone.0060283) PMID: [23593187](https://pubmed.ncbi.nlm.nih.gov/23593187/)
3. Hamad HM. Cancer initiatives in Sudan. *Annals of oncology: official journal of the European Society for Medical Oncology / ESMO*. 2006; 17 Suppl 8:viii32–viii6.
4. Shastry BS. SNP alleles in human disease and evolution. *Journal of human genetics*. 2002; 47(11):561–6. doi: [10.1007/s100380200086](https://doi.org/10.1007/s100380200086) PMID: [12436191](https://pubmed.ncbi.nlm.nih.gov/12436191/)
5. Zheng SL, Sun J, Wiklund F, Smith S, Stattin P, Li G, et al. Cumulative association of five genetic variants with prostate cancer. *The New England journal of medicine*. 2008; 358(9):910–9. doi: [10.1056/NEJMoa075819](https://doi.org/10.1056/NEJMoa075819) PMID: [18199855](https://pubmed.ncbi.nlm.nih.gov/18199855/)
6. Cheng HL, Liu YF, Su CW, Su SC, Chen MK, Yang SF, et al. Functional genetic variant in the Kozak sequence of WW domain-containing oxidoreductase (WWOX) gene is associated with oral cancer risk. *Oncotarget*. 2016; 7(43):69384–96. Epub 2016/09/23. doi: [10.18632/oncotarget.12082](https://doi.org/10.18632/oncotarget.12082) PMID: [27655721](https://pubmed.ncbi.nlm.nih.gov/27655721/)
7. Chou YE, Hsieh MJ, Hsin CH, Chiang WL, Lai YC, Lee YH, et al. CD44 gene polymorphisms and environmental factors on oral cancer susceptibility in Taiwan. *PloS one*. 2014; 9(4):e93692. Epub 2014/04/05. doi: [10.1371/journal.pone.0093692](https://doi.org/10.1371/journal.pone.0093692) PMID: [24699672](https://pubmed.ncbi.nlm.nih.gov/24699672/)
8. Chu YH, Tzeng SL, Lin CW, Chien MH, Chen MK, Yang SF. Impacts of microRNA gene polymorphisms on the susceptibility of environmental factors leading to carcinogenesis in oral cancer. *PloS one*. 2012; 7(6):e39777. Epub 2012/07/05. doi: [10.1371/journal.pone.0039777](https://doi.org/10.1371/journal.pone.0039777) PMID: [22761899](https://pubmed.ncbi.nlm.nih.gov/22761899/)
9. Chung TT, Pan MS, Kuo CL, Wong RH, Lin CW, Chen MK, et al. Impact of RECK gene polymorphisms and environmental factors on oral cancer susceptibility and clinicopathologic characteristics in Taiwan. *Carcinogenesis*. 2011; 32(7):1063–8. Epub 2011/05/14. doi: [10.1093/carcin/bgr083](https://doi.org/10.1093/carcin/bgr083) PMID: [21565829](https://pubmed.ncbi.nlm.nih.gov/21565829/)
10. Lin CW, Yang SF, Chuang CY, Lin HP, Hsin CH. Association of matrix metalloproteinase-11 polymorphisms with susceptibility and clinicopathologic characteristics for oral squamous cell carcinoma. *Head & neck*. 2015; 37(10):1425–31. Epub 2014/05/20.
11. Su CW, Huang YW, Chen MK, Su SC, Yang SF, Lin CW. Polymorphisms and Plasma Levels of Tissue Inhibitor of Metalloproteinase-3: Impact on Genetic Susceptibility and Clinical Outcome of Oral Cancer. *Medicine*. 2015; 94(46):e2092. Epub 2015/11/19. doi: [10.1097/MD.0000000000002092](https://doi.org/10.1097/MD.0000000000002092) PMID: [26579821](https://pubmed.ncbi.nlm.nih.gov/26579821/)
12. Su SC, Hsieh MJ, Liu YF, Chou YE, Lin CW, Yang SF. ADAMTS14 Gene Polymorphism and Environmental Risk in the Development of Oral Cancer. *PloS one*. 2016; 11(7):e0159585. Epub 2016/07/28. doi: [10.1371/journal.pone.0159585](https://doi.org/10.1371/journal.pone.0159585) PMID: [27463966](https://pubmed.ncbi.nlm.nih.gov/27463966/)

13. Goos JA, Coupe VM, Diosdado B, Delis-Van Diemen PM, Karga C, Belien JA, et al. Aurora kinase A (*AURKA*) expression in colorectal cancer liver metastasis is associated with poor prognosis. *British journal of cancer*. 2013; 109(9):2445–52. doi: [10.1038/bjc.2013.608](https://doi.org/10.1038/bjc.2013.608) PMID: [24104968](https://pubmed.ncbi.nlm.nih.gov/24104968/)
14. Wang C, Yan Q, Hu M, Qin D, Feng Z. Effect of *AURKA* Gene Expression Knockdown on Angiogenesis and Tumorigenesis of Human Ovarian Cancer Cell Lines. *Targeted oncology*. 2016.
15. Treekitkarmongkol W, Katayama H, Kai K, Sasai K, Jones JC, Wang J, et al. Aurora kinase-A overexpression in mouse mammary epithelium induces mammary adenocarcinomas harboring genetic alterations shared with human breast cancer. *Carcinogenesis*. 2016.
16. Li Y, Zhang J. *AURKA* is a predictor of chemotherapy response and prognosis for patients with advanced oral squamous cell carcinoma. *Tumour biology: the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2015; 36(5):3557–64.
17. Ewart-Toland A, Briassouli P, de Koning JP, Mao JH, Yuan J, Chan F, et al. Identification of *Stk6/STK15* as a candidate low-penetrance tumor-susceptibility gene in mouse and human. *Nature genetics*. 2003; 34(4):403–12. doi: [10.1038/ng1220](https://doi.org/10.1038/ng1220) PMID: [12881723](https://pubmed.ncbi.nlm.nih.gov/12881723/)
18. Sen S, Zhou H, White RA. A putative serine/threonine kinase encoding gene *BTAK* on chromosome 20q13 is amplified and overexpressed in human breast cancer cell lines. *Oncogene*. 1997; 14(18):2195–200. doi: [10.1038/sj.onc.1201065](https://doi.org/10.1038/sj.onc.1201065) PMID: [9174055](https://pubmed.ncbi.nlm.nih.gov/9174055/)
19. Bischoff JR, Anderson L, Zhu Y, Mossie K, Ng L, Souza B, et al. A homologue of *Drosophila aurora* kinase is oncogenic and amplified in human colorectal cancers. *The EMBO journal*. 1998; 17(11):3052–65. doi: [10.1093/emboj/17.11.3052](https://doi.org/10.1093/emboj/17.11.3052) PMID: [9606188](https://pubmed.ncbi.nlm.nih.gov/9606188/)
20. Zhou H, Kuang J, Zhong L, Kuo WL, Gray JW, Sahin A, et al. Tumour amplified kinase *STK15/BTAK* induces centrosome amplification, aneuploidy and transformation. *Nature genetics*. 1998; 20(2):189–93. doi: [10.1038/2496](https://doi.org/10.1038/2496) PMID: [9771714](https://pubmed.ncbi.nlm.nih.gov/9771714/)
21. Lee CP, Chiang SL, Lee CH, Tsai YS, Wang ZH, Hua CH, et al. *AURKA* Phe31Ile polymorphism interacted with use of alcohol, betel quid, and cigarettes at multiplicative risk of oral cancer occurrence. *Clinical oral investigations*. 2015; 19(8):1825–32. doi: [10.1007/s00784-015-1432-5](https://doi.org/10.1007/s00784-015-1432-5) PMID: [25697104](https://pubmed.ncbi.nlm.nih.gov/25697104/)
22. Dai ZJ, Kang HF, Wang XJ, Shao YP, Lin S, Zhao Y, et al. Association between genetic polymorphisms in *AURKA* (rs2273535 and rs1047972) and breast cancer risk: a meta-analysis involving 37,221 subjects. *Cancer cell international*. 2014; 14(1):91. doi: [10.1186/s12935-014-0091-y](https://doi.org/10.1186/s12935-014-0091-y) PMID: [25253995](https://pubmed.ncbi.nlm.nih.gov/25253995/)
23. Pickhard A, Siegl M, Baumann A, Huhn M, Wirth M, Reiter R, et al. The response of head and neck squamous cell carcinoma to cetuximab treatment depends on Aurora kinase A polymorphism. *Oncotarget*. 2014; 5(14):5428–38. doi: [10.18632/oncotarget.2117](https://doi.org/10.18632/oncotarget.2117) PMID: [24980817](https://pubmed.ncbi.nlm.nih.gov/24980817/)
24. Chuang TP, Wang JY, Jao SW, Wu CC, Chen JH, Hsiao KH, et al. Over-expression of *AURKA*, *SKA3* and *DSN1* contributes to colorectal adenoma to carcinoma progression. *Oncotarget*. 2016.
25. Sillars-Hardebol AH, Carvalho B, Tijssen M, Belien JA, de Wit M, Delis-van Diemen PM, et al. *TPX2* and *AURKA* promote 20q amplicon-driven colorectal adenoma to carcinoma progression. *Gut*. 2012; 61(11):1568–75. doi: [10.1136/gutjnl-2011-301153](https://doi.org/10.1136/gutjnl-2011-301153) PMID: [22207630](https://pubmed.ncbi.nlm.nih.gov/22207630/)
26. Davidson KC, Adams AM, Goodson JM, McDonald CE, Potter JC, Berndt JD, et al. *Wnt/beta-catenin* signaling promotes differentiation, not self-renewal, of human embryonic stem cells and is repressed by *Oct4*. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109(12):4485–90. doi: [10.1073/pnas.1118777109](https://doi.org/10.1073/pnas.1118777109) PMID: [22392999](https://pubmed.ncbi.nlm.nih.gov/22392999/)
27. Dar AA, Belkhiri A, El-Rifai W. The aurora kinase A regulates *GSK-3beta* in gastric cancer cells. *Oncogene*. 2009; 28(6):866–75. doi: [10.1038/onc.2008.434](https://doi.org/10.1038/onc.2008.434) PMID: [19060929](https://pubmed.ncbi.nlm.nih.gov/19060929/)
28. Xu LZ, Long ZJ, Peng F, Liu Y, Xu J, Wang C, et al. Aurora kinase A suppresses metabolic stress-induced autophagic cell death by activating *mTOR* signaling in breast cancer cells. *Oncotarget*. 2014; 5(17):7498–511. doi: [10.18632/oncotarget.2241](https://doi.org/10.18632/oncotarget.2241) PMID: [25115395](https://pubmed.ncbi.nlm.nih.gov/25115395/)
29. Jia L, Lee HS, Wu CF, Kundu J, Park SG, Kim RN, et al. *SMAD4* suppresses *AURKA*-induced metastatic phenotypes via degradation of *AURKA* in a *TGFbeta*-independent manner. *Molecular cancer research: MCR*. 2014; 12(12):1779–95. doi: [10.1158/1541-7786.MCR-14-0191](https://doi.org/10.1158/1541-7786.MCR-14-0191) PMID: [25061104](https://pubmed.ncbi.nlm.nih.gov/25061104/)
30. Kao SY, Chen YP, Tu HF, Liu CJ, Yu AH, Wu CH, et al. Nuclear *STK15* expression is associated with aggressive behaviour of oral carcinoma cells in vivo and in vitro. *The Journal of pathology*. 2010; 222(1):99–109. doi: [10.1002/path.2737](https://doi.org/10.1002/path.2737) PMID: [20597069](https://pubmed.ncbi.nlm.nih.gov/20597069/)
31. Guo XG, Zheng L, Feng WB, Xia Y. The *AURKA* gene rs2273535 polymorphism contributes to breast carcinoma risk—meta-analysis of eleven studies. *Asian Pacific journal of cancer prevention: APJCP*. 2014; 15(16):6709–14. Epub 2014/08/30. PMID: [25169513](https://pubmed.ncbi.nlm.nih.gov/25169513/)
32. Ju H, Cho H, Kim YS, Kim WH, Ihm C, Noh SM, et al. Functional polymorphism 57Val>Ile of aurora kinase A associated with increased risk of gastric cancer progression. *Cancer letters*. 2006; 242(2):273–9. doi: [10.1016/j.canlet.2005.11.015](https://doi.org/10.1016/j.canlet.2005.11.015) PMID: [16412566](https://pubmed.ncbi.nlm.nih.gov/16412566/)

33. Kao SY, Lim E. An overview of detection and screening of oral cancer in Taiwan. *The Chinese journal of dental research: the official journal of the Scientific Section of the Chinese Stomatological Association (CSA)*. 2015; 18(1):7–12. Epub 2015/03/31.
34. Ko YC, Huang YL, Lee CH, Chen MJ, Lin LM, Tsai CC. Betel quid chewing, cigarette smoking and alcohol consumption related to oral cancer in Taiwan. *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*. 1995; 24(10):450–3.
35. Liu TY, Chen CL, Chi CW. Oxidative damage to DNA induced by areca nut extract. *Mutation research*. 1996; 367(1):25–31. PMID: [8596543](#)
36. Kuo MY, Jeng JH, Chiang CP, Hahn LJ. Mutations of Ki-ras oncogene codon 12 in betel quid chewing-related human oral squamous cell carcinoma in Taiwan. *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*. 1994; 23(2):70–4.
37. Jin YT, Tsai ST, Wong TY, Chen FF, Chen RM. Studies on promoting activity of Taiwan betel quid ingredients in hamster buccal pouch carcinogenesis. *European journal of cancer Part B, Oral oncology*. 1996; 32B(5):343–6. PMID: [8944839](#)
38. Wang CK, Peng CH. The mutagenicities of alkaloids and N-nitrosoguvacoline from betel quid. *Mutation research*. 1996; 360(3):165–71. PMID: [8692215](#)
39. Lee-Chen SF, Chen CL, Ho LY, Hsu PC, Chang JT, Sun CM, et al. Role of oxidative DNA damage in hydroxychavicol-induced genotoxicity. *Mutagenesis*. 1996; 11(5):519–23. PMID: [8921515](#)
40. Miao X, Sun T, Wang Y, Zhang X, Tan W, Lin D. Functional STK15 Phe31Ile polymorphism is associated with the occurrence and advanced disease status of esophageal squamous cell carcinoma. *Cancer research*. 2004; 64(8):2680–3. PMID: [15087379](#)
41. Sun T, Miao X, Wang J, Tan W, Zhou Y, Yu C, et al. Functional Phe31Ile polymorphism in Aurora A and risk of breast carcinoma. *Carcinogenesis*. 2004; 25(11):2225–30. doi: [10.1093/carcin/bgh244](#) PMID: [15271856](#)
42. Dicioccio RA, Song H, Waterfall C, Kimura MT, Nagase H, McGuire V, et al. STK15 polymorphisms and association with risk of invasive ovarian cancer. *Cancer epidemiology, biomarkers & prevention: a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2004; 13(10):1589–94.
43. Cox DG, Hankinson SE, Hunter DJ. Polymorphisms of the *AURKA* (STK15/Aurora Kinase) Gene and Breast Cancer Risk (United States). *Cancer causes & control: CCC*. 2006; 17(1):81–3.
44. Ewart-Toland A, Dai Q, Gao YT, Nagase H, Dunlop MG, Farrington SM, et al. Aurora-A/STK15 T+91A is a general low penetrance cancer susceptibility gene: a meta-analysis of multiple cancer types. *Carcinogenesis*. 2005; 26(8):1368–73. doi: [10.1093/carcin/bgi085](#) PMID: [15802297](#)
45. Hienonen T, Salovaara R, Mecklin JP, Jarvinen H, Karhu A, Aaltonen LA. Preferential amplification of *AURKA* 91A (Ile31) in familial colorectal cancers. *International journal of cancer*. 2006; 118(2):505–8. doi: [10.1002/ijc.21344](#) PMID: [16049965](#)
46. Matarasso N, Bar-Shira A, Rozovski U, Rosner S, Orr-Urtreger A. Functional analysis of the Aurora Kinase A Ile31 allelic variant in human prostate. *Neoplasia*. 2007; 9(9):707–15. PMID: [17898866](#)
47. Taylor NJ, Bensen JT, Poole C, Troester MA, Gammon MD, Luo J, et al. Genetic variation in cell cycle regulatory gene *AURKA* and association with intrinsic breast cancer subtype. *Molecular carcinogenesis*. 2015; 54(12):1668–77. doi: [10.1002/mc.22238](#) PMID: [25328151](#)
48. Chiu CC, Chen BH, Hour TC, Chiang WF, Wu YJ, Chen CY, et al. Betel quid extract promotes oral cancer cell migration by activating a muscarinic M4 receptor-mediated signaling cascade involving SFKs and ERK1/2. *Biochemical and biophysical research communications*. 2010; 399(1):60–5. doi: [10.1016/j.bbrc.2010.07.042](#) PMID: [20638369](#)