
ERRATUM Open Access

## Erratum to: Oncolytic virus efficiency inhibited growth of tumour cells with multiple drug resistant phenotype in vivo and in vitro

Elena P. Goncharova<sup>1</sup>, Julia S. Ruzhenkova<sup>1</sup>, Ivan S. Petrov<sup>1,3</sup>, Sergey N. Shchelkunov<sup>2</sup> and Marina A. Zenkova<sup>1\*</sup>

## Erratum to: J Transl Med (2016) 14:241 DOI 10.1186/s12967-016-1002-x

Unfortunately, the original version of this article [1] contained an error. Figures 2 and 7 were the incorrect versions. They have been corrected in the original article and are also included correctly in this erratum.




**Fig. 2** Development of LIVP-GFP infection in various tumour cells. Development of LIVP-GFP infection in RLS (*green circles*), RLS-40 (*red circles*), KB-3-1 (*violet squares*), KB-8-5 (*black line with open squares*) and melanoma B-16 (*blue triangles*) tumour cells at MOI of 1 (**a**) and MOI of 10 (**b**). Cells were incubated with virus for 1 h, washed with PBS and incubated up to the analysis in IMDM supplemented with 2 % FBS. Viral titre was measured by PFU assay. Data of three independent experiments are presented

<sup>&</sup>lt;sup>1</sup> Institute of Chemical Biology and Fundamental Medicine SB RAS, 8, Lavrentiev Ave., Novosibirsk 630090, Russian Federation Full list of author information is available at the end of the article



<sup>\*</sup>Correspondence: marzen@niboch.nsc.ru



**Fig. 7** Effect of LIVP-GFP treatment on the immune responses of tumour-bearing mice. **a** Summary data showing a significant increase of the number of IFN-γ-ransecreting splenocytes in LIVP-GFP-treated mice (n = 6) with RLS-40 or with melanoma B-16 tumours. Comparison of immune-related proteins GMC-SF (**b**) and IL-6 (**c**) in the blood serum of RLS-40 bearing mice during the treatment with LIVP-GFP (the experimental scheme was shown in Fig. 5a): red circles and blue squares for RLS-40 bearing mice treated with LIVP-GFP and PBS, respectively; black triangles healthy mice receiving PBS. The levels of cytokines in the blood serum were measured by ELISA. For each day, the value of mean  $\pm$  SEM is shown

## **Author details**

<sup>1</sup> Institute of Chemical Biology and Fundamental Medicine SB RAS, 8, Lavrentiev Ave., Novosibirsk 630090, Russian Federation. <sup>2</sup> Institute of Cytology and Genetics SB RAS, Novosibirsk, Russian Federation. <sup>3</sup> Department of Biochemistry, Biocenter, University of Wuerzburg, Am Hubland, 97074 Würzburg, Germany.

The online version of the original article can be found under doi:10.1186/s12967-016-1002-x.

Published online: 27 September 2016

## Reference

 Goncharova EP, Ruzhenkova JS, Petrov IS, Shchelkunov SN, Zenkova MA. J Transl Med. 2016;14:241. doi:10.1186/s12967-016-1002-x.