



Bisphenol A derivatives act as novel coactivator-binding inhibitors for estrogen receptor β

Received for publication, May 10, 2021, and in revised form, September 1, 2021 Published, Papers in Press, September 6, 2021,
<https://doi.org/10.1016/j.jbc.2021.101173>

Masaki Iwamoto^{1,‡}, Takahiro Masuya^{1,‡}, Mari Hosose¹, Koki Tagawa¹, Tomoka Ishibashi¹, Keitaro Suyama¹,
Takeru Nose¹, Eiji Yoshihara^{2,3,4}, Michael Downes², Ronald M. Evans², and Ayami Matsushima^{1,*} 

From the ¹Department of Chemistry, Faculty of Science, Kyushu University, Fukuoka, Japan; ²Gene Expression Laboratory, Salk Institute for Biological Studies, La Jolla, California, USA; ³Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, California, USA; and ⁴David Geffen School of Medicine at University of California, Los Angeles, Los Angeles, California, USA

Edited by Henrik Dohlman

Bisphenol A and its derivatives are recognized as endocrine disruptors based on their complex effects on estrogen receptor (ER) signaling. While the effects of bisphenol derivatives on ER α have been thoroughly evaluated, how these chemicals affect ER β signaling is less well understood. Herein, we sought to identify novel ER β ligands using a radioligand competitive binding assay to screen a chemical library of bisphenol derivatives. Many of the compounds identified showed intriguing dual activities as both ER α agonists and ER β antagonists. Docking simulations of these compounds and ER β suggested that they bound not only to the canonical binding site of ER β but also to the coactivator binding site located on the surface of the receptor, suggesting that they act as coactivator-binding inhibitors (CBIs). Receptor–ligand binding experiments using WT and mutated ER β support the presence of a second ligand–interaction position at the coactivator-binding site in ER β , and direct binding experiments of ER β and a coactivator peptide confirmed that these compounds act as CBIs. Our study is the first to propose that bisphenol derivatives act as CBIs, presenting critical insight for the future development of ER signaling–based drugs and their potential to function as endocrine disruptors.

Estrogen receptors (ERs) are members of the nuclear receptor family of transcription factors that directly bind to consensus nucleotide sequences to induce gene transcription. Forty-eight human nuclear receptors have been identified, including those for sex steroid hormones, glucocorticoids, retinoids, and vitamin D (1, 2), with many of these receptors recognized as therapeutic targets for a wide range of diseases (3). In particular, ERs are major drug targets for breast cancer (4) and menopausal disorders. Two ER isoforms exist, ER α and ER β , that have high amino acid similarity in both the DNA-binding domains and ligand-binding domains (LBDs) (5). Many ER α and/or ER β -associated gene disruption experiments have been reported (6). Female mice lacking ER α are infertile,

whereas male mice exhibit decreased fertility (7). Disruption of ER α in female mice leads to hypoplastic uteri, and ER α -disrupted female mice do not respond to estradiol treatments. ER β KO mice present with less-severe phenotypes than those with ER α KO, although ER β -disrupted female mice are subfertile predominantly because of reduced ovarian efficiency (8). Moreover, ER α and ER β double-KO mice show normal reproductive tract development during the prepubertal period. However, those animals present with similar features to ER α KO mice during adulthood. Furthermore, this diagnostic phenotype indicates that ER β plays a role in oocyte progression in the postnatal ovary (9, 10). Both ER α and ER β are activated by endogenous estrogens; however, their expression patterns and actions are different (11), with each receptor assumed to have specific biological functions.

A growing body of work in laboratory animals supports bisphenol A (BPA) as an endocrine-disrupting chemical (EDC) (12) that has adverse effects on not only the female reproductive system but also on the brain and immune system (13). BPA is used extensively as a raw material for making polycarbonate plastics and epoxy resins. However, its likely adverse effects on humans, especially infants and fetuses, have recently led to BPA being phased out of polycarbonate plastic and resin production (14). Various BPA derivatives have been developed to create more firm and stable plastics and resins, and these derivatives are now preferred as raw materials (15) (Fig. 1). However, BPA analogs have already been detected in the environment (15, 16). Fluorine-containing BPA, that is, bisphenol AF (BPAF, 2,2-Bis(4-hydroxyphenyl)hexafluoropropane, Chemical Abstracts Service [CAS] No. 1478-61-1), is seen as a practical alternative to BPA, despite reported estrogenic activity in MCF-7 breast cancer cells (17). Eight BPA derivatives, including BPAF, have been detected in sediments collected from industrialized areas (18) and indoor dust (19). In addition, BPA analogs have been found in urine samples from individuals living close to a BPAF-manufacturing plant (20) and a municipal solid waste incineration plant (21). Chlorine-containing BPA, that is, bisphenol C (BPC, also known as bisphenol C2 or bisphenol Cl2, 1,1-dichloro-2,2-bis(4-

[‡] These authors contributed equally to this work.

* For correspondence: Ayami Matsushima, ayami@chem.kyushu-univ.jp.

Dual role of bisphenols as ER α agonists and ER β antagonists




Figure 1. Structures of BPA derivatives selected via screening using an ER β binding assay. Chemical structures of E2, 4OHT, and 20 BPA-related compounds exhibited stronger binding abilities than BPA; BPC had the highest binding ability to ER β . Fluorine-containing BPA derivatives, that is, 9,9-bis(4-hydroxyphenyl)fluorene and 9,9-bis(4-hydroxy-3-methylphenyl)fluorene, exerted stronger binding abilities than did BPA. 4OHT, 4-hydroxytamoxifen; BPA, bisphenol A; BPC, bisphenol C; E2, 17- β estradiol; ER, estrogen receptor.

hydroxyphenyl)ethylene, CAS No. 14868-03-2), is a beneficial substrate for polymer production because of the high thermal stability of BPC-containing polycarbonate (22–24). Notably, BPC is structurally similar to two banned pesticides

dichlorodiphenyltrichloroethane (1,1'-(2,2,2-trichloroethylidene)bis(4-chlorobenzene), CAS No. 50-29-3) and methoxychlor (1,1'-(2,2,2-trichloroethylidene)bis(4-methoxybenzene), CAS No. 72-43-5) (25, 26). Based on its high affinity

for endogenous ERs in MCF-7 cells (27), BPC was considered but ultimately not included in the list of *in vitro* endocrine disruptors by the Interagency Coordinating Committee on the Validation of Alternative Methods (ICCVAM) (NIH Publication No: 03-4503) in 2003. Historically, the designation of 2,2-bis(4-hydroxy-3-methylphenyl) propane (CAS No. 79-97-0, which does not have chlorine atoms) as BPC has led to some confusion in the literature; however, chlorine-containing BPC has been detected in human breast milk (28).

ER α and/or ER β are major targets of EDCs that interfere with their estrogen-responsive signaling pathways (29). Human ER α and ER β have almost identical DNA-binding domains, differing by only two amino acids, and both receptors bind the same estrogen-response elements in transcriptional control regions. Although ER α and ER β also have similar LBDs, they have some distinctive features in terms of ligand selectivity and target gene regulation (30). Endogenous estrogen, 17- β estradiol (E2), binds to ER α slightly stronger than to ER β . Similarly, BPA binds ER α with higher affinity than ER β , although its binding abilities are much weaker than those of E2. In contrast, BPAF and BPC display higher affinity for both ER α and ER β than BPA, with a preference for ER β over ER α binding. BPAF and BPC show antagonistic activity against ER β in reporter gene assays using HeLa cells (31, 32). BPAF and BPC show much stronger antagonist activity for ER β than ER α , (32, 33). While crystal structures have provided insight into ER α activation/inactivation mediated by BPAF and BPC binding (32, 33), the structural changes induced by the strong antagonistic activity of BPAF and BPC against ER β are not well established. Recently, we found that the bisphenol moiety is a privileged structure for ER α . Here, we describe the biphasic binding of BPAF and BPC to ER β and propose a novel two-site binding model of the ER β -BPC complex, based on the crystal structure of 4-hydroxytamoxifen (4OHT) bound to ER β . This

is the first study to mechanistically associate the antagonistic actions of EDCs with interactions at the coactivator-binding site, thereby providing insight into developing safer raw materials that do not exhibit endocrine-disrupting features.

Results

The bisphenol scaffold binds both ER α and ER β

We screened a library of 119 bisphenol derivatives and related compounds using a radioligand competitive binding assay with tritium-labeled E2 ($[^3\text{H}]$ E2) for ER β . Some of these bisphenol derivatives have been detected in human biological samples (16). The CAS registry numbers (RNs), common names, and IUPAC names are provided in Table S1. We found 18 bisphenol derivatives with similar or stronger ER β binding than BPA (Table 1 and Fig. S1). BPC showed the strongest ER β (IC₅₀ of 2.99 nM) and highest ER α (IC₅₀ of 2.81 nM) binding affinity of the derivatives examined. The second strongest ER β binding was seen with compound No. 2 (4,4'-(1,3-dimethylbutylidene)bisphenol; IC₅₀ of 16.1 nM), although higher affinity was measured with ER α (IC₅₀ of 5.75 nM). 4,4'-(1,3-Dimethylbutylidene)bisphenol, 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane (HPTE) (3), and BPAF showed comparable binding ability to ER β (IC₅₀ of ~18 nM). Contrary to the results for 4,4'-(1,3-dimethylbutylidene)bisphenol (2), HPTE (3) and BPAF were preferential ER β ligands, displaying three times stronger binding to ER β than ER α . Although bisphenol Z (5), 4,4'-(2-ethylhexylidene)bisphenol (6), and 4,4'-(2-hydroxybenzylidene)-bis(2,3,6-trimethylphenol) (7) showed similar results to BPAF, they bound more strongly to ER α . The majority of the chemicals tested elicited comparable binding to both ER α and ER β . Of the 18 derivatives with similar or stronger ER β binding compared with BPA, 14 showed slightly stronger binding

Table 1
Receptor binding affinity (mean \pm SD) of BPA derivatives for ER β

Compound No.	Chemicals	Binding affinity (IC ₅₀ , nM)	
		ER β	ER α (34)
E2	Estradiol	2.17 \pm 0.6	0.88 \pm 0.13
1	Bisphenol C	2.99 \pm 1.0	2.81 \pm 0.61
4OHT	4-Hydroxytamoxifen	4.66 \pm 1.5	2.85 \pm 0.20
2	4,4'-(1,3-Dimethylbutylidene)bisphenol	16.1 \pm 6.1	5.75 \pm 1.92
3	2,2-Bis(<i>p</i> -hydroxyphenyl)-1,1,1-trichloroethane (HPTE)	18.1 \pm 4.9	59.1 \pm 1.5
4	Bisphenol AF	18.9 \pm 8.4	53.4 \pm 7.3
5	Bisphenol Z	21.6 \pm 1.9	56.9 \pm 0.6
6	4,4'-(2-Ethylhexylidene)bisphenol	25.9 \pm 8.5	18.5 \pm 6.7
7	4,4'-(2-Hydroxybenzylidene)-bis(2,3,6-trimethylphenol)	41.5 \pm 2.0	12.3 \pm 7.3
8	Bisphenol B	79.8 \pm 12.6	195 \pm 44
9	1,1-Bis(4-hydroxy-3-methylphenyl)cyclohexane	132 \pm 6.5	38.6 \pm 7.2
10	Bisphenol M	148 \pm 80	56.8 \pm 11.7
11	Bisphenol AP	158 \pm 33	259 \pm 41
12	α , α' -Tris(4-hydroxyphenyl)-1-ethyl-4-isopropylbenzene	212 \pm 36	61.7 \pm 10.4
13	2,2-Bis(3-amino-4-hydroxyphenyl)hexafluoropropane	224 \pm 113	334 \pm 112
14	9,9-Bis(4-hydroxyphenyl)fluorene	325 \pm 60	2230 \pm 202
15	9,9-Bis(4-hydroxy-3-methylphenyl)fluorene	405 \pm 108	321 \pm 103
16	Bisphenol P	607 \pm 28	176 \pm 35
17	2,2-Bis[4-(4-aminophenoxy)phenyl]hexafluoropropane	609 \pm 81	1030 \pm 375
18	2,2-Bis(4-hydroxy-3-methylphenyl)propane	744 \pm 429	368 \pm 22
19	Bisphenol A	900 \pm 70	1780 \pm 764
20	α , α' -Bis(4-hydroxy-3,5-dimethylphenyl)-1,4-diisopropylbenzene	>10,000	733 \pm 628

Receptor binding affinity was evaluated by competitive binding assay using [^3H] 17 β -estradiol as a radioligand.

Dual role of bisphenols as ER α agonists and ER β antagonists

abilities to ER α than ER β (Table 1). We reported that 18 bisphenol derivatives bound to ER α more strongly than did BPA (34). Bulky functional groups at their sp³-carbon connecting two phenol groups were beneficial for ER β binding, similar to the results previously observed for ER α (34). However, ER β binding abilities did not precisely correlate with those of ER α . Fluorene derivatives, 9,9-bis(4-hydroxyphenyl)fluorene (14) and 9,9-bis(4-hydroxy-3-methylphenyl)fluorene (15) not only bound to ER α (34, 35) but also to ER β (35), with their ER β binding ability stronger than that of BPA. Bisphenol derivatives possessing halogen atoms between two phenol groups, especially chlorine-containing derivatives, showed strong ER β binding.

To gain insight into the differences observed in ER β and ER α binding, we compared the ligand-binding cavities in the deposited ER β and ER α LBD crystal structures. The sizes of the canonical binding pockets were calculated for 45 ER α and 25 ER β structures in their active conformations using MOE SiteFinder function, and the amino acid residues surrounding the bound ligands identified (Tables S2 and S3). The average ER β pocket was smaller than for ER α (430.9 Å³ and 369.3 Å³ for ER α and ER β , respectively; Fig. 2A). The typical ligand-binding pockets of each receptor in the active conformation are illustrated (Fig. 2, C and D). Moreover, the average size of the ligand-binding pocket in E2-bound ER α and ER β structures was 419.4 Å³ and 385.0 Å³, respectively, and in

genistein-bound ER α and ER β structures was 475.9 Å³ and 375.8 Å³, respectively. Although these results suggested that ER α is able to accept larger ligands than ER β , the amino acid residues surrounding the ligands differ slightly. Some of the smaller ligands fit more adequately into the ER β than the ER α ligand-binding pocket.

BPC and BPAF bind but fail to activate ER β

Reporter assays using HeLa cells were performed to evaluate ER β transcriptional activity induced by BPA, BPC, BPAF, and 17 bisphenol derivatives (Fig. 2B). The detailed dose-dependent transcriptional activity of each compound is shown in Figure S2. The statistical significance and the value of maximum fold induction of each compound are summarized in Table S4. BPA elicited the strongest ER β agonistic activity of the derivatives, with the activity at 10 μM comparable with that seen with the endogenous ligand E2 despite its affinity being 400 times weaker than that of E2. 4,4'-(1,3-Dimethylbutylidene)bisphenol (2) and bisphenol B (8) achieved ~50% of BPA-induced transcriptional activity at the highest concentration of 10 μM. While compound 2, found as an impurity in industrial-grade BPA, has been shown to function as an ER α agonist in yeast-two hybrid assays (36), our results reveal a high affinity for and functional activation of ER β . Compounds 2 and 8 are structurally similar to BPA,




Figure 2. Differential activities of BPA derivatives on ER α and ER β . A, ligand-binding pocket volumes from ER α (open circles) and ER β (filled circles) calculated from crystal structures in the presence of activating ligands; average volumes indicated by red lines. B, top 20 BPA derivatives binding to ER β induced partial agonistic activity against ER β . C, the ligand-binding pockets of ER α (PDB ID: 1QKU) and (D) ER β (3OLL) are illustrated in gray; estradiol is bound as the ligand. E, sixteen chemicals, including tricyclic bisphenols, inhibited more than half of the 10 nM E2-induced transcriptional activity. BPA, bisphenol A; BPC, bisphenol C; E2, 17- β estradiol; ERs, estrogen receptors.

possessing one methyl group on the sp³-carbon that bridges the two phenol groups, suggesting that this conformation is beneficial for ER β activation. BPC, HTPe, BPAF, bisphenol Z, 1,1-bis(4-hydroxy-3-methylphenyl)cyclohexane (9), 9,9-bis(4-hydroxy-3-methylphenyl)fluorene (15), and 2,2-bis(4-hydroxy-3-methylphenyl)propane (18) functioned as partial agonists, inducing 20% to 30% of the E2-induced transcriptional activity. The transcriptional activity of BPC, HTPe, and BPAF was consistent with a previous report investigating ER α and ER β , in which these compounds elicited weaker activity against ER β than ER α (32, 33). Surprisingly, 4,4'-(2-ethylhexylidene)bisphenol (6), 4,4'-(2-hydroxybenzylidene)-bis(2,3,6-trimethylphenol) (7), bisphenol M (10), α , α , α' -tris(4-hydroxyphenyl)-1-ethyl-4-isopropylbenzene (12), bisphenol P (16), and α , α' -bis(4-hydroxy-3,5-dimethylphenyl)-1,4-diisopropylbenzene (20) showed no agonist activity against ER β . These findings contrast with ER α , where the majority of bisphenol derivatives with strong binding affinity also showed strong agonistic activity (34).

BPA derivatives function as ER β antagonists

The finding that many BPA derivatives with high binding affinities showed almost no agonist activity suggested that they function as ER β antagonists. To explore this possibility, the inhibitory effects of the BPA derivatives (100 nM, 1 μ M, 10 μ M) against 10 nM E2-induced ER β activation were measured (Fig. 2E). The statistical significance and transcriptional activity of each compound at the highest effective inhibitory concentration are summarized in Table S5. BPC showed the strongest antagonistic activity, with additional halogen-containing bisphenols (*i.e.*, HTPe, and BPAF), also elicited antagonistic activities, consistent with previous reports (31–33). 4,4'-(1,3-Dimethylbutylidene)bisphenol (2), which had the second strongest binding ability and partial agonist activity compared with BPA, showed weak antagonist activity, contrasting with its reported ER α agonism. Bisphenol B (8) showed similar weak antagonist activity, with both bisphenol B (8) and 4,4'-(1,3-dimethylbutylidene)bisphenol (2) inhibiting 50% of BPA-induced activation. Tricycle bisphenols (*i.e.*, bisphenol M (10), α , α , α' -tris(4-hydroxyphenyl)-1-ethyl-4-isopropylbenzene (12), bisphenol P (16), and α , α' -bis(4-hydroxy-3,5-dimethylphenyl)-1,4-diisopropylbenzene (20)) showed antagonistic activity, presumably through the disruption of the active conformation, as reported for ER α (34). While demonstrating no agonist activity, 4,4'-(2-ethylhexylidene)bisphenol (6) and 4,4'-(2-hydroxybenzylidene)-bis(2,3,6-trimethylphenol) (7) suppressed 90% of E2-induced activation at the 10 μ M concentration. Interestingly, the fluorene derivative, 9,9-bis(4-hydroxy-3-methylphenyl)fluorene (15), functioned as a weak antagonist, demonstrating that fluorene derivatives 14 and 15 can exhibit both ER β and ER α antagonistic activities (34, 35). With the exception of the tricyclic bisphenols, these findings indicate that most bisphenol derivatives with strong ER β binding functioned as antagonists, although they showed only agonist activities to ER α (34).

Docking analysis predicts BPC binding to the surface of ER β

To investigate the contrasting actions of BPA derivatives as ER β antagonists and ER α agonists, we performed docking simulations using the LBD of human ER β and BPC, the strongest binder among the BPA derivatives examined using a competitive binding assay with [³H]E2. Possible ligand-binding sites in 38 deposited ER β crystal structures were identified using MOE SiteFinder, a program for binding-site analysis equipped in the Molecular Operating Environment (MOE). Canonical as well as putative binding sites were ranked according to propensity for ligand binding (PLB), a specific parameter in MOE SiteFinder (37). Consistently, the top five predicted sites in each structure were the canonical ligand-binding sites. Interestingly, an actual surface 4OHT-binding site close to the hydrophobic groove for the coactivator recognition surface of ER β (PDB ID: 2FSZ) was ranked 11th in the PLB order. Moreover, this location was a predicted binding site on all antagonist-bound ER β structures, based on PLB. Notably, this second site was not predicted as a binding site on over half of the agonist-bound structures (Table S6). These predictions suggest that ER β antagonism induced by BPC and other BPA derivatives may be due to inhibition of coactivator recruitment. Next, we performed a docking simulation for ER β LBD and BPC using both its canonical and second binding sites as target rooms. BPC was able to fit and bind in both rooms, with one of its chlorine atoms interacting with the tryptophan residue (Trp335) on helix 5 *via* halogen interaction (Fig. 3, A and B). The obtained model structure suggested that BPC binding to the second binding site prevented recruitment of coactivators for gene transcriptions, similar to 4OHT (Fig. 3, C and D). We hypothesized that the binding affinity of BPA derivatives to this coactivator binding site would correlate with antagonistic activity. To explore this notion, docking simulations were performed for each BPA derivatives (Fig. S3), and the free energy of ligand binding evaluated using a docking simulation and the GBVI/WSA dG scoring function (larger negative scores indicate more stable ligand/receptor complexes) (38). Correlation of the GBVI/WSA dG scores with the extent of antagonism (reported as the % inhibition of 10 nM E2 induced transcriptional activity) revealed a linear relationship (correlation coefficient of – 0.83), suggesting that inhibition of coactivator recruitment underlies the antagonism of ER β by BPA derivatives (Fig. 3E).

Binding of the coactivator peptide is reduced by BPC

Ligand binding induces a conformation change in the ER β LBD that facilitates its translocation to the cell nucleus and the subsequent recruitment of coactivator proteins. To explore the effects of BPC on ER β activation, surface plasmon resonance experiments were performed to measure the direct binding of the coactivator peptide derived from human nuclear receptor coactivator 1, also known as steroid receptor coactivator (SRC1). Consistently, the E2 ligand increased SRC1 peptide binding to ER β -LBD (K_d 3.3 \pm 0.6 μ M and 9.1 \pm 0.7 μ M with and without E2, respectively; Fig. 3G). Notably, SRC1 peptide binding was reduced in the presence of BPC (K_d 16.4 \pm 0.9 μ M; Fig. 3H).

Dual role of bisphenols as ER α agonists and ER β antagonists




Figure 3. ER β harbors two ligands in its LBD. *A*, two BPC bound to ER β during the docking simulation. The canonical binding site is indicated in gray; the second binding site, located on the surface of the receptor, is shown in magenta. The activation helix, H12, is indicated in magenta. *B*, chlorine, a halogen atom of BPC, interacted with the Trp335 side chain via halogen interaction in the second binding site. BPC and 4OHT are illustrated in blue and gray, respectively, in the stick model. *C*, superimposition of the calculated BPC-bound ER β structure (blue) and its agonist form with the nuclear receptor coactivator 1, SRC1 (green, PDB ID: 3OLL). SRC1 is indicated as a red α -helix, H12 of its agonist form is indicated in purple, BPC is illustrated in blue, and 4OHT is shown in gray. BPC clashed with the amino acid residues on H12 in the ER β agonist form; therefore, BPC prevented the ER β activation. BPC and 4OHT disrupted the SRC1 binding due to steric hindrance of the amino acid residues shown in the red stick models. *D*, in ER β -agonist form, amino acid residues surrounding Trp335 within 4.5 Å on H12 are shown in the purple stick model, while leucine residues on the SRC1 LXXLL motif are indicated via the red stick model. *E* and *F*, correlation of the calculated binding scores and inhibitory activity for ER β . Inhibitory activity is defined as the ratio of chemicals inhibiting transcriptional activity induced by 10 nM E2. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. *G* and *H*, dose response of SRC1 peptide binding to ER β LBD in the presence of (G) 10 μ M E2 or (H) 10 μ M BPC. 4OHT, 4-hydroxytamoxifen; BPC, bisphenol C; E2, 17- β estradiol; ERs, estrogen receptors; LBD, ligand-binding domain; SRC1, steroid receptor coactivator.

Biphasic 4OHT binding indicative of two ER β -binding sites

To further support the presence of a second ligand-binding site, competitive binding assays were performed using BPA, BPC, and BPAF and tritium-labeled 4OHT ($[^3\text{H}]4\text{OHT}$) (Fig. 4A). Notably, a biphasic dose–response curve was observed for BPC (18.1 nM and 2281 nM IC_{50}) that was not evident in the $[^3\text{H}]E2$ competitive analyses. Similarly, BPAF displayed a biphasic binding curve, albeit with weaker binding at both the high- and low-affinity sites than BPC. Moreover, 4OHT showed a biphasic curve, consistent with the 4OHT/ER β crystal structure (PDB: ID 2FSZ). In contrast, BPA, which did not elicit antagonistic activity, showed a sigmoidal curve

indicative of a single ligand-binding site. Interestingly, the trifluorine substitution of the methyl groups in BPAF increased ER β binding ~ 50 -fold compared with BPA. These results confirmed the presence of two distinguishable binding sites for BPC and BPAF on ER β . In contrast, the typical sigmoidal curves seen in E2 competitive binding assays using $[^3\text{H}]4\text{OHT}$ and $[^3\text{H}]E2$ are indicative of single ligand-binding site.

Trp335 is required for biphasic ligand binding

The docking simulations suggested that hydrophobic interactions between the BPA derivatives and the indole group of Trp335 were required for ER β binding and identified a




Figure 4. Binding properties and transcriptional features of BPAF and BPC showed the importance of ER β W335 for their receptor binding and activation. *A*, detailed competitive binding curves of BPA, BPAF, BPC, and 4OHT using [3 H]4OHT illustrated a biphasic binding curve, in which chemicals compete with [3 H]4OHT in two binding sites on WT ER β . *B*, ER β (W335A) competitive binding assays showed typical sigmoidal binding curves. *C*, the reporter gene assay indicated that BPAF and BPC induced weak transcriptional activity in WT ER β , whereas E2 and BPA showed strong transcriptional activity. *D*, ER β (W335A) lost E2 or BPA-induced transcriptional activity, indicating that Trp335 substitution disrupted active conformation. *E*, in ER β agonist form, amino acid residues surrounding Trp335 within 4.5 Å are represented as green and purple stick models. (PDB ID: 3OLL). [3 H]4OHT, tritium-labeled 4OHT; BPAF, 2,2-Bis(4-hydroxyphenyl)hexafluoropropane; BPC, bisphenol C; E2, 17- β estradiol; ERs, estrogen receptors.

potential halogen interaction between the chlorine atom of BPC and the indole ring. To determine the contributions of these putative interaction to BPC binding, the corresponding tryptophan was mutated to alanine (A). Saturation binding assays revealed a typical sigmoidal dose–response curve and a K_d of 23.1 nM for E2 against ER β (W335A), indicating preservation of the canonical binding site (Fig. S4A).

Competitive binding assays confirmed two 4OHT-binding sites in ER β , with K_d values of 4.6 nM and 53.1 nM. In contrast, a single binding site was evident in ER β (W335A) (K_d 34.2 nM) (Fig. S4B). Similarly, the biphasic binding of BPC and BPAF was lost in the ER β (W335A) mutant (Fig. 4, A and B). The IC_{50} values of 4OHT, BPC, and BPAF were 106 ± 51 nM, 691 ± 29 nM, and 1249 ± 579 nM, respectively. BPA illustrated a typical sigmoidal competitive dose–response curve against ER β (W335A), similar to the result against ER β . These results indicated that replacing Trp for Ala compromises the second 4OHT and BPA derivatives binding site on the surface of the ER β LBD.

W335A reduces ER β transcription activity

Reporter assays revealed that E2-induced transcriptional activation was markedly reduced by the tryptophan to alanine

substitution in ER β (Fig. 4, C and D). Given that E2 binding ability was retained, this is consistent with reduced coactivator binding. Indeed, in the active conformation, Trp335 interacts with Leu491, Met494, and Leu495 on H12 (Fig. 4E). Supporting this notion, the SRC1 peptide bound poorly to ER β (W335A), as measured by surface plasmon resonance experiments using Biacore T100 (Fig. S5). These results indicated that Trp335 on the ER β coactivator-binding site plays an important role, not only in interacting with bisphenol derivatives but also in recruiting coactivators on the surface of ER β by stabilizing H12 in its active conformation.

Discussion

Here, we report the ER β transcriptional activities of BPA derivatives including BPC and BPAF using a combination of receptor binding and reporter assays. Of note, 18 derivatives bound ER β with higher affinity than BPA. The binding abilities of these BPA derivatives are stronger than those of known environmental chemicals such as dichlorodiphenyltrichloroethane, nonylphenol, phytoestrogens, and dioxins (39). Unexpectedly, our results clearly showed that many BPA derivatives function as ER β antagonists, contrasting with their previously reported ER α agonism. Docking simulations indicated that

Dual role of bisphenols as ER α agonists and ER β antagonists

BPA derivatives bind to a second site located near the coactivator-binding site on the surface of ER β -LBD that requires interactions with Trp335. Mutation of tryptophan to alanine led to the loss of this low-affinity binding site in ER β . These results indicated that some BPA derivatives act as antagonists, although most of EDCs, including BPA, are assumed ER agonists. We previously reported that most of the BPA derivatives examined in this study act as weak agonists for ER α . The results obtained in this study demonstrate the importance of screening for both agonist and antagonist activity, especially against ER β .

We previously reported that tricyclic bisphenols, that is, bisphenol M, α , α , α' -tris(4-hydroxyphenyl)-1-ethyl-4-isopropylbenzene, bisphenol P, and α,α' -Bis(4-hydroxy-3,5-dimethylphenyl)-1,4-diisopropylbenzene, act as antagonists against ER α because of the steric hindrance caused by the third aromatic ring structure (34). This study showed that this feature is also valid for ER β ; tricyclic bisphenols act as antagonists not only for ER α but also ER β . In addition to tricyclic bisphenols, many BPA derivatives, including BPAF and BPC, elicit antagonist activity. Our finding for BPAF and BPC are consistent with reports that both chemicals showed partial agonism for ER α and antagonism for ER β (31, 32, 40, 41).

Several ER α - or ER β -specific agonists have been reported, including propyl pyrazole triol that selectively binds to and transcriptionally activates ER α (42). The first chemical shown to function as an ER α agonist and ER β antagonist is HPTE, a metabolite of the banned pesticide, methoxychlor [1,1,1-trichloro-2,2-bis(4-methoxyphenyl)ethane] (43, 44). Accumulated knowledge gained from protein crystal structures emphasize the importance of halogens in receptor–ligand interactions (45, 46). We found that in addition to the halogen containing BPAF and BPC, many BPA derivatives display ER α agonist activities similar to HPTE. These results indicate the complexity of establishing the mechanisms of action of environmental chemicals that activate or suppress the physiological functions of one or more nuclear receptors. In particular, antagonist activities might be overlooked if both binding affinity and transcriptional activity are not determined, as environmental chemicals are typically categorized based on the ability to active ERs.

Recent studies have indicated the value of small molecules that bind to coactivator protein-binding sites on nuclear receptors (47). Coactivator-binding inhibitors (CBIs) have been developed for ERs, an androgen receptor, a progesterone receptor, a vitamin D receptor, a thyroid hormone receptor, a pregnane X receptor, a retinoid X receptor, and peroxisome proliferator-activated receptors (48–51). This study is the first to conclude that EDCs can function as CBIs for ER β , indicating the importance of assessing both agonist and antagonist activities of these chemicals.

In summary, we showed that tricyclic bisphenols elicit antagonistic activity against both ER α and ER β . Our results also indicate that many next-generation bisphenols are agonists and antagonists of ER α and ER β . Mutagenesis of an ER β surface amino acid indicated that these next-generation bisphenols act as CBIs. While *in silico* docking analyses

support this mechanism of action, future crystallographic studies will be required to provide more direct information on CBIs. This study highlights the mechanistic complexity of the next-generation of bisphenols acting as EDCs.

Experimental procedures

Chemicals

E2 (CAS RN 50-28-2, >98.9%) was obtained from Research Biochemicals International. 4OHT (CAS RN 68047–06–3, >98%) and HPTE (CAS RN 2971–36–0, >98.9%) were obtained from Sigma-Aldrich Inc. 4,4'-dihydroxydiphenylmethane (bisphenol F, CAS RN 620-92-8, >99.0%) and hexestrol (CAS RN 84-16-2, >99.0%) were obtained from FUJIFILM Wako Pure Chemical Corporation; the remaining 117 chemicals were purchased from Tokyo Chemical Industry Co, Ltd. Dimethyl sulfoxide, used to dissolve each compound in a 10 mM stock solution, was obtained from Sigma-Aldrich. [^3H]E2 (4458 GBq/mmol) and [^3H]4OHT (2960 GBq/mmol) were purchased from PerkinElmer.

ER β expression and purification

The LBD of ER β (amino acids 263–530) was expressed as a GST-fused protein for receptor-binding assays. Human ER β cDNA was obtained from OriGene Technologies. The cDNA of ER β -LBD was amplified using PCR and subcloned into a pGEX-6p-1 expression vector. The expression of GST-fused ER β -LBD was induced by 1 mM IPTG in *Escherichia coli* BL21 α at 16 °C for overnight. The resulting crude protein was affinity-purified using Glutathione-Sepharose 4B (Cytiva), followed by gel filtration using a Sephadex G-10 column (Cytiva).

Radioligand-binding assay

Radioligand-binding assays for ER β and ER β (W335A) were performed mainly according to a previously reported method (31, 34). Saturation binding assays were conducted with [^3H]E2 or [^3H]4OHT using GST-ER β -LBD or GST-ER β (W335A)-LBD to evaluate the binding ability of radiolabeled compounds. The reaction mixtures of each LBD (20 ng) and a series of concentrations of [^3H]E2 (0.01–10 nM) or [^3H]4OHT (0.1–30 nM) were incubated in a total volume of 100 μl of the binding buffer (10 mM Tris-buffered saline (pH 7.4), 1 mM EGTA, 1 mM EDTA, 10% glycerol, 0.5 mM PMSF, 0.2 mM leupeptin, and 1 mM sodium vanadate (V)) at 20 °C for 2 h, to analyze total binding. Corresponding reaction mixtures, containing 10 μM nonlabeled E2 or 4OHT, were incubated to detect each nonspecific binding. [^3H]E2 or [^3H]4OHT-specific binding was evaluated by subtracting the obtained radioactivity values of total binding from the those of nonspecific binding. After successive incubation with 100 μl of 0.4% dextran-coated charcoal (DCC) (Sigma-Aldrich) in PBS (pH 7.4) on ice for 10 min, free radioligands bound to DCC were removed using a vacuum filtration system with a 96-well filtration plate (MultiScreenHTS HV, 0.45-mm pore size, Merck KGaA) for the bound/free separation. The radioactivity of each eluent was measured using a liquid scintillation counter (LS6500;

Beckman Coulter) and Clear-sol I (Nacalai Tesque Inc). Calculated specific binding of [3 H]E2 was assessed using Scatchard plot analysis (52). Competitive binding assays were performed to evaluate the binding ability of each test compound using [3 H]E2, for a library screening or detailed BPA binding assay. Each compound was dissolved in dimethyl sulfoxide to prepare a 1.0 mM stock solution and further diluted to prepare serial dilutions (10^{-12} M to 10^{-5} M) in the binding buffer. To assess their binding abilities, each compound was incubated with GST-ER β -LBD or GST-ER β (W335A)-LBD (20 ng) and radiolabeled ligand (5 nM of [3 H]E2 or 5 nM of [3 H]4OHT, final concentration) for 2 h at 20 °C. Bound/free separation was performed as described above, and the radioactivity was determined using a MicroBeta microplate counter (PerkinElmer Inc). The IC $_{50}$ value of each test compound was calculated from the dose–response curves generated *via* nonlinear regression analysis using Prism software (GraphPad Software Inc).

Luciferase reporter gene assay

Transcriptional activities of ER β and ER β (W335A) were measured as previously reported (31, 34). HeLa cells were maintained in Eagle's minimum essential medium (Nissui Pharmaceutical Co, Ltd) supplemented with DCC-treated fetal bovine serum (10%, v/v) at 37 °C under 5% CO $_2$. To evaluate agonistic activity, HeLa cells were seeded at a density of 5×10^5 cells per 60-mm dish and cultured for 24 h, followed by transfection of the reporter plasmid (3 μ g, pGL4.23/3 \times ERE) and each expression plasmid (1 μ g, pcDNA3.1/ER β or pcDNA3.1/ER β (W335A)) using Lipofectamine LTX with Plus Reagent (Thermo Fisher Scientific, Inc), according to the manufacturer's instructions. After incubation for 24 h, cells were harvested and seeded onto 96-well plates at 5×10^4 cells/well, and then treated with a series of the test compounds (10^{-12} M to 10^{-5} M, final concentration) diluted with 1% bovine serum albumin/PBS (v/v). After a 24-h incubation, luciferase activity was measured using the ONE-Glo Luciferase Assay System (Promega Co) on an EnSpire multimode plate reader (PerkinElmer, Inc). To analyze antagonistic activity, serial concentrations of test compounds (10^{-12} M to 10^{-5} M) were treated in the presence of 10 nM E2, which normally induces full transcriptional activity levels in transiently expressed ER β .

Docking simulation of each antagonist onto the ER β LBD

Three-dimensional coordinates of the compounds were obtained from the Cambridge Structural Database (CSD-Core, The Cambridge Crystallographic Data Centre). Ligand IDs of compounds utilized for docking simulations are summarized in Table S7. For the compounds with no corresponding entry in the CSD System, 3D coordinates were constructed *in silico* using Gaussian 16 (Gaussian, Inc), with the basis set of 6–31G. Docking simulations for the ligand–ER β complex were performed using a Dock functions in the MOE package (Chemical Computing Group); the free energy of each complex was evaluated according to its GBVI/WSA dG score (38).

Ligand-binding cavity volumes of the deposited crystal structures were analyzed and calculated using the MOE SiteFinder function in MOE.

Binding analysis of ER β LBD and SRC1 peptide by surface plasmon resonance

The anti-GST antibody was immobilized on a Sensor Chip CM5 (Cytiva) using Amine Coupling kit (Cytiva) and GST Capture kit (Cytiva) according to the manufacturer's instruction for Biacore T100 instrument (Cytiva). The binding of SRC1 peptide (amino acids 685–697; ERHKILHRLIQEG) to the ER β -LBD was analyzed by capturing GST-ER β -LBD on the sensor chip and injecting SRC1 peptide with E2 or BPC. The peptide was synthesized using the ABI 433A peptide synthesizer (Applied Biosystems) by the solid-phase method with Fmoc chemistry. GST-ER β -LBD (50 μ g/ml) was incubated with 10 μ M E2 or 10 μ M BPC for 1 h and captured at 25 °C with a flow rate of 5 μ l/min on the sensor chip. Binding between SRC1 peptide and ER β -LBD was analyzed using HBS-EP+ buffer (0.01 M HEPES, pH 7.4, 0.15 M NaCl, 3 mM EDTA, and 0.05% (w/v) Surfactant P20) as a running buffer under the following conditions: contact time 120 s, flow rate 30 μ l/min, and dissociation time 180 s. The sensor chip was recovered by 10 mM Gly-HCl (pH 2.0) with a flow rate of 20 μ l/min and a contact time of 120 s. The data obtained were analyzed using the Biacore T100 evaluation software.

Statistical analysis

Significance of the data between experimental groups was determined using unpaired *t*-tests. Data are presented as the mean \pm SD, and *p* values are summarized in supplementary tables.

Data availability

All data needed to evaluate the conclusions in the article are present in the article and/or the supporting information.

Supporting information—This article contains supporting information.

Acknowledgments—We appreciate R.T. Yu and A.R. Atkins (Salk Institute for Biological Studies) for helpful suggestions and discussions, and L. Ong and C. Brondos for administrative assistance. We appreciate Y. Shimohigashi (Kyushu University) for providing the chemical library and X. Liu (Kyushu University) for providing the ER β -mutated plasmid. We thank the RIKEN BRC Cell Bank and Cell Resource Center for Biomedical Research, Institute of Development, Aging and Cancer, Tohoku University, for providing the HeLa cells. This work was supported in part by a grant from Izumi Science and Technology Foundation.

Author contributions—M. I., T. M., M. H., K. T., T. I., K. S., T. N., and A. M. data curation; M. I., T. M., M. H., K. T., T. I., and A. M. formal analysis; E. Y., M. D., R. M. E., and A. M. writing—review and editing; R. M. E. and A. M. funding acquisition; A. M. methodology; A. M. conceptualization; M. I., T. M., K. T., T. I. and A. M. investigation; A. M. writing—original draft, A. M. project administration.

Dual role of bisphenols as ER α agonists and ER β antagonists

Funding and additional information—This work was supported by a grant-in-aid for scientific research JSPS KAKENHI JP17H01881, JP18K19147, 18KK0320, and 20H00635, to A. M. R. M. E. is supported by the NOMIS Foundation—Science of Health. M. D. and R. M. E. were supported in part by the National Institute of Environmental Health Sciences of the NIH under Award Number P42ES010337. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest—R. M. E. holds the March of Dimes Chair in Molecular and Developmental Biology at the Salk Institute. All other authors declare that they have no conflicts of interest with the contents of this article.

Abbreviations—The abbreviations used are: [^3H]4OHT, tritium-labeled 4OHT; [^3H]E2, tritium-labeled E2; 4OHT, 4-hydroxytamoxifen; BPA, bisphenol A; BPAF, 2,2-Bis(4-hydroxyphenyl)hexafluoropropane; BPC, bisphenol C; CAS, Chemical Abstracts Service; CBIs, coactivator-binding inhibitors; DCC, dextran-coated charcoal; E2, 17- β estradiol; EDC, endocrine-disrupting chemical; ERs, estrogen receptors; HPTE, 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane; LBDs, ligand-binding domains; MOE, Molecular Operating Environment; PLB, propensity for ligand binding; RNs, registry numbers; SRC1, steroid receptor coactivator.

References

1. Nuclear Receptors Nomenclature Committee (1999) A unified nomenclature system for the nuclear receptor superfamily. *Cell* **97**, 161–163
2. Evans, R. M., and Mangelsdorf, D. J. (2014) Nuclear receptors, RXR, and the Big Bang. *Cell* **157**, 255–266
3. Moore, J. T., Collins, J. L., and Pearce, K. H. (2006) The nuclear receptor superfamily and drug discovery. *ChemBioChem* **1**, 504–523
4. Renoir, J.-M., Marsaud, V., and Lazennec, G. (2013) Estrogen receptor signaling as a target for novel breast cancer therapeutics. *Biochem. Pharmacol.* **85**, 449–465
5. Kuiper, G., Enmark, E., PeltoHuikko, M., Nilsson, S., and Gustafsson, J. A. (1996) Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 5925–5930
6. Hewitt, S. C., Harrell, J. C., and Korach, K. S. (2005) Lessons in estrogen biology from knockout and transgenic animals. *Annu. Rev. Physiol.* **67**, 285–308
7. Kregel, J. H., Hodgins, J. B., Couse, J. F., Enmark, E., Warner, M., Mahler, J. F., Sar, M., Korach, K. S., Gustafsson, J. A., and Smithies, O. (1998) *Proc. Natl. Acad. Sci. U. S. A.* **90**, 11162–11166
8. Kregel, J. H., Hodgins, J. B., Couse, J. F., Enmark, E., Warner, M., Mahler, J. F., Sar, M., Korach, K. S., Gustafsson, J. A., and Smithies, O. (1998) Generation and reproductive phenotypes of mice lacking estrogen receptor beta. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 15677–15682
9. Couse, J. F., Curtis Hewitt, S., Bunch, D. O., Sar, M., Walker, V. R., Davis, B. J., and Korach, K. S. (1999) Postnatal sex reversal of the ovaries in mice lacking estrogen receptors α and β . *Science* **286**, 2328–2331
10. Dupont, S., Krust, A., Gansmuller, A., Dierich, A., Chambon, P., and Mark, M. (2000) Effect of single and compound knockouts of estrogen receptors α (ER α) and β (ER β) on mouse reproductive phenotypes. *Development* **127**, 4277–4291
11. Uhlén, M., Fagerberg, L., Hallström, B. M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., Olsson, I. M., Edlund, K., Lundberg, E., Navani, S., Szijgyarto, C. A. K., et al. (2015) Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419
12. Dodds, E. C., and Lawson, W. (1936) Synthetic estrogenic agents without the phenanthrene nucleus. *Nature* **137**, 996
13. Richter, C. A., Birnbaum, L. S., Farabollini, F., Newbold, R. R., Rubin, B. S., Talsness, C. E., Vandenbergh, J. G., Walser-Kuntz, D. R., and vom Saal, F. S. (2007) *In vivo* effects of bisphenol A in laboratory rodent studies. *Reprod. Toxicol.* **24**, 199–224
14. Usman, A., and Ahmad, M. (2016) From BPA to its analogues: Is it a safe journey? *Chemosphere* **158**, 131–142
15. Pelch, K., Wignall, J. A., Goldstone, A. E., Ross, P. K., Blain, R. B., Shapiro, A. J., Holmgren, S. D., Hsieh, J.-H., Svoboda, D., Auerbach, S. S., Parham, F. M., Masten, S. A., Walker, V., Rooney, A., and Thayer, K. A. (2019) A scoping review of the health and toxicological activity of bisphenol A (BPA) structural analogues and functional alternatives. *Toxicology* **424**, 152235
16. Program, N. N. T., Pelch, K. E., Wignall, J. A., Goldstone, A. E., Ross, P. K., Blain, R. B., Shapiro, A. J., Holmgren, S. D., Hsieh, J.-H., Svoboda, D., Auerbach, S. S., Parham, F. M., Masten, S. A., and Thayer, K. A. (2017) *NTP research report on biological activity of bisphenol A (BPA) structural analogues and functional alternatives*. Research Report 4, National Toxicology Program, Research Triangle Park, NC
17. Pérez, P., Pulgar, R., Olea-Serrano, F., Villalobos, M., Rivas, A., Metzler, M., Pedraza, V., and Olea, N. (1998) The estrogenicity of bisphenol A-related diphenylalkanes with various substituents at the central carbon and the hydroxy groups. *Environ. Health Perspect.* **106**, 167–174
18. Liao, C., Liu, F., Moon, H.-B., Yamashita, N., Yun, S., and Kannan, K. (2012) Bisphenol analogues in sediments from industrialized areas in the United States, Japan, and Korea: Spatial and temporal distributions. *Environm. Sci. Technol.* **46**, 11558–11565
19. Liao, C., Liu, F., Guo, Y., Moon, H.-B., Nakata, H., Wu, Q., and Kannan, K. (2012) Occurrence of eight bisphenol analogues in indoor dust from the United States and several Asian countries: Implications for human exposure. *Environm. Sci. Technol.* **46**, 9138–9145
20. Yang, Y., Guan, J., Yin, J., Shao, B., and Li, H. (2014) Urinary levels of bisphenol analogues in residents living near a manufacturing plant in south China. *Chemosphere* **112**, 481–486
21. Lu, S., Yang, D., Ge, X., Li, L., Zhao, Y., Li, C., Ma, S., and Yu, Y. (2020) The internal exposure of phthalate metabolites and bisphenols in waste incineration plant workers and the associated health risks. *Environ. Int.* **145**, 106101
22. Dobkowski, Z. (1984) Procedure for evaluation of the Mark-Houwink constants. *J. Appl. Polym. Sci.* **29**, 2683–2694
23. Factor, A., and Orlando, C. M. (1980) Polycarbonates from 1,1-dichloro-2,2-bis(4-hydroxyphenyl)ethylene and bisphenol A: A highly flame-resistant family of engineering thermoplastics. *J. Polym. Sci. Polym. Chem.* **18**, 579–592
24. Jurs, J. L., and Tour, J. M. (2003) Novel flame retardant polyarylethers: Synthesis and testing. *Polymer* **44**, 3709–3714
25. Glustrom, L. W., Mitton-Fry, R. M., and Wuttke, D. S. (2002) Re: 1,1-Dichloro-2,2-bis-(*p*-chlorophenyl)ethylene and polychlorinated biphenyls and breast cancer: Combined analysis of five U.S. studies. *J. Natl. Cancer Instit.* **94**, 1337–1338. author reply 1338–1339
26. Matsushima, A. (2018) A novel action of endocrine-disrupting chemicals on wildlife; DDT and its derivatives have remained in the environment. *Int. J. Mol. Sci.* **19**, 1377
27. Stoessel, S., and Leclercq, G. (1986) Competitive binding assay for estrogen receptor in monolayer culture: Measure of receptor activation potency. *J. Steroid Biochem.* **25**, 677–682
28. Deceuninck, Y., Bichon, E., Marchand, P., Boquien, C.-Y., Legrand, A., Boscher, C., Antignac, J. P., and Le Bizec, B. (2015) Determination of bisphenol A and related substitutes/analogues in human breast milk using gas chromatography-tandem mass spectrometry. *Anal. Bioanal. Chem.* **407**, 2485–2497
29. Shanle, E. K., and Xu, W. (2011) Endocrine disrupting chemicals targeting estrogen receptor signaling: Identification and mechanisms of action. *Chem. Res. Toxicol.* **24**, 6–19
30. Kuiper, G. G., Carlsson, B., Grandien, K., Enmark, E., Häggblad, J., Nilsson, S., and Gustafsson, J. A. (1997) Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* **138**, 863–870
31. Matsushima, A., Liu, X., Okada, H., Shimohigashi, M., and Shimohigashi, Y. (2010) Bisphenol AF is a full agonist for the estrogen receptor ER α but

- a highly specific antagonist for ER β . *Environ. Health Perspect.* **118**, 1267–1272
32. Delfosse, V., Grimaldi, M., Pons, J.-L., Boulahtouf, A., Le Maire, A., Cavailles, V., Labesse, G., Bourguet, W., and Balaguer, P. (2012) Structural and mechanistic insights into bisphenols action provide guidelines for risk assessment and discovery of bisphenol A substitutes. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14930–14935
 33. Delfosse, V., Grimaldi, M., Cavailles, V., Balaguer, P., and Bourguet, W. (2014) Structural and functional profiling of environmental ligands for estrogen receptors. *Environ. Health Perspect.* **122**, 1306–1313
 34. Masuya, T., Iwamoto, M., Liu, X., and Matsushima, A. (2019) Discovery of novel oestrogen receptor α agonists and antagonists by screening a revisited privileged structure moiety for nuclear receptors. *Sci. Rep.* **9**, 9954
 35. Zhang, Z., Hu, Y., Guo, J., Yu, T., Sun, L., Xiao, X., Zhu, D., Nakanishi, T., Hiromori, Y., Li, J., Fan, X., Wan, Y., Cheng, S., Li, J., Guo, X., *et al.* (2017) Fluorene-9-bisphenol is anti-oestrogenic and may cause adverse pregnancy outcomes in mice. *Nat. Commun.* **8**, 14585
 36. Terasaki, M., Shiraiishi, F., Nishikawa, T., Edmonds, J. S., Morita, M., and Makino, M. (2005) Estrogenic activity of impurities in industrial grade bisphenol A. *Environm. Sci. Technol.* **39**, 3703–3707
 37. Soga, S., Shirai, H., Kobori, M., and Hirayama, N. (2007) Identification of the druggable concavity in homology models using the PLB index. *J. Chem. Inf. Model.* **47**, 2287–2292
 38. Corbeil, C. R., Williams, C. I., and Labute, P. (2012) Variability in docking success rates due to dataset preparation. *J. Comput. Aided Mol. Des.* **26**, 775–786
 39. Kuiper, G., Lemmen, J. G., Carlsson, B., Corton, J. C., Safe, S. H., van der Saag, P. T., van der Brug, B., and Gustafsson, J.-Å. (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* **139**, 4252–4263
 40. Kojima, H., Takeuchi, S., Sanoh, S., Okuda, K., Kitamura, S., Uramaru, N., Sugihara, K., and Yoshinari, K. (2019) Profiling of bisphenol A and eight its analogues on transcriptional activity via human nuclear receptors. *Toxicology* **413**, 48–55
 41. Li, Y., Perera, L., Coons, L. A., Burns, K. A., Tyler Ramsey, J., Pelch, K. E., Houtman, R., van Beuningen, R., Teng, C. T., and Korach, K. S. (2018) Differential *in vitro* biological action, coregulator interactions, and molecular dynamic analysis of bisphenol A (BPA), BPAF, and BPS ligand-ER α complexes. *Environ. Health Perspect.* **126**, 017012
 42. Stauffer, S. R., Coletta, C. J., Tedesco, R., Nishiguchi, G., Carlson, K., Sun, J., Katzenellenbogen, B. S., and Katzenellenbogen, J. A. (2000) Pyrazole ligands: Structure - affinity/activity relationships and estrogen receptor- α -selective agonists. *J. Med. Chem.* **43**, 4934–4947
 43. Gaido, K. W., Leonard, L. S., Maness, S. C., Hall, J. M., McDonnell, D. P., Saville, B., and Safe, S. (1999) Differential interaction of the methoxychlor metabolite 2,2-bis-(p-hydroxyphenyl)-1,1,1-trichloroethane with estrogen receptors α and β . *Endocrinology* **140**, 5746–5753
 44. Gaido, K. W., Maness, S. C., McDonnell, D. P., Dehal, S. S., Kupfer, D., and Safe, S. (2000) Interaction of methoxychlor and related compounds with estrogen receptor alpha and beta, and androgen receptor: Structure-activity studies. *Mol. Pharmacol.* **58**, 852–858
 45. Cavallo, G., Metrangolo, P., Milani, R., Pilati, T., Priimagi, A., Resnati, G., and Terraneo, G. (2016) The halogen bond. *Chem. Rev.* **116**, 2478–2601
 46. Bauzá, A., Mooibroek, T. J., and Frontera, A. (2015) The bright future of unconventional σ/π -hole interactions. *ChemPhysChem* **16**, 2496–2517
 47. Skowron, K. J., Booker, K., Cheng, C., Creed, S., David, B. P., Lazzara, P. R., Lian, A., Siddiqui, Z., Speltz, T. E., and Moore, T. W. (2019) Steroid receptor/coactivator binding inhibitors: An update. *Mol. Cell. Endocrinol.* **493**, 110471
 48. Shapiro, D. J., Mao, C., and Cherian, M. T. (2011) Small molecule inhibitors as probes for estrogen and androgen receptor action. *J. Biol. Chem.* **286**, 4043–4048
 49. Hughes, T. S., Giri, P. K., de Vera, I. M. S., Marciano, D. P., Kuruvilla, D. S., Shin, Y., Blayo, A.-L., Kamenecka, T. M., Burris, T. P., Griffin, P. R., and Kojetin, D. J. (2014) An alternate binding site for PPAR γ 3 ligands. *Nat. Commun.* **5**, 1–13
 50. Staudinger, J. L. (2019) Clinical applications of small molecule inhibitors of Pregnane X receptor. *Mol. Cell. Endocrinol.* **485**, 61–71
 51. Tice, C. M., and Zheng, Y.-J. (2016) Non-canonical modulators of nuclear receptors. *Bioorg. Med. Chem. Lett.* **26**, 4157–4164
 52. Scatchard, G. (1949) The attractions of proteins for small molecules and ions. *Proc. Natl. Acad. Sci. U. S. A.* **51**, 660–672