

Draft Whole-Genome Sequence of *Serratia marcescens* Strain RM66262, Isolated from a Patient with a Urinary Tract Infection

Roberto E. Bruna,^a Santiago Revale,^b Eleonora García Vescovi,^a Javier F. Mariscotti^a

Instituto de Biología Molecular y Celular de Rosario (IBR-CONICET), Departamento de Microbiología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina^a; Instituto de Agrobiotecnología Rosario (INDEAR)-CONICET, Rosario, Argentina^b

E.G.V. and J.F.M. contributed equally to this work.

***Serratia marcescens* strains are ubiquitous bacteria isolated from environmental niches and also constitute emergent nosocomial opportunistic pathogens. Here, we report on the draft genome sequence of *S. marcescens* strain RM66262, which was isolated from a patient with urinary tract infection in the Bacteriology Service of the Rosario National University, Rosario, Argentina.**

Received 13 October 2015 Accepted 17 October 2015 Published 3 December 2015

Citation Bruna RE, Revale S, García Vescovi E, Mariscotti JF. 2015. Draft whole-genome sequence of *Serratia marcescens* strain RM66262, isolated from a patient with a urinary tract infection. *Genome Announc* 3(6):e01423-15. doi:10.1128/genomeA.01423-15.

Copyright © 2015 Bruna et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to Eleonora García Vescovi, garciavescovi@ibr-conicet.gov.ar, or Javier F. Mariscotti, mariscotti@ibr-conicet.gov.ar.

Serratia marcescens is an opportunistic human pathogen associated with urinary and respiratory tract as well as wound and eye infections, endocarditis, osteomyelitis, meningitis, and septicemia. Immunocompromised people and newborns are the most affected hosts. The incidence of *S. marcescens* infection has increased over the last years, mainly due to the acquisition of multiple antibiotic resistances (1–4). Despite its clinical prevalence, the factors and mechanisms that contribute to *Serratia* pathogenesis remain unclear. *S. marcescens* strain RM66262 is a nonpigmented clinical isolate from a patient with urinary tract infection from the Bacteriology Service of the Facultad de Ciencias Bioquímicas y Farmacéuticas of the Rosario National University, Rosario, Argentina (5).

In previous reports from our group, we demonstrated that this strain of *S. marcescens*, using an *in vitro* infection model of nonphagocytic cells, provokes an extracellular induction of autophagy mediated by Sh1A pore-forming toxin, and then is able to internalize, survive, and replicate inside large membrane-bound compartments that recruit prototypical markers of autophagosomes (6–8). Earlier work from our group pointed to the *S. marcescens* Rcs system as a key player in the regulation of the expression of virulence determinants of the bacterium (5, 9). We also found that our strain RM66262 produces outer membrane vesicles (OMVs), which package and deliver a specific cargo of potential virulence determinants (10).

In this study, we present the draft genome sequence of *S. marcescens* strain RM66262. Genomic DNA was isolated from an overnight-grown liquid culture, using the Wizard genomic DNA purification kit (Promega). DNA was quantified using the Ultraspec 2000 spectrophotometer (Pharmacia Biotech). The genome sequence was obtained on an Illumina HiSeq 1500 instrument at INDEAR (Argentina), using a whole-genome shotgun strategy with 2 × 100-bp reads. Overall genome coverage of ~900-fold was obtained. The A5-miseq pipeline (11, 12) was used to perform read trimming and correction, contig assembly, crude scaffolding, misassembly correction, and final scaffolding.

The genome of *S. marcescens* strain RM66262 has 4,882,260 bp, with a total G+C content of 60.1%. Genome annotation was performed using the NCBI Prokaryotic Genome Automatic Annotation Pipeline (PGAAP) (13). The *S. marcescens* strain RM66262 genome has 4,520 genes; among the identified genes 4,366 are protein-coding sequence genes (CDSs), and 53 are pseudogenes. The genome also has 10 rRNA genes (5S, 16S, and 23S) and 79 tRNAs genes. The RAST server (14) predicted coding sequences belonging to 561 subsystems.

The information provided in the genome sequence of *S. marcescens* strain RM66262 will enable further studies to understand the mechanisms of pathogenesis, characterize virulence factors, and analyze gene expression regulation of this opportunistic human pathogen.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number JWLO00000000. The version described in this paper is the first version, JWLO01000000, and consists of sequences JWLO01000001 to JWLO01000019.

ACKNOWLEDGMENT

This work was supported by grant PICT-2012-1403 (to E.G.V.) from Agencia Nacional de Promoción Científica (ANPCyT), Argentina.

REFERENCES

1. Grimont PAD, Grimont F. 1978. The genus *Serratia*. *Annu Rev Microbiol* 32:221–248. <http://dx.doi.org/10.1146/annurev.mi.32.100178.001253>.
2. Hejazi A, Falkiner FR. 1997. *Serratia marcescens*. *J Med Microbiol* 46:903–912. <http://dx.doi.org/10.1099/00222615-46-11-903>.
3. Hejazi A, Auckan HM, Falkiner FR. 2000. Epidemiology and susceptibility of *Serratia marcescens* in a large general hospital over an 8-year period. *J Hosp Infect* 45:42–46. <http://dx.doi.org/10.1053/jhin.1999.0722>.
4. Mahlen SD. 2011. *Serratia* infections: from military experiments to current practice. *Clin Microbiol Rev* 24:755–791. <http://dx.doi.org/10.1128/CMR.00017-11>.
5. Castelli ME, Fedrigo GV, Clementín AL, Ielmini MV, Feldman MF, García Vescovi EG. 2008. Enterobacterial common antigen integrity is a checkpoint for flagellar biogenesis in *Serratia marcescens*. *J Bacteriol* 190:213–220. <http://dx.doi.org/10.1128/JB.01348-07>.

6. Fedrigo GV, Campoy EM, Di Venanzio G, Colombo MI, García Vescovi E. 2011. *Serratia marcescens* is able to survive and proliferate in autophagic-like vacuoles inside non-phagocytic cells. *PLoS One* 6:e24054. <http://dx.doi.org/10.1371/journal.pone.0024054>.
7. Barchiesi J, Castelli ME, Di Venanzio G, Colombo MI, García Vescovi E. 2012. The PhoP/PhoQ system and its role in *Serratia marcescens* pathogenesis. *J Bacteriol* 194:2949–2961. <http://dx.doi.org/10.1128/JB.06820-11>.
8. Di Venanzio G, Stepanenko TM, García Vescovi E. 2014. *Serratia marcescens* ShlA pore-forming toxin is responsible for early induction of autophagy in host cells and is transcriptionally regulated by RcsB. *Infect Immun* 82:3542–3554. <http://dx.doi.org/10.1128/IAI.01682-14>.
9. Castelli ME, Vescovi EG. 2011. The Rcs signal transduction pathway is triggered by enterobacterial common antigen structure alterations in *Serratia marcescens*. *J Bacteriol* 193:63–74. <http://dx.doi.org/10.1128/JB.00839-10>.
10. McMahon KJ, Castelli ME, García Vescovi EG, Feldman MF, García Vescovi E. 2012. Biogenesis of outer membrane vesicles in *Serratia marcescens* is thermoregulated and can be induced by activation of the Rcs phosphorelay system. *J Bacteriol* 194:3241–3249. <http://dx.doi.org/10.1128/JB.00016-12>.
11. Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. An integrated pipeline for de novo assembly of microbial genomes. *PLoS One* 7:e42304. <http://dx.doi.org/10.1371/journal.pone.0042304>.
12. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <http://dx.doi.org/10.1093/bioinformatics/btu661>.
13. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. National Center for Biotechnology Information, Bethesda, MD.
14. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.