

RESEARCH

Open Access



Home treatment and use of informal market of pharmaceutical drugs for the management of paediatric malaria in Cotonou, Benin

Edwige Apetoh^{1,2*}, Marina Tilly¹, Carine Baxerres^{1,3} and Jean-Yves Le Hesran¹

Abstract

Background: Malaria is the main cause of hospital admissions in Benin and a leading cause of death in childhood. Beside consultations, various studies have underlined the management of the disease through home treatment. The medicines used can be purchased in informal market of pharmaceutical drugs (IMPD) without prescription or any involvement of healthcare professional. Pharmaceutical drugs are sold by informal private vendors, who operate at any time in the immediate environment of the patients. The present study was conducted in Cotonou to study the health-seeking behaviour of caregivers to treat malaria in children under 12 years old. Factors associated with malaria home treatment and drugs purchase in IMPD were studied.

Methods: A cross-sectional study was carried out among 340 children's caregivers who were interviewed about their socio-demographic characteristics and their care-seeking behaviour during the most recent episode of malaria in their children under 12. Medicines used and purchase place were also collected. Multivariate logistic regression model was used to determine factors associated with malaria home treatment and drug purchase in IMPD.

Results: Beyond all the 340 caregivers, 116 (34%) consulted healthcare professional, 224 (66%) home treat the children, among whom 207 (61%) gave pharmaceutical drugs and 17 (5%) gave traditional remedies to children. Malaria home treatment was associated with family size, health insurance (OR = 0.396, 95% CI 0.169–0.928), and wealth quintiles where home treatment was less used by the richest (OR = 0.199, 95% CI 0.0676–0.522) compared to those in the poorest quintile. The caregivers age group 30–39 years was associated to the use of IMPD (OR = 0.383, 95% CI 0.152–0.964), the most economically wealthy people were less likely to use IMPD (wealth quintile richest: OR = 0.239, 95% CI 0.064–0.887; wealth quintile fourth OR = 0.271, 95% CI 0.100–0.735) compared to those in the poorest quintile. All caregivers who benefited from health insurance did not use IMPD.

Conclusion: This study highlights the link between worse economic conditions and accessibility to medical care as one of the main factors of malaria home treatment and drug purchase in IMPD, even if those two phenomena need to be understood apart.

Keywords: Benin, Cotonou, Home treatment, Informal market of pharmaceutical drugs, Malaria, Parallel market, Paediatric malaria, Self-medication

*Correspondence: eapetoh@gmail.com

¹ Institut de recherche pour le développement, Unité mixte de recherche 216: Mères et enfants face aux infections tropicales, Université Paris-Descartes, 4 Avenue de l'Observatoire, 75006 Paris, France
Full list of author information is available at the end of the article



Background

Malaria is the leading cause of morbidity among adults, and mortality among children under five in Benin. It accounted for 40% of outpatient consultations and 25% of hospital admissions in 2016 [1]. In the context of the extended resistance to chloroquine and sulfadoxine–pyrimethamine (SP), the malaria management policy in Benin has been modified in 2004 by introducing into the national malaria coordination programme the use of artemisinin-based combination therapy (ACT) to treat non-complicated malaria [2]. This change was made 2 years before the official recommendation of the World Health Organization (WHO) to most sub-Saharan African countries [3]. In 2011, in Benin recommended malaria diagnosis [rapid diagnostic test (RDT) or microscopy] for all patients, as mandated by the diagnostic guidelines, and treatment being given only to patients with a positive test result [4].

Early diagnosis and prompt treatment hugely reduces progression of the illness to severe state and prevents death [5]. However, in Benin like in other sub-Saharan Africa countries, caregivers first manage malaria in children at home, and when the symptoms persist a health-care professional is consulted [6–9].

In Benin, malaria RDT is performed only after consulting health-care professional. Thus, home treatment of malaria in children is based on perceived symptoms by caregivers. It is well recognized that home treatment of malaria presents risk of false diagnosis and increased delay in presentation at health-care centres [6]. Home treatment was operationally defined as the administration of treatment (pharmaceutical drugs or traditional remedies) at home by the caregiver on his own initiative or on the advice of a relative, as opposed to treatment recommended by health-care professional on the present episode of disease [10]. Home treatment risks are increased in environments where people have access to a wide range of pharmaceutical drugs, even for drugs normally requiring prescription by a healthcare professional. Drugs can be bought in private or public formal sectors (pharmacies, health centres, private warehouses), but also in informal market of pharmaceutical drugs (IMPD). The selling of informal pharmaceutical drugs is a widespread practice since at least the 1980's in several African countries, particularly in the French-speaking ones [11–13].

Informal refers to activities involving the sale and purchase of drugs outside the legislative and administrative framework imposed by a country's government and biomedical health system. Pharmaceutical drugs are sold by informal private vendors, who operate as shop keepers, traders, itinerant drug sellers, and drug wholesalers [14–16]. This informal practice is taking place in a

context marked in part by the lack of rigidity in executive, administrative or other measures, including penalties against sellers and distributors [17, 18]. In Benin, the procedures against IMPD have been mentioned in the national pharmaceutical policy [19]. On a related subject, a national counterfeit medicines committee was created in 2012 and actions, including awareness campaigns and repression through the seizure and destruction of drugs on the informal market were conducted. Since February 2017, an unprecedented repression was organized by the president of the Republic of Benin.

The last quantitative survey on the use of IMPD was conducted in Cotonou, the biggest town of Benin, in 2003, 15 years ago. The survey aimed to determine the percent of the population which at least once have purchased pharmaceutical drugs in IMPD. The percent reported was 72%, all diseases taken into account [20].

Cotonou is the city in which IMPD is mostly present due to the biggest West African international market named “Dantokpa”, where drugs are sold wholesale and retail [16]. In the order to describe malaria management in households in this environment, a household cross-sectional study was conducted. The study aimed to (i) describe the management of the most recent episode of malaria among children under twelve, (ii) document the pharmaceutical drugs used by caregivers to treat children, (iii) describe factors associated with malaria home treatment, and (iv) describe factors associated with the use of IMPD when home treatment was practiced.

Methods

Study area

This study was carried out before the 09th edition of Operation PANGEA IX in 2017, coordinated by Interpol with a view to curbing the informal circuit of pharmaceutical drugs in Benin. The study was conducted in Cotonou, the economic capital and biggest city of Benin. The country is divided in twelve departments and Cotonou forms the department of Littoral on its own. The general population of Cotonou was estimated at 679,012 in 2013, with 211,192 children under 12 years old [21]. Cotonou is a coastal city, limited to the west by the Atlantic department and to the east by Ouémé department. The Atlantic ocean is its southern boundary and the Nokoué lake is its northern boundary. This geographic situation makes malaria endemic in the Cotonou area. The primary malaria vector is *Anopheles gambiae* sensu stricto [4]. Severe malaria was the leading cause of death (24%) among children under 5 in hospitals in 2016 [22].

In Benin, the use of ACT to treat mild malaria was recommended by the health ministry of Benin since 2004 [2]. The National Malaria Control Programme of Benin recommended in 2011 new guidelines for effective

malaria prevention and treatment [4]. Thus, patients of all ages should receive a confirmed malaria diagnosis [with either a rapid diagnostic test (RDT) or microscopy] before receiving treatment. Treatment is free of charge for pregnant women and children under 5 attending public health facilities, and the ACT subsidy system has been implemented in the public sector for other categories of patients. The subsidized ACT, artemether and lumefantrine, can be made available through different trade names, including Coartem® or Lumartem®. Prevention strategies involve the use of insecticide-treated mosquito nets (ITNs), indoor residual spraying (IRS), and intermittent preventive treatment of pregnant women (IPTp).

Study population

Caregivers were identified in households and asked about the treatment-seeking behaviour for the most recent malaria episode in a child under 12 years under their care. As self-medication is virtually non-existent in this age group, caregivers have full picture of pharmaceutical drugs consumed by children. This hypothesis was consolidated by a previous study. Geissler et al. [23] studied self-medication practice among schoolchildren aged 11–17 years in western Kenya, and reported that self-medication decreases with a younger age. Twenty percent of the 15–17 years old were practicing self-medication without parental or adult involvement. The prevalence was 14% among the 13–14 years old, and 9% among the 11–13 years old.

Study design and sample size

A cross-sectional study was conducted in Cotonou, during April–May 2016. The numbers of caregivers to interview was calculated by the Schwartz’ formula [24]:

$$n = \frac{(z^2)(\hat{p})(1 - \hat{p})}{e^2},$$

where *z* represents the z-score fixed at 1.96; *e* indicates the desired margin of error fixed at 0.05 [25], and \hat{p} the proportion of children under 12 which have already contracted malaria. This latter proportion was unknown, and was fixed to 50% in order to maximize the sample size [26]. A total of 384 caregivers of children under 12 should be included in the study.

Because of the absence of relevant data to perform a simple random sampling, a GPS sampling method was used to access caregivers. The sampling method consisted of random selection of a calculated number of GPS points into the study area. Buildings identified by GPS points were visited for the survey. Every GPS point may not necessarily identify a household because of empty spaces (park, field, swamps, or puddles) and uninhabited

buildings (commercial, company or administrative buildings, buildings under construction or abandoned). Simulations performed determined that two-thirds of GPS points will identify a building. Considering that 1 out of 2 households would include at least one child less than 12 years of age, twice as many households should be visited. The number of GPS points to randomize to reach 384 caregivers of children under 12 years old was calculated with the formula:

$$n_{GPS} = n \times 1/p_{child} \times 1/p_{building} \times 1/p_{non-response},$$

where *n* was the calculated number of children to include in the study, *p_{child}* the proportion (1/2) of households with at least one child under 12, *p_{building}* the proportion (2/3) of GPS points which identifies a building, and *p_{non-response}* the assumed proportion (1/10) of non-response [26] *n_{GPS}* was 1300.

A total of 421 households contained at least a child under 12 and caregivers were interviewed about the most recent episode of malaria in a child under their care. A total of 340 (81%) caregivers were able to provide information on the health-care seeking behaviour and were included in the study. Forty-nine (12%) declared that the child has never contracted malaria, 23 (5%) declared that they practised prevention, so that children had not contracted malaria in the recent past, and 9 (2%) caregivers declared they did not remember.

Collected data

Questionnaire for data collection was translated into the local dialect “fon”, the most common spoken in Cotonou by non-French speakers. In the households, caregivers were interviewed. A questionnaire was administered by investigators who graduated in sociology with substantial knowledge of the field surveys.

Questions were about the health-seeking behaviour during the most recent malaria episode among their children under 12 in the household. The episode date was collected and caregivers were asked to cite any symptoms which led them to recognize that the child had malaria. Afterward, the first action done by caregivers after they recognized symptoms were collected. Action could then be either home treatment or consultation with a healthcare professional. Information were collected on the type of treatments used at home or prescribed (pharmaceutical drugs or traditional remedies), treatment name and galenic form of pharmaceutical drugs. When home treatment was practised, the purchase place of pharmaceutical drugs was collected. Purchase places were categorized into IMPD and formal market.

A first binary outcome variable was home treatment of malaria, defined in this paper as treatment

(pharmaceutical drugs or traditional remedies) administration at home by the caregiver on his own initiative or on the advice of a relative, as opposed to treatment recommended by health-care professional. For this purpose, used leftover medicines having been advised by pharmaceutical drugs vendors (pharmacist or informal vendor) were considered as home treatment.

A second binary outcome variable was drug purchase in IMPD or in formal market when home treatment was practised. Independent variables were, age, education level, marital status, religion, health insurance, family size, and presence of relatives living in the same inner court. A set of 17 variables on the household possession and housing characteristics (nature of habitat, access to resources and property owned by the household) were collected and facilitated the construction of a wealth index by using principal component analyses, like in the demographic and health survey (DHS) [27]. The wealth index was categorized into quintiles in ascending order of poorest, fourth, middle, second, and richest. However, the variables used in this study were not exactly similar to those of the demographic and health survey, and, therefore, the wealth quintile used in this study cannot strictly be compared to DHS data.

Data analysis

Data were analysed using STATA version 13 statistical software (Stata Corp., TX USA). Descriptive analyses were performed to characterize socio-demographic aspects of caregivers, and management of the disease. Median and length were calculated for continuous variables and percentages for categorical variables. To assess the significance of differences between proportions the Chi square test or Fisher's exact test, were used as appropriate.

Forgetting the name of drugs used over time can be a consequence of a memory time-related bias. To identify any possible memory time-related bias, the date of the most recent episode was quantified and grouped by months, and Chi square tests were conducted based on information related to the forgetfulness of the used treatments names, and secondly with therapeutic classes of cited drugs.

Bivariate logistic regression analyses were performed on the one hand with home treatment as the outcome, and in the other hand with drug purchase in IMPD as the outcome. For each outcome, all significant variables in the bivariate analyses at $p < 0.20$ were included in a multivariate logistic regression model. Odds ratios (OR) were reported with 95% confidence intervals (CI). Statistical significance was set at $p < 0.05$.

Ethical considerations

The study has been approved by the Ministry of Higher Education and Scientific Research of Benin; Ethics Committee CER-ISBA –FAVORABLE ADVICE N° 30. Each interviewed person was informed about the objectives of the study, the types of collected data and were asked to sign a document of informed consent.

Results

Among the 340 caregivers, 27% reported that children had malaria during the month preceding the interview, 36% had malaria in the last 1–3 months, 15% within the last 4–6 months, and 22% declared that the most recent episode dates back more than 6 months. The median time was 2 months. There was no evidence for memory time-related bias; neither for forgetfulness of the used treatment names, nor for the medicines therapeutic class (see Additional file 1). Based on these results, all data were included in subsequent analyses.

Socio-demographic characteristics of caregivers

Table 1 presents the socio-demographic characteristics of caregivers and data on the management of the disease. Of the 340 interviewed caregivers, 75% were female and 80% were in a marital or non-marital relationship. Median age was 38 (range 18–83 years), 23% reported having received no formal education, 53% had primary education, 38% had attended secondary education and 9% had tertiary level education. Most caregivers were Christian (72%). Regarding family size, 28% of caregivers lived in households composed of at least three persons (four persons was the average size of households in Cotonou, according to the last census [21]), and 57% lived in households composed of 4–8 persons and 15% up to 8 persons. Thirty-one percent of the families lived in an inner court together with relatives. Thirty-eight percent of caregivers belonged to the upper third quintile, which characterizes the most economically wealthy people. Only 9% of caregivers reported that they subscribed to a health insurance. Seventy-nine percent of caregivers who brought their children to a healthcare professional reported that RDT or microscopy was done to confirm malaria.

Treatment-seeking behaviour and illness history

Table 2 presents the malaria health-seeking behaviours of caregivers and the symptoms in the child they recognized as malaria symptoms. To treat malaria, 116 (34%) caregivers indicated they brought the child to a healthcare professional, whereas 224 (66%) home-treated children. In the case of home-treatment, 203 (90%) reported they gave only pharmaceutical drugs to their child, while 17 (8%) reported they gave only herbal remedies. Only 4 (2%) caregivers declared giving

Table 1 Socio-demographic characteristics of caregivers of under twelve children in Cotonou, Benin, May 2016 (N = 340)

Categories	Subcategories	Frequencies (n)	Percentages (%)
Age group (years)	< 30	63	18
	30–39	128	38
	40–49	68	20
	≥ 50	81	24
Gender	Female	255	75
	Male	85	25
Marital status	In relationship	273	80
	Single	67	20
Education	None	79	23
	Primary	100	53
	Secondary	130	38
	Tertiary	31	9
Religion	Christian	246	72
	Islam	47	14
	Other	47	14
Family size	< 4	38	28
	4–8	252	57
	≥ 9	50	15
Relative in the inner court	No	233	69
	Yes	107	31
Health insurance	No	310	81
	Yes	30	9
Wealth quintiles	Quintile 1 (richest)	57	17
	Quintile 2 (fourth)	70	21
	Quintile 3 (middle)	68	20
	Quintile 4 (second)	75	22
	Quintile 5 (poorest)	70	21

both at the same time. As these later caregivers gave pharmaceutical drugs to the child, they were analysed together with those who used pharmaceutical drugs.

When asked about the symptoms that made them think that the child has malaria, at least five percent of caregivers reported fever (89%), loss of appetite (20%), vomiting (19%), weakness (17%), chills (10%), headache (9%), coughing (10%), and dark urine (6%). Forty-two percent of the caregivers cited only one symptom (30% fever), indicating they relied on a single symptom to conclude the child had malaria, 39% relied on two symptoms, and 19% on more than two symptoms. For the remainder of the analyses, cited symptoms were grouped into seven categories. The first category was “fever, headache, dizziness”, the second includes “body aches, chills, weak”, the third “bitter taste in the mouth, no appetite”, the fourth “vomiting, diarrhoea”, the fifth “dark urine, yellow eyes”, the sixth “cough”, and the last category regroups other symptoms which are rarely mentioned (“prickly, sores in the mouth, anaemia”).

Factor associated with the malaria home treatment

Table 3 presents factors associated with malaria home treatment in bivariate and multivariate analyses. In bivariate analyses, caregivers who practised home treatment were significantly different from those who did not, in terms of age, education, family size, possession of health insurance, wealth quintiles and the fact that the identified symptoms were either “fever, headache, dizziness” and/or “vomiting, diarrhoea”.

In multivariate analysis, caregivers who lived in a family composed of more than eight persons were more likely to home treat malaria in children, but the difference was only statistically significant when compared to family size in the 4–8 group (OR = 0.399, 95% CI 0.180–0.885). When caregivers had health insurance, they were less likely to home treat children (OR = 0.396, 95% CI 0.169–0.928) compared to those who did not have it. Caregivers belonging to the richest wealth quintiles were less likely to treat children at home (OR = 0.199, 95% CI 0.0676–0.522) compared to those in the poorest quintile. Home

Table 2 Health-seeking behaviours of caregivers to treat malaria in children under twelve and malaria symptoms recognized in Cotonou, Benin, May 2016 (N = 340)

Categories	Subcategories	Frequencies (n)	Percentages (%)
Treatment-seeking	Healthcare professional	116	34
	Home treatment with pharmaceutical drugs	207	61
	Home treatment with traditional remedies	17	5
Fever, headache, dizziness	No	30	8
	Yes	310	91
Body aches, chills, weak	No	248	73
	Yes	92	27
Bitter taste in the mouth, no appetite	No	268	79
	Yes	72	21
Vomiting, diarrhoea	No	265	78
	Yes	75	22
Dark urine, yellow eyes	No	312	92
	Yes	28	8
Cough	No	310	91
	Yes	30	9
Other: prickly, sores in the mouth, anaemia	No	328	96
	Yes	12	4
Number of symptoms (md[length] = 2 [1,6])	1	141	42
	2	134	39
	≥ 3	65	19
Use of IMPD for home treatment with pharmaceutical drugs ^a (n = 207)	No	123	59
	Yes	84	41

^a Based on the number (n = 207) of caregivers who practised home treatment with pharmaceutical drugs

treatment were less practised by those who reported “vomiting, diarrhoea” (OR = 0.374, 95% CI 0.196–0.712) than those who reported other symptoms.

Factors associated with the medicines purchased in IMPD

Table 4 presents factors associated with the use of IMPD in bivariate and multivariate analyses. It is important to underline that any caregiver who declared to have health insurance did not buy drugs in IMPD in this study.

In bivariate analysis, caregivers who purchased medicines in IMPD significantly differed from those who went to formal market in terms of age, education, wealth quintiles, and when reporting “fever, headache, dizziness”, “vomiting, diarrhoea”.

In multivariate analyses, caregivers belonging to age groups 30–39 (OR = 0.383, 95% CI 0.152–0.964) and economically wealthy people (wealth quintile richest: OR = 0.239, 95% CI 0.064–0.887; wealth quintile fourth OR = 0.271, 95% CI 0.100–0.735) were less likely to purchase drug in IMPD compared to those in the poorest quintile. When reporting “fever, headache, dizziness”, caregivers were more likely to go to IMPD.

Types of drugs cited

Sixty-seven (20%) caregivers reported they forgot the name of the drugs used, 45 (38%) of whom consulted a healthcare professional, and 22 (11%) administered home treatment. Table 5 presents medicines reported by the 256 caregivers when a healthcare professional was consulted versus home treatment, and Table 6 presents location of treatment purchase in the case of home treatment.

Different therapeutic classes were cited. Seventy-one percent of caregivers cited anti-malarial drugs, 61% antipyretics (analgesics or anti-inflammatory drugs), 16% antibiotics, 13% supplements (vitamins or anti-anaemics), and 6% anthelmintic drugs. Other therapeutic classes, including decongestants, cough or cold drugs, antidiarrhoeal drugs and antihistamines were also mentioned, but rarely. A total of 92 different trade names of pharmaceutical drugs were cited. The most cited drugs (by at least 5% of caregivers) were Paracetamol (44%), Quinine (32%), Efferalgan® 10%, Amoxicillin (9%) and Lufanter® (6%), which is the most cited ACT. Coartem® (4 home treatments and 2 prescriptions) and Lumartem (2 home treatments and 1 prescription) were rarely mentioned.

Table 3 Factors associated with the practice of home treatment, in Cotonou, Benin, May 2016 (N = 340)

Covariates	Categories	Subcategories	Home treatment n (%)	Total n	Bivariate			Multivariate		
					OR	95% IC	p-value	OR	95% IC	p-value
Age		< 30	40 (18)	63	1			1		
		30–39	87 (39)	128	1.220	0.647–2.298	0.538	1.382	0.678–2.815	0.372
		40–49	45 (20)	68	1.125	0.548–2.306	0.748	1.057	0.479–2.328	0.890
		≥ 50	52 (23)	81	1.031	0.519–2.045	0.930	0.959	0.447–2.054	0.915
Gender		Female	168 (75)	255	1					
		Male	56 (25)	85	1	0.595–1.678	1.000			
Marital status		Single	41 (18)	67	1					
		In relationship	183 (82)	273	1.354	0.782–2.341	0.278			
Education		None	58 (26)	79	1			1		
		Primary	68 (30)	100	0.769	0.400–1.477	0.431	0.699	0.236–2.069	0.518
		Secondary	79 (35)	130	0.560	0.304–1.033	0.064	0.627	0.230–1.709	0.362
		Tertiary	19 (9)	31	0.573	0.238–1.379	0.214	0.535	0.206–1.385	0.198
Religion		Christian	164 (73)	246	1					
		Islam	27 (12)	47	0.675	0.357–1.275	0.226			
		Other	33 (15)	47	1.178	0.597–2.324	0.635			
Family size		< 4	25 (11)	38	0.542	0.210–1.398	0.205	0.396	0.139–1.125	0.082
		4–8	160 (72)	252	0.490	0.239–1.004	0.051	0.399	0.180–0.885	0.024
		9–22	39 (17)	50	1					
Relative in the inner court		No	154 (69)	233	1					
		Yes	70 (31)	107	0.970	0.599–1.571	0.903			
Health insurance		No	211 (94)	310	1			1		
		Yes	13 (6)	30	0.358	1.677–0.767	0.008	0.396	0.169–0.9284	0.033
Wealth quintiles		Quintile 5 (poorest)	57 (25)	70	1			1		
		Quintile 4 (second)	50 (22)	75	0.456	0.211–0.985	0.046	0.470	0.207–1.065	0.071
		Quintile 3 (middle)	46 (21)	68	0.476	0.211–0.985	0.066	0.428	0.184–0.994	0.049
		Quintile 2 (fourth)	44 (20)	70	0.385	0.178–0.836	0.016	0.472	0.198–1.121	0.089
		Quintile 1 (richest)	27 (12)	57	0.205	0.092–0.454	0.000	0.199	0.076–0.522	0.001
Fever, headache, dizziness		No	23 (10)	30	1			1		
		Yes	201 (90)	310	0.561	0.233–1.349	0.197	0.477	0.176–1.286	0.144
Body aches, chills, weak		No	163 (73)	248	1					
		Yes	61 (27)	92	1.026	0.618–1.701	0.920			
Bitter taste in the mouth, no appetite		No	182 (81)	268	1					
		Yes	42 (19)	72	0.725	0.421–1.246	0.245			
Vomiting, diarrhoea		No	187 (83)	265	1			1		
		Yes	37 (17)	75	0.406	0.240–0.685	0.001	0.374	0.196–0.712	0.003
Dark urine, yellow eyes		No	205 (92)	312	1					
		Yes	19 (8)	28	1.101	0.482–2.519	0.818			
Cough		No	207 (92)	310	1					
		Yes	17 (8)	30	0.650	0.304–1.391	0.268			
Other (prickly, sores in the mouth, anaemia)		No	217 (89)	328	1					
		Yes	7 (3)	12	0.716	0.222–2.307	0.576			
Number of symptoms					0.691	0.534–0.894	0.005	0.864	0.629–1.186	0.368

Statistically significant p values are in italic

Home treatment versus prescription by a healthcare professional

Caregivers who brought their child to a healthcare professional more frequently reported the use of anti-malarial

drugs (89%), compared to those who home treat children (65%) ($p < 0.001$). According to the type of the anti-malarial drug used, ACT was used in the same proportion in the case of home treatment and prescription (30%),

Table 4 Factors associated with drug purchase in IMPD for home treatment (N = 207) in Cotonou, Benin, May 2016

Covariates	Categories	Subcategories	IMPD n (%)	Total n	Bivariate			Multivariate		
					OR	95% IC	p-value	OR	95% IC	p-value
Age		< 30	21 (25)	36	1			1		
		30–39	27 (33)	80	0.363	0.162–0.816	0.014	0.383	0.152–0.964	0.042
		40–49	18 (21)	44	0.494	0.202–1.209	0.123	0.517	0.190–1.1403	0.195
		≥ 50	18 (21)	47	0.443	0.182–1.075	0.072	0.470	0.170–1.298	0.145
Gender		Female	60 (71)	154	1					
		Male	24 (29)	53	1.296	0.690–2.435	0.419			
Marital status		Single	15 (18)	37	1					
		In relationship	69 (82)	170	1.001	0.485–2.067	0.996			
Education		None	29 (35)	52	1			1		
		Primary	26 (31)	61	0.589	0.279–1.246	0.165	0.726	0.305–1.729	0.470
		Secondary	25 (30)	75	0.396	0.191–0.821	0.013	0.563	0.242–1.312	0.184
		Tertiary	4 (5)	19	0.211	0.061–0.724	0.013	0.419	0.094–1.857	0.253
Religion		Christian	59 (70)	153	1					
		Islam	11 (13)	26	1.168	0.502–2.715	0.718			
		Other	14 (17)	28	1.593	0.709–3.578	0.259			
Family size		< 4	6 (26)	21	1					
		4–8	64 (76)	150	1.860	0.684–5.059	0.224			
		> 8	14 (17)	36	1.590	0.498–5.074	0.433			
Relative in the inner court		No	56 (67)	143	1					
		Yes	28 (33)	64	1.208	0.665–2.195				
Wealth quintiles		Quintile 5 (poorest)	30 (36)	48	1			1		
		Quintile 4 (second)	23 (27)	45	0.142	0.274–1.433	0.269	0.860	0.350–2.108	0.742
		Quintile 3 (middle)	15 (18)	44	0.200	0.132–0.729	0.007	0.500	0.198–1.261	0.142
		Quintile 2 (fourth)	11 (13)	44	0.310	0.814–0.491	0.000	0.271	0.100–0.735	0.010
		Quintile 1 (richest)	5 (6)	26	0.627	0.045–0.445	0.001	0.239	0.064–0.887	0.032
Fever, headache, dizziness		No	2 (2)	18	1			1		
		Yes	82 (98)	189	6.130	1.370–27.417	0.018	9.384	1.807–48.730	0.008
Body aches, chills, weak		No	62 (74)	151	1					
		Yes	22 (26)	56	0.928	0.496–1.738	0.817			
Bitter taste in the mouth, no appetite		No	65 (77)	168	1					
		Yes	19 (23)	39	1.505	0.747–3.032	0.252			
Vomiting, diarrhoea		No	67 (80)	176	1			1		
		Yes	17 (20)	31	1.975	0.914–4.266		2.606	0.928–7.315	0.069
Dark urine, yellow eyes		No	79 (94)	191	1					
		Yes	5 (6)	16	0.644	0.215–1.927	0.432			
Cough		No	78 (93)	191	1					
		Yes	6 (7)	16	0.869	0.303–2.490				
Other: prickly, sores in the mouth, anaemia		No	82 (98)	202	1					
		Yes	2 (2)	5	0.975	0.159–5.967	0.979			
Number of symptoms					1.419	0.968–2.078	0.072	1.577	0.725–1.846	0.539

Statistically significant p values are in italic

IMPD Informal market of pharmaceutical drugs

while quinine was more often prescribed ($p < 0.001$) and chloroquine was more often used for home treatment (< 0.001).

A higher percentage of caregivers who treated children at home gave antipyretics (68%) than those who brought

their child to a healthcare professional (42%) ($p < 0.001$). Antibiotics were more often cited when healthcare professionals were consulted (25%) than when home treatment was used (13%) ($p < 0.05$).

Table 5 Pharmaceutical drugs used by caregivers to treat malaria in children under 12 in Cotonou, Benin, May 2016 (N = 256): home treatment versus healthcare professional

Variates		Healthcare professional	Home treatment	Chi square test
Categories	Subcategories	n = 71 (%)	n = 185 (%)	p-value
Therapeutic classes	Anti-malarial	63 (89)	120 (65)	<i>0.000</i>
	Antipyretic	30 (42)	125 (68)	<i>0.000</i>
	Anthelmintic	4 (6)	11 (6)	0.924
	Antibiotic	18 (25)	24 (13)	<i>0.017</i>
	Supplements	13 (18)	26 (14)	0.828
Anti-malarial	ACT	21 (30)	55 (30)	0.980
	SP	2 (3)	7 (4)	0.707
	Quinine	42 (59)	45 (24)	<i>0.000</i>
	Chloroquine	0 (0)	11 (6)	<i>0.036</i>
	Other anti-malarial	1 (1)	4 (2)	0.696
Galenic form	Tablet/pill	28 (39)	167 (90)	<i>0.000</i>
	Injection/infusion	39 (55)	1 (1)	<i>0.000</i>
	Syrup	25 (35)	29 (16)	0.019
	Other	7 (10)	7 (4)	0.157
Number of different therapeutic classes	1	22 (31)	84 (45)	<i>0.036</i>
	2	25 (35)	77 (42)	0.348
	≥ 3	24 (34)	24 (13)	<i>0.000</i>

Statistically significant p values are in italic

ACT artemisinin-based combination therapy, SP sulfadoxine–pyrimethamine

Table 6 Purchase place of drugs used by caregivers who practised home treatment to treat malaria in children under 12 (N = 185)

Variates		Formal Market	IMPD	Chi square
Categories	Subcategories	n = 118 (%)	n = 67 (%)	p-value
Therapeutic classes	Anti-malarial	81 (69)	39 (58)	0.153
	Antipyretic	71 (60)	54 (81)	<i>0.004</i>
	Anthelmintic	4 (3)	7 (10)	0.101
	Antibiotic	13 (11)	11 (16)	0.293
	Supplements	10 (8)	16 (24)	<i>0.004</i>
Anti-malarial	ACT	44 (37)	11 (16)	<i>0.003</i>
	SP	6 (5)	1 (1)	0.425
	Quinine	24 (20)	21 (31)	0.094
	Chloroquine	3 (3)	8 (12)	<i>0.019</i>
	Other anti-malarial	4 (3)	0 (0)	0.298
Galenic form	Tablet/pill	101 (76)	66 (93)	<i>0.003</i>
	Injection/infusion	1 (1)	0 (0)	1
	Syrup	25 (19)	4 (6)	<i>0.010</i>
	Other	6 (5)	1 (1)	0.425
Number of different therapeutic classes	1	60 (51)	24 (36)	<i>0.048</i>
	2	46 (39)	31 (46)	0.334
	≥ 3	12 (10)	12 (18)	0.132

Statistically significant p values are in italic

IMPD Informal market of pharmaceutical drugs, ACT artemisinin-based combination therapy, SP sulfadoxine–pyrimethamine

The most frequently therapeutic class used was anti-malarials when a healthcare professional had been seen ($p < 0.001$), and antipyretics in case of home treatment ($p < 0.001$). Concerning therapeutic classes used together, the pair anti-malarials-antipyretics was the most frequently used in home treatment, whereas anti-malarials-antibiotics was the most frequently prescribed. Concerning the galenic form, parenteral administration ($p < 0.001$) and syrup ($p < 0.05$) were used more in prescriptions, and tablet ($p < 0.001$) was preferred in the case home treatment.

IMPD versus formal market

The places of purchase were recorded only in the case of home treatment, and not after consulting a healthcare professional. Table 5 presents data on locations provided by 185 caregivers who bought a pharmaceutical treatment. Various commercial drug names were cited, underlining the availability of a wide variety of pharmaceutical drugs available for purchase without prescription (see Additional file 2 for the name of pharmaceutical drugs cited).

There was no difference in the use of anti-malarials in relation to the purchase place. However, ACT ($p < 0.05$) was more often purchased in formal market, whereas chloroquine ($p < 0.02$) was more often purchased in IMPD. When drugs were used alone, anti-malarials ($p < 0.001$) was the most often purchased in the formal market. Caregivers who went to IMPD, were more likely to purchase more than two different therapeutic classes of drugs ($p < 0.01$). Concerning the galenic form, tablets ($p < 0.01$) were purchased more frequently in IMPD, and syrup ($p < 0.05$) in the formal market.

Discussion

While home treatment is common it carries risks, whether individuals use medicines inappropriately, or whether its practice increases the delay in consulting a healthcare professional to receive appropriate care [28, 29]. Prompt diagnosis and treatment is the most effective means of preventing a mild case of malaria from developing into severe disease with increased risk of death [30]. In sub-Saharan African countries, home treatment has been widely documented as a common practice to treat illnesses, including malaria [13, 16, 31, 32]. Aside from the delay and the inappropriate use of medicines, home treatment presents risk in environment in which people can purchase medicines that are not subject to any government control (such as in the IMPD). This study sought to characterize the practice of malaria home treatment in children under 12 years old, and the use of the IMPD among caregivers who home treat malaria.

Malaria home treatment

In this study, the prevalence of malaria home treatment (66%) is fairly close to the one previously reported (69.7%) by Agueh et al. [31] who considered it across all ages in Comé, a semi-rural area, 80 km far from Cotonou. This similarity is surprising. It would have been expected that home treatment was less frequent when treating malaria in children, knowing that on the one hand malaria treatment is free of charge for children under the age of five, and on the other hand the living environment were different in the two areas in terms of the health-care structures availability. However, as malaria is endemic in Benin, the similar prevalence can result from the fact that symptoms associated with malaria are very common and that people believe to know how to treat them. Consequently home-treatment is first used before any other action, including consulting health-care professional [33]. However, when considering the factors associated with malaria home treatment in this study (i.e. more than 8 members in the family, not having health insurance, belonging in the poorest quintile), the problem seems to rather be the lack of financial means prevailing in both rural and urban areas. Indeed, Agueh et al. also reported that the poorest quintile was associated with the highest home treatment rate. These results confirm those reported by several other studies showing that economic hardship was one of the main reasons cited by persons who home treated, both for malaria or other diseases [20, 34].

Following the Bamako Initiative, Benin, like other African countries, introduced out-of-pocket payments for health [35]. Healthcare that was hitherto free, became payable by the population. In Benin, it has been reported that 75% to 80% of households directly pay medical fees, and 76% of health spending is spent on pharmaceuticals and other medical goods [36]. This can cause households to incur catastrophic expenditures, which in turn can push them into poverty [37]. As this study shows that only very few caregivers declared to possess health insurance, obviously the financial expenditures increase with the number of persons in a family without health insurance, and could explain why the size of the household is associated with the practice of home treatment.

The proportion of caregivers who declared to have health insurance was very low (9%), even though health insurances have been operating in Benin since the 1990s [38], mainly for salaried employees, officials and army personnel, but not for informal workers. Unfortunately, health insurance's contribution in relieving the financial burden on households remains low [38]. The majority covers only small risks (primary curative consultation, pre/post-natal consultation, normal childbirth, essential

drugs) that are supported at the level of dispensaries, district health centres and communal health centres. In 2008, the health ministry launched the process of setting up a Universal Health Insurance Plan (Régime d'Assurance Maladie Universelle, RAMU), which would offer effective protection against the risk of illness to every Beninese whatever his social condition [35]. However, this project did not succeed, and was replaced in 2016 by the Insurance for Strengthening Human Capital [Assurance pour le Renforcement du Capital Humain (ARCH)] which will have to identify the essential dimensions of the Universal Health Coverage recommended by the WHO [36]. The fact that people who got health insurance practised less home treatment than those who got none, should encourage the government of Benin in the effective implementation of the ARCH.

Caregivers who declared that symptoms exhibited by the child were “vomiting, diarrhoea” practised home treatment less frequently. Diarrhoea was poorly known as a malaria symptom and was rarely mentioned in Tanzania [39] and Nigeria [40]. Furthermore, the proportions of respondents who perceived vomiting as a symptom of malaria varies across studies, from low (8.8%) in Lagos, Nigeria among pregnant women [41], to varying from 17.6% to 32.4% in households in four selected districts of the Central African Republic [42]. The fact that, caregivers who declared “vomiting, diarrhoea” practised home treatment less frequently might suggested that these symptoms were not always considered as symptoms malaria symptoms in also this study, thus necessitate referral to a health-care professional for a differential diagnosis.

Caregivers who home treat children more often remembered the name of medicines they used. This raises the question whether caregivers were being careful with prescriptions made by a healthcare professional. A study conducted on the Kenya coast reported that, among mothers who bring children to a healthcare professional, about 55% of them did not understand their prescription, yet none asked for clarification as they believed that health workers had no time for such explanations, and could become harsh when asked too many questions [43].

Informal market of pharmaceutical drugs

The prevalence of the use of IMPD when home treatment was practised (41%) is lower in this study than in the study of Come (83.7%). This can be explained by environmental and economic factors. Results show that economic factors are associated with IMPD use. People from rural areas are generally poorer than those from urban areas. In the department Mono, where Come is located, the indicator of human poverty is almost twice that of

Cotonou [21]. Additionally, the density of pharmacies is much higher in Cotonou (2017: 97 pharmacies over a surface of 79 km²) than in the Mono department (7 over a surface area of 1605 km²) [21, 22].

Without considering events that occurred after this study, the Ministry of Health of Benin conducted awareness campaigns in 2007 to curb IMPD in Benin, and since 2012, a strong appeal is made annually to inform populations about risks associated with consuming IMPD's drugs. Yet, after these measures, the prevalence of IMPD use still seems high. More than one in three caregivers buy drugs in IMPD for home treatment of children under twelve.

The poorest people were the most likely to purchase drugs in the IMPD, as reported in Nigeria [44] and previously in Cotonou [20]. People without a fixed monthly income more frequently bought drugs in the IMPD, while a lower proportion of people with a higher income (more than 100,000 FCFA) buy IMPD drugs. However, some caregivers of the richest quintile purchased drug in IMPD, underlining that it is not exclusively a financial problem. Furthermore, the lack of difference in IMPD usage by caregivers belonging to the middle quintile and the poorest quintile adds complexity. It should also be noted that informal vendors of pharmaceutical drugs are usually part of the neighbourhood. As shown by Baxerres [16], each inhabitant of Cotonou potentially has an informal vendor of pharmaceutical drugs in his social or family circle (a friend of parents, a sister, a mother-in-law, a sister-in-law). The routine and trusting bonds which are being established between them could be the reason.

Pharmaceutical drugs used

Various therapeutic classes of drugs were administered to children. Antipyretics were the most frequently used therapeutic class for home treatment in this study. This may not be surprising, considering that fever is the main symptom associated with malaria. However, fever can be also associated with virus infections that can be treated successfully with antipyretics. Therefore, some caregivers choose to use antipyretic drugs and not anti-malarial drugs to treat children. Fever can transiently decrease after treatment, but home treatment increases the delay of seeking appropriate health care, and may increase the risk of disease progression into severe malaria.

In this study, ACT was used in the same proportions both in the case of home treatment and by prescription. However, the first anti-malarial drug used for home treatment was a artemisinin-based combination, while quinine was the first prescribed drug. Parenteral administration was most frequently mentioned for prescriptions, supposing that the disease seemed severe and considering that caregivers who consulted a healthcare

professional more often cited “vomiting, diarrhoea” as symptoms. The fact that health professionals prescribed antibiotics and anti-malarials together could highlight a lack of diagnosis (infection or malaria). Chloroquine was not prescribed, in compliance with recommendations, but continues to be used for home treatment in Cotonou. Use of chloroquine was also reported in Comé and other countries like Ghana or Nigeria [13, 45], where drug resistance has been confirmed, and where the use of ACT was implemented as a first-line drug to treat uncomplicated malaria.

Synthesis

Children home treatment is not country specific. One important motivation for home treatment is the need for quick relief. Moreover, home treatment is prevalent when the illness is deemed to not require medical consultation or when it is considered as a familiar problem for which the drug is already known [6, 28, 46]. The same reasons prevail in low-income and high-income countries. But, as this study shows, the burden of economic constraints is more felt in low-income countries [47] and limited accessibility to medical care [13, 17]. Limited financial means are also a reason to purchase pharmaceutical drugs in IMPDs [14].

Affordability and the possibility to purchase quantities of drugs adapted to specific needs are additional reasons that may push people toward IMPDs [14, 48]. Besides affordability, people can be motivated by various factors, such as the availability of sellers irrespective of the time of the day, and the ability to purchase ‘prescription-only drugs’ without prescription [14, 49].

Another important point is that vendors often live in close proximity to the customers, sharing the same way of life [16, 49].

Based on Ministry of Health information, Benin has an average of 7.3 qualified health professionals (doctors, nurses and midwives) per 10,000 inhabitants. Cotonou is the city with the highest ratio, 29 qualified health professionals per 10,000 inhabitants [22], when the WHO recommends 23 qualified health professionals per 10,000 inhabitants as needed to provide the most essential maternal and child care [50]. Question about the lack of human resource does not arise in Cotonou. Factors identified in this study may differ from other city where there is a lack of health facilities. Several studies show that the distance to the health facility is a predictor of home treatment [51, 52], of access appropriate and effective health care [53] and also of childhood mortality [54]. It was found in this study that financial means was a factor associated with children home treatment. The financial means were also found

in area where there is a lack of health facilities [31]. The conclusion is that distance from facility is not the main reason of the practice of home treatment in region that faces the lack of health facilities. Indeed, persons who live far from health facility have to spend money on transportation in addition to hospital expenses.

Authors reported that the continued existence of self-medication and then home treatment is favoured by the IMPD [11, 14] as people have access to the same drugs than in formal market, but at cheaper prices. Caregivers who purchased drugs in IMPD purchased more medicines than those who went to formal markets.

Information about the place of purchase was not asked in the case of consultation with a healthcare professional. Indeed, it would be interesting to study whether, after prescription, people still go to purchase drugs in IMPDs, as it was previously observed in Cotonou in a qualitative survey conducted from informal pharmaceutical drugs vendors [16].

Limitations

Data were obtained from 340 children out of the 384 children expected. In this case, there is a loss of statistical power. Information collected was based on a past disease consequently the memory work was asked to the caregivers. Results must be considered while taking into account these elements. Nevertheless, sensitive analyses were done, and allowed to conclude there was no evidence for memory time-related bias.

Conclusion

This study shows that malaria home treatment and use of informal drug market are at the centre of the health-care practice of people in Cotonou, Benin. As patients who consult health-care professional also purchase prescribed drug in the IMPD, these two phenomena must be considered separately.

On IMPD, a part of the medicines originate from the formal pharmaceutical sector in Benin, but the majority are coming informally from Ghana and Nigeria where they are legally distributed. Due to different drug distribution policies between countries, they are considered as false medicine in Benin. On February 24, 2017 as part of the 9th edition of Operation “PANGEA IX coordinated by Interpol, in partnership with the World Customs Organization (WCO), in the main popular market of Cotonou, 80 tons of medicines were seized and 109 people arrested. It is obvious that this action largely slowed down the informal market but, at the same time, reduced the capacity of the population to purchase medicine at low price. New studies are needed to

better understand how people presently acquire pharmaceutical drugs, mainly for self-medication or home treatment.

Additional files

Additional file 1. Sensitivity analysis based on chi-square test to highlight a possible memory bias related to time.

Additional file 2. Trade names of pharmaceutical drugs used by caregivers to treat children under twelve in Cotonou, Benin, May 2016.

Abbreviations

ACT: artemisinin-based combination therapy; GPS: Global Positioning System; IMPD: informal market of pharmaceutical drugs; OR: odds ratio; WHO: World Health Organization.

Authors' contributions

All authors contributed significantly to the work. EA participated in the design of the study, conducted the field work, analysed and interpreted data, drafted and revised the manuscript. MT participated to data management. CB conceived the study. JYLH conceived the study, participated in work conception and data collection in the field, and revised the manuscript. All authors read and approved the final manuscript.

Author details

¹ Institut de recherche pour le développement, Unité mixte de recherche 216: Mères et enfants face aux infections tropicales, Université Paris-Descartes, 4 Avenue de l'Observatoire, 75006 Paris, France. ² Ecole doctorale Pierre Louis de santé publique, ED 393 Epidémiologie et Sciences de l'Information Biomédicale, Paris, France. ³ Centre Norbert Elias EHESS-Campus Marseille La Vieille Charité, 2 Rue de la Charité, 13002 Marseille, France.

Acknowledgements

We would like to thank Dr. Friso PALSTRA for his important contributions to the review of this document.

Competing interests

The authors report no competing interests. The authors alone are responsible for the content and the writing of the paper.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The study has been approved by the Ministry of Higher Education and Scientific Research of Benin; Ethics Committee CER-ISBA –FAVORABLE ADVICE N° 30. Each interviewed person was informed about the objectives of the study, the types of collected data and signed a document of informed consent.

Funding

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement n°337372.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 4 July 2018 Accepted: 3 October 2018

Published online: 10 October 2018

References

1. Ministère de la Santé. Rapport annuel d'activités 2014 du Programme National de Lutte contre le Paludisme. Cotonou, République du Bénin; 2015. http://www2.sante.gouv.bj/documents/PNLP/Rapport_annuel_PNLP_2014%20final.pdf. Accessed 24 Apr 2018.
2. Ogouyemi-Hounto A, Kinde-Gazard D, Nahum A, Abdillahi A, Massoug-bodji A. Management of malaria in Benin: evaluation of the practices of healthcare professionals following the introduction of artemisinin derivatives. *Med Trop (Mars)*. 2009;69:561–4 (in French).
3. WHO. Malaria including proposal for establishment of World Malaria Day; 2007. <https://afro.who.int/publications/wma6018-malaria-including-proposal-establishment-world-malaria-day>. Accessed 25 Sept 2018.
4. Ministry of Health. Malaria Operational Plan FY 2018. Cotonou, Republic of Benin; 2018. <https://www.pmi.gov/docs/default-source/default-document-library/malaria-operational-plans/fy-2018/fy-2018-benin-malaria-operational-plan.pdf?sfvrsn=5h>. Accessed 20 Feb 2018.
5. Mitiku I, Assefa A. Caregivers' perception of malaria and treatment-seeking behaviour for under five children in Mandura District, West Ethiopia: a cross-sectional study. *Malar J*. 2017;16:144.
6. Orimadegun AE, Ilesanmi KS. Mothers' understanding of childhood malaria and practices in rural communities of Ise-Orun, Nigeria: implications for malaria control. *J Fam Med Prim Care*. 2015;4:226–31.
7. Romay-Barja M, Jarrin I, Ncogo P, Nseng G, Sagrado MJ, Santana-Morales MA, et al. Rural-urban differences in household treatment-seeking behaviour for suspected malaria in children at Bata District, Equatorial Guinea. *PLoS ONE*. 2015;10:e0135887.
8. Nwaneri DU, Sadoh AE, Ibadin MO. Impact of home-based management on malaria outcome in under-fives presenting in a tertiary health institution in Nigeria. *Malar J*. 2017;16:187.
9. Malik EM, Hanafi K, Ali SH, Ahmed ES, Mohamed KA. Treatment-seeking behaviour for malaria in children under five years of age: implication for home management in rural areas with high seasonal transmission in Sudan. *Malar J*. 2006;5:60.
10. Tipke M, Louis VR, Yé M, De Allegri M, Beiersmann C, Sié A, et al. Access to malaria treatment in young children of rural Burkina Faso. *Malar J*. 2009;8:266.
11. WHO. Practical chemotherapy of malaria: report of a WHO scientific group. Geneva, World Health Organization; 1990. <http://www.who.int/iris/handle/10665/39778>. Accessed 13 Dec 2017.
12. Baxerres C, Le Hesran J-Y. Where do pharmaceuticals on the market originate? An analysis of the informal drug supply in Cotonou, Benin. *Soc Sci Med*. 2011;73:1249–56.
13. Ezenduka CC, Ogbonna BO, Ekwunife OI, Okonta MJ, Esimone CO. Drugs use pattern for uncomplicated malaria in medicine retail outlets in Enugu urban, southeast Nigeria: implications for malaria treatment policy. *Malar J*. 2014;13:243.
14. Van Der Geest S. Self-care and the informal sale of drugs in South Cameroon. *Soc Sci Med*. 1987;25:293–305.
15. Goodman C, Brieger W, Unwin A, Mills A, Meek S, Greer G. Medicine sellers and malaria treatment in sub-Saharan Africa: what do they do and how can their practice be improved? *Am J Trop Med Hyg*. 2007;77(Suppl 6):203–18.
16. Baxerres C. Du médicament informel au médicament libéralisé: une anthropologie du médicament pharmaceutique au Bénin. Paris: Archives Contemporaines; 2008.
17. Ouattara A. Achat de médicaments de la rue en Afrique: essai de compréhension d'un comportement « irrationnel ». *Mark Manag*. 2011;9:59–73.
18. Du Fassin D. clandestin à l'officieux. Les réseaux de vente illicite des médicaments au Sénégal. *Cahier d'études Africaines*. 1985;25:161–77.
19. Ministère de la santé du Bénin, Direction des pharmacies et du médicament. Politique Pharmaceutique Nationale. Cotonou, République du Bénin; 2008. <http://apps.who.int/medicinedocs/fr/m/abstract/Js21742fr/>. Accessed 27 Apr 2018.
20. Abdoulaye I, Chastanier H, Azondekon A, Dansou A, Bruneton C. Survey on the illicit drug market in Cotonou, Benin in March 2003. *Med Trop (Mars)*. 2006;66:573–6 (in French).
21. Institut National de la Statistique et de l'Analyse Economique (INSAE). Principaux indicateurs sociodémographiques et économiques du département du Littoral; 2016. <https://www.insae-bj.org/images/docs/insae-statistiques/enquetes-recensements/Recensement-General-de-la-Popul>

- ation-et-de-Habitation/Principaux%20indicateurs%20RGP4/Principaux%20indicateurs%20Littoral%20Final.pdf. Accessed 15 June 2017.
22. Institut National de la Statistique et de l'Analyse Economique (INSAE). Annuaire des statistiques sanitaires 2016; 2017. http://www.sante.gouv.bj/IMG/pdf/annuaire_2016_vf.pdf. Accessed 25 Mar 2017.
 23. Geissler PW, Nokes K, Prince RJ, Odhiambo RA, Aagaard-Hansen J, Ouma JH. Children and medicines: self-treatment of common illnesses among Luo schoolchildren in western Kenya. *Soc Sci Med*. 1982;2000(50):1771–83.
 24. Schawrtz D. Méthodes statistiques à l'usage des médecins et des biologistes. Paris: Flammarion Médecins Sciences; 1996.
 25. Ardilly P. Les Techniques de Sondage. Paris: Editions TECHNIP; 2006.
 26. United Nations. Designing household survey samples: practical guidelines. New York: United Nations; 2008.
 27. Institut National de la Statistique et de l'Analyse Economique (INSAE), ICF International. Enquête Démographique et de Santé 2011-2012; 2013. https://www.insae-bj.org/images/docs/insae-statistiques/enquetes-recensements/Enquete%20Demographique%20et%20de%20Sante/2011-2012/EDS_2012_Rapport_final-11-15-2013.pdf. Accessed 2 Mar 2017.
 28. Hughes CM, McElroy JC, Fleming GF. Benefits and risks of self medication. *Drug Saf*. 2001;24:1027–37.
 29. Lee C-H, Chang F-C, Hsu S-D, Chi H-Y, Huang L-J, Yeh M-K. Inappropriate self-medication among adolescents and its association with lower medication literacy and substance use. *PLoS ONE*. 2017;12:e0189199.
 30. WHO. World malaria report 2017. Geneva, World Health Organization; 2017. <http://www.who.int/malaria/media/world-malaria-report-2017/en/>. Accessed 26 Feb 2018.
 31. Agueh V, Badet ME, Jerome CS, Paraiso MN, Azandjeme CS, Metonnou C, et al. Prevalence and determinants of antimalarial self-medication in Southern Benin. *Int J Trop Dis Health*. 2016;18:1–11.
 32. Briggs MA, Kalolella A, Bruxvoort K, Wiegand R, Lopez G, Festo C, et al. Prevalence of malaria parasitemia and purchase of artemisinin-based combination therapies (ACTs) among drug shop clients in two regions in Tanzania with ACT subsidies. *PLoS ONE*. 2014;9:e94074.
 33. Kpatchavi AC. Savoirs, maladie et thérapie en Afrique de l'Ouest: Pour une anthropologie du paludisme chez les Fgn et Waci du Bénin. Éditions Ablodé; 2011.
 34. Fenny AP, Asante FA, Enemark U, Hansen KS. Malaria care seeking behavior of individuals in Ghana under the NHIS: are we back to the use of informal care? *BMC Public Health*. 2015;15:370.
 35. Ridde V, Girard J-E. Douze ans après l'initiative de Bamako: constats et implications politiques pour l'équité d'accès aux services de santé des indigents africains. *Santé Publique*. 2004;16:37–51.
 36. Ministère de la Santé du Bénin. Processus de mise en place d'un Régime d'Assurance Maladie Universelle (RAMU); 2011. http://www.coopami.org/fr/countries/countries/benin/social_protection/pdf/social_protection_01.pdf. Accessed 19 June 2018.
 37. Wagstaff A, Flores G, Hsu J, Smits M-F, Chepynoga K, Buisman LR, et al. Progress on catastrophic health spending in 133 countries: a retrospective observational study. *Lancet Glob Health*. 2017;6:169–79.
 38. Ministère de la Santé du Bénin. Régime d'Assurance Maladie Universelle (RAMU); 2011. <http://www.sante.gouv.bj/spip.php?article76>. Accessed 29 May 2018.
 39. Mayala BK, Fahey CA, Wei D, Zinga MM, Bwana VM, Mlacha T, et al. Knowledge, perception and practices about malaria, climate change, livelihoods and food security among rural communities of central Tanzania. *Infect Dis Poverty*. 2015;4:21.
 40. Okeke TA. Improving malaria recognition, treatment and referral practices by training caretakers in rural Nigeria. *J Biosoc Sci*. 2010;42:325–39.
 41. Iriemenam NC, Dosunmu AO, Oyibo WA, Fagbenro-Beyioku AF. Knowledge, attitude, perception of malaria and evaluation of malaria parasitaemia among pregnant women attending antenatal care clinic in metropolitan Lagos, Nigeria. *J Vector Borne Dis*. 2011;48:12–7.
 42. Serengbe GB, Moyen J-M, Fioboy R, Beyam EN, Kango C, Bangue C, et al. Knowledge and perceptions about malaria in communities in four districts of the Central African Republic. *BMC Res Notes*. 2015;8:162.
 43. Foster S. Treatment of malaria outside the formal health services. *J Trop Med Hyg*. 1995;98:29–34.
 44. Isiguzo C, Anyanti J, Ujuju C, Nwoko E, De La Cruz A, Schatzkin E, et al. Presumptive treatment of malaria from formal and informal drug vendors in Nigeria. *PLoS ONE*. 2014;9:e110361.
 45. Ameh S, Welaga P, Kabiru CW, Ndifon W, Ikpeme B, Nsan E, et al. Factors associated with appropriate home management of uncomplicated malaria in children in Kassena-Nankana district of Ghana and implications for community case management of childhood illness: a cross-sectional study. *BMC Public Health*. 2015;15:458.
 46. Coulomb A, Baumelou A. Situation de l'automédication en France et perspectives d'évolution : marché, comportements, positions des acteurs. La Documentation française; 2007. <http://www.bdsp.ehesp.fr/Base/353391/>. Accessed 20 Apr 2018.
 47. Brutus L, Fleuret S, Guienne V. Se soigner par soi-même. Recherche interdisciplinaire sur l'automédication. Paris: CNRS Edition Alpha; 2017.
 48. Angbo-Effi KO, Kouassi DP, Yao GHA, Douba A, Secki R, Kadjo A. Facteurs déterminant la consommation des médicaments de la rue en milieu urbain. *Santé Publique*. 2012;23:455–64.
 49. Baxerres C, Le Hesran J-Y. Le marché parallèle du médicament en milieu rural au Sénégal: Les atouts d'une offre de soins populaire (Note de recherche). *Anthropol Sociétés*. 2006;30:219–30.
 50. WHO. World Health Statistics 2014. Geneva, World Health Organization; 2014. http://www.who.int/gho/publications/world_health_statistics/2014/en/. Accessed 27 June 2018.
 51. Ocan M, Bwanga F, Bbosa GS, Bagenda D, Waako P, Ogwal-Okeng J, et al. Patterns and predictors of self-medication in northern Uganda. *PLoS ONE*. 2014;9:e92323.
 52. Begashaw B, Tessema F, Gesesew HA. Health care seeking behavior in Southwest Ethiopia. *PLoS ONE*. 2016;11:e0161014.
 53. Mwaliko E, Downing R, O'Meara W, Chelagat D, Obala A, Downing T, et al. "Not too far to walk": the influence of distance on place of delivery in a western Kenya health demographic surveillance system. *BMC Health Serv Res*. 2014;14:212.
 54. Schoeps A, Gabrysch S, Niamba L, Sié A, Becher H. The effect of distance to health-care facilities on childhood mortality in rural Burkina Faso. *Am J Epidemiol*. 2011;173:492–8.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

