

A convenient Simple Method for Synthesis of Meta-iodobenzylguanidine (MIBG)

Zahra Sheikholislam^a, Zohreh Soleimani^a, Abolghasem Moghimi^b and Soraya Shahhosseini^{a*}

^aPharmaceutical Chemistry Department, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran, P O Box: 14155-6153. ^bDepartment of Chemistry, University of Imam Hossein.

Abstract

Radioiodinated meta-iodobenzylguanidine (MIBG) is one of the important radiopharmaceuticals in Nuclear Medicine. [^{123/131}I] MIBG is used for imaging of Adrenal medulla, studying heartsympathetic nerves, treatment of pheochromocytoma and neuroblastoma. For clinical application, radioiodinated MIBG is prepared through isotopic exchange method, which includes replacement of radioactive iodine in a nucleophilic substitution reaction with cold iodine (¹²⁷I). The unlabelled MIBG hemisulfate is synthesized by the procedure described by Wieland *et al.* (1980). The availability of a more practical and cost-effective procedure for MIBG preparation encouraged us to study the MIBG synthesis methods. In this study the preparation of MIBG through different methods were evaluated and a new method, which is one step, simple and cost-effective is introduced. The method has ability to be scaled up for production of unlabelled MIBG.

Keywords: Meta-iodobenzylguanidine; Neuroendocrine tumors; Sympathetic neurons; Radioiodinated MIBG.

Introduction

Meta-iodobenzylguanidine (MIBG) a structural and functional analogue of norepinephrine (NE), is transported, stored, and released in the adrenergic neuron by the same mechanisms as NE which includes neuronal (uptake-1) and extra neuronal (uptake-2, myocyte) specific membrane transport process and passive diffusion in and from both cellular compartments. MIBG unlike NE is a stable and non-metabolizable compound in vivo, resistant to monoamine oxidase (MAO) and catechol-o-methyl transferase (COMT) enzyme systems. Studies have shown that the norepinephrine transporter (NET) is highly over expressed in

neuroendocrine tumors such as neuroblastoma, pheochromocytoma, paraganglioma, and carcinoid. Since MIBG is a substrate for NET, therefore, MIBG and its radioiodinated forms accumulate in neuroendocrine tumors. In clinic, radioiodinated MIBG is used for treatment and diagnosis of neuroendocrine tumors. Accumulation of radioiodinated MIBG in adrenergic neurons clinically used for non-invasive assessment of sympathetic neuronal functions (1-9).

Although MIBG has been radiolabelled with radioisotopes such as ²¹¹At, ¹⁸F, ⁷⁶Br none of these radiopharmaceuticals are in clinic yet (10-13). The most important radioisotopes for radiolabeling MIBG are still iodine radioisotopes (^{123/131}I). Radioiodinated MIBG is an important diagnostic and therapeutic radiopharmaceutical in clinic. For clinical application, radioiodinated

* Corresponding author:

E-mail: s_shahoseini@sbm.ac.ir

MIBG is prepared through isotopic exchange method (14-16). In this method, radioactive iodine ($^{123/131}\text{I}$) in a nucleophilic substitution reaction replaces cold iodine (^{127}I) in MIBG molecule. The unlabelled MIBG hemisulfate is synthesized by the procedure described by Wieland *et al.* (1980). The availability of a more practical procedure for MIBG preparation encouraged us to study the MIBG synthesis methods. In this study, MIBG was synthesized through 4 different methods. In each method, the product was characterized by melting point and spectroscopic methods (IR, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$, Mass). The purity was determined using Reverse phase HPLC and TLC. The spectra data for in-house MIBG was in agreement with structure of MIBG. The purity of MIBG hemisulfate synthesized using meta-iodobenzylamine and S-ethylisothiuronium (method four) was found to be comparable with authentic reference MIBG (Sigma). The synthesis compare to Wieland procedure, which is used for MIBG synthesis, has fewer steps. It is simple and because of production of ethanethiol as a by-product is cost-effective. The new method has the ability to be replaced by traditional methods for preparation of unlabelled MIBG.

Experimental

All chemicals were purchased from Merck except meta-iodobenzylamine hydrochloride (MIBA HCl) and reference MIBG which were purchased from Sigma-Aldrich.

Four different methods were used for synthesizing MIBG.

Method 1: MIBG hemisulfate was synthesized according to the procedure described by Wieland *et al.* (1980). Briefly, a mixture of MIBA HCl (1 mmole) and cyanamide (1.5 mmole) was refluxed 4 hrs at 100°C. The resulting oily product was dissolved in 0.5 mL water and 0.5 mL KHCO_3 (100 mg/0.5 mL) was added slowly with stirring. The precipitated MIBG bicarbonate was filtered by vacuum, washed by cold water, and dried in vacuum oven (M.P 121-124°C, yield: 60%). To a mixture of MIBG bicarbonate (0.6 mmole) in 2 mL water, 0.3 mL 2 N H_2SO_4 was added slowly. The MIBG hemisulfate was dissolved in water by heat and crystallized upon cooling

at room temperature. The crystals was filtered by vacuum, washed by cold water, and dried in vacuum oven. Re-crystallization was done in a mixture of water-ethanol (50:50) (2). M.P 166-167.5°C, IR (KBr, cm^{-1}): 3302 (NH), 3138 (NH), 1656 and 1537 (C=N), 1057 (S=O), 785 and 686 (1, 3-disubstituted benzene). $^1\text{H-NMR}$ ($\text{D}_2\text{O}/500\text{ MHz}$): δ 7.1 (t, 1H, phenyl H5), 7.3 (d, 1H, phenyl H6), 7.5 (m, 2H, phenyl H2, H4), 4.3 (s, 1H, CH_2 Benzyl) ppm. $^{13}\text{C-NMR}$ ($\text{D}_2\text{O}/125\text{ MHz}$): δ 40 (CH_2 -Benzyl), 95 (C3-phenyl), 130 (C6-phenyl), 135 (C5-phenyl), 137 (C4-phenyl), 140 (C2- phenyl), 142 (C1 phenyl), 158 (C-Imine) ppm. MS (ESI): 276 ($\text{M}+\text{H}^+$).

Method 2: To a solution of 3.3 g sodium (Na) in 20 mL super dry ethanol, 2 mmole guanidine HCl was added, and refluxed for 15 min. After filtration of sodium chloride, 2 mmole meta-iodobenzylbromide in anhydrous ethanol was added to filtrate and refluxed for 36 hrs. Solvent was removed by rotary evaporation. MIBG was crystallized in methanol (17-18). Yield was 43%. IR (KBr, cm^{-1}): 3367 and 3162 (NH), 1636 and 1560 (C=N), 801 and 737 (1,3-disubstituted benzene). $^1\text{H-NMR}$ ($\text{CDCl}_3/500\text{ MHz}$): δ 7 (t, 1H, phenyl H5), 7.1 (d, 1H, phenyl H6), 7.3 (m, 2H, phenyl H2, H4), 4 (s, 1H, CH_2 Benzyl) ppm. $^{13}\text{C-NMR}$ ($\text{CDCl}_3/125\text{ MHz}$): δ 40 (CH_2 -Benzyl), 93 (C3- phenyl), 130 (C6- phenyl), 135 (C5- phenyl), 135 (C4- phenyl), 139 (C2-phenyl), 142 (C1- phenyl), 158 (C-Imine) ppm. MS (ESI): 276 ($\text{M}+\text{H}^+$).

Method 3: A mixture of 0.8 mmole cyanoguanidine with 2 mmole MIBA HCl was heated for 6 h at 200°C while purging with nitrogen gas. MIBG was crystallized upon cooling (19). Yield was 60.5%. IR (KBr, cm^{-1}): 3298 and 3011(NH), 1685 and 1628 (C=N), 841 and 697 (1,3-disubstituted benzene). $^1\text{H-NMR}$ ($\text{CDCl}_3/500\text{ MHz}$): δ 7 (t, 1H, phenyl H5), 7.3 (d, 1H, phenyl H6), 7.5 (m, 2H, phenyl H2, H4), 3.9 (s, 1H, CH_2 Benzyl) ppm. $^{13}\text{C-NMR}$ ($\text{CDCl}_3/125\text{ MHz}$): δ 40(CH_2 -Benzyl), 95 (C3-phenyl), 125 (C6-phenyl), 130 (C5-phenyl), 135 (C4- phenyl), 137 (C2- phenyl), 139 (C1-phenyl), 160 (C-Imine) ppm. MS (ESI): 276 ($\text{M}+\text{H}^+$).

Method 4: To a mixture of 0.7 mmole MIBA HCl in 15 mL water, 0.3 mmole S-ethylisothiuronium sulfate [$\text{NH}_2\text{C}(\text{NH})$




Figure 1. Scheme for the synthesis of MIBG from meta-iodobenzylbromide with guanidine.

$S\text{-Et}^+]_2\text{SO}_4^{2-}$ was added gradually. The mixture was stirred 5 hrs at room temperature. Water was removed by rotary evaporation and the precipitate was crystallized in acetone. Re-crystallization was done on acetone. Ethanethiol a valuable by-product of reaction was collected during the reaction. Yield was 63%. IR (KBr, cm^{-1}): 3339 and 3274 (NH), 1643 and 1662 (C=N), 1092 (S=O), 887 and 701 (1,3-disubstituted benzene). ^{13}C -NMR ($\text{D}_2\text{O}/125\text{ MHz}$): δ 40 (CH_2 -Benzyl), 95 (C3- phenyl), 125 (C6- phenyl), 127 (C5- phenyl), 130 (C4- phenyl), 133 (C2-phenyl), 135 (C1- phenyl), 165 (C-Imine) ppm. MS (ESI): 276 ($\text{M}+\text{H}^+$).

Synthesis of S-ethylisothiuronium sulfate $[\text{NH}_2\text{C}(\text{NH})\text{S-Et}]_2^+\text{SO}_4^{2-}$ was done based on method of Osklo-Salmani A, 2003. Briefly, a reaction mixture of 0.15 mole diethylsulfate and 0.2 mole thiourea was stirred at 100°C for 2.5 hrs and then at 140°C for an additional 1.5 hrs. Upon cooling, S-ethylisothiuronium sulfate was obtained with melting point $219\text{-}222^\circ\text{C}$ and 93% yield (20). IR (KBr, cm^{-1}): 3385, 3140, 2820, 2125, 1620, 1405, 1380, 1330, 1230, 1080, 1000, 920, 870. ^{13}C -NMR (DMSO-d_6 , $\text{D}_2\text{O}/125\text{ MHz}$): δ 172.1 (C=N), 25.9 (SCH_2), 13.9 (CH_3) ppm.

Quality control of synthesized MIBG was done through RP-HPLC and TLC based on United States Pharmacopeia USP (21, 2008). HPLC (Merck-Hitachi, L7100, L7420): C18 Chro.Monolithic 100-4.6 mm, mobile phase: 3.04 g triethylamine per 1 liter of Water: Acetonitrile (90:10), pH 4. Flow rate: 1.5 mL/min, 229 nm. TLC: Silica gel, HCCl_3 : CH_3OH (95:5).

Results and Discussion

Radioiodinated MIBG is one of the

important radiopharmaceuticals in Nuclear Medicine. $^{123/131}\text{I}$ MIBG is used for imaging of Adrenal medulla, studying heart sympathetic nerves, treatment of pheochromocytoma and neuroblastoma. For clinical application, radioiodinated MIBG is prepared by exchange of radioactive iodine with non-radioactive one using isotope exchange method.

The method used for synthesis of unlabelled MIBG is based on Wieland's procedure (2). In this method, the reaction of MIBA HCl and cyanamide in the presence of KHCO_3 , results in MIBG bicarbonate formation. By adding H_2SO_4 to MIBG bicarbonate, MIBG hemisulfate is crystallized. A few times re-crystallization in a mixture of water:ethanol (50:50) is used for purification of MIBG hemisulfate.

The reaction of alkyl halide with appropriate amine was used for preparation of MIBG in second method. Guanidine HCl in a solution of Na in ethanol makes a good base to react with m-iodobenzylbromide to produce MIBG (Figure 1). Resulting MIBG was crystallized on methanol. Yield was 43%. This method does not have any advantages to Wieland procedure.

The reaction of cyanoguanidine with ammonium salts, which results in guanidinium salts, was used in third method for MIBG preparation. Mixture of Cyanoguanidine and MIBA HCl was heated up 200°C for 6 hrs (Figure 2). Crystals of MIBG was formed upon cooling. The yield was 60.5%. Since no solvent is used in this reaction, there is no step to eliminate solvent which is advantageous of this method.

The reaction of MIBA HCl with salt of S-ethylisothiuronium sulfate produced MIBG sulfate in one step. Yield was 63%. Ethanethiol, the by-product, was collected which is used for synthesis of other chemicals. In this method,




Figure 2. Scheme for the synthesis of MIBG from meta-iodobenzylamine with cyanoguanidine.

MIBG hemisulfate is formed directly without adding H_2SO_4 .

Comparing the methods, method 2 and 3 are similar and do not have any specific advantages to Wieland procedure which is routinely used to synthesis MIBG sulfate. Method four is a new method which is reported for first time. In this method, MIBG hemisulfate is prepared through one step. The by-product of reaction is easily collected and is used in industry for synthesizing other chemicals. Considering yield, cost, steps, and simplicity the new method looks promising to be replaced instead of Wieland procedure.

In summary, a convenient and simple method for the synthesis of MIBG was developed. The method has ability to be scaled up for production of unlabelled MIBG.

Acknowledgment

This study was supported financially by a grant from Vice-Chancellor's office for Research and Technology affairs, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

References

- (1) Wieland DM, Swanson DP, Brown EL and Beierwaltes WH. Imaging the adrenal medulla with an ^{131}I -labeled antiadrenergic agent. *J. Nucl. Med.* (1979) 20: 155-158.
- (2) Wieland DM, W JL, Brown LE, Mangner TJ, Swanson DP and Beierwaltes WH. Radiolabeled adrenergic neuron-blocking agents: adrenomedullary imaging with ^{131}I -iodobenzylguanidine. *J. Nucl. Med.* (1980) 21: 349-353.
- (3) Wieland DM, Mangner TJ, Inbasekaran MN, Brown LE and Wu JL. Adrenal medulla imaging agents: a structure-distribution relationship study of radiolabeled aralkylguanidines. *J. Med. Chem.* (1984) 27: 149-155.
- (4) Glowniak JV, Turner FE, Gray LL, Palac RT, Lagunas-Solar MC and Woodward WR. ^{131}I -metaiodobenzylguanidine imaging of the heart in idiopathic congestive cardiomyopathy and cardiac transplants. *J. Nucl. Med.* (1989) 30: 1182-1191.
- (5) Wafelman AR, Hoefnagel CA and Beijnen JH. Radiochemical stability during infusion of ^{131}I -labelled meta-iodobenzylguanidine for therapeutic use. *Nucl. Med Commun.* (1993) 14: 819-822.
- (6) Hanson MW, Feldman JM, Beam CA, Leight GS and Coleman RE. Iodine-131 labeled metaiodobenzylguanidine scintigraphy and biochemical analysis in suspected pheochromocytoma. *Arch. Intern. Med.* (1991) 151: 1397-1402.
- (7) DeGrado TR, Zalutsky MR and Vaidyanathan G. Uptake mechanisms of ^{123}I -MIBG in isolated rat heart. *Nucl. Med. Biol.* (1995) 22: 1-12.
- (8) Trendelenburg U. Extraneuronal uptake and metabolism of catecholamines as a site of loss. *Life Sci.* (1978) 22: 1217-1222.
- (9) Vaidyanathan G, Affleck DJ, Alston KL, Zhao XG, Hens M, Hunter DH, Babich J and Zalutsky MR. A kit method for the high level synthesis of [^{211}At] MABG. *Bioorganic and Medicinal Chemistry.* (2007) 15: 3430-3436.
- (10) Berry CR, Garg PK, Zalutsky MR. Coleman RE and DeGrado TR. Uptake and retention kinetics of para fluorine-18 fluorobenzylguanidine in isolated rat heart. *J. Nucl. Med.* (1996) 37: 2011-2016.
- (11) Garg P. K, Garg S, Zalutsky M. R. Synthesis and preliminary evaluation of para and meta 18-fluorobenzylguanidine. *Nucl. Med. Biol.* (1994) 21: 97-103.
- (12) Vaidyanathan G, Affleck DJ, Alston KL, Zhao XG, Hens M, Hunter DH, Babich J and Zalutsky M R. A kit method for high level synthesis of ^{211}At -MABG. *Bioorganic and Medicinal Chemistry.* (2007) 15: 3430-3436.
- (13) Loc'h C, Mardon K, Valette H, Brutesco C, Merlet P, Syrota A and Maziere B. preparation and pharmacological characterization of ^{76}Br -MIBG. *Nucl. Med. Biol.* (1994) 21: 49-55.
- (14) Mock BH and Weiner RF. Simplified solid state labeling of ^{123}I -MIBG. *Applied Radiation and Isotopes.* (1988) 39: 939-942.

- (15) Prabhakar G, Mehra K.S, Ramamoorthy N, Chattopadhyay S, Oommen R, Narasimhan S, Nair NA and Gunasekaran S. Evaluation of radioiodination of MIBG catalysed by in situ generated Cu(I) and directly added Cu(II). *Applied Radiation and Isotopes*. (1999) 50: 1011-1014.
- (16) Donovan A C and Valliant J F. A convenient solution phase method for the preparation of MIBG in high effective specific activity. *Nuclear Medicine and Biology* (2008) 35: 741-746.
- (17) Willson FGI, Wheeler TS. Benzyl aniline. Organic synthesis. (1941) Coll. Vol. 1, P: 102-104; (1928) Coll. Vol. 8, P: 38.
- (18) Cymerman CJ, Yong J. 1-benzyl piperazine. Organic synthesis. (1973) Coll. Vol. 5, P: 88; (1962) 42: 19.
- (19) Lecher H, Graf F. 1923, Ber. 56 B, 1326-1330. C.A., 1923; 17: 3319.
- (20) Osklo-Salmani A. Synthesis of alkylguanidines. Master of Sciences thesis in organic chemistry. Chemistry department, Faculty of basic Sciences, Imam Hossein University. (2003): 45-60.

This article is available online at <http://www.ijpr.ir>
