

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Methoxybenzamidineium hydrogen sulfate

Simona Irrera and Gustavo Portalone*

Chemistry Department, 'Sapienza' University of Rome, P. le A. Moro, 5, I-00185 Rome, Italy

Correspondence e-mail: g.portalone@caspur.it


Received 23 October 2012; accepted 25 October 2012

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.039; wR factor = 0.114; data-to-parameter ratio = 15.7.

The title salt, $\text{C}_8\text{H}_{11}\text{N}_2\text{O}^+\cdot\text{HSO}_4^-$, has been synthesized by the reaction between 4-methoxybenzamidine and sulfuric acid. The asymmetric unit comprises a nonplanar 4-methoxybenzamidineium cation and one hydrogen sulfate anion. In the cation, the amidinium group has two identical C—N bonds [1.306 (2) and 1.308 (2) Å], and its plane forms a dihedral angle of 6.49 (8)° with the mean plane of the benzene ring. The ionic components are associated in the crystal *via* N—H⁺··O⁻, resulting in chains running approximately along the *b*-axis direction which are interconnected by O—H··O⁻ hydrogen bonds.

Related literature

For the biological and pharmacological relevance of benzamidine, see: Powers & Harper (1999); Grzesiak *et al.* (2000). For structural analysis of proton-transfer adducts containing molecules of biological interest, see: Portalone (2011a); Portalone & Irrera (2011). For the supramolecular association in proton-transfer adducts containing benzamidineium cations, see: Portalone (2010, 2011b, 2012); Irrera & Portalone (2012); Irrera *et al.* (2012). For hydrogen-bond motifs, see Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_8\text{H}_{11}\text{N}_2\text{O}^+\cdot\text{HSO}_4^-$
 $M_r = 248.26$
 Monoclinic, $P2_1/c$

$a = 14.2608$ (14) Å
 $b = 10.1844$ (9) Å
 $c = 7.5723$ (9) Å

$\beta = 94.206$ (10)°
 $V = 1096.83$ (19) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.30$ mm⁻¹
 $T = 298$ K
 $0.31 \times 0.25 \times 0.15$ mm

Data collection

Oxford Diffraction Xcalibur S CCD diffractometer
 Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2006)
 $T_{\min} = 0.912$, $T_{\max} = 0.956$

16529 measured reflections
 2626 independent reflections
 2133 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.114$
 $S = 1.04$
 2626 reflections
 167 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O3—H3A···O2 ⁱ	0.78 (3)	1.79 (3)	2.562 (2)	172 (3)
N1—H1A···O4	0.86 (2)	2.10 (2)	2.938 (2)	163 (2)
N1—H1B···O5 ⁱⁱ	0.84 (2)	2.10 (2)	2.884 (2)	154 (2)
N2—H2A···O5	0.84 (2)	2.07 (3)	2.907 (2)	177 (2)
N2—H2B···O4 ⁱⁱⁱ	0.86 (3)	2.22 (3)	2.965 (2)	145 (3)

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: CrysAlis CCD (Oxford Diffraction, 2006); cell refinement: CrysAlis RED (Oxford Diffraction, 2006); data reduction: CrysAlis RED; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ5020).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Grzesiak, A., Helland, R., Smalas, A. O., Krowarsch, D., Dadlez, M. & Otlewski, J. (2000). *J. Mol. Biol.* **301**, 205–217.
- Irrera, S., Ortaggi, G. & Portalone, G. (2012). *Acta Cryst.* **C68**, o447–o451.
- Irrera, S. & Portalone, G. (2012). *Acta Cryst.* **E68**, o3083.
- Oxford Diffraction (2006). *CrysAlis CCD and CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Portalone, G. (2010). *Acta Cryst.* **C66**, o295–o301.
- Portalone, G. (2011a). *Chem. Cent. J.* **5**, 51.
- Portalone, G. (2011b). *Acta Cryst.* **E67**, o3394–o3395.
- Portalone, G. (2012). *Acta Cryst.* **E68**, o268–o269.
- Portalone, G. & Irrera, S. (2011). *J. Mol. Struct.* **991**, 92–96.
- Powers, J. C. & Harper, J. W. (1999). *Proteinase Inhibitors*, edited by A. J. Barrett & G. Salvesen, pp. 55–152. Amsterdam: Elsevier.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o3244 [doi:10.1107/S1600536812044327]

4-Methoxybenzamidine hydrogen sulfate

Simona Irrera and Gustavo Portalone

Comment

In our search for new salt compounds as part of investigation of the functionality of biological systems (Portalone, 2011*a*; Portalone & Irrera, 2011), we have prepared 4-methoxybenzamidine hydrogen sulfate, (I), which was obtained by a reaction between 4-methoxybenzamide (4-amidinoanisole) and sulfuric acid.

Benzamide derivatives, which have shown strong biological and pharmacological activity (Powers & Harper, 1999; Grzesiak *et al.*, 2000), are being used in our group as bricks for supramolecular construction (Portalone, 2010, 2011*b*, 2012). Indeed, these molecules are strong Lewis bases and their cations can be easily anchored onto numerous inorganic and organic anions and polyanions, largely because of the presence of four potential donor sites for hydrogen-bonding.

The asymmetric unit of (I) comprises one non-planar 4-methoxybenzamidine cation and one hydrogen sulfate anion (Fig. 1). In the cation the amidinium group forms dihedral angle of 6.49 (8)° with the mean plane of the phenyl ring, which is close to the values observed in protonated benzamidine ions (14.4 (1) - 32.7 (1)°, Portalone, 2010, 2012; Irrera *et al.*, 2012). The lack of planarity in all these systems is obviously caused by steric hindrances between the H atoms of the aromatic ring and the amidine moiety. This conformation is rather common in benzamidine-containing small-molecule crystal structures, with the only exception of benzamidine diliturate, where the benzamidine cation is planar (Portalone, 2010). The pattern of bond lengths and bond angles of the 4-methoxybenzamidine cation agrees with that reported in previous structural investigations (Irrera *et al.*, 2012; Portalone, 2010, 2012; Irrera & Portalone, 2012). In particular the amidinium group, true to one's expectations, features identical C—N bonds within experimental error [1.306 (2) and 1.308 (2) Å], evidencing the delocalization of the π electrons and double-bond character.

Bond lengths in the slightly distorted tetrahedral hydrogen sulfate anion indicate the position of the H atom. There are three short S—O bonds of 1.4504 (13), 1.4442 (13) and 1.4464 (13) Å to terminal atoms O2, O4 and O5, respectively, and one longer bond of 1.5470 (15) Å to atom O3, which is bound to atom H3A.

The ionic components of compound (I) are joined by two $N^+—H\cdots O^-$ (\pm) hydrogen bonds (Table 1) to form ionic dimers with graph-set motif $R^2_2(8)$ (Bernstein *et al.*, 1995). Analysis of the crystal packing of (I), (Fig. 2), shows that four $N^+—H\cdots O^-$ hydrogen bonds link the molecular components into a mono-dimensional structure. As previously mentioned, each subunit, built from the ion pairs of the asymmetric unit, forms $R^2_2(8)$ dimers *via* the bidentate interaction of the N—H and S—O groups. Adjacent ion pairs are then linked together by way of the remaining two $N^+—H\cdots O^-$ hydrogen bonds to form $R^2_4(8)$, resulting in chains running approximately along crystallographic *b* axis. These chains are then interconnected by means of the only $O—H\cdots O$ hydrogen bond present in the structure.

Experimental

4-Methoxybenzamide (0.1 mmol, Fluka at 96% purity) was dissolved without further purification in 6 ml of hot water and heated under reflux for 3 h. While stirring, H_2SO_4 (2 mol L^{-1}) was added dropwise until pH reached 2. After cooling the solution to an ambient temperature, colourless crystals suitable for single-crystal X-ray diffraction were grown by

slow evaporation of the solvent after one week.

Refinement

All H atoms were identified in difference Fourier maps, but for refinement all C-bound H atoms were placed in calculated positions, with C—H = 0.93 Å (phenyl) and 0.97 Å (methyl), and refined as riding on their carrier atoms. The U_{iso} values were kept equal to $1.2U_{\text{eq}}(\text{C, phenyl})$, and to $1.5U_{\text{eq}}(\text{C, methyl})$. Positional and thermal parameters of H atoms of the amidinium group and of the hydrogen sulfate ion were freely refined, giving N—H distances in the range 0.84 (2)–0.86 (2) Å and an O—H distance equal to 0.78 (3) Å.

Computing details

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduction: *CrysAlis RED* (Oxford Diffraction, 2006); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).





Figure 1

The asymmetric unit of (I), showing displacements ellipsoids drawn at the 50% probability level. The asymmetric unit was selected so that the two ions are linked by $\text{N}^+ \cdots \text{H} \cdots \text{O}^-$ hydrogen bonds. H atoms are shown as small spheres of arbitrary radii. Hydrogen bonding is indicated by dashed lines.


Figure 2

Crystal packing diagram for (I), viewed approximately down the *c* axis. Displacements ellipsoids are at the 50% probability level. H atoms are shown as small spheres of arbitrary radii. For the sake of clarity, H atoms not involved in hydrogen bonding (dashed lines) have been omitted.

4-Methoxybenzamidinium hydrogen sulfate

Crystal data

$C_8H_{11}N_2O^+ \cdot HSO_4^-$

$M_r = 248.26$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.2608 (14) \text{ \AA}$

$b = 10.1844 (9) \text{ \AA}$

$c = 7.5723 (9) \text{ \AA}$

$\beta = 94.206 (10)^\circ$

$V = 1096.83 (19) \text{ \AA}^3$

$Z = 4$

$F(000) = 520$

$D_x = 1.503 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$

Cell parameters from 4647 reflections

$\theta = 2.9\text{--}29.7^\circ$

$\mu = 0.30 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Tablets, colourless

$0.31 \times 0.25 \times 0.15 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur S CCD
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: $16.0696 \text{ pixels mm}^{-1}$

ω and φ scans

Absorption correction: multi-scan

(*CrysAlis RED*; Oxford Diffraction, 2006)

$T_{\min} = 0.912$, $T_{\max} = 0.956$

16529 measured reflections

2626 independent reflections

2133 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 28.0^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -18 \rightarrow 18$

$k = -13 \rightarrow 13$

$l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.114$
 $S = 1.04$
 2626 reflections
 167 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0582P)^2 + 0.2957P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. Absorption correction: [CrysAlis RED (Oxford Diffraction, 2006); empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm]

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.16129 (3)	0.16203 (4)	0.84884 (6)	0.03448 (15)
O2	-0.20315 (10)	0.11649 (13)	1.00664 (17)	0.0514 (4)
O3	-0.24735 (9)	0.20809 (15)	0.7284 (2)	0.0498 (4)
H3A	-0.2287 (18)	0.259 (3)	0.662 (4)	0.064 (8)*
O4	-0.09972 (9)	0.27365 (12)	0.87891 (18)	0.0467 (3)
O5	-0.11706 (9)	0.05608 (12)	0.75864 (19)	0.0485 (3)
O1	0.49301 (10)	0.19362 (16)	0.4667 (3)	0.0684 (5)
N1	0.07593 (12)	0.28008 (15)	0.6992 (3)	0.0452 (4)
H1A	0.0189 (18)	0.280 (2)	0.730 (3)	0.053 (6)*
H1B	0.1027 (17)	0.354 (2)	0.693 (3)	0.061 (7)*
N2	0.07598 (14)	0.06125 (17)	0.6593 (3)	0.0595 (5)
H2A	0.0212 (18)	0.058 (2)	0.692 (3)	0.057 (7)*
H2B	0.107 (2)	-0.009 (3)	0.647 (4)	0.087 (9)*
C1	0.21780 (12)	0.18063 (15)	0.6067 (2)	0.0346 (4)
C2	0.26115 (13)	0.30028 (17)	0.5855 (3)	0.0465 (5)
H2	0.2275	0.3770	0.6020	0.056*
C3	0.35275 (14)	0.30917 (18)	0.5407 (3)	0.0497 (5)
H3	0.3806	0.3910	0.5288	0.060*
C4	0.40279 (13)	0.19663 (19)	0.5136 (3)	0.0465 (5)
C5	0.36028 (15)	0.0754 (2)	0.5315 (3)	0.0595 (6)
H5	0.3937	-0.0010	0.5123	0.071*
C6	0.26906 (14)	0.06751 (18)	0.5775 (3)	0.0493 (5)
H6	0.2413	-0.0143	0.5892	0.059*

C7	0.12029 (12)	0.17365 (15)	0.6575 (2)	0.0365 (4)
C8	0.54205 (16)	0.3152 (3)	0.4575 (4)	0.0749 (8)
H8A	0.5094 (10)	0.3716 (13)	0.369 (2)	0.112*
H8B	0.6058 (12)	0.2988 (4)	0.425 (3)	0.112*
H8C	0.5445 (13)	0.3583 (13)	0.573 (2)	0.112*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0360 (3)	0.0244 (2)	0.0440 (3)	-0.00176 (15)	0.00969 (18)	-0.00018 (16)
O2	0.0718 (9)	0.0368 (7)	0.0477 (8)	-0.0093 (6)	0.0198 (7)	0.0026 (6)
O3	0.0352 (7)	0.0531 (8)	0.0616 (9)	-0.0021 (6)	0.0082 (6)	0.0112 (7)
O4	0.0456 (7)	0.0321 (7)	0.0627 (9)	-0.0101 (5)	0.0068 (6)	-0.0029 (6)
O5	0.0488 (7)	0.0302 (6)	0.0688 (9)	0.0022 (5)	0.0193 (6)	-0.0041 (6)
O1	0.0351 (8)	0.0586 (9)	0.1145 (14)	0.0038 (7)	0.0259 (8)	-0.0054 (9)
N1	0.0333 (8)	0.0306 (8)	0.0735 (12)	-0.0025 (6)	0.0154 (8)	-0.0090 (7)
N2	0.0389 (10)	0.0285 (8)	0.1137 (17)	-0.0032 (7)	0.0232 (10)	-0.0039 (9)
C1	0.0304 (8)	0.0282 (8)	0.0454 (9)	0.0011 (6)	0.0031 (7)	-0.0015 (7)
C2	0.0367 (9)	0.0277 (8)	0.0763 (14)	0.0020 (7)	0.0133 (9)	-0.0030 (8)
C3	0.0395 (10)	0.0335 (9)	0.0775 (14)	-0.0050 (8)	0.0137 (9)	-0.0031 (9)
C4	0.0309 (9)	0.0457 (10)	0.0639 (12)	0.0020 (8)	0.0097 (8)	-0.0032 (9)
C5	0.0448 (11)	0.0352 (10)	0.1007 (17)	0.0089 (8)	0.0210 (11)	-0.0056 (10)
C6	0.0423 (10)	0.0277 (8)	0.0793 (14)	0.0008 (7)	0.0132 (9)	-0.0048 (9)
C7	0.0328 (9)	0.0283 (8)	0.0483 (10)	-0.0003 (6)	0.0023 (7)	-0.0020 (7)
C8	0.0381 (12)	0.0773 (17)	0.112 (2)	-0.0104 (11)	0.0238 (13)	-0.0012 (15)

Geometric parameters (Å, °)

S1—O4	1.4442 (13)	C1—C6	1.391 (2)
S1—O5	1.4464 (13)	C1—C7	1.471 (2)
S1—O2	1.4504 (13)	C2—C3	1.377 (3)
S1—O3	1.5470 (15)	C2—H2	0.9300
O3—H3A	0.78 (3)	C3—C4	1.374 (3)
O1—C4	1.360 (2)	C3—H3	0.9300
O1—C8	1.427 (3)	C4—C5	1.386 (3)
N1—C7	1.306 (2)	C5—C6	1.374 (3)
N1—H1A	0.86 (2)	C5—H5	0.9300
N1—H1B	0.84 (2)	C6—H6	0.9300
N2—C7	1.308 (2)	C8—H8A	0.9739
N2—H2A	0.84 (2)	C8—H8B	0.9739
N2—H2B	0.86 (3)	C8—H8C	0.9739
C1—C2	1.381 (2)		
O4—S1—O5	112.37 (8)	C4—C3—H3	120.2
O4—S1—O2	113.83 (8)	C2—C3—H3	120.2
O5—S1—O2	111.75 (8)	O1—C4—C3	124.73 (18)
O4—S1—O3	107.52 (8)	O1—C4—C5	115.75 (17)
O5—S1—O3	107.60 (8)	C3—C4—C5	119.52 (17)
O2—S1—O3	103.05 (9)	C6—C5—C4	120.42 (17)
S1—O3—H3A	106.6 (19)	C6—C5—H5	119.8

C4—O1—C8	118.01 (17)	C4—C5—H5	119.8
C7—N1—H1A	123.0 (15)	C5—C6—C1	120.68 (17)
C7—N1—H1B	119.6 (16)	C5—C6—H6	119.7
H1A—N1—H1B	117 (2)	C1—C6—H6	119.7
C7—N2—H2A	120.2 (16)	N1—C7—N2	118.74 (18)
C7—N2—H2B	118.6 (19)	N1—C7—C1	120.48 (15)
H2A—N2—H2B	120 (2)	N2—C7—C1	120.77 (16)
C2—C1—C6	117.86 (16)	O1—C8—H8A	109.5
C2—C1—C7	120.84 (15)	O1—C8—H8B	109.5
C6—C1—C7	121.29 (15)	H8A—C8—H8B	109.5
C3—C2—C1	121.85 (16)	O1—C8—H8C	109.5
C3—C2—H2	119.1	H8A—C8—H8C	109.5
C1—C2—H2	119.1	H8B—C8—H8C	109.5
C4—C3—C2	119.65 (17)		
C6—C1—C2—C3	-1.4 (3)	C3—C4—C5—C6	-0.5 (4)
C7—C1—C2—C3	179.01 (19)	C4—C5—C6—C1	0.0 (4)
C1—C2—C3—C4	0.9 (3)	C2—C1—C6—C5	0.9 (3)
C8—O1—C4—C3	4.9 (3)	C7—C1—C6—C5	-179.5 (2)
C8—O1—C4—C5	-176.0 (2)	C2—C1—C7—N1	-6.3 (3)
C2—C3—C4—O1	179.1 (2)	C6—C1—C7—N1	174.06 (19)
C2—C3—C4—C5	0.1 (3)	C2—C1—C7—N2	172.7 (2)
O1—C4—C5—C6	-179.6 (2)	C6—C1—C7—N2	-6.9 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H3A \cdots O2 ⁱ	0.78 (3)	1.79 (3)	2.562 (2)	172 (3)
N1—H1A \cdots O4	0.86 (2)	2.10 (2)	2.938 (2)	163 (2)
N1—H1B \cdots O5 ⁱⁱ	0.84 (2)	2.10 (2)	2.884 (2)	154 (2)
N2—H2A \cdots O5	0.84 (2)	2.07 (3)	2.907 (2)	177 (2)
N2—H2B \cdots O4 ⁱⁱⁱ	0.86 (3)	2.22 (3)	2.965 (2)	145 (3)

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x, y+1/2, -z+3/2$; (iii) $-x, y-1/2, -z+3/2$.