

A Thermostable β -Glucuronidase Obtained by Directed Evolution as a Reporter Gene in Transgenic Plants

Ai-Sheng Xiong^{1*}, Ri-He Peng², Jing Zhuang³, Jian-Min Chen⁴, Bin Zhang², Jian Zhang^{3*}, Quan-Hong Yao^{2*}

1 State Key Laboratory of Crop Genetics and Germplasm Enhancement, College of Horticulture, Nanjing Agricultural University, Nanjing, China, **2** Biotechnology Research Institute, Shanghai Academy of Agricultural Sciences, Shanghai, China, **3** Alberta Innovates-Technology Futures, Vegreville, Alberta, Canada, **4** College of Bioscience and Biotechnology, Yangzhou University, Yangzhou, China

Abstract

A β -glucuronidase variant, GUS-TR3337, that was obtained by directed evolution exhibited higher thermostability than the wild-type enzyme, GUS-WT. In this study, the utility of GUS-TR3337 as an improved reporter was evaluated. The corresponding *gus-tr3337* and *gus-wt* genes were independently cloned in a plant expression vector and introduced into *Arabidopsis thaliana*. With 4-MUG as a substrate, plants containing the *gus-wt* gene showed no detectable β -glucuronidase activity after exposure to 60°C for 10 min, while those hosting the *gus-tr3337* gene retained 70% or 50% activity after exposure to 80°C for 10 min or 30 min, respectively. Similarly, *in vivo* β -glucuronidase activity could be demonstrated by using X-GLUC as a substrate in transgenic *Arabidopsis* plants hosting the *gus-tr3337* gene that were exposed to 80°C for up to 30 min. Thus, the thermostability of GUS-TR3337 can be exploited to distinguish between endogenous and transgenic β -glucuronidase activity, which is a welcome improvement in its use as a reporter.

Citation: Xiong A-S, Peng R-H, Zhuang J, Chen J-M, Zhang B, et al. (2011) A Thermostable β -Glucuronidase Obtained by Directed Evolution as a Reporter Gene in Transgenic Plants. PLoS ONE 6(11): e26773. doi:10.1371/journal.pone.0026773

Editor: Annalisa Pastore, National Institute for Medical Research, Medical Research Council, London, United Kingdom

Received: August 3, 2011; **Accepted:** October 3, 2011; **Published:** November 9, 2011

Copyright: © 2011 Xiong et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The research is supported by the Priority Academic Program Development of Jiangsu Higher Education Institutions; National key Project of Transgenic Crops of China (2009ZX08002-011B); International Scientific and Technological Cooperation (2010DFA62320). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Jing Zhuang and Jian Zhang are employees of Alberta Innovates-Technology Futures. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLoS ONE policies on sharing data and materials, as detailed online in the guide for authors.

* E-mail: xiongaisheng@njau.edu.cn (A-SX); Jian.Zhang@albertainnovates.ca (JZ); yaoquanhong@saas.sh.cn (Q-HY)

Introduction

There are about fifty reporter genes that have proved useful in transgenic plant research, based on their efficiency, scientific applications and commercialization. Reporter genes serve as indicators to study transgenic events, for example by facilitating visual identification of successfully transformed entities in a large background of non-transformed material. Although many reporter genes have been proposed, only a few of them have been commercialized [1]. β -glucuronidase (GUS) [2,3], green fluorescent protein (GFP) [4,5], luciferase (LUC) [6,7,8] are three important reporters that have a proven track record in transgenic plant research.

The advantage of GFP and LUC as reporters includes the possibility of monitoring in live tissues and in real time [1,4,8]. These two reporters allow continuous monitoring of the gene activity even through the developmental stages of the transgenic plants [1,5,9]. However, there are several disadvantages in the use of these two reporter genes. For example, the half life of LUC is short (a few hours) in plants and the process of catalyzing the release of oxyluciferin is slow [6,7,10]. In case of GFP, there may be interference due to a high background of auto-fluorescence. Another major drawback in the use GFP as a reporter is that its detection requires relatively expensive instrumentation, namely the fluorescence microscope.

The bacterial enzyme GUS encoded by the *E. coli gusA* gene is to date the most widely used as a reliable transgenic reporter system

in transgenic plant research [1]. GUS has some advantages as a reporter in transgenic plant research and commercialization. First, in genetically modified plants *E. coli* GUS can be regarded as safe for environment and consumers [11]. Second, in transgenic plants, overexpressed GUS is very stable and shows no toxicity in plant growth and development. In addition, detection of GUS does not require specialized equipment as it can easily be observed after staining with substrate such as 5-bromo-4-chloro-3-indolyl- β -D-glucuronic acid (X-GLUC). Thus, the GUS reporter system is a simple, reliable and cost-effective method to be used in transgenic studies [1,12]. Despite these benefits of the *gusA* gene, false-positives are difficult to eliminate in some plants due to endogenous GUS activity [13]. Increasing the thermal stability of GUS through directed evolution *in vitro* could potentially eliminate false-positives and dramatically increase its veracity as a reporter.

The methodology of *in vitro* directed evolution has been approved in the laboratory [14,15,16]. Directed evolution has provided significant advances in many fields, such as for biocatalysis with industrially important enzymes [17,18,19,20,21,22], plant improvement [23,24,25,26], or in vaccines and pharmaceuticals [27,28,29].

Directed evolution *in vitro* via DNA shuffling, is an important method for generating proteins with improved functions. Stemmer introduced the method of *in vitro* DNA shuffling for the formation of recombinant genes from a set of parental genes [30,31]. We have previously reported the directed evolution of a GUS variant,

GUS-TR3337, which is significantly more resistant to high temperatures than wild-type enzyme, GUS-WT [32,33]. Here, our objective was to test the utility of GUS-TR3337 as an improved reporter gene in a transgenic plant.

In order to study the activity of the thermostable GUS in plants, it was introduced into *Arabidopsis thaliana* by the floral dip method. The transgenic *Arabidopsis* plants containing the wild-type *gus-wt* gene (Genbank No. [AF485783](#)), lost most of the GUS activity within 5 min of exposure to 60°C. In contrast, *Arabidopsis* plants transformed with the *gus-tr3337* gene (GenBank No. [DQ513152](#)), retained the activity when heated upto 80°C for as long as 30 min. We confirm here that the mutant thermostable GUS gene, *gus-tr3337* is an improved reporter for transgenic research.

Materials and Methods

Chemicals, enzymes, plasmids and yeast strains

All reagents were purchased from Sigma Chemical Co. Ltd (St Louis, MO, USA) and Sinopharm Chemical Reagent Co. Ltd (Shanghai, China). Restriction enzymes were purchased from Promega (Madison, USA). High-fidelity DNA polymerase Pyrobest™ was purchased from Takara Co., Ltd (Dalian, China). DNA sequencing kit was purchased from Applied Systems Company (Foster City, CA, USA). QIAquick™ gel extraction kit was purchased from Qiagen (Stanford, CA, USA). X-GLUC and plasmid pBI121, containing the *gusA* gene, were purchased from Clontech (Palo Alto, CA, USA). *E. coli* strain DH5 α used for amplification of plasmid DNA was purchased from Stratagene (La Jolla, CA, USA).

Construction of plant expression vectors and plant transformation

The original *gus-wt* and mutant *gus-tr3337* genes were independently cloned in the binary vector pYF7716 that was constructed by our research team. This vector contains double-cauliflower mosaic virus 35S promoter, to direct transcription of the transgene as well as that of the intron-containing neomycin phosphotransferase II gene (*km*). In addition, the vector carries the omega enhancer and the nopaline synthase terminator, that were fused to the tobacco SAR (scaffold attachment region) (Figure S1). The *gus-wt* and *gus-tr3337* constructs generated in this vector were introduced into *Agrobacterium tumefaciens* GV3101 by electroporation. *A. thaliana* cv. Columbia was transformed by the floral dip method as described previously [34]. The presence of the *gus* transgenes was confirmed via PCR amplification of the genes in *Arabidopsis* plants using the following specific primers pairs: GUS-WTF1 (5'-ATGTTACGTCCTGTAGAAACCCCAACC-3') and GUS-WTR1 (5'-TCATTGTTTGCCTCCCTGCTGCGGTTT-TTC-3'), for the *gus-wt* gene, and GUS-TRF1 (5'-TCACGCCG-TACG TTATTGCCGGG-3') and GUS-TRR1 (5'-TTGCT-ACCGCAACGCGTAATATG-3') for the *gus-tr3337* gene. The PCR conditions were: denaturation for 10 min at 94°C, followed by 30 cycles, each of 40 s at 94°C, 30 s at 58°C, 2 min at 72°C; and then extension for 10 min at 72°C. Standard DNA manipulation methods were used in plasmid construction and *E. coli* transformation [35].

Histochemical analysis of GUS activity in transgenic *Arabidopsis* plants

For the analysis of expression of the *gus* transgenes, seeds of the transgenic *Arabidopsis* plants were sterilized with 1% (v/v) NaOCl solution and sown in pots containing a 1:1:1 mixture of peat moss/perlite/vermiculite. Pots were kept in a controlled-environment growth chamber at 22°C, programmed for a photoper-

iod of 16 h-day under cool white fluorescent light and 8 h-dark. High temperature treatments were performed on whole two-week old seedlings by exposure to temperatures of 70°C, 80°C, 30°C for different times (10 min, 20 min, 30, min). The heat treatment was given by placing a single seedling in a 0.6 mL capacity Eppendorf™ tube containing 150 μ L ddH₂O, and then holding the tube at the desired heating regime in a PCR System PTC-200, using the lid heating function to avoid the water evaporation. Then, the water was removed. Based on the apparent optimal temperature of β -glucuronidase activity tested in transgenic *Arabidopsis* plants, the plant samples were incubated at 45°C for transgenic *gus-wt Arabidopsis* samples or 60°C for transgenic *gus-tr3337 Arabidopsis* samples, with 1 mg/mL X-GLUC in a GUS activity buffer (50 mM sodium phosphate, pH 7.0; 1 mM EDTA; 0.1% Triton X-100; 0.5 mM K₃[Fe(CN)₆]; 0.5 mM K₄[Fe(CN)₆]·3H₂O). Histochemical staining for GUS activity was performed essentially as described by Jefferson et al. [2] and Sudan et al. [36] for most experiments. The stained samples were immersed in 70% ethanol to remove chlorophyll. The ethanol solution was changed three times at 1-h intervals.

GUS enzyme kinetics by fluorometric assay in transgenic *Arabidopsis* plants

The GUS activity in plant samples was measured by a fluorometric assay using DyNA Quant 200™ Fluorometer (Hoefel Pharmacia Biotech, San Francisco, CA, USA) as described by Martin et al. [37] and Fior et al. [38] for most experiments. Standards were prepared with different concentrations of 4-methylumbelliferone sodium salt (4-MU, Sigma Chemical Co. Ltd., No. M1508): 1 μ M, 1 mM and 100 mM, in 0.2 M sodium carbonate. 4-Methylumbelliferyl- β -D-glucuronide (4-MUG, Sigma Chemical Co. Ltd., No. M5664), 2 mM, was added to sample as substrate. GUS extraction buffer (50 mM sodium phosphate, pH 7.0; 10 mM β -mercaptoethanol; 10 mM EDTA and 0.1% Triton X-100) was used as reaction medium.

Extracts of whole three-week-old seedlings in GUS extraction buffer were used for the analysis of enzyme kinetics. Seedlings were frozen by immersion in liquid nitrogen and ground with a pestle and mortar in the presence of liquid nitrogen. GUS extraction buffer, 500 μ L, was added and the plant material was vigorously ground to a finely pulverized powder. The extract thus obtained was centrifuged at 8,000 $\times g$ for 1 min at 4°C. The supernatant was recovered and diluted one-hundred fold with GUS extraction buffer in order to decrease the effect of any potential indigenous GUS inhibitors [39]. The diluted extract, 10 μ L was added to 4-MUG, 90 μ L (2 mM), to initiate the reaction. After 1 h the reaction was stopped by the addition of 900 μ L of 0.2 M Na₂CO₃, and fluorescence was measured using a DyNA Quant 200™ spectrofluorometer.

The optimal pH for the activity of enzymes GUS-WT and GUS-TR3337 from transgenic *Arabidopsis* plants was determined at 37°C at every half point between pH values 2.5 and 13.0. The activity of both enzymes from transgenic *Arabidopsis* plants was measured at 5-degree intervals between 25°C and 90°C to determine the optimum temperature at which maximum activity was retained. The thermostability of the two enzymes from transgenic *Arabidopsis* plants was determined by measuring the residual activity at 37°C after subjecting the samples to temperatures of 60°C, 70°C, 80°C and 90°C for different times (10 min, 20 min and 30 min), followed by chilling on ice. All reactions were performed in triplicate. Kinetic parameters were determined as described in previous reports [32,40,41].

Results

Over-expression of *gus-wt* and *gus-tr3337* in *Arabidopsis*

The cloned *gus-wt* or *gus-tr3337* genes under the control of the CaMV 35S promoter were independently introduced into *Arabidopsis* (Figure S1). The transgenic *Arabidopsis* plants thus recovered showed high levels of expression of the two genes (Figure S2). There were no noticeable differences in the morphology or growth among the above transgenic plants and the parental *Arabidopsis* plants (data not shown).

Thermostability of the mutant GUS-TR3337 in transgenic *Arabidopsis* plant

High temperature treatments were performed with two-week-old whole seedlings to evaluate the thermostability of the mutant GUS-TR3337 in transgenic *Arabidopsis* plants. Seedlings of wild-type *Arabidopsis* could not be stained by X-GLUC when they were not or subjected to 60°C for 5 min (Figure 1). Seedlings containing the wild-type *gus-wt* gene or the mutant *gus-tr3337* gene exhibited stable GUS activity when they were not subjected to high temperature treatments (Figures 2A and 3A). The seedlings expressing the *gus-wt* gene were stained weakly by X-GLUC after being subjected to 60°C for 5 min or 10 min (Figures 2B and 2C) and the enzyme appeared to be inactivated by treatments for longer duration or at higher temperature (Figures 2D–F).

On the opposite, seedlings expressing the mutant *gus-tr3337* gene exhibited stable GUS activity when treated at 60°C and 70°C for 10 min, 20 min or 30 min (Figures 3C–E and F–H), and even at 80°C for 10 min, 20 min or 30 min (Figures 3I–K). However, treatment at 90°C for 10 min or longer appeared to inactivate the GUS-TR3337 enzyme (Figures 3H–I).

Characterization of the GUS-TR3337 activity in transgenic *Arabidopsis* plants

Three-week-old transgenic *Arabidopsis* seedlings containing the mutant *gus-tr3337* gene or the wild-type *gus-wt* gene were used to characterize the GUS-TR3337 and GUS-WT enzymes in seedling extracts. Three representative seedlings each (P1, P2 and P3) expressing either of transgenes were tested using 4-MUG as substrate at 37°C. A single activity peak was found at pH 6.5 in all six transgenic *Arabidopsis* lines expressing *gus-tr3337* or *gus-wt* gene (Figure 4A).




Figure 1. Histochemical staining of GUS activity in wild-type *Arabidopsis* seedlings. Seedlings were subjected to heat treatment and then stained with X-GLUC 37°C overnight. A: Control (untreated); B: 60°C for 5 min.

doi:10.1371/journal.pone.0026773.g001

The apparent optimal temperature of GUS activity tested at pH 7.0 in extracts of transgenic *Arabidopsis* plants over-expressing *gus-wt* gene was 45°C, while that of transgenic *Arabidopsis* plants over-expressing *gus-tr3337* gene was 60°C (Figure 4B). The GUS-WT activity in transgenic *Arabidopsis* lines at 45°C was about 30% higher than that at 35°C. The GUS-TR3337 activity in transgenic *Arabidopsis* lines at 60°C was >2 times of its activity at 35°C.

The GUS-WT activity in the extracts of transgenic *Arabidopsis* decreased significantly after being subjected to 60°C for 10 min, and was undetectable after treatment at 70°C or higher temperatures for 10 min or longer (Figure 5). In contrast, GUS-TR3337 activity in the transgenic *Arabidopsis* plants was much more resistant to heat treatment. Nearly 90% and 70% of GUS activity was retained when plants were heated at 70°C for 10 min or 30 min, and about 70% and 50% was retained when heated at 80°C for 10 min or 30 min.

Similarly, when heated at 70°C for 10 min or 30 min, they retained nearly 90% and 70% of this activity, respectively; and when heated at 80°C for 10 min or 30 min, retained about 70% and 50%, of this activity, respectively. The GUS-TR3337 activity decreased significantly following at heat treatment 90°C for 10 min and was undetectable if subjected to 90°C or higher temperatures for 30 min or longer (Figure 5).

Discussion

Various reporter genes encoding GFP, GAL, LUC, GUS, or oxalate oxidase (OxO) have been used in transgenic plant research or crop development, and have been assessed for the efficiency, scientific application and commercialization. Many of these genes have specific limitations or have not been sufficiently tested to merit their widespread use [1]. There are some reports on improved reporter genes that allow easy detection or a wide range of applications and that are generated by directing their evolution in the laboratory. Cramer and his colleagues used DNA shuffling to obtain GFP variants with 40 times increased fluorescence intensity [42]. A mutant GFP with stronger fluorescence intensity was selected by directed evolution in conjunction with a functional salvage screen [43].

The GUS enzyme encoded by *gusA* gene of *E. coli* is still to date the most widely used reporter gene in plants. Some limitations of using GUS as a reporter are the activity assay that is destructive to the plants and the false positive events due to endogenous GUS or GUS-like enzymes activity in some plants [44]. Hu et al. reported that there were intrinsic GUS-like activities in some seed plants [13]. Some novel GUS or GUS-like genes were also isolated and characterized from plants, such as *Scutellaria baicalensis Georgi* [45]. GUS-like enzymes were known to be present in microorganisms [2,3,46,47], animals [48,49] and plants [11,13,36,39,50], such as *Scutellaria baicalensis* [51], rye [52], and maize [53]. Sudan and his colleagues demonstrated that GUS is ubiquitously present in plants [36]. Over the past three decades, numerous reports have illustrated that there was GUS background activity in different plant species, including some model plants, such as *Arabidopsis thaliana*, *Oryza sativa*, *Nicotiana tabacum* and *Zea mays* [36,38,54,55,56].

Therefore, efforts have been focused on the suppression of the background activity [57,58,59]. As reported by Kosugi et al., the addition of methanol to 20% of the final volume during GUS staining, or during fluorimetric assays, eliminates the endogenous activities found in many plants [59]. The methanol appears to have no effect on the bacterial GUS encoded by the *gusA* from *E. coli*. However, methanol is not always effective in suppressing the endogenous GUS activities, and moreover, it is harmful to the experimentalists.




Figure 2. Histochemical staining of GUS activity in transgenic *Arabidopsis* seedlings hosting the *gus-wt* gene. Seedlings were subjected to various heat treatments and then stained at with X-GLUC 45°C overnight. (A) Control (untreated); (B) 60°C for 5 min; (C) 60°C for 10 min; (D) 60°C for 20 min; (E) 70°C for 10 min; (F) 70°C for 20 min.

doi:10.1371/journal.pone.0026773.g002

The *E. coli* GUS system is a good model system for the study of directed evolution *in vitro*. Matsumura et al. have directed the evolution of a GUS variant that is significantly more resistant to both glutaraldehyde and formaldehyde than the wild-type enzyme [40]. This GUS variant could resolve the problem of loss of GUS activity in transgenic plants during tissue fixation by glutaraldehyde or formaldehyde. After four cycles of screening the best variant was more active than the wild-type enzyme, and retained function at 70°C, whereas the wild-type enzyme lost activity at 65°C [41]. DNA shuffling (sexual recombination) and a histochemical screen have also been used to direct the evolution of *E. coli* GUS variants with improved β -glucuronidase activity [60]. The same model evolutionary system was employed to test the efficiency of several other techniques [61]. One of the GUS variants was obtained by screening for increased p-nitrophenyl- β -d-xylopyranoside (pNP-xyI) activity [62]. Although there have been a few studies on directed evolution of GUS to obtain an improved reporter, none of them have been in transgenic plants.

We obtained a temperature resistant *gusA* variant *gus-tr3337* (GenBank No. [DQ513151](#)) by employing a high throughput system for directed evolution *in vitro* [32,33]. The mutant enzyme maintained its activity even when the nitrocellulose filter containing the variant colony was heated at 100°C for 30 min.

Sequencing of the *gus-tr3337* revealed six mutations: Q493R, T509A, M532T, N550S, G559S and N566S that are present in GUS-TR3337, indicating that these were the key amino acids needed to confer its high thermostability. The purified GUS-TR3337 protein retained most of its activity when heated at 80°C for 10 min, while the wild-type GUS protein lost its activity completely after 10 min at 70°C. With pNPG as a substrate, the purified GUS-TR3337 protein retained 75% or 48% of its activity when heated at 80°C for 10 min or 30 min, respectively [33]. In the present study, with 4-MUG as substrate, transgenic *Arabidopsis* plants hosting the *gus-tr3337* gene retained about 70% and 50% of GUS activity when heated at 80°C for 10 min or 30 min, respectively. With pNPG as substrate, a single activity peak had been recorded at pH 6.5 at 37°C with the purified GUS-WT and GUS-TR3337 enzymes [33]. Similar results were obtained in the present study on GUS enzymes in extracts from transgenic *Arabidopsis* plants overexpressing the *gus-tr3337* gene and the *gus-wt* gene. The apparent optimal temperature for GUS-WT and GUS-TR3337 tested at pH 6.5 was 45°C and 65°C, respectively with pNPG as substrate; while temperature optima for the two enzymes expressed in *Arabidopsis* plants, as tested with 4-MUG as a substrate at pH 7.0 was 45°C and 60°C respectively. Thus, the apparent temperature optimum of GUS-TR3337 expressed in transgenic




Figure 3. Histochemical staining of GUS activity in transgenic *Arabidopsis* seedlings hosting the *gus-tr3337* gene. Seedlings were subjected to various heat treatments and then stained with X-GLUC 60°C overnight. (A) Control (untreated); (B) 60°C for 5 min; (C–E) 60°C for 10 min, 20 min or 30 min respectively; (F–H) 70°C for 10 min, 20 min or 30 min, respectively; (I–K) 80°C for 10 min, 20 min or 30 min, respectively; (L–N) 90°C for 10 min, 20 min or 30 min, respectively.
doi:10.1371/journal.pone.0026773.g003

plants was 5°C lower than that of purified enzyme, under the conditions of the assays. We also repeated the characterization (determination of the optimal pH, temperature and of residual activity after heat treatment) of the purified enzymes of GUS-TR3337 and GUS-WT by comparing pNPG and MUG as substrates and found that the two substrates yield similar results (Figures S3 and S4).

Although the GUS system is the most popular transgenic reporter gene for plants [1], there were some limitations of the use of GUS system as a reporter. Other potential problems with GUS, as discussed by Taylor [44], include its false positive, particularly in those plants with endogenous high endogenous GUS or GUS-like activity that interferes with the histochemical detection of

transgenic GUS. In the present study, transgenic *Arabidopsis* plants with the *gus-tr3337* gene (GenBank No. [DQ513152](#)) retained GUS enzyme activity in plant extracts as well *in vivo* even after being heated at 80°C for up to 30 min, while plants with the *gus-wt* gene lost both *in vitro* and *in vivo* enzyme activity almost completely after 10 min at 60°C. Based on the staining with 1 mg/mL X-GLUC, the apparent temperature optimum for GUS-TR3337 activity was found to be 60°C. The transgenic *Arabidopsis* plant hosting the *gus-tr3337* gene could be easily stained with X-GLUC after heated at 80°C for 10 min to 30 min. The thermostability of GUS-TR3337 can be exploited to minimize the recovery of false positives. Some limitations of GUS as a reporter could be overcome by using this thermostable GUS gene (*gus-tr3337*). Staining with X-GLUC after




Figure 4. pH- and temperature-dependence of GUS activity in transgenic *Arabidopsis* plants hosting the *gus-tr3337* gene or the *gus-wt* gene. P1, P2, P3 are triplicates. (A) Enzymatic activity in seedling extracts was measured in buffers at the indicated pH; (B) Enzymatic activity in seedling extracts was measured in phosphate buffer at pH 7.0 at the indicated temperatures. Values are mean \pm S.D. of triplicates.
doi:10.1371/journal.pone.0026773.g004




Figure 5. Thermostability of GUS in transgenic *Arabidopsis* plants. P1-, P2-, P3-GUS-TR3337: transgenic *Arabidopsis* plant lines hosting the mutant *gus-tr3337* gene; P1-, P2-, P3-GUS-WT: transgenic *Arabidopsis* plant lines hosting the wild-type *gus-wt* gene. Seedlings were subjected to heat treatments at the indicated temperature and duration and the residual GUS activity was measured in the plant extracts at 37°C. Values are mean \pm S.D. of triplicates.

doi:10.1371/journal.pone.0026773.g005

heat treatment at 70°C for 30 min or at 80° for 10–30 min will allow to distinguish the GUS activity associated to the transgenic *gus-tr3337* expression from the endogenous one, responsible of false positives, in the plants where the endogenous Gus or GUS-like activity is not thermostable. In conclusion, the use of *gus-tr3337* transgene can enhance the precision, sensitivity, and reliability of *gus* gene as a reporter.

Supporting Information

Figure S1 Schematic diagram of the vectors used in this study. The original (*gus-wt*) and mutant (*gus-tr3337*) genes was cloned into the binary vector pYF7716, at the *Bam* HI and *Sac* I restriction enzyme sites, under the control of double CaMV 35S promoter (D35S). For steady transmission of *gus-wt* and mutant *gus-tr3337*, two scaffold attachment regions (SAR) were fused upstream of the D35S promoter and downstream of the Nos-Terminator (Nos-T). (BMP)

Figure S2 Identification of T₁ generation of transgenic plants. (A) Screening of T₁ generation seeds on kanamycin

containing plates; (B) Screening of T₁ transgenic seedlings after staining for GUS activity. (TIF)

Figure S3 pH- and temperature-dependence of GUS activity in purified protein with 4-MUG as a substrate. (A) Enzymatic activity was measured at the indicated pH; (B) Enzymatic activity was measured at the indicated temperatures. Values are mean \pm S.D. of triplicates. (BMP)

Figure S4 Thermostability of GUS in purified protein with 4-MUG as a substrate. Proteins were subjected to heat treatments at the indicated temperature and duration and the residual GUS activity was measured at 37°C. Values are mean \pm S.D. of triplicates. (BMP)

Author Contributions

Conceived and designed the experiments: ASX QHY J. Zhang. Performed the experiments: ASX J. Zhuang BZ. Contributed reagents/materials/analysis tools: QHY RHP JMC. Wrote the paper: ASX. Revised the paper: ASX QHY. All authors read and approved the final manuscript.

References

- Miki B, McHugh S (2004) Selectable marker genes in transgenic plants: applications, alternatives and biosafety. *J Biotechnol* 107: 193–232.
- Jefferson RA, Kavanaugh TA, Bevan MW (1987) GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker for higher plants. *EMBO J* 6: 3901–3907.
- Jefferson RA (1989) The GUS reporter gene system. *Nature* 342: 837–838.
- Stewart CN (2001) The utility of green fluorescent protein in transgenic plants. *Plant Cell Rep* 20: 376–382.
- Haseloff J, Siemering KR (2006) The uses of green fluorescent protein in plants. *Methods Biochem Anal* 47: 259–284.
- Millar AJ, Short SR, Chua NH, Kay SA (1992) A novel circadian phenotype based on firefly luciferase expression in transgenic plants. *Plant Cell* 4: 1075–1087.
- Millar AJ, Short SR, Hiratsuka K, Chua NH, Kay SA (1992) Firefly luciferase as a reporter of regulated gene expression in higher plants. *Plant Mol Biol Rep* 10: 324–333.
- Hastings JW, Morin JG (2006) Photons for reporting molecular events: green fluorescent protein and four luciferase systems. *Methods Biochem Anal* 47: 15–38.
- Verhees J, van der Krol AR, Vreugdenhil D, van der Plas LH (2002) Characterization of gene expression during potato tuber development in individuals and populations using the luciferase reporter system. *Plant Mol Biol* 50: 653–665.
- van Leeuwen W, Hagendoorn MJM, Ruttink T, van Poecke R, van der Plas LHW, et al. (2000) The use of the luciferase reporter system for in planta gene expression studies. *Plant Mol Biol Rep* 18: 143–144.

11. Gilissen LJ, Metz PL, Stiekema WJ, Nap JP (1998) Biosafety of *E. coli* beta-glucuronidase (GUS) in plants. *Transgenic Res* 7: 157–163.
12. Mantis J, Tague RW (2000) Comparing the utility of β -glucuronidase and green fluorescent protein for detection of weak promoter activity in *Arabidopsis thaliana*. *Plant Mol Biol Rep* 18: 319–330.
13. Hu C, Chee PP, Robert H, Chesney, James H, Zhou JH, et al. (1990) Intrinsic GUS-like activities in seed plants. *Plant Cell Rep* 9: 1–5.
14. Dougherty MJ, Arnold FH (2009) Directed evolution: new parts and optimized function. *Curr Opin Biotechnol* 20: 486–491.
15. Shivange AV, Marienhagen J, Mundhada H, Schenk A, Schwaneberg U (2009) Advances in generating functional diversity for directed protein evolution. *Curr Opin Chem Biol* 13: 19–25.
16. Xiong AS, Yao QH, Peng RH, Cheng ZM (2010) Directed in vitro evolution of reporter genes based on semi-rational design and high throughput screening. *In Vitro Mutagenesis Protocols*. 3rd edition. Edited by Jeff Braman. Springer, Humana Press Inc. Totowa, NJ, USA.
17. Arnold FH, Volkov AA (1999) Directed evolution of biocatalysts. *Curr Opin Chem Biol* 3: 54–59.
18. Chirumamilla RR, Muralidhar R, Marchant R, Nigam P (2001) Improving the quality of industrially important enzymes by directed evolution. *Mol Cell Biochem* 224: 159–168.
19. Zhao H, Chockalingam K, Chen Z (2002) Directed evolution of enzymes and pathways for industrial biocatalysis. *Curr Opin Biotechnol* 13: 104–110.
20. Alexeeva M, Carr R, Turner NJ (2003) Directed evolution of enzymes: new biocatalysts for asymmetric synthesis. *Org Biomol Chem* 1: 4133–4137.
21. Otten LG, Quax WJ (2005) Directed evolution: selecting today's biocatalysts. *Biomol Eng* 22: 1–9.
22. Tian YS, Peng RH, Xu J, Zhao W, Gao F, et al. (2010) Mutations in two amino acids in phylls from *Aspergillus niger* 113 improve its phytase activity. *World J Microbiol Biotechnol* 26: 903–907.
23. Lassner M, Bedbrook J (2001) Directed molecular evolution in plant improvement. *Curr Opin Plant Biol* 4: 152–156.
24. Dixon DP, McEwen AG, Laphorn AJ, Edwards R (2003) Forced evolution of a herbicide detoxifying glutathione transferase. *J Biol Chem* 278: 23930–23935.
25. Castle LA, Siehl DL, Gorton R, Patten PA, Chen YH, et al. (2004) Discovery and directed evolution of a glyphosate tolerance gene. *Science* 304: 1151–1154.
26. Zhou M, Xu H, Wei X, Ye Z, Wei L, et al. (2006) Identification of a glyphosate-resistant mutant of rice 5-enolpyruvylshikimate 3-phosphate synthase using a directed evolution strategy. *Plant Physiol* 40: 184–195.
27. Chartrain M, Salmon PM, Robinson DK, Buckland BC (2000) Metabolic engineering and directed evolution for the production of pharmaceuticals. *Curr Opin Biotechnol* 11: 209–214.
28. Locher CP, Paldhungat M, Whalen RG, Punnonen J (2005) DNA shuffling and screening strategies for improving vaccine efficacy. *DNA Cell Biol* 24: 256–263.
29. Whalen RG, Kaiwar R, Soong NW, Punnonen J (2001) DNA shuffling and vaccines. *Curr Opin Mol Ther* 3: 31–36.
30. Stemmer WP (1994) Rapid evolution of a protein in vitro by DNA shuffling. *Nature* 370: 389–391.
31. Stemmer WP (1994) DNA shuffling by random fragmentation and reassembly: in vitro recombination for molecular evolution. *Proc Natl Acad Sci USA* 91: 10747–10751.
32. Xiong AS, Peng RH, Liu JG, Zhuang J, Qiao YS, et al. (2007) High efficiency and throughput system in directed evolution in vitro of reporter gene. *Appl Microbiol Biotechnol* 74: 160–168.
33. Xiong AS, Peng RH, Cheng ZM, Li Y, Liu JG, et al. (2007) Concurrent mutations in six amino acids in beta-glucuronidase improve its thermostability. *Protein Eng Des Sel* 20: 319–325.
34. Zhang X, Henriques R, Lin SS (2006) Agrobacterium-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nat Protoc* 1: 641–646.
35. Sambrook J, Russell DW (2001) *Molecular Cloning: A laboratory Manual*. (Cold Spring Harbor laboratory Press, Cold Spring Harbor, New York, USA.
36. Sudan C, Prakash S, Bhomkar P, Jain S, Bhalla-Sarin N (2006) Ubiquitous presence of β -glucuronidase (GUS) in plants and its regulation in some model plants. *Planta* 224: 853–864.
37. Martin T, Wohner RV, Hummel S, Willmitzer L, Frommer WB (1992) The GUS reporter system as a tool to study plant gene expression. In *GUS Protocols* Gallagher SR, ed. San Diego: Academic Press. pp 23–43.
38. Fior S, Vianelli A, Gerola PD (2009) A novel method for fluorometric continuous measurement of β -glucuronidase (GUS) activity using 4-methylumbelliferyl- β -D-glucuronide (MUG) as substrate. *Plant Sci* 176: 130–135.
39. Fior S, Gerola PD (2009) Impact of ubiquitous inhibitors on the GUS gene reporter system: evidence from the model plants *Arabidopsis*, tobacco and rice and correction methods for quantitative assays of transgenic and endogenous GUS. *Plant Methods* 5: 19.
40. Matsumura I, Wallingford JB, Surana NK, Vize PD, Ellington AD (1999) Directed evolution of the surface chemistry of the reporter enzyme beta-glucuronidase. *Nat Biotechnol* 17: 696–701.
41. Flores H, Ellington AD (2002) Increasing the thermal stability of an oligomeric protein, beta-glucuronidase. *J Mol Biol* 315: 325–337.
42. Cramer A, Whitehorn EA, Tate E, Stemmer WP (1996) Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat Biotechnol* 14: 315–319.
43. Nam SH, Oh KH, Kim GJ, Kim HS (2003) Functional tuning of a salvaged green fluorescent protein variant with a new sequence space by directed evolution. *Protein Eng* 16: 1099–1105.
44. Taylor CB (1997) Promoter fusion analysis: an insufficient measure of gene expression. *Plant Cell* 9: 273–275.
45. Sasaki K, Taura F, Shoyama Y, Morimoto S (2000) Molecular characterization of a novel beta-glucuronidase from *Scutellaria baicalensis georgi*. *J Biol Chem* 275: 27466–27472.
46. Levvy GA, Marsh CA (1959) Preparation and properties of β -glucuronidase. *Adv Carbohydr Chem* 14: 381–428.
47. Hawkesworth G, Draser BS, Hill MJ (1971) Intestinal bacteria and the hydrolysis of glycosidic bonds. *J Med Microbiol* 4: 451–459.
48. Langley SD, Wilson SD, Gros AS, Warner CK, Finnerty V (1983) A genetic variant of β -glucuronidase in *Drosophila melanogaster*. *J Biol Chem* 258: 7416–7424.
49. Sebastiano M, D'Alessio M, Bazzicalupo P (1986) β -glucuronidase mutants of the nematode *Caenorhabditis elegans*. *Genetics* 112: 459–468.
50. Plegt L, Bino J (1989) β -glucuronidase activity during development of male gametophyte from transgenic and non-transgenic plants. *Mol Gen Genet* 216: 321–327.
51. Lewy GA (1954) Baicalinase, a plant β -glucuronidase. *Biochem J* 56: 462–469.
52. Anhalt S, Weissenbock G (1992) Subcellular localization of luteolin glucuronides and related enzymes in rye mesophyll. *Planta* 187: 83–88.
53. Muhitch MJ (1998) Characterization of pedicel β -glucuronidase activity in developing maize (*Zea mays*) kernels. *Physiol Plant* 104: 423–430.
54. Bekkaoui F, Pilon M, Laine E, Raju DSS, Crosby WL, et al. (1988) Transient gene expression in electroporated *Picea glauca* protoplasts. *Plant Cell Rep* 7: 481–484.
55. Martin T, Schmidt R, Altman T, Frommer WB (1990) Non-destructive assay system for detection of β -glucuronidase activity in higher plants. *Plant Mol Biol Rep* 10: 37–46.
56. Woznaik CA, Owens LD (1994) Native β -glucuronidase activity in sugarbeet (*Beta vulgaris*). *Physiol Plant* 90: 763–771.
57. Alwen A, Moreno RMB, Vicente O, Heberle-Bors E (1992) Plant endogenous activity: how to avoid interference with the use of *E. coli* β -glucuronidase as a reporter gene in transgenic plants. *Transgenic Res* 1: 63–70.
58. Hodal L, Bocharde A, Neilson JE, Mattson O, Okkels FT (1992) Detection, expression and specific elimination of endogenous β -glucuronidase activity in transgenic and non-transgenic plants. *Plant Sci* 87: 115–122.
59. Kosugi S, Ohashi Y, Nakajima K, Arai Y (1990) An improved assay for β -glucuronidase in transformed cells: Methanol almost completely suppresses a putative endogenous β -glucuronidase activity. *Plant Sci* 70: 133–140.
60. Matsumura I, Ellington AD (2001) *In vitro* evolution of betaglucuronidase into a beta-galactosidase proceeds through nonspecific intermediates. *J Mol Biol* 305: 331–339.
61. Rowe LA, Geddie ML, Alexander OB, Matsumura I (2003) A comparison of directed evolution approaches using the beta-glucuronidase model system. *Mol Biol* 332: 851–860.
62. Geddie ML, Matsumura I (2004) Rapid evolution of beta-glucuronidase specificity by saturation mutagenesis of an active site loop. *J Biol Chem* 279: 26462–8.