



OPEN

## Ratio of venous-to-arterial $\text{PCO}_2$ to arteriovenous oxygen content difference during regional ischemic or hypoxic hypoxia

Jihad Mallat<sup>1,2,3</sup>✉ & Benoit Vallet<sup>4</sup>

The purpose of the study was to evaluate the behavior of the venous-to-arterial  $\text{CO}_2$  tension difference ( $\Delta\text{PCO}_2$ ) over the arterial-to-venous oxygen content difference ( $\Delta\text{O}_2$ ) ratio ( $\Delta\text{PCO}_2/\Delta\text{O}_2$ ) and the difference between venous-to-arterial  $\text{CO}_2$  content calculated with the Douglas' equation ( $\Delta\text{CCO}_{2D}$ ) over  $\Delta\text{O}_2$  ratio ( $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$ ) and their abilities to reflect the occurrence of anaerobic metabolism in two experimental models of tissue hypoxia: ischemic hypoxia (IH) and hypoxic hypoxia (HH). We also aimed to assess the influence of metabolic acidosis and Haldane effects on the  $\text{PCO}_2/\text{CO}_2$  content relationship. In a vascularly isolated, innervated dog hindlimb perfused with a pump-membrane oxygenator system, the oxygen delivery ( $\text{DO}_2$ ) was lowered in a stepwise manner to decrease it beyond critical  $\text{DO}_2$  ( $\text{DO}_{2\text{crit}}$ ) by lowering either arterial  $\text{PO}_2$  (HH-model) or flow (IH-model). Twelve anesthetized and mechanically ventilated dogs were studied, 6 in each model. Limb  $\text{DO}_2$ , oxygen consumption ( $\text{VO}_2$ ),  $\Delta\text{PCO}_2/\Delta\text{O}_2$ , and  $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$  were obtained every 15 min. Beyond  $\text{DO}_{2\text{crit}}$   $\text{VO}_2$  decreased, indicating dysoxia.  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$  increased significantly only after reaching  $\text{DO}_{2\text{crit}}$  in both models. At  $\text{DO}_{2\text{crit}}$   $\Delta\text{PCO}_2/\Delta\text{O}_2$  was significantly higher in the HH-model than in the IH-model ( $1.82 \pm 0.09$  vs.  $1.39 \pm 0.06$ ,  $p = 0.002$ ). At  $\text{DO}_{2\text{crit}}$   $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$  was not significantly different between the two groups ( $0.87 \pm 0.05$  for IH vs.  $1.01 \pm 0.06$  for HH,  $p = 0.09$ ). Below  $\text{DO}_{2\text{crit}}$  we observed a discrepancy between the behavior of the two indices. In both models,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  continued to increase significantly (higher in the HH-model), whereas  $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$  tended to decrease to become not significantly different from its baseline in the IH-model. Metabolic acidosis significantly influenced the  $\text{PCO}_2/\text{CO}_2$  content relationship, but not the Haldane effect.  $\Delta\text{PCO}_2/\Delta\text{O}_2$  was able to depict the occurrence of anaerobic metabolism in both tissue hypoxia models. However, at very low  $\text{DO}_2$  values,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  did not only reflect the ongoing anaerobic metabolism; it was confounded by the effects of metabolic acidosis on the  $\text{CO}_2$ -hemoglobin dissociation curve, and then it should be interpreted with caution.

### Abbreviations

$\text{CO}_2$	Carbon dioxide
$\text{VO}_2$	Oxygen consumption
$\text{VCO}_2$	Carbon dioxide production
$\text{DO}_2$	Oxygen delivery
RQ	Respiratory quotient
$\Delta\text{PCO}_2$	Venous-to-arterial carbon dioxide tension difference
$\text{CCO}_2$	$\text{CO}_2$ content
$\Delta\text{CCO}_2$	Venous-to-arterial carbon dioxide content difference
$\text{CCvCO}_2$	Venous $\text{CO}_2$ content
$\text{CCaCO}_2$	Arterial $\text{CO}_2$ content
$\Delta\text{O}_2$	Arterial-to-venous oxygen content difference
$\text{PaCO}_2$	Partial arterial carbon dioxide tension

<sup>1</sup>Department of Critical Care Medicine, Critical Care Institute, Cleveland Clinic Abu Dhabi, Al Maryah Island, Abu Dhabi, UAE. <sup>2</sup>Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH, USA. <sup>3</sup>Normandy University, UNICAEN, ED 497, Caen, France. <sup>4</sup>EA2694, University of Lille, Lille, France. ✉email: mallatjihad@gmail.com

PvCO <sub>2</sub>	Partial venous carbon dioxide tension
SvO <sub>2</sub>	Venous oxygen saturation
SaO <sub>2</sub>	Arterial oxygen saturation
PaO <sub>2</sub>	Partial arterial oxygen tension
PvO <sub>2</sub>	Partial venous oxygen tension
Hb	Hemoglobin

In a landmark study, Vallet et al. demonstrated the determinant role of blood flow in the tissue hypoxia-induced increased venous-to-arterial CO<sub>2</sub> tension difference ( $\Delta\text{PCO}_2$ )<sup>1</sup>. Their data supported the hypothesis that increases in the venous PCO<sub>2</sub> are primarily a function of changes in regional blood flow, independently of the degree of hypoxia. Gutierrez G has confirmed this conclusion in a mathematical model of tissue-to-blood CO<sub>2</sub> exchange during hypoxia<sup>2</sup>. In these previous publications, the behavior of  $\Delta\text{PCO}_2$  over the arterial-to-venous oxygen content difference ( $\Delta\text{O}_2$ ) ratio ( $\Delta\text{PCO}_2/\Delta\text{O}_2$ ), and the difference between venous-to-arterial CO<sub>2</sub> content ( $\Delta\text{CCO}_2$ ) over  $\Delta\text{O}_2$  ratio ( $\Delta\text{CCO}_2/\Delta\text{O}_2$ ) in a model of progressive tissue hypoxia generated by reducing either flow [ischemic hypoxia (IH)] or arterial oxygen tension [hypoxic hypoxia (HH)], were not investigated<sup>1,2</sup>.

Several clinical studies<sup>3–7</sup> have shown that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio, taken as a surrogate of respiratory quotient (RQ), was associated with elevated lactate levels and oxygen supply dependency considered, in those studies, as indices of global anaerobic metabolism in critically ill patients with tissue hypoperfusion. However, in an experimental study, Dubin et al. found that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio was a poor indicator of anaerobic metabolism in the hemodilution model of tissue hypoxia, where anemia was associated with preserved blood flow<sup>8</sup>. Similarly, other authors suggested that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio might not rise during tissue hypoxia conditions when associated with normal/high blood flow because venous blood flow seemed to guarantee a sufficient clearance of CO<sub>2</sub> generated by the anaerobic metabolism<sup>9</sup>. Thus, it is unclear if the  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio would be able to depict the presence of anaerobic metabolism in patients with maintained blood flow (cardiac output).

Furthermore, one estimates that the  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio might be affected by other factors than anaerobic metabolism by influencing the relationship between CO<sub>2</sub> content (CCO<sub>2</sub>) and PCO<sub>2</sub>. Indeed, metabolic acidosis can change the PCO<sub>2</sub>/CCO<sub>2</sub> relationship so that PCO<sub>2</sub> is higher for a given CCO<sub>2</sub>. Low oxygen saturation, by promoting more CO<sub>2</sub> binding to hemoglobin (Haldane effect), increases the CCO<sub>2</sub> for a given PCO<sub>2</sub><sup>10</sup>. It is not completely clear to what extent these factors would impact the PCO<sub>2</sub>/CCO<sub>2</sub> relationship and influence the  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio. Answering this question would help to define the applicability of this ratio in different clinical situations.

Therefore, we used, in secondary analysis, the original study published by Vallet et al.<sup>1</sup> with the aim to assess the behavior of  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio,  $\Delta\text{CCO}_2/\Delta\text{O}_2$  ratio, and their components in the regional model of progressive tissue hypoxia generated by IH or HH<sup>1</sup>. We also investigated the metabolic acidosis (pH) and Haldane effects on the PCO<sub>2</sub>/CCO<sub>2</sub> relationship. Since the flow was maintained unchanged in the HH model, we hypothesized that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_2/\Delta\text{O}_2$  ratios might not be able to detect the occurrence of anaerobic metabolism as the sustained blood flow would be sufficient to wash out the CO<sub>2</sub> generated by hypoxic cells in that model.

## Methods

**Animal preparation.** The original study was approved by the University of Alabama at Birmingham Institutional Animal Care and Use Committee. The study is reported in accordance with the ARRIVE guidelines. All experiments were performed in accordance with relevant guidelines and regulations. Twelve dogs of either sex and mixed breed were used<sup>1</sup>. All animals were anesthetized with intravenous 30 mg/kg of sodium phenobarbital and mechanically ventilated with a Harvard animal respirator at 10 breaths/min. Lamps suspended above the operating table were used to maintain core temperature near 37 °C. Tidal volume was varied to maintain systemic arterial PCO<sub>2</sub> between 30 and 35 mmHg. The ventilator setting was kept unchanged during the rest of the experiment. A 20 mg of succinylcholine chloride was given intramuscularly and a continuous infusion (0.1 mg/mL/min) was begun. Anesthesia depth was checked regularly by vigorous toe pinching, and additional anesthetic was given if systemic blood pressure and heart rate responded.

Catheters were inserted into the pulmonary artery (through the internal jugular vein) and common carotid artery for continuous measurements of vascular pressures and blood sampling. Arterial inflow (Q) and venous outflow from the left hindlimb were isolated, as previously described<sup>1,11</sup> (Supplemental Digital Content 1, Appendix). A roller occlusive pump directed blood flow from the right hindlimb femoral artery to the femoral artery of the vascularly isolated left hindlimb. A sampling port and pressure transducer were placed in this circuit proximal to the limb. A membrane oxygenator (model 0800-2A, Sci Med) was interposed in the perfusion circuit. A gas flow mixer (model GF-3, Cameron Instruments) supplied O<sub>2</sub>, N<sub>2</sub>, and CO<sub>2</sub> to the oxygenator, as needed, to produce normoxia or hypoxia with normocapnia in the blood supply to the hindlimb. A water bath warmed the oxygenator so that perfusion to the isolated hindlimb was at 37 °C after heat loss through the tubing.

**Measurements.** Blood samples from the carotid, femoral, and pulmonary arteries and femoral vein were obtained simultaneously. Blood gas tensions and pH were measured in an acid–base analyzer (ABL-30, Radiometer, Westlake, OH) at 37 °C and later corrected to esophageal temperature at the time of sampling. Oxygen saturation was measured with a co-oximeter calibrated for dog blood (IL-282, Instrumentation Lab, Lexington, MA). Arterial oxygen content was calculated as  $\text{CaO}_2$  (mL) =  $1.34 \times \text{Hb}$  (g/dL)  $\times \text{SaO}_2 + 0.0031 \times \text{PaO}_2$  (mmHg), where SaO<sub>2</sub> is the oxygen saturation of arterial blood, Hb the hemoglobin concentration, and PaO<sub>2</sub> the arterial oxygen tension. Hindlimb venous oxygen content was calculated as  $\text{CvO}_2$  (mL) =  $1.34 \times \text{Hb}$  (g/dL)  $\times \text{SvO}_2 + 0.0031 \times \text{PvO}_2$  (mmHg), where PvO<sub>2</sub> is the hindlimb venous oxygen tension, and SvO<sub>2</sub> is the hindlimb venous oxygen saturation.  $\Delta\text{O}_2$  was calculated as  $\text{CaO}_2 - \text{CvO}_2$ . Hindlimb VO<sub>2</sub> ( $\dot{\text{V}}\text{O}_2$ ) was calculated

as the product of Q (leg blood flow) and  $\Delta O_2$ . Hindlimb oxygen delivery ( $DO_2$ ) was calculated by using the formula:  $DO_2$  (mL/min) =  $CaO_2 \times Q \times 10$ . Hindlimb oxygen extraction (OE) was defined as:  $OE = \dot{V}O_2/DO_2$ .

$\Delta PCO_2$  was calculated as the difference between the hindlimb venous carbon dioxide tension ( $PvCO_2$ ) and hindlimb arterial  $PCO_2$  ( $PaCO_2$ ). In the original study, the hindlimb difference between venous-to-arterial  $CO_2$  content ( $CvCO_2 - CaCO_2$ ) was calculated with the McHardy equation (as proposed by Neviere et al.<sup>12</sup>):  $\Delta CCO_2 = 11.02 \times [(PvCO_2)^{0.396} - (PaCO_2)^{0.396}] - (15 - Hb) \times 0.015 \times (PvCO_2 - PaCO_2) - (95 - SaO_2) \times 0.064$ . However, the most used equation to calculate the blood  $CO_2$  content is the Douglas equation<sup>13</sup>, which includes pH:

$$\begin{aligned} \text{Blood } CO_{2D} \text{ content [blood Douglas } CCO_2 \text{ (mL)]} \\ = \text{Plasma } CCO_2 \times [1 - 0.0289 \times (Hb)/(3.352 - 0.456 \times SO_2) \times (8.142 - pH)] \end{aligned}$$

where plasma  $CCO_2 = 2.226 \times S \times \text{plasma } PCO_2 \times (1 + 10^{pH - pK'})$ ,  $CCO_2$  is  $CO_2$  content,  $SO_2$  is oxygen saturation,  $S$  is the plasma  $CO_2$  solubility coefficient, and  $pK'$  is the apparent  $pK$ .

$S$  and  $pK'$  were calculated as follow:

$$S = 0.0307 + [0.00057 \times (37 - T)] + [0.00002 \times (37 - T)^2]$$

and

$$pK' = 6.086 + [0.042 \times (7.4 - pH)] + [(38 - T) \times \{0.00472 + [0.00139 \times (7.4 - pH)]\}]$$

where  $T$  is the temperature expressed as  $^{\circ}C$ .

The difference between venous-to-arterial  $CCO_2$  calculated with the Douglas equation was:  $\Delta CCO_{2D} = CvCO_{2D} - CaCO_{2D}$ .

To investigate the metabolic acidosis and Haldane effects on the  $PCO_2/CCO_2$  relationship, default (Def) values of blood  $CCO_2$  were calculated with the Douglas's equation by using only the resting values of pH and  $SvO_2$  for each dog as following:  $DefpH - \Delta CCO_{2D} = DefpH - CvCO_{2D} - DefpH - CaCO_{2D}$ , and  $DefSvO_2 - \Delta CCO_{2D} = DefSvO_2 - CvCO_{2D} - DefSvO_2 - CaCO_{2D}$ .

Leg blood flow,  $DO_2$ , and  $\dot{V}O_2$  were reported per kilogram of muscle mass.

We also calculated the hindlimb  $\Delta PCO_2/\Delta O_2$ ,  $\Delta CCO_2/\Delta O_2$ , and  $\Delta CCO_{2D}/\Delta O_2$  ratios.

**Experimental protocol.** The experimental model was already described previously<sup>1</sup>. After all pressures and flows were stable for at least 30 min, the experiment began with a 30-min control period, during which measurements were obtained every 15 min. In the progressive ischemic hypoxia (IH) group, Q was then decreased every 15 min to produce Q values of ~60, 45, 40, 30, 20, 15, and 10 mL/kg/min. In the hypoxic hypoxia (HH) group, Q was set at 60 mL/kg/min and limb  $DO_2$  was reduced by decreasing arterial  $PO_2$  from 100 to ~15 mmHg (i.e.,  $CaO_2$  of 17 to 2 mL O<sub>2</sub>/100 mL) in eight steps at 15-min intervals. A flow rate of 60 mL/kg/min was chosen for progressive hypoxia because it is within the range of resting blood flow to normal skeletal muscle and for the practical reason that a moderate flow was necessary to achieve the desired low  $PO_2$  values using the membrane oxygenator. Oxygen and  $CO_2$ -derived variables were determined every 15 min, 13 min after the change in hindlimb arterial flow or  $PO_2$ .

For each experiment, regression lines were fitted to the delivery independent and dependent portions of the delivery-uptake curve using a dual-line, least squares method<sup>14</sup>. The intercept of these two lines defined the critical  $DO_2$  ( $DO_{2crit}$ ), that is, the delivery at which  $\dot{V}O_2$  began to fall with any further decline in  $DO_2$ .

**Statistical analysis.** All data are expressed as mean  $\pm$  SEM after assessed for normality using the Kolmogorov-Smirnov test.

Comparisons of data within and between groups were performed using a mixed ANOVA. Post-hoc paired and unpaired  $t$  tests were used, as appropriate, for one-time comparisons. The Bonferroni method was used to adjust for multiple comparisons.


Statistical analysis was performed using GraphPad Prism 6.0 software for windows (San Diego, California, USA).  $p < 0.006$  and  $p < 0.007$  were considered statistically significant for the between-group and within-group (with the baseline) comparisons, respectively. All reported  $p$  values are two-sided.

## Results


Systemic hemodynamics and oxygen-derived variables remain unchanged throughout the protocol with no differences between the IH and HH models (Supplemental Digital Content 2, Table S1).

In both groups, the  $\dot{V}O_2/DO_2$  graph depicts the typical biphasic relationship (Supplemental Digital Content 3, Figure S1). There was no statistically significant difference between the mean  $DO_{2crit}$  in the HH and IH models ( $6.9 \pm 0.6$  vs.  $6.0 \pm 0.5$  mL/kg/min,  $p = 0.28$ , respectively).  $SvO_2$  at  $DO_{2crit}$  was not statistically different between the two groups ( $25 \pm 1.7\%$  in HH vs.  $26 \pm 1.5\%$  in IH,  $p = 0.66$ ). However, for the lower  $DO_2$  values,  $SvO_2$  was significantly higher in the IH model than in the HH group (Supplemental Digital Content 4, Figure S2).  $EO_2$  at  $DO_{2crit}$  was significantly higher in the IH group than in the HH model ( $74 \pm 2\%$  vs.  $60 \pm 4\%$ ,  $p = 0.01$ ) and increased continuously and similarly in both groups (Supplemental Digital Content 5, Figure S3).  $\Delta PCO_2$  risen significantly in the IH model and did not change in the HH model (Supplemental Digital Content 6, Figure S4).

**Time course of venous-to-arterial  $CCO_2$  difference.**  $\Delta CCO_2$  calculated with the McHardy equation increased progressively along with the decrease in  $DO_2$  in the IH group but remained unchanged and even significantly decreased at the lowest  $DO_2$  value on the HH group (Fig. 1A). At  $DO_{2crit}$ ,  $\Delta CCO_2$  was significantly



**Figure 1.** Hindlimb venous-to-arterial CO<sub>2</sub> content difference ( $\Delta\text{CCO}_2$ ) calculated with McHardy equation (A) and with Douglas equation ( $\Delta\text{CCO}_{2D}$ ) (B) as a function of hindlimb oxygen delivery ( $\text{DO}_2$ ) for ischemic hypoxia model (IH) and hypoxic hypoxia model (HH). \* $p < 0.006$  vs. HH, # $p < 0.007$  vs. baseline, mixed ANOVA.



**Figure 2.** Hindlimb venous pH as a function of hindlimb oxygen delivery ( $\text{DO}_2$ ) for ischemic hypoxia model (IH) and hypoxic hypoxia model (HH). \* $p < 0.006$  vs. HH, # $p < 0.007$  vs. baseline, mixed ANOVA.


higher in the IH group than in the HH group ( $7.5 \pm 0.66$  vs.  $4.6 \pm 0.5$  mL,  $p = 0.006$ , respectively), and it was significantly different from the baseline only in the IH group ( $p = 0.0023$ ).

$\Delta\text{CCO}_{2D}$  calculated with the Douglas equation, in the IH group, increased with the decrease in  $\text{DO}_2$  down to  $\text{DO}_{2\text{crit}}$ . However, beyond  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{CCO}_{2D}$  started to decrease with the further decline in  $\text{DO}_2$  to become not significantly different from its baseline value at the lowest value of  $\text{DO}_2$  (Fig. 1B). In the HH group,  $\Delta\text{CCO}_{2D}$  had the same pattern as  $\Delta\text{CCO}_2$  calculated with the McHardy equation (Fig. 1A,B), which remained unchanged in parallel with the decreases in  $\text{DO}_2$  to become significantly lower than its baseline ( $p < 0.001$ ) only at the end of the experiment. At  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{CCO}_{2D}$  was greater in the IH group compared to the HH group ( $11.0 \pm 0.88$  vs.  $7.0 \pm 0.56$  mL,  $p = 0.003$ , respectively), and it was significantly higher than its baseline value ( $p < 0.001$ ) only in the IH group (Fig. 1B).


**pH and Haldane effects on the  $\text{PCO}_2/\text{CCO}_2$  relationship.** Hindlimb venous pH ( $\text{pH}_v$ ) remained unchanged with the decline in  $\text{DO}_2$  down to  $\text{DO}_{2\text{crit}}$  in both groups (Fig. 2). However, beyond  $\text{DO}_{2\text{crit}}$ ,  $\text{pH}_v$  decreased significantly only in the IH group and remained stable in the HH group (Fig. 2).

The venous  $\text{CCO}_2$  calculated, with the Douglas equation, by acknowledging the changes in  $\text{pH}_v$  ( $\text{CvCO}_{2D}$ ) increased first with the rise in  $\text{PvCO}_2$ , but then after, it stabilized despite further increases in  $\text{PvCO}_2$ , due to the fall in  $\text{pH}_v$ . Eventually, despite the continuously increasing  $\text{PvCO}_2$ ,  $\text{CvCO}_{2D}$  decreased due to the marked decline in  $\text{pH}_v$  (Fig. 3). On the contrary, there was almost a linear increase in  $\text{DefpH} - \text{CvCO}_{2D}$  (without accounting for the changes in  $\text{pH}_v$ ) with the increase in  $\text{PvCO}_2$  (Fig. 4). Also,  $\text{DefpH} - \Delta\text{CCO}_{2D}$  increased linearly with the decreases in  $\text{DO}_2$  in the IH group, while it remained unchanged in the HH group (Supplemental Digital Content 7, Figure S5).

The relationship between  $\text{PvCO}_2$  and  $\text{CCO}_2$  calculated without accounting for the changes in  $\text{SvO}_2$  was the same as that if we acknowledged the variations in  $\text{SvO}_2$  (Supplemental Digital Content 8, Figure S6).



**Figure 3.** Hindlimb venous  $\text{CO}_2$  content ( $\text{CCO}_{2\text{D}}$ ) as a function of hindlimb venous  $\text{PCO}_2$  for  $\text{CCO}_2$  calculated with accounting for pH changes (with\_pH) and without accounting for pH changes (without\_pH) using Douglas equation ( $\text{CCO}_{2\text{D}}$ ). \* $p < 0.006$  vs. HH, # $p < 0.007$  vs. baseline, mixed ANOVA.



**Figure 4.** Hindlimb venous-to-arterial  $\text{PCO}_2$  difference ( $\Delta\text{PCO}_2$ ) over the arterial-to-venous  $\text{O}_2$  difference ( $\Delta\text{O}_2$ ) ratio ( $\Delta\text{PCO}_2/\Delta\text{O}_2$ ) as a function of hindlimb oxygen delivery ( $\text{DO}_2$ ) for ischemic hypoxia model (IH) and hypoxic hypoxia model (HH). At  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  was significantly higher in HH model ( $1.82 \pm 0.09$ ) than IH model ( $1.39 \pm 0.06$ ). \* $p < 0.006$  vs. HH, # $p < 0.007$  vs. baseline, mixed ANOVA.

**Time course of  $\Delta\text{PCO}_2/\Delta\text{O}_2$ ,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$ , and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratios.**  $\Delta\text{O}_2$  increased significantly in the IH and decreased in the HH in parallel with the decreases in  $\text{DO}_2$  (Supplemental Digital Content 9, Figure S7).

At  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio was significantly higher in the HH group than in the IH group ( $1.82 \pm 0.09$  mmHg/mL vs.  $1.39 \pm 0.06$  mmHg/mL,  $p = 0.002$ , respectively). In both groups,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio increased significantly only after reaching  $\text{DO}_{2\text{crit}}$  (Fig. 4). Also, the increase in  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio was significantly higher in the HH than in the IH group.


$\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio increased after  $\text{DO}_{2\text{crit}}$  was reached in both groups, with a trend to decrease by the end of the experiment in the HH group (Supplemental Digital Content 10, Figure S8). At  $\text{DO}_{2\text{crit}}$ , there was no significant difference between the two groups (IH:  $0.59 \pm 0.02$  vs. HH:  $0.67 \pm 0.03$ ,  $p = 0.05$ ).

In both groups,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio increased significantly after reaching  $\text{DO}_{2\text{crit}}$ . However, in the HH group, at lower values of  $\text{DO}_2$ ,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio started to decline but remained significantly higher than its baseline value. In the IH group, beyond  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio began to decrease at a higher value of  $\text{DO}_2$  than in the HH group, to become not significantly different from its baseline value at the end of the experiment (Fig. 5). At  $\text{DO}_{2\text{crit}}$ ,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  was not significantly different between the two groups ( $0.87 \pm 0.05$  for IH vs.  $1.01 \pm 0.06$  for HH,  $p = 0.09$ ).

In both groups,  $\text{DefpH-}\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  (without accounting for pH changes) increased similarly and linearly in parallel with the decrease in  $\text{DO}_2$  (Supplemental Digital Content 11, Figure S9). The increase in  $\text{DefpH-}\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  in IH occurred before reaching  $\text{DO}_{2\text{crit}}$ .

## Discussion

The main findings of our study were that: (1) in both groups,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  as well as  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$ , and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  increases significantly in parallel with the decreases in  $\text{DO}_2$  only after reaching  $\text{DO}_{2\text{crit}}$ ; (2) beyond  $\text{DO}_{2\text{crit}}$ , the time course of  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio was different from that of  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  or  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio, in both



**Figure 5.** Hindlimb venous-to-arterial CO<sub>2</sub> content difference calculated with Douglas equation ( $\Delta\text{CCO}_{2D}$ ) over the arterial-to-venous O<sub>2</sub> difference ( $\Delta\text{O}_2$ ) ratio ( $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$ ) as a function of hindlimb oxygen delivery ( $\text{DO}_2$ ) for ischemic hypoxia model (IH) and hypoxic hypoxia model (HH). At  $\text{DO}_{2\text{crit}}$ , there was no significantly difference between HH model ( $1.01 \pm 0.06$ ) and IH model ( $0.87 \pm 0.05$ ). \* $p < 0.006$  vs. HH, # $p < 0.007$  vs. baseline, mixed ANOVA.

groups; (3) metabolic acidosis, but not Haldane effect influenced significantly the  $\text{PCO}_2/\text{CCO}_2$  relationship explaining the discrepancy between  $\Delta\text{PCO}_2$  and  $\Delta\text{CCO}_{2D}$ ; (4) the method of  $\text{CCO}_2$  calculation had a considerable impact on the results and yielded different conclusions.

Anaerobic metabolism occurrence is usually due to cellular hypoxia<sup>15</sup>. Whenever oxygen delivery decreases relative to demand, and the compensatory mechanism is exhausted, extra-mitochondrial anaerobic glycolysis occurs, and lactic acidosis develops<sup>16</sup>. We aimed to investigate if  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_{2D}/\Delta\text{O}_2$  could reflect the development of anaerobic metabolism in two regional models of tissue hypoxia: IH, where the oxygen delivery progressively decreased by decreasing the blood flow, and HH, where the blood flow was maintained unchanged, and the oxygen delivery was reduced by decreasing the arterial oxygen content.

In experimental conditions of tissue hypoxia, the drop in  $\text{VO}_2$  leads to decreased total  $\text{VCO}_2$  generation, mainly related to the decrease in aerobic CO<sub>2</sub> production. However, under situations of hypoxia, tissue CO<sub>2</sub> increases as hydrogen ions generated by anaerobic sources of energy (hydrolysis of high-energy phosphates) are buffering by bicarbonate existing in the cells (anaerobic CO<sub>2</sub> production)<sup>17</sup>. Therefore,  $\text{VCO}_2$  being reduced less than  $\text{VO}_2$ , the RQ ( $\text{VCO}_2/\text{VO}_2$ ) should increase. Accordingly, the increase in RQ has been shown to be a useful marker of global tissue hypoxia<sup>18,19</sup>. Indeed, Groeneveld et al.<sup>18</sup> observed, in an experimental model of a graded increase in positive end-expiratory pressure-induced a decrease in cardiac output and oxygen delivery in pigs, that the decline in  $\text{VCO}_2$  (by  $21 \pm 2\%$ ) was less than in  $\text{VO}_2$  (by  $27 \pm 2\%$ ).

However, airway RQ measurement necessitates a specific monitoring device (indirect calorimetry) that many hospitals might not have. Recently, there has been a growing interest in the  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio as a surrogate of the RQ to detect the development of global anaerobic metabolism in critically ill patients<sup>3-7</sup>. Indeed, several studies found an association between increased  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio and hyperlactatemia<sup>5</sup> and decreased lactate clearance<sup>6,7</sup>, which were taken as markers of anaerobic metabolism activation. We<sup>4</sup> and other authors<sup>3</sup> have also shown that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio had an excellent ability to detect the presence of  $\text{VO}_2/\text{DO}_2$  dependency phenomenon, better than central venous oxygen saturation and blood lactate levels, in septic shock patients. Recently, Mesquida et al.<sup>20</sup> reported an association between  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio and ICU mortality in septic shock patients. In contrast, in other studies,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  was unable to predict hyperlactatemia, poor lactate clearance, or  $\text{VO}_2/\text{DO}_2$  dependency and was not associated with outcome in septic shock or cardiac surgery patients<sup>9,21-23</sup>. Thus, the relationship between  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and the presence of tissue hypoxia is controversial.

Indeed, the use of  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio as a surrogate of RQ supposes that the  $\text{PCO}_2/\text{CCO}_2$  relationship is quasi-linear, which may be true over the physiological range of  $\text{PCO}_2$ <sup>24</sup>. However, this relationship can be influenced by the degree of metabolic acidosis<sup>25</sup>, hematocrit<sup>26</sup>, and oxygen saturation (Haldane effect)<sup>8,27</sup>, and it becomes nonlinear if these factors change<sup>28</sup>. Indeed, severe metabolic acidosis, low hematocrit, and high oxygen saturation can increase  $\text{PCO}_2$  for a given  $\text{CCO}_2$  since less CO<sub>2</sub> is bound to hemoglobin<sup>8</sup>. Thus,  $\Delta\text{PCO}_2$  and  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio might be increased due to several factors unrelated to the blood flow and anaerobic metabolism. We found that metabolic acidosis influenced the  $\text{PCO}_2/\text{CCO}_2$  relationship significantly. Indeed, when the changes in pH<sub>v</sub> were ignored, the  $\text{PCO}_2/\text{CCO}_2$  relationship was almost linear (Fig. 3). However,  $\text{CCO}_2$  was not linearly related to  $\text{PCO}_2$  when the changes in pH were acknowledged. In fact,  $\text{PCO}_2$  and  $\text{CCO}_2$  changed in opposite directions as metabolic acid was added to the blood by the hypoxic cells (Fig. 3). That is because metabolic acidosis causes plasma and red blood cell  $\text{CCO}_2$  and bicarbonates to decrease<sup>29</sup>. In our study, the Haldane effect did not influence the  $\text{PCO}_2/\text{CCO}_2$  relationship as the latter was the same, taking into account or not for the changes in  $\text{SvO}_2$  (Supplemental Digital Content 8, Figure S6).

Our findings suggest that, in situations with moderate/severe metabolic acidosis, an elevated  $\Delta\text{PCO}_2$  might not reflect only low or inadequate blood flow but could also be ascribed to modifications of the  $\text{CO}_2$ –hemoglobin dissociation curve. Our results are in line with previous studies. Indeed, Sun et al.<sup>29</sup> found that, in healthy subjects, during heavy exercise, changes in pH had a significant influence on the  $\text{PCO}_2/\text{CCO}_2$  relationship with  $\text{CCO}_2$  not linearly related to  $\text{PCO}_2$  and even varied in opposite directions after the lactic acidosis threshold was reached. However, in that study, changes in  $\text{SO}_2$  (Haldane effect) had a minor influence on the  $\text{PCO}_2/\text{CCO}_2$  relationship. Also, in septic shock patients, Mesquida et al.<sup>20</sup> observed that pH was the only best predictor of the discrepancy found between  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$ ; venous oxygen saturation (Haldane effect) had a minimal effect.

We observed that  $\Delta\text{PCO}_2/\Delta\text{O}_2$ , and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  significantly increased at  $\text{DO}_{2\text{crit}}$  and not before (Figs. 4 and 5), suggesting that these variables were able to depict the occurrence of oxygen supply dependency ( $\text{DO}_{2\text{crit}}$ ) in both IH and HH groups. The increases in these variables were mainly due to the decline in  $\Delta\text{O}_2$  in the HH group and the rise in  $\Delta\text{PCO}_2$  and  $\Delta\text{CCO}_2$  in the IH group induced by the decrease in blood flow. In contrast, in an experimental study of hemodilution model of tissue hypoxia, Dubin et al.<sup>8</sup> found that  $\Delta\text{PCO}_2/\Delta\text{O}_2$  significantly increased before the fall in  $\text{VO}_2$  and the sharp increase in RQ (measured by indirect calorimetry), and thus, it was a misleading indicator of anaerobic metabolism. The authors explained this finding by the effects of low hemoglobin on the  $\text{CO}_2$ –hemoglobin dissociation curve<sup>8</sup>. However, it is hard to compare these results together as the two tissue hypoxia models (HH and hemodilution) are different. Indeed, the effects of anemia on the  $\text{CO}_2$ –hemoglobin dissociation curve could be different from that of the low oxygen saturation (Haldane effect). Also, the magnitude of the decrease in venous oxygen saturation would be much more pronounced in the HH model, where the flow was maintained constant, than in the hemodilution model, where cardiac output increased by 126%<sup>8</sup>. Beyond  $\text{DO}_{2\text{crit}}$ , we observed a discrepancy between the evolutions of  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  in both groups (Figs. 4 and 5). That might be explained by the different behavior of  $\Delta\text{PCO}_2$  and  $\Delta\text{CCO}_{2\text{D}}$  at lower  $\text{DO}_2$  values. Indeed, in the IH group, these two variables changed in opposite directions:  $\Delta\text{PCO}_2$  continued to increase, whereas  $\Delta\text{CCO}_{2\text{D}}$  fell caused by metabolic acidosis (decreases in bicarbonate levels). In the HH model,  $\Delta\text{PCO}_2$  remained unchanged, whereas  $\Delta\text{CCO}_{2\text{D}}$  decreased at lower  $\text{DO}_2$  values (Fig. 1B and Supplemental Digital Content 6, Figure S4). Therefore, below  $\text{DO}_{2\text{crit}}$  and at very low  $\text{DO}_2$  values,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  ratio is confounded by the changes in the  $\text{CO}_2$ –hemoglobin curve induced by metabolic acidosis, and it does not reliably reflect the oxygen supply dependency phenomenon and the activation of anaerobic metabolism, especially in the IH tissue hypoxia model. However, in clinical practice, in such cases with very low  $\text{DO}_2$ , the clinical diagnosis of tissue hypoxia would be obvious without the need for such markers.

It is worth to note that the method of calculation of the difference in  $\text{CCO}_2$  matters as the McHardy equation<sup>12</sup>, and Douglas equation<sup>13</sup> yielded different findings (Figs. 5 and Supplemental Digital Content 10, Figure S8). However, we think that the Douglas equation is much more used in research papers, and more accurate as it accounts for much more factors such as pH.

There is no reported data, in the literature, on the behavior of  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio beyond  $\text{DO}_{2\text{crit}}$  at very low  $\text{DO}_2$  values. This ratio tended to decrease in both tissue hypoxia models, even in the presence of anaerobic  $\text{CO}_2$  production. It is possible that in case of advanced tissue hypoxia with massive decreases in  $\text{VO}_2$ , the anaerobic sources of  $\text{CO}_2$  becoming much less important than the dramatically decreased aerobic ones leading to a reduction in  $\text{VCO}_2/\text{VO}_2$  ratio.

We acknowledge several limitations to our study. First, our study was a secondary analysis that is subject to inherent limitations. Second, computation of  $\text{CCO}_2$  is subject to an important potential risk of measurement errors due to the number of variables included in the equation<sup>30</sup> that might amplify during the calculation of  $\Delta\text{CCO}_{2\text{D}}$ . Nevertheless,  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratio was already shown to be associated with mortality in septic shock patients<sup>9</sup>, suggesting that the influence of measurement errors might be limited.

## Conclusions

In both IH and HH regional models of tissue hypoxia,  $\Delta\text{PCO}_2/\Delta\text{O}_2$  and  $\Delta\text{CCO}_{2\text{D}}/\Delta\text{O}_2$  ratios both widened significantly only at the beginning of oxygen supply dependency. The hypoxic tissue hypoxia model yielded higher increases in  $\Delta\text{PCO}_2/\Delta\text{O}_2$  than the IH model. At advanced stages of tissue hypoxia (very low  $\text{DO}_2$ ),  $\Delta\text{PCO}_2/\Delta\text{O}_2$  did not only reflect the ongoing anaerobic metabolism, but it was confounded by the effects of metabolic acidosis on the  $\text{CO}_2$ –hemoglobin dissociation curve, and then it should be interpreted with caution. For clinical practice, in severe metabolic acidosis situations, elevated  $\Delta\text{PCO}_2$  may not reflect the degree of tissue hypoperfusion. In these cases, calculating the difference in  $\text{CCO}_2$  with the Douglas equation is advisable.

Received: 21 March 2021; Accepted: 28 April 2021

Published online: 13 May 2021

## References

- Vallet, B., Teboul, J. L., Cain, S. & Curtis, S. Venoarterial  $\text{CO}_2$  difference during regional ischemic or hypoxic hypoxia. *J. Appl. Physiol.* **89**, 1317–1321 (2000).
- Gutierrez, G. A mathematical model of tissue–blood carbon dioxide exchange during hypoxia. *Am. J. Respir. Crit. Care Med.* **169**, 525–533 (2004).
- Monnet, X. *et al.* Lactate and venoarterial carbon dioxide difference/arterial–venous oxygen difference ratio, but not central venous oxygen saturation, predict increase in oxygen consumption in fluid responders. *Crit. Care Med.* **41**, 1412–1420 (2013).
- Mallat, J. *et al.* Ratios of central venous-to-arterial carbon dioxide content or tension to arteriovenous oxygen content are better markers of global anaerobic metabolism than lactate in septic shock patients. *Ann. Intensive Care* **6**, 10 (2016).
- Mekontso-Dessap, A. *et al.* Combination of venoarterial  $\text{PCO}_2$  difference with arteriovenous  $\text{O}_2$  content difference to detect anaerobic metabolism in patients. *Intensive Care Med.* **28**, 272–277 (2002).

6. Mesquida, J. *et al.* Central venous-to-arterial carbon dioxide difference combined with arterial-to-venous oxygen content difference is associated with lactate evolution in the hemodynamic resuscitation process in early septic shock. *Crit. Care* **19**, 126 (2015).
7. He, H. W., Liu, D. W., Long, Y. & Wang, X. T. High central venous-to-arterial CO<sub>2</sub> difference/arterial–central venous O<sub>2</sub> difference ratio is associated with poor lactate clearance in septic patients after resuscitation. *J. Crit. Care* **31**, 76–81 (2016).
8. Dubin, A. *et al.* Venoarterial PCO<sub>2</sub>-to-arteriovenous oxygen content difference ratio is a poor surrogate for anaerobic metabolism in hemodilution: An experimental study. *Ann. Intensive Care* **7**, 65 (2017).
9. Ospina-Tascón, G. A. *et al.* Combination of arterial lactate levels and venous–arterial CO<sub>2</sub> to arterial–venous O<sub>2</sub> content difference ratio as markers of resuscitation in patients with septic shock. *Intensive Care Med.* **41**, 796–805 (2015).
10. Teboul, J. L. & Scheeren, T. Understanding the Haldane effect. *Intensive Care Med.* **43**, 91–93 (2016).
11. Cain, S. M. & Chapler, C. K. Oxygen extraction by canine hindlimb during hypoxic hypoxia. *J. Appl. Physiol.* **46**, 1023–1028 (1979).
12. Nevriere, R. *et al.* Carbon dioxide rebreathing method of cardiac output measurement during acute respiratory failure in patients with chronic obstructive pulmonary disease. *Crit. Care Med.* **22**, 81–85 (1994).
13. Douglas, A. R., Jones, N. L. & Reed, J. W. Calculation of whole blood CO<sub>2</sub> content. *J. Appl. Physiol.* **65**, 473–477 (1988).
14. Samsel, R. & Schumacker, P. T. Determination of the critical O<sub>2</sub> delivery from experimental data: Sensitivity to error. *J. Appl. Physiol.* **64**, 2074–2082 (1988).
15. Cohen, P. J. The metabolic function of oxygen and biochemical lesions of hypoxia. *Anesthesiology* **37**, 148–177 (1972).
16. Ronco, J. J. *et al.* Identification of the critical oxygen delivery for anaerobic metabolism in critically ill septic and nonseptic humans. *JAMA* **270**, 1724–1730 (1993).
17. Mallat, J., Lemyze, M., Tronchon, L., Vallet, B. & Thevenin, D. Use of venous-to-arterial carbon dioxide tension difference to guide resuscitation therapy in septic shock. *World J. Crit. Care Med.* **5**, 47–56 (2016).
18. Groeneveld, A. B., Vermeij, C. G. & Thijs, L. G. Arterial and mixed venous blood acid-base balance during hypoperfusion with incremental positive end-expiratory pressure in the pig. *Anesth. Analg.* **73**, 576–582 (1991).
19. Cohen, I. L., Sheikh, F. M., Perkins, R. J., Feustel, P. J. & Foster, E. D. Effect of hemorrhagic shock and reperfusion on the respiratory quotient in swine. *Crit. Care Med.* **23**, 545–552 (1995).
20. Mesquida, J. *et al.* Respiratory quotient estimations as additional prognostic tools in early septic shock. *J. Clin. Monit. Comput.* **32**, 1065–1072 (2018).
21. Muller, G. *et al.* Prognostic significance of central venous-to-arterial carbon dioxide difference during the first 24 hours of septic shock in patients with and without impaired cardiac function. *Br. J. Anaesth.* **119**, 239–248 (2017).
22. Abou-Arab, O. *et al.* The ratios of central venous to arterial carbon dioxide content and tension to arteriovenous oxygen content are not associated with overall anaerobic metabolism in postoperative cardiac surgery patients. *PLoS ONE* **13**, e0205950 (2018).
23. Fischer, M. O. *et al.* Assessment of macro- and micro-oxygenation parameters during fractional fluid infusion: A pilot study. *J. Crit. Care* **40**, 91–98 (2017).
24. Cavaliere, F. *et al.* Comparison of two methods to assess blood CO<sub>2</sub> equilibration curve in mechanically ventilated patients. *Respir. Physiol. Neurobiol.* **146**, 77–83 (2005).
25. Jakob, S. M., Groeneveld, A. B. & Teboul, J. L. Venous–arterial CO<sub>2</sub> to arterial–venous O<sub>2</sub> difference ratio as a resuscitation target in shock states?. *Intensive Care Med.* **41**, 91–93 (2015).
26. Chiarla, C. *et al.* Significance of hemoglobin concentration in determining blood CO<sub>2</sub> binding capacity in critical illness. *Respir. Physiol. Neurobiol.* **172**, 32–36 (2010).
27. Jakob, S. M., Kosonen, P., Ruokonen, E., Parviainen, I. & Takala, J. The Haldane effect—An alternative explanation for increasing gastric mucosal PCO<sub>2</sub> gradients?. *Br. J. Anaesth.* **83**, 740–746 (1999).
28. Mchardy, G. J. The relationship between the differences in pressure and content of carbon dioxide in arterial and venous blood. *Clin. Sci.* **32**, 299–309 (1967).
29. Sun, X. G., Hansen, J. E., Stringer, W. W., Ting, H. & Wasserman, K. Carbon dioxide pressure-concentration relationship in arterial and mixed venous blood during exercise. *J. Appl. Physiol.* **90**, 1798–1810 (2001).
30. Mallat, J. *et al.* Repeatability of blood gas parameters, PCO<sub>2</sub> gap, and PCO<sub>2</sub> gap to arterial-to-venous oxygen content difference in critically ill adult patients. *Medicine (Baltimore)* **94**, e415 (2015).

## Author contributions

J.M., and B.V. designed the study. J.M. conducted statistical analyses. J.M. and B.V. participated in manuscript writing and reviewing. All authors read and approved the final manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-89703-5>.

**Correspondence** and requests for materials should be addressed to J.M.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021