

Review Article

Molecular Fingerprints to Identify *Candida* Species

Claudia Spampinato^{1,2} and Darío Leonardi^{3,4}

¹ Departamento de Química Biológica, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario (UNR), Suipacha 531, 2000 Rosario, Argentina

² Centro de Estudios Fotosintéticos y Bioquímicos (CEFOTBI, UNR-CONICET), Suipacha 531, 2000 Rosario, Argentina

³ Departamento de Tecnología Farmacéutica, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario (UNR), Suipacha 531, 2000 Rosario, Argentina

⁴ Instituto de Química Rosario (IQUIR, UNR-CONICET), Suipacha 531, 2000 Rosario, Argentina

Correspondence should be addressed to Claudia Spampinato; spampinato@cefobi-conicet.gov.ar

Received 9 April 2013; Revised 30 May 2013; Accepted 6 June 2013

Academic Editor: Yoko Tabe

Copyright © 2013 C. Spampinato and D. Leonardi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A wide range of molecular techniques have been developed for genotyping *Candida* species. Among them, multilocus sequence typing (MLST) and microsatellite length polymorphisms (MLP) analysis have recently emerged. MLST relies on DNA sequences of internal regions of various independent housekeeping genes, while MLP identifies microsatellite instability. Both methods generate unambiguous and highly reproducible data. Here, we review the results achieved by using these two techniques and also provide a brief overview of a new method based on high-resolution DNA melting (HRM). This method identifies sequence differences by subtle deviations in sample melting profiles in the presence of saturating fluorescent DNA binding dyes.

1. Introduction

Candida species are opportunistic pathogens which can cause diseases ranging from mucosal infections to systemic mycoses depending on the vulnerability of the host. The major pathogen worldwide is *Candida albicans* [1, 2]. This fungus is detected in the body microbiota of healthy humans [3] and accounts for 75% of the organisms residing in the oral cavity [4]. It is diploid and has a largely clonal mode of reproduction. However, it can undergo considerable genetic variability either by gene regulation and/or genetic changes including chromosomal alterations, mutations, and loss of heterozygosity (LOH). In fact, LOH events lead to *MTL* homozygosity [5], azole resistance [6–8] and microevolution during infection [9–11], passage through a mammalian host [12], or *in vitro* exposure to physiologically relevant stresses [13].

Non-*albicans* *Candida* species such as *Candida glabrata*, *Candida parapsilosis*, *Candida tropicalis*, *Candida krusei*, and *Candida dubliniensis* are also found with increasing frequency [14–17]. *C. glabrata* has been reported to be the

second etiologic agent, after *C. albicans*, of superficial and invasive candidiasis in adults in the United States [18, 19], whereas, in Europe and Latin America, *C. parapsilosis* is the specie responsible for approximately 45% of all cases of candidemia [14, 20].

The ability to discriminate *Candida* isolates at the molecular level is crucial to better understand the spread of these species, particularly in hospitals and to assist in an early diagnosis and initiation of the appropriate antifungal therapy as these organisms show a range of susceptibilities to existing antifungal drugs. *C. albicans*, *C. parapsilosis*, and *C. tropicalis* remain susceptible to polyenes, azoles, and echinocandins [21]. However, *C. glabrata* and *C. krusei* show reduced triazole susceptibility [22, 23]. In addition, the majority of clade 1 isolates of *C. albicans* are less susceptible to flucytosine [24]. The faster and more accurate the species and strains can be identified, the greater the impact in the patient clinical response is. Several methods, such as pulsed-field gel electrophoresis, restriction enzyme analysis, Southern-blot assays, random amplified polymorphic DNA, and amplified fragment length

TABLE 1: International consensus gene set used for *C. albicans* MLST analysis.

Locus	Chromosome	Gene product	Primers	Sequenced fragment size (bp)
CaAAT1a	2	Aspartate aminotransferase	F: ACTCAAGCTAGATTTTTGGC R: CAGCAACATGATTAGCCC	349
CaACCI	R	Acetyl-coenzyme A carboxylase	F: GCAAGAGAAATTTTAATTCAATG R: TTCATCAACATCATCCAAGTG	407
CaADPI	1	ATP-dependent permease	F: GAGCCAAGTATGAATGATTTG R: TTGATCAACAAACCCGATAAT	443
CaPMIb	2	Mannose phosphate isomerase	F: ACCAGAAATGGCCATTGC R: GCAGCCATGCATTCAATTAT	375
CaSYA1	6	Alanyl-RNA synthetase	F: AGAAGAATTGTTGCTGTTACTG R: GTTACCTTTACCACCAGCTTT	391
CaVPS13	4	Vacuolar protein sorting protein	F: TCGTTGAGAGATATTCGACTT R: ACGGATGGATCTCCAGTCC	403
CaZWF1b	1	Glucose-6-phosphate dehydrogenase	F: GTTCATTTGATCCTGAAGC R: GCCATTGATAAGTACCTGGAT	491

F and R indicate forward and reverse primers, respectively.

polymorphism, were used to track differences among *Candida* isolates [25, 26]. However, these approaches have limitations such as time consuming, use of radioactive elements, poor reproducibility, and/or discriminatory power [25, 26]. In the present review, we summarize the most exact and/or recent DNA-based techniques developed for a better understanding of the epidemiology of *Candida* species. The availability of the *C. albicans* genome sequence [27–29] facilitated studies in comparative genomics and genome evolution.

2. Multilocus Sequence Typing

The multilocus sequence typing (MLST) is based on the analysis of nucleotide sequences of internal regions of various independent housekeeping genes. MLST studies for *C. albicans*, *C. glabrata*, *C. tropicalis*, *C. krusei*, and *C. dubliniensis* have been reported (reviewed in [30]). MLST of *C. albicans* was introduced during the early 2000s [31, 32]. On the basis of a collaborative work, an international consensus set of seven genes for *C. albicans* MLST have been proposed [33]. This gene set includes *AAT1a*, *ACCI*, *ADPI*, *MPIb*, *SYA1*, *VPS13*, and *ZWF1b* (Table 1). *MPIb* has been renamed *PMII* [34]. Table 1 also shows primers for the amplification and sequencing of the seven gene fragments.

MLST system has proved to be a useful method for epidemiological differentiation of *C. albicans* clinical isolates [31, 32]. Indeed, isolations of *C. albicans* strains recovered from human patients seem to be specific to the patient but not associated with different anatomical sources or hospital origin [9, 10, 35, 36]. MLST studies also revealed a population structure with five major clades of closely related strain types (numbered 1, 2, 3, 4, and 11) plus various minor clades [37]. Clades do not represent cryptic species as genetic exchange between and within clades is limited [38]. Clade 1 is particularly rich in flucytosine-resistant isolates [39, 40]. All clade 1 flucytosine-resistant isolates carry a point mutation (R101C) in the *FUR1* gene which encodes uridine phosphoribosyl transferase [40].

A potential weakness of the *C. albicans* international standard gene set is that three of the chromosomes are not

represented and two gene pairs are located on the same chromosome (Table 1). In order to include highly informative polymorphisms, a MLST-biased single nucleotide polymorphism (SNP) microarray has been developed [41]. This system which includes 7 loci from the consensus scheme and 12 additional discrete loci located at intervals along the 8 chromosomes may provide a basis for a standardized system.

MLST schemes have been also reported for *C. glabrata* [42]. This typing system is based on fragments of six genes ([42], Table 2). Utilizing this MLST method, several studies have described the population structure of geographically diverse collections of *C. glabrata* isolates [43–45]. Recent MLST analysis of 230 isolates of *C. glabrata* from five populations that differed both geographically and temporally confirmed that the six unlinked loci provide genotypic diversity and differentiation among isolates of this species [46]. MLST studies also revealed that *C. glabrata* strains causing bloodstream infections have similar population structures and fluconazole susceptibilities compared to those normally residing in/on the host [47]. When susceptibility testing of colonizing isolates while receiving azole therapy was studied, MLST revealed the occurrence of resistance development far more frequently in *C. glabrata* than in any other species [48]. This resistance to azole prophylaxis has led to an increased use of echinocandin for primary therapy of *C. glabrata* infections. However, decreased susceptibility to echinocandin drugs can be observed among *C. glabrata* isolates with mutations in the *FKS1* and *FKS2* genes. These genes encode Fks1p and Fks2p subunits of the 1,3- β -glucan synthase complex, which synthesizes the principal cell wall component β -1,3-glucan, target of echinocandin drugs. In light of this, MLST analysis performed on isolates with *FKS* mutations indicated that the predominant S663P mutation in the *FKS2* gene was not due to the clonal spread of a single resistant phenotype [49].

The MLST system for *C. tropicalis* comprises six housekeeping genes ([52], Table 2). Data indicate that *C. tropicalis* phylogenetically resembles *C. albicans* [53]. Both are diploid organisms, exhibit a predominant clonal mode of reproduction, and support high level of recombination events, which

TABLE 2: Summary of loci used for individual MLST schemes. Data for *C. dubliniensis*, *C. glabrata*, *C. krusei*, and *C. tropicalis* are from McManus et al. [50], Dodgson et al. [42], Jacobsen et al. [51], and Tavanti et al. [52], respectively.

Species	Locus	Gene product	Primers	Sequenced fragment size (bp)	Genotypes/site
<i>C. dubliniensis</i>	<i>CdAAT1a</i>	Aspartate aminotransferase	F: ATCAAACACTACTAAATTTTGAC R: CGGCAACCAIAGATTAGCCC	373	1.25
	<i>CdACC1</i>	Acetyl-coenzyme A carboxylase	F: GCCAGAGAAAATTTTGATCCAATGT R: TTCATCAACATCATCCAAGTG	407	1.33
	<i>CdADP1</i>	ATP-dependent permease	F: GAGCCAAGTATGAATGACTTG R: TTGATCAACAACAAACCCGATAAT	443	1.2
	<i>CdPMB</i>	Mannose phosphate isomerase	F: ACCAGAAATGGCC R: GCAGCCATACATTCAATTAT	375	3.5
	<i>CdRPN2</i>	26S proteasome regulatory subunit	F: TTTATGCACTGCTGGTACTACTGATG R: TAACCCCAATACTCAAAGCAGCAGCCT	302	1
	<i>CdSYA1</i>	Alanyl-RNA synthetase	F: AGAAGAATAGTTGCTCTTACTG R: GTTGCCCTTACCACCAGCTTT	391	1
	<i>CdVPS13</i>	Vacuolar protein sorting 13	F: CGTTGAGAGATATTCGACTT R: ACGGATCGATCGCCAATCC	403	1.33
	<i>CdZWF1b</i>	Glucose-6-phosphate dehydrogenase	F: GTTTCATTTGATCCTGAAGC R: GCCATTGATAAGTACCTGGAT	491	0.86
	<i>CgFKS</i>	1,3- β -glucan synthase	F: GTCAAATGCCACAACAACACCT R: AGCACTTCAGCAGGCTTCAG	589	1.27
	<i>CgLEU2</i>	3-Isopropylmalate dehydrogenase	F: TTTCTTGTATCCTCCCATTTGTTCA R: ATAGGTAAAGGTGGTGTGTGTGC	512	1
	<i>CgNMT1</i>	Myristoyl-coenzyme A, protein N-myristoyltransferase	F: GCCGGTGTGGTGTGCCTGCTC R: CGTTACTGCGGTGCTCGGTGTGC	607	0.81
	<i>CgTRP1</i>	Phosphoribosyl-anthranilate isomerase	F: AATGTTCAGCGTTTTTGT R: GACCAGTCCAGCTCTTTCAC	419	1.08
	<i>CgUGP1</i>	UTP-glucose-1-phosphate uridylyltransferase	F: TTTCAACACCGACAAGGACACAGA R: TCGGACTTCACTAGCAGCAAAATCA	616	0.75
<i>CgURA3</i>	Orotidine-5'-phosphate decarboxylase	F: AGCGAATTTGTAAGTTGGTTGA R: AATTCGGTTGTAAGATGATGTTC	602	0.68	

TABLE 2: Continued.

Species	Locus	Gene product	Primers	Sequenced fragment size (bp)	Genotypes/site	
<i>C. krusei</i>	<i>CkADE2</i>	Phosphoribosylaminoimidazole carboxylase	F: GTCACTTCACAGTTTGAAGC R: ACACCATCTAAAGTAGAGCC	470	2.33	
	<i>CkHIS3</i>	Imidazole glycerol phosphate dehydratase	F: GGAGGGACATATCACTGCC R: AATCTTTAATTGCCAAAGCC	400	1.75	
	<i>CkLEU2</i>	3-Isopropylmalate dehydrogenase	F: CTGTGAGACCAGAACAGGGG R: GCAGAGCCACCCAAAGTCTCC	619	1.89	
	<i>CkLYS2</i>	L-Amino adipate-semialdehyde dehydrogenase	F: ATCTGAGAAAGCAGTTGGCGC R: AGACTTGTAAAGAAATTATCCC	441	1.90	
	<i>CkNMT1</i>	Myristoyl-coenzyme A, protein N-myristoyltransferase	F: CTGATGAAAGAAATCACCCG R: GCTTGATATCATCTTTGTCC	537	2.00	
	<i>CkTRP1</i>	Phosphoribosyl-anthranilate isomerase	F: AGCTATGTGGAGCAAAGAGG R: ACATCAACGGCCACAACACC	380	2.00	
<i>C. tropicalis</i>	<i>CtCLI</i>	Isocitrate lyase	F: CAACAGATTGGTTGGCCATCAGAGC R: CGAAGTCATCAACACAGCCCAAAGCAG	447	0.71	
	<i>CtMDR1</i>	Multidrug resistance protein	F: TGTGGCATTACCCCTTCTT R: TGGAGCACCCAAACAATGGGA	425	1.67	
	<i>CtSAPT2</i>	Secreted aspartic protease 2	F: CAACGATCGTGGTGTCTG R: CACTGGTAGCTGAAGGAG	525	0.51	
	<i>CtSAPT4</i>	Secreted aspartic protease 4	F: TGCTTCTCCTACAACCTCACCTCC R: ATTCCCATGACTCCCTGAGCAACA	390	0.90	
	<i>CtXYR1</i>	D-xylose reductase I or II	F: AGTTGGTTTTCGGATGTTG R: TCGTAAATCAAAGCACCAGT	370	3.00	
	<i>CtZWF1</i>	Glucose-6-phosphate dehydrogenase	F: GGTGCTTCAGGAGATTTAGC R: ACCTTCAGTACCAAAAAGCTTC	520	0.94	

F and R indicate forward and reverse primers, respectively. Genotypes/site indicate the ratio of genotypes to SNPs.

mimic sexual reproduction processes [53]. However, unlike *C. albicans* [35], *C. tropicalis* shows a clonal cluster enriched with isolates with fluconazole resistant or “trailing growth” phenotypes [54]. The term “trailing growth” describes the growth that some isolates exhibit at drug concentrations above the minimum inhibitory concentration (MIC) after 48 h of incubation, although isolates appear fluconazole susceptible after 24 h of incubation. However, Wu et al. [55] reported that *C. tropicalis* isolates were unrelated to the fluconazole resistance pattern, suggesting that the antifungal resistance may develop geographically. Association between the MLST type of each isolate and flucytosine resistance has also been observed [40, 56, 57]. It is interesting that MLST genotypes were only distantly related, thus indicating that flucytosine resistant strains emerged independently in different geographic areas [56].

MLST gene sets for *C. krusei* and *C. dubliniensis* have also been described [50, 51]. Characteristics of the housekeeping loci used for these species are described in Table 2.

3. Microsatellite Length Polymorphisms Analysis

Microsatellite length polymorphisms (MLP) analysis identifies microsatellite instability. Microsatellites, also called simple sequence repeats (SSRs) or short tandem repeats (STRs), are tandem repeat nucleotides comprising 1–6 bp dispersed throughout the genome. These sequences undergo considerable length variations due to DNA polymerase slippage and as a consequence are highly mutagenic [58]. In *Candida* species, this technique has been applied for strain typing [43–45, 59–63], analysis of population structure [64, 65], and epidemiological studies [57, 61, 66–68]. For *C. albicans*, several polymorphic microsatellite loci have been identified (Table 3 and references therein). They were located in the promoter sequence of the elongation factor 3 (*EF3*) [60, 69], in coding regions of extracellular-signal-regulated kinase gene (*ERK1*) [70], downstream of coding sequences of cell division cycle protein (*CDC3*) [59, 60, 71] and imidazole glycerol phosphate dehydratase genes (*HIS3*) [60] and in noncoding regions (CARABEME, CAI, CAIII, CAV, CAVI, and CAVII) [44, 66, 72]. These markers were used alone or in combination. The best discriminatory powers (DPs) obtained were 0.998 for CAI, CAIII, and CAVI [44] and 0.999 for *EF3*, CAREBEME, *CDC3*, *HIS3*, *KRE6*, *LOC4 (MREII)*, *ZNF1*, CAI, CAIII, CAV, and CAVII [73]. The DP estimates the method ability to differentiate between two unrelated strains. A high DP value (close to 1) indicates that the typing method is able to distinguish each member of a strain population from all other members of that population [74]. It is noteworthy to mention that CA markers were specific for *C. albicans* [44, 66]. In fact, CA microsatellites were named after *C. albicans* and numbered according to the order of the analysis [44]. These markers are highly polymorphic since they are located outside known coding regions, thus being under inconsequential selective pressures. Recently, an allelic *CDC3* ladder has been developed for interlaboratory comparison of *C. albicans* genotyping data [75]. This ladder proved to

be important as an internal standard for a correct allele assignment.

Genotyping systems based on SSR markers have been also described for *C. glabrata*. In 2005, Foulet et al. [67] adopted three polymorphic microsatellite markers located upstream of the mitochondrial RNase P precursor (*RP2*), metallothionein 1 (*MTI*), and $\delta 5,6$ -sterol desaturase (*ERG3*) genes to generate a rapid strain typing method with a DP of 0.84. These markers were specific for *C. glabrata* isolates. Addition of three new microsatellite markers (GLM4, GLM5, and GLM6) generated a typing system with a DP value of 0.941 [76]. However, by combining only 4 microsatellite markers (*MTI*, *ERG3*, GLM4, and GLM5), authors achieved a DP value of 0.949. A different set of six different microsatellite markers located in noncoding regions (Cg4, Cg5, and Cg6) and in coding regions (Cg7, Cg10, and Cg11) have been described [68], although the highest DP value, 0.902, was reached by using a combination of only four markers (Cg4, Cg5, Cg6, and Cg10). Another research group adopted eight polymorphic microsatellite markers distributed among different chromosomes [77]. This method has a DP value of 0.97, making it suitable for tracing strains. Studies using this system indicate that *C. glabrata* is a persistent colonizer of the human tract, where it appears to undergo microevolution [78].

A highly polymorphic CKTNR locus for molecular strain typing of *C. krusei* has been identified [43]. Such locus consists of CAA repeats interspersed with CAG and CAT trinucleotides. Analysis of the CKTNR allele distribution suggested that the reproductive mode of *C. krusei* is mainly clonal [43].

MLP analysis also proved to be a reproducible method for molecular genotyping of *C. parapsilosis* [79]. Seven polymorphic loci containing dinucleotide repeats, most of them located in noncoding regions, were analyzed. The DP calculated for such loci was 0.971. These microsatellites were not amplified with DNA from single representatives of related species, *Candida orthopsilosis* and *Candida metapsilosis* [79]. Recently, another research group conducted *C. parapsilosis* typing studies using one of the previously reported marker (locus B, [79]) and three additional new microsatellite loci located outside known coding regions [80]. This multilocus analysis resulted in a DP of 0.99. These markers were also specific for the molecular typing of *C. parapsilosis* since no amplification products were obtained with DNA of *C. orthopsilosis* and *C. metapsilosis*.

4. High-Resolution DNA Melting

High-resolution DNA melting (HRM) is a novel technique for SNPs genotyping and for the identification of new genetic variants in real time (Figure 1). First, a PCR method is used to amplify specific DNA polymorphic regions in the presence of saturating DNA fluorophores [81]. The dye does not interact with single-stranded DNA but binds to double-stranded DNA, resulting in a bright structure. After PCR amplification, at the beginning of the HRM analysis, the fluorescence is high. As DNA samples are heated up, the double-stranded DNA dissociates releasing the dye which leads to a decrease in

TABLE 3: Description of SSRs used for MLPs analysis of *C. albicans*.

Locus	Ch	Gene product	Repeat motif	Location	DP	Primers	References
1	EF3	Elongation factor 3	(TTC) _n (TTTC) _n	Upstream region	0.88 ⁽¹⁾ 0.97 ⁽¹⁺³⁺⁴⁾ 0.999 ^(ALL except 2+12)	F: TTTCTCTTCCCTTTCATATAGAA R: GGATTCACATAGCAGCAGACA	[59, 60, 64, 69, 73]
2	ERK1	Extracellular-signal-regulated kinase	(CAGGCT) _n (CAAAGCT) _n --(CAA) _n --(GCCGCA) _n --(CTT) _n	Coding region	nr	F: CGACCACCGTCATCAATAGAAATCG R: CGTTGAATGAAACTTGACGAGGGG	[64, 70]
3	CARE-BEME	nc	(GAA) _n	Noncoding region	0.999 ^(ALL except 2+12)	F: GAATCATGAAACAGAAACTG R: TGGGTGAAGGATAATCTGCA	[72, 73]
4	CDC3	Cell division cycle protein	(AGTA) _n	Downstream region	0.97 ⁽¹⁺³⁺⁴⁾ 0.999 ^(ALL except 2+12)	F: CAGATGATTTTTGTATGAGAAGAA R: CAGTCACAAGATTAATAATGTTC AAG	[60, 73]
5	HIS3	Imidazole glycerol phosphate dehydratase	(ATTT) _n	Downstream region	0.97 ⁽¹⁺³⁺⁴⁾ 0.999 ^(ALL except 2+12)	F: TGGCAAAAATGATATTTCCAA R: TACACTATGCCCCAAACACA	[60, 73]
6	KRE6	β-1,6-Glucan synthesis	(AAT) _n	Coding region	0.999 ^(ALL except 2+12)	F: CAAGCTTATAGTGGCTACTA R: CCAAACATGATACATCTCG	[64, 73]
7	LOC4 (MREH1)	Double-strand break repair protein	(GAA) _n	Coding region	0.999 ^(ALL except 2+12)	F: GTAATGATTACGGCAATGAC R: AGAACGACGIGTACTATTTGG	[64, 73]
8	ZNF1	Zinc finger transcription factor	(CAA) _n	Coding region	0.999 ^(ALL except 2+12)	F: CCATTACAGCTGAACCAGCGAGGG R: CGCTAGGTAACCTACAGATTGTGGC	[64, 73]
9	CAI	nc	(CAA) _n --(CAA) _n	Noncoding region	0.967 ⁽⁹⁾ 0.998 ⁽⁹⁺¹⁰⁺¹²⁾ 0.999 ^(ALL except 2+12)	F: ATGCCATTGAGTGGAAATTGG R: AGTGGCTTGTGTGGGTTTT	[44, 66, 73]
10	CAIII	nc	(GAA) _n	Noncoding region	0.853 ⁽¹⁰⁾ 0.998 ⁽⁹⁺¹⁰⁺¹²⁾ 0.999 ^(ALL except 2+12)	F: TTGGAATCACTTCACCAGGA R: TTTCCTGGCATCAGTATCA	[44, 73]
11	CAV	nc	(ATT) _n	Noncoding region	0.853 ⁽¹¹⁾ 0.999 ^(ALL except 2+12)	F: TGCCAAATCTTGAGATACAAGTG R: CTGTCTCTCTTGTCTTAAATTG	[44, 73]
12	CAVI	nc	(TAAA) _n	Noncoding region	0.853 ⁽¹²⁾ 0.998 ⁽⁹⁺¹⁰⁺¹²⁾	F: ACAATTAAGAAAATGGATTTTAGTCAG R: TGCTGGTCTGCTGGTATTA	[44]
13	CAVII	nc	(CAAAT) _n	Noncoding region	0.670 ⁽¹³⁾ 0.999 ^(ALL except 2+12)	F: GGGGATAGAAAATGGCATCAA R: TGTGAAAACAATTCCTCCTTGC	[44, 73]

Discriminatory power (DP) based on one or various loci is indicated in brackets according to the row number. Dashed line indicates various nucleotides that separate different microsatellites in the sequence. Ch: chromosome; nc: not corresponding; nr: not reported.




FIGURE 1: Schematic representation of HRM analysis for SNPs genotyping. Arrows indicate the positions of the primers for allele amplification of a region harboring a SNP. The DNA fluorophore has a bright fluorescence when intercalated to double-stranded DNA (black circle) and low fluorescence in the unbound state (gray circles). Mismatched nucleotides are shown as diagonally broken lines. PCR products from homozygous wild type (solid lines), heterozygous mutant (dotted lines), and homozygous mutant (dashed lines) were analyzed by normalized melting curves (a), derivate melting curves (b), and difference plots (c).

the fluorescence intensity (Figure 1(a)). The observed melting temperature (T_m) and the shape of the melt curve are characteristics of the specific sequence of the fragment (primarily the GC content and the length). Data can also easily be interpreted by derivative melting curves (Figure 1(b)) and by plotting the fluorescence difference between a sample and a selected control at each temperature (Figure 1(c)) [81]. Some recent studies used HRM to differentiate clinical *Candida* species [82–84]. HRM has been proven to be a sensitive, reproducible, and inexpensive tool for a clinical laboratory but exhibits low DP values. DP for *CDC3*, *EF3*, and *HIS3* markers was 0.77 [84]. However, HRM can be used along other genotyping methods to increase the resolving power. In fact, the combination of HRM with MLP and SNaPshot minisequencing of the *CDC3* locus provided a DP value of 0.88 [83].

5. Conclusions

The development of DNA sequence-based technologies led to a great progress in understanding the epidemiology of clinical isolates of *Candida* species. Both MLST and MLP analysis offer a number of technical advantages over conventional typing methods including extremely high DP values and

reproducibility, ease of use, and rapid reliable data. The selection of the technique depends on the purpose of the study, the accessibility of genotypic strains archives, the time available to complete the analysis, and the cost. MLST remains the most reliable method for the assessment of population structure, diversity, and dynamics among *C. albicans*, whereas MLP analysis is most suitable for a rapid and less expensive study of a limited number of isolates.

Acknowledgments

This paper was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) to Claudia Spampinato (PICT 0458) and Darío Leonardi (PICT 2643), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) to Claudia Spampinato (PIP 0018), and Universidad Nacional de Rosario (UNR) to Claudia Spampinato (BIO 221) and Darío Leonardi (BIO 328). Both authors are members of the researcher career of CONICET.

References

[1] M. A. Pfaller and D. J. Diekema, “Epidemiology of invasive candidiasis: a persistent public health problem,” *Clinical Microbiology Reviews*, vol. 20, no. 1, pp. 133–163, 2007.

- [2] J. Kim and P. Sudbery, "Candida albicans, a major human fungal pathogen," *Journal of Microbiology*, vol. 49, no. 2, pp. 171–177, 2011.
- [3] S. K. Mehta, D. A. Stevens, S. K. Mishra, F. Feroze, and D. L. Pierson, "Distribution of *Candida albicans* genotypes among family members," *Diagnostic Microbiology and Infectious Disease*, vol. 34, no. 1, pp. 19–25, 1999.
- [4] M. A. Ghannoum, R. J. Jurevic, P. K. Mukherjee et al., "Characterization of the oral fungal microbiome (mycobiome) in healthy individuals," *PLoS Pathogens*, vol. 6, no. 1, Article ID e1000713, 2010.
- [5] C. M. Hull, R. M. Raisner, and A. D. Johnson, "Evidence for mating of the 'asexual' yeast *Candida albicans* in a mammalian host," *Science*, vol. 289, no. 5477, pp. 307–310, 2000.
- [6] A. Coste, A. Selmecki, A. Forche et al., "Genotypic evolution of azole resistance mechanisms in sequential *Candida albicans* isolates," *Eukaryotic Cell*, vol. 6, no. 10, pp. 1889–1904, 2007.
- [7] A. T. Coste, M. Karababa, F. Ischer, J. Bille, and D. Sanglard, "TAC1, transcriptional activator of CDR genes, is a new transcription factor involved in the regulation of *Candida albicans* ABC transporters *CDR1* and *CDR2*," *Eukaryotic Cell*, vol. 3, no. 6, pp. 1639–1652, 2004.
- [8] A. Coste, V. Turner, F. Ischer et al., "A mutation in Taclp, a transcription factor regulating *CDR1* and *CDR2*, is coupled with loss of heterozygosity at chromosome 5 to mediate antifungal resistance in *Candida albicans*," *Genetics*, vol. 172, no. 4, pp. 2139–2156, 2006.
- [9] M.-E. Bougnoux, D. Diogo, N. François et al., "Multilocus sequence typing reveals intrafamilial transmission and microevolutions of *Candida albicans* isolates from the human digestive tract," *Journal of Clinical Microbiology*, vol. 44, no. 5, pp. 1810–1820, 2006.
- [10] F. C. Odds, A. D. Davidson, M. D. Jacobsen et al., "Candida albicans strain maintenance, replacement, and microvariation demonstrated by multilocus sequence typing," *Journal of Clinical Microbiology*, vol. 44, no. 10, pp. 3647–3658, 2006.
- [11] A. Forche, G. May, and P. T. Magee, "Demonstration of loss of heterozygosity by single-nucleotide polymorphism microarray analysis and alterations in strain morphology in *Candida albicans* strains during infection," *Eukaryotic Cell*, vol. 4, no. 1, pp. 156–165, 2005.
- [12] A. Forche, P. T. Magee, A. Selmecki, J. Berman, and G. May, "Evolution in *Candida albicans* populations during a single passage through a mouse host," *Genetics*, vol. 182, no. 3, pp. 799–811, 2009.
- [13] A. Forche, D. Abbey, T. Pisithkul et al., "Stress alters rates and types of loss of heterozygosity in *Candida albicans*," *MBio*, vol. 2, no. 4, 2011.
- [14] A. M. Tortorano, C. Kibbler, J. Peman, H. Bernhardt, L. Klingspor, and R. Grillot, "Candidaemia in Europe: epidemiology and resistance," *International Journal of Antimicrobial Agents*, vol. 27, no. 5, pp. 359–366, 2006.
- [15] J. K. Chow, Y. Golan, R. Ruthazer et al., "Factors associated with candidemia caused by non-albicans *Candida* species versus *Candida albicans* in the intensive care unit," *Clinical Infectious Diseases*, vol. 46, no. 8, pp. 1206–1213, 2008.
- [16] J. D. Sobel, "The emergence of non-albicans *Candida* species as causes of invasive candidiasis and candidemia," *Current Infectious Disease Reports*, vol. 8, no. 6, pp. 427–433, 2006.
- [17] T. Y. Tan, A. L. Tan, N. W. S. Tee, L. S. Y. Ng, and C. W. J. Chee, "The increased role of non-albicans species in candidaemia: results from a 3-year surveillance study," *Mycoses*, vol. 53, no. 6, pp. 515–521, 2010.
- [18] A. Dongari-Bagtzoglou, P. Dwivedi, E. Ioannidou, M. Shaqman, D. Hull, and J. Burleson, "Oral *Candida* infection and colonization in solid organ transplant recipients," *Oral Microbiology and Immunology*, vol. 24, no. 3, pp. 249–254, 2009.
- [19] D. L. Horn, D. Neofytos, E. J. Anaissie et al., "Epidemiology and outcomes of candidemia in 2019 patients: data from the prospective antifungal therapy alliance registry," *Clinical Infectious Diseases*, vol. 48, no. 12, pp. 1695–1703, 2009.
- [20] B. Almirante, D. Rodríguez, M. Cuenca-Estrella et al., "Epidemiology, risk factors, and prognosis of *Candida parapsilosis* bloodstream infections: case-control population-based surveillance study of patients in Barcelona, Spain, from 2002 to 2003," *Journal of Clinical Microbiology*, vol. 44, no. 5, pp. 1681–1685, 2006.
- [21] M. A. Pfaller, P. G. Pappas, and J. R. Wingard, "Invasive fungal pathogens: current epidemiological trends," *Clinical Infectious Diseases*, vol. 43, 1, pp. S3–S14, 2006.
- [22] M. A. Pfaller, D. J. Diekema, L. Steele-Moore et al., "Twelve years of fluconazole in clinical practice: global-trends in species distribution and fluconazole susceptibility of bloodstream isolates of *Candida*," *Clinical Microbiology and Infection*, vol. 10, 1, pp. 11–23, 2004.
- [23] M. A. Pfaller, D. J. Diekema, R. N. Jones et al., "International surveillance of bloodstream infections due to *Candida* species: frequency of occurrence and in vitro susceptibilities to fluconazole, ravuconazole, and voriconazole of isolates collected from 1997 through 1999 in the SENTRY antimicrobial surveillance program," *Journal of Clinical Microbiology*, vol. 39, no. 9, pp. 3254–3259, 2001.
- [24] A. R. Dodgson, K. J. Dodgson, C. Pujol, M. A. Pfaller, and D. R. Soll, "Clade-specific flucytosine resistance is due to a single nucleotide change in the *FUR1* gene of *Candida albicans*," *Antimicrobial Agents and Chemotherapy*, vol. 48, no. 6, pp. 2223–2227, 2004.
- [25] F. Saghrouni, J. B. Abdeljelil, J. Boukadida, and M. B. Said, "Molecular methods for strain typing of *Candida albicans*: a review," *Journal of Applied Microbiology*, vol. 114, no. 6, pp. 1559–1574.
- [26] D. R. Soll, "The ins and outs of DNA fingerprinting the infectious fungi," *Clinical Microbiology Reviews*, vol. 13, no. 2, pp. 332–370, 2000.
- [27] T. Jones, N. A. Federspiel, H. Chibana et al., "The diploid genome sequence of *Candida albicans*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 19, pp. 7329–7334, 2004.
- [28] B. R. Braun, M. van het Hoog, C. d'Enfert et al., "A human-curated annotation of the *Candida albicans* genome," *PLoS Genetics*, vol. 1, no. 1, pp. 0036–0057, 2005.
- [29] M. Van het Hoog, T. J. Rast, M. Martchenko et al., "Assembly of the *Candida albicans* genome into sixteen supercontigs aligned on the eight chromosomes," *Genome Biology*, vol. 8, no. 4, article R52, pp. 1–11, 2007.
- [30] F. C. Odds and M. D. Jacobsen, "Multilocus sequence typing of pathogenic *Candida* species," *Eukaryotic Cell*, vol. 7, no. 7, pp. 1075–1084, 2008.
- [31] M.-E. Bougnoux, S. Morand, and C. D'Enfert, "Usefulness of multilocus sequence typing for characterization of clinical isolates of *Candida albicans*," *Journal of Clinical Microbiology*, vol. 40, no. 4, pp. 1290–1297, 2002.

- [32] A. Tavanti, N. A. R. Gow, S. Senesi, M. C. J. Maiden, and F. C. Odds, "Optimization and validation of multilocus sequence typing for *Candida albicans*," *Journal of Clinical Microbiology*, vol. 41, no. 8, pp. 3765–3776, 2003.
- [33] M.-E. Bougnoux, A. Tavanti, C. Bouchier et al., "Collaborative consensus for optimized multilocus sequence typing of *Candida albicans*," *Journal of Clinical Microbiology*, vol. 41, no. 11, pp. 5265–5266, 2003.
- [34] M. B. Arnaud, M. C. Costanzo, M. S. Skrzypek et al., "Sequence resources at the *Candida* genome database," *Nucleic Acids Research*, vol. 35, no. 1, pp. D452–D456, 2007.
- [35] K.-W. Chen, Y.-C. Chen, H.-J. Lo et al., "Multilocus sequence typing for analyses of clonality of *Candida albicans* strains in Taiwan," *Journal of Clinical Microbiology*, vol. 44, no. 6, pp. 2172–2178, 2006.
- [36] D. A. Da Matta, A. S. Melo, A. L. Colombo, J. P. Frade, M. Nucci, and T. J. Lott, "Candidemia surveillance in Brazil: evidence for a geographical boundary defining an area exhibiting an abatement of infections by *Candida albicans* group 2 strains," *Journal of Clinical Microbiology*, vol. 48, no. 9, pp. 3062–3067, 2010.
- [37] F. C. Odds, "Molecular phylogenetics and epidemiology of *Candida albicans*," *Future Microbiology*, vol. 5, no. 1, pp. 67–79, 2010.
- [38] M.-E. Bougnoux, C. Pujol, D. Diogo, C. Bouchier, D. R. Soll, and C. d'Enfert, "Mating is rare within as well as between clades of the human pathogen *Candida albicans*," *Fungal Genetics and Biology*, vol. 45, no. 3, pp. 221–231, 2008.
- [39] F. C. Odds, M.-E. Bougnoux, D. J. Shaw et al., "Molecular phylogenetics of *Candida albicans*," *Eukaryotic Cell*, vol. 6, no. 6, pp. 1041–1052, 2007.
- [40] A. Tavanti, A. D. Davidson, M. J. Fordyce, N. A. R. Gow, M. C. J. Maiden, and F. C. Odds, "Population structure and properties of *Candida albicans*, as determined by multilocus sequence typing," *Journal of Clinical Microbiology*, vol. 43, no. 11, pp. 5601–5613, 2005.
- [41] T. J. Lott and R. T. Scarborough, "Development of a MLST-biased SNP microarray for *Candida albicans*," *Fungal Genetics and Biology*, vol. 45, no. 6, pp. 803–811, 2008.
- [42] A. R. Dodgson, C. Pujol, D. W. Denning, D. R. Soll, and A. J. Fox, "Multilocus sequence typing of *Candida glabrata* reveals geographically enriched clades," *Journal of Clinical Microbiology*, vol. 41, no. 12, pp. 5709–5717, 2003.
- [43] R. Shemer, Z. Weissman, N. Hashman, and D. Kornitzer, "A highly polymorphic degenerate microsatellite for molecular strain typing of *Candida krusei*," *Microbiology*, vol. 147, no. 8, pp. 2021–2028, 2001.
- [44] P. Sampaio, L. Gusmão, A. Correia et al., "New microsatellite multiplex PCR for *Candida albicans* strain typing reveals microevolutionary changes," *Journal of Clinical Microbiology*, vol. 43, no. 8, pp. 3869–3876, 2005.
- [45] J.-M. Costa, O. Eloy, F. Botterel, G. Janbon, and S. Bretagne, "Use of microsatellite markers and gene dosage to quantify gene copy numbers in *Candida albicans*," *Journal of Clinical Microbiology*, vol. 43, no. 3, pp. 1387–1389, 2005.
- [46] T. J. Lott, J. P. Frade, and S. R. Lockhart, "Multilocus sequence type analysis reveals both clonality and recombination in populations of *Candida glabrata* bloodstream isolates from U.S. surveillance studies," *Eukaryotic Cell*, vol. 9, no. 4, pp. 619–625, 2010.
- [47] T. J. Lott, J. P. Frade, G. M. Lyon, N. Iqbal, and S. R. Lockhart, "Bloodstream and non-invasive isolates of *Candida glabrata* have similar population structures and fluconazole susceptibilities," *Medical Mycology*, vol. 50, no. 2, pp. 136–142, 2012.
- [48] P. A. Mann, P. M. McNicholas, A. S. Chau et al., "Impact of antifungal prophylaxis on colonization and azole susceptibility of *Candida* species," *Antimicrobial Agents and Chemotherapy*, vol. 53, no. 12, pp. 5026–5034, 2009.
- [49] A. J. Zimbeck, N. Iqbal, A. M. Ahlquist et al., "FKS mutations and elevated echinocandin MIC values among *Candida glabrata* isolates from U.S. population-based surveillance," *Antimicrobial Agents and Chemotherapy*, vol. 54, no. 12, pp. 5042–5047, 2010.
- [50] B. A. McManus, D. C. Coleman, G. Moran et al., "Multilocus sequence typing reveals that the population structure of *Candida dubliniensis* is significantly less divergent than that of *Candida albicans*," *Journal of Clinical Microbiology*, vol. 46, no. 2, pp. 652–664, 2008.
- [51] M. D. Jacobsen, N. A. R. Gow, M. C. J. Maiden, D. J. Shaw, and F. C. Odds, "Strain typing and determination of population structure of *Candida krusei* by multilocus sequence typing," *Journal of Clinical Microbiology*, vol. 45, no. 2, pp. 317–323, 2007.
- [52] A. Tavanti, A. D. Davidson, E. M. Johnson et al., "Multilocus sequence typing for differentiation of strains of *Candida tropicalis*," *Journal of Clinical Microbiology*, vol. 43, no. 11, pp. 5593–5600, 2005.
- [53] M. D. Jacobsen, A. D. Davidson, S.-Y. Li, D. J. Shaw, N. A. R. Gow, and F. C. Odds, "Molecular phylogenetic analysis of *Candida tropicalis* isolates by multi-locus sequence typing," *Fungal Genetics and Biology*, vol. 45, no. 6, pp. 1040–1042, 2008.
- [54] H.-H. Chou, H.-J. Lo, K.-W. Chen, M.-H. Liao, and S.-Y. Li, "Multilocus sequence typing of *Candida tropicalis* shows clonal cluster enriched in isolates with resistance or trailing growth of fluconazole," *Diagnostic Microbiology and Infectious Disease*, vol. 58, no. 4, pp. 427–433, 2007.
- [55] Y. Wu, H. Zhou, J. Wang et al., "Analysis of the clonality of *Candida tropicalis* strains from a general hospital in Beijing using multilocus sequence typing," *PLoS One*, vol. 7, Article ID e47767, 2012.
- [56] K.-W. Chen, Y.-C. Chen, Y.-H. Lin, H.-H. Chou, and S.-Y. Li, "The molecular epidemiology of serial *Candida tropicalis* isolates from ICU patients as revealed by multilocus sequence typing and pulsed-field gel electrophoresis," *Infection, Genetics and Evolution*, vol. 9, no. 5, pp. 912–920, 2009.
- [57] M. Desnos-Ollivier, S. Bretagne, C. Bernède et al., "Clonal population of flucytosine-resistant *Candida tropicalis* from blood cultures, Paris, France," *Emerging Infectious Diseases*, vol. 14, no. 4, pp. 557–565, 2008.
- [58] E. J. Oliveira, J. G. Pádua, M. I. Zucchi, R. Vencovsky, and M. L. C. Vieira, "Origin, evolution and genome distribution of microsatellites," *Genetics and Molecular Biology*, vol. 29, no. 2, pp. 294–307, 2006.
- [59] F. Dalle, N. Franco, J. Lopez et al., "Comparative genotyping of *Candida albicans* bloodstream and nonbloodstream isolates at a polymorphic microsatellite locus," *Journal of Clinical Microbiology*, vol. 38, no. 12, pp. 4554–4559, 2000.
- [60] F. Botterel, C. Desterke, C. Costa, and S. Bretagne, "Analysis of microsatellite markers of *Candida albicans* used for rapid typing," *Journal of Clinical Microbiology*, vol. 39, no. 11, pp. 4076–4081, 2001.
- [61] O. Eloy, S. Marque, F. Botterel et al., "Uniform distribution of three *Candida albicans* microsatellite markers in two French ICU populations supports a lack of nosocomial cross-contamination," *BMC Infectious Diseases*, vol. 6, article no. 162, 2006.

- [62] D. Garcia-Hermoso, O. Cabaret, G. Lecellier et al., "Comparison of microsatellite length polymorphism and multilocus sequence typing for DNA-based typing of *Candida albicans*," *Journal of Clinical Microbiology*, vol. 45, no. 12, pp. 3958–3963, 2007.
- [63] F. Stéphan, M. Sialou Bah, C. Desterke et al., "Molecular diversity and routes of colonization of *Candida albicans* in a surgical intensive care unit, as studied using microsatellite markers," *Clinical Infectious Diseases*, vol. 35, no. 12, pp. 1477–1483, 2002.
- [64] R. E. Fundyga, T. J. Lott, and J. Arnold, "Population structure of *Candida albicans*, a member of the human flora, as determined by microsatellite loci," *Infection, Genetics and Evolution*, vol. 2, no. 1, pp. 57–68, 2002.
- [65] T. J. Lott, R. E. Fundyga, M. E. Brandt et al., "Stability of allelic frequencies and distributions of *Candida albicans* microsatellite loci from U.S. population-based surveillance isolates," *Journal of Clinical Microbiology*, vol. 41, no. 3, pp. 1316–1321, 2003.
- [66] P. Sampaio, L. Gusmão, C. Alves, C. Pina-Vaz, A. Amorim, and C. Pais, "Highly polymorphic microsatellite for identification of *Candida albicans* strains," *Journal of Clinical Microbiology*, vol. 41, no. 2, pp. 552–557, 2003.
- [67] F. Foulet, N. Nicolas, O. Eloy et al., "Microsatellite marker analysis as a typing system for *Candida glabrata*," *Journal of Clinical Microbiology*, vol. 43, no. 9, pp. 4574–4579, 2005.
- [68] F. Grenouillet, L. Millon, J.-M. Bart et al., "Multiple-locus variable-number tandem-repeat analysis for rapid typing of *Candida glabrata*," *Journal of Clinical Microbiology*, vol. 45, no. 11, pp. 3781–3784, 2007.
- [69] S. Bretagne, J.-M. Costa, C. Besmond, R. Carsique, and R. Calderone, "Microsatellite polymorphism in the promoter sequence of the elongation factor 3 gene of *Candida albicans* as the basis for a typing system," *Journal of Clinical Microbiology*, vol. 35, no. 7, pp. 1777–1780, 1997.
- [70] D. Metzgar, D. Field, R. Haubrich, and C. Wills, "Sequence analysis of a compound coding-region microsatellite in *Candida albicans* resolves homoplasies and provides a high-resolution tool for genotyping," *FEMS Immunology and Medical Microbiology*, vol. 20, no. 2, pp. 103–109, 1998.
- [71] F. Dalle, L. Dumont, N. Franco et al., "Genotyping of *Candida albicans* oral strains from healthy individuals by polymorphic microsatellite locus analysis," *Journal of Clinical Microbiology*, vol. 41, no. 5, pp. 2203–2205, 2003.
- [72] F. V. Lunel, L. Licciardello, S. Stefani et al., "Lack of consistent short sequence repeat polymorphisms in genetically homologous colonizing and invasive *Candida albicans* strains," *Journal of Bacteriology*, vol. 180, no. 15, pp. 3771–3778, 1998.
- [73] C. L. ̀ollivier, C. Labruère, A. Jebrane et al., "Using a Multi-Locus Microsatellite Typing method improved phylogenetic distribution of *Candida albicans* isolates but failed to demonstrate association of some genotype with the commensal or clinical origin of the isolates," *Infection, Genetics and Evolution*, vol. 12, pp. 1949–1957, 2012.
- [74] P. R. Hunter, "Reproducibility and indices of discriminatory power of microbial typing methods," *Journal of Clinical Microbiology*, vol. 28, no. 9, pp. 1903–1905, 1990.
- [75] D. Garcia-Hermoso, D. M. MacCallum, T. J. Lott et al., "Multi-center collaborative study for standardization of *Candida albicans* genotyping using a polymorphic microsatellite marker," *Journal of Clinical Microbiology*, vol. 48, no. 7, pp. 2578–2581, 2010.
- [76] S. Abbes, H. Sellami, A. Sellami et al., "*Candida glabrata* strain relatedness by new microsatellite markers," *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 31, pp. 83–91, 2012.
- [77] S. Brisse, C. Pannier, A. Angoulvant et al., "Uneven distribution of mating types among genotypes of *Candida glabrata* isolates from clinical samples," *Eukaryotic Cell*, vol. 8, no. 3, pp. 287–295, 2009.
- [78] A. Enache-Angoulvant, M. Bourget, S. Brisse et al., "Multi-locus microsatellite markers for molecular typing of *Candida glabrata*: application to analysis of genetic relationships between bloodstream and digestive system isolates," *Journal of Clinical Microbiology*, vol. 48, no. 11, pp. 4028–4034, 2010.
- [79] B. A. Lasker, G. Butler, and T. J. Lott, "Molecular genotyping of *Candida parapsilosis* group I clinical isolates by analysis of polymorphic microsatellite markers," *Journal of Clinical Microbiology*, vol. 44, no. 3, pp. 750–759, 2006.
- [80] R. Sabino, P. Sampaio, L. Rosado, D. A. Stevens, K. V. Clemons, and C. Pais, "New polymorphic microsatellite markers able to distinguish among *Candida parapsilosis sensu stricto* isolates," *Journal of Clinical Microbiology*, vol. 48, no. 5, pp. 1677–1682, 2010.
- [81] M. Erali, K. V. Voelkerding, and C. T. Wittwer, "High resolution melting applications for clinical laboratory medicine," *Experimental and Molecular Pathology*, vol. 85, no. 1, pp. 50–58, 2008.
- [82] S. Arancia, S. Sandini, F. De Bernardis, and D. Fortini, "Rapid, simple, and low-cost identification of *Candida* species using high-resolution melting analysis," *Diagnostic Microbiology and Infectious Disease*, vol. 69, no. 3, pp. 283–285, 2011.
- [83] J.-M. Costa, D. Garcia-Hermoso, M. Olivi et al., "Genotyping of *Candida albicans* using length fragment and high-resolution melting analyses together with minisequencing of a polymorphic microsatellite locus," *Journal of Microbiological Methods*, vol. 80, pp. 306–309, 2010.
- [84] S. Gago, B. Lorenzo, A. Gomez-Lopez, I. Cuesta, M. Cuenca-Estrella, and M. J. Buitrago, "Analysis of strain relatedness using High Resolution Melting in a case of recurrent candiduria," *BMC Microbiology*, vol. 13, article 13, 2013.