

ORAL PRESENTATION**Open Access**

The Trunk Appearance Perception Scale (TAPS): a new tool to evaluate subjective impression of trunk deformity in patients with idiopathic scoliosis

J Sánchez Raya^{1*}, J Bagó¹, FJ Sánchez Perez-Grueso², JM Climent³

From 8th International Conference on Conservative Management of Spinal Deformities and SOSORT 2011
Annual Meeting
Barcelona, Spain. 19-21 May 2011

Background

Outcome assessment in idiopathic scoliosis should probably include patients' perception of their trunk deformity in addition to self-image. This can be accomplished with the Walter Reed Visual Assessment Scale (WRVAS) [1-4]. Nevertheless, this instrument has some shortcomings. These considerations prompted us to design the Trunk Appearance Perception Scale (TAPS).

Material and methods

Patients with idiopathic scoliosis and no prior surgical treatment were included. Each patient completed the TAPS and SRS-22 questionnaire and underwent a complete radiographic study of the spine [5,6]. The magnitude of the upper thoracic, main thoracic, and thoracolumbar/lumbar structural curves was recorded. The TAPS includes 3 sets of figures that depict the trunk from 3 viewpoints: looking toward the back, looking toward the head with the patient bending over and looking toward the front. Drawings are scored from 1 (greatest deformity) to 5 (smallest deformity), and a mean score is obtained.

Results

A total of 186 patients (86% females), with a mean age of 17.8 years participated. The mean of the largest curve (CMAX) was 40.2°. The median of TAPS sum score was 3.6. The floor effect was 1.6% and ceiling effect 3.8%. Cronbach's alpha coefficient was 0.89; the ICC for the mean sum score was 0.92. Correlation coefficient of the TAPS mean sum and CMAX was -0.55 ($P < 0.01$). Correlation coefficients between TAPS mean sum score and

SRS-22 scales were all statistically significant, ranging from 0.45 to 0.52 ($P < 0.05$).

Conclusions

The TAPS is a valid instrument for evaluating the perception patients have of their trunk deformity.

Author details

¹Hospital de la Vall d' Hebron, Barcelona, Spain. ²Hospital de La Paz, Madrid, Spain. ³Hospital General de Alicante, Spain.

Published: 27 January 2012

References

1. Sanders JO, Polly DW, Cats-Baril W, Jones JA, Lenke LG, O'Brien MF, et al: Analysis of patient and parent assessment of deformity in idiopathic scoliosis using the Walter-Reed Visual Assessment Scale. *Spine* 2003, 28:2158-2163.
2. Sanders JO, Harrast JJ, Kuklo TR, Polly DW, Bridwell KH, Diab M, et al: The Spinal Appearance Questionnaire. Results of reliability, validity, and responsiveness testing in patients with idiopathic scoliosis. *Spine* 2007, 32(24):2719-2722.
3. Pineda S, Bagó J, Climent JM, Gilperez C: Validity of the Walter Reed Visual Assessment Scale to measure subjective perception of spine deformity in patients with idiopathic scoliosis. *Scoliosis* 2006, 1:18.
4. Bagó J, Climent JM, Pineda S, Gilperez C: Further evaluation of the Walter Reed Visual Assessment Scale: correlation with curve pattern and radiological deformity. *Scoliosis* 2007, 2:12.
5. Asher MA, Lai SM, Burton D, Manna B: Discrimination validity of the scoliosis research society-22 patient questionnaire: relationship to idiopathic scoliosis curve pattern and curve size. *Spine* 2003, 28(1):74-78.
6. Bagó J, Climent JM, Ey A, Perez-Grueso FJ, Izquierdo E: The Spanish version of the SRS-22 patient questionnaire for idiopathic scoliosis: transcultural adaptation and reliability analysis. *Spine* 2004, 29(15):1676-1680.

doi:10.1186/1748-7161-7-S1-O1

Cite this article as: Sánchez Raya et al.: The Trunk Appearance Perception Scale (TAPS): a new tool to evaluate subjective impression of trunk deformity in patients with idiopathic scoliosis. *Scoliosis* 2012 7(Suppl 1):O1.

¹Hospital de la Vall d' Hebron, Barcelona, Spain

Full list of author information is available at the end of the article