

POSTER PRESENTATION

Open Access

A cortical attractor network with dynamic synapses

Pradeep Krishnamurthy^{1*}, Gilad Silberberg², Anders Lansner^{1,3}

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Neocortical inhibitory interneurons play a critical role in shaping the network activity patterns by directly controlling the firing rates of pyramidal cells (PC) [1]. Evidences are accumulating for the possible role of Martinotti cells (MC), which are dendrite-targeting interneurons that receive strongly facilitating synapses from PC, as opposed to basket cells (BC) that are soma targeting and receive strongly depressing synapses [2]. We have previously developed a network model of neo-

cortical layers 2/3 [3] and we here extend this set-up to explore the possible division of labour between basket and Martinotti cells. We used single-compartment cells taken from Pospischill et al. [4] and implemented in NEURON [5]. Short-term depression and facilitation were incorporated for all glutamatergic and GABAergic synapses according to the formalism of Tsodyks & Markram [6] with parameters tuned from traces provided by Silberberg et al. [2]. We commenced with




Figure 1 (A) PC-MC microcircuitry showing the disynaptic pathway between the PC mediated by MC showing frequency dependent recruitment and frequency dependent disynaptic inhibition of PC. (B) Raster plot (top) and average firing rate (bottom) of all the cells. The cells are colour-coded: PC (blue), BC (red) and MC (green).

* Correspondence: pkri@csc.kth.se

¹Department of Numerical analysis and Computer Science, Stockholm University, 114 21 Stockholm, Sweden

Full list of author information is available at the end of the article

reproducing in our model the PC – MC microcircuit, as previously described by Silberberg & Markram [2], and reproduced (a) frequency dependent disynaptic inhibition of PC and (b) frequency dependent recruitment of MC (Figure 1A). Thereafter, we integrated this microcircuit into our cortical network model to study the effects of MC on the attractor dwell time while the network is spontaneously hopping between the attractor states (stored memories) in the absence of external input. Raster plot and average firing rate (Figure 1B) show that BC that receive depressing synapses has a high firing rate at the beginning of the attractor state which then tapers off. On the other hand, MC that receive facilitating synapses display a late onset of activation and tend to terminate an ongoing attractor state. Cortex is provided with many mechanisms, e.g. spike frequency adaptation, synaptic depression of PC-PC synapses and late firing MC, to control its activity levels and termination of attractors. However, our simulations show that MC inhibition could be a dominating factor, the high divergence of MC to PC connections also assists this.

Author details

¹Department of Numerical analysis and Computer Science, Stockholm University, 114 21 Stockholm, Sweden. ²Nobel Institute of Neurophysiology, Department of Neuroscience, Karolinska Institute, Stockholm, Sweden.
³Department of Computational Biology, Royal Institute of Technology (KTH), 114 21 Stockholm, Sweden.

Published: 18 July 2011

References

1. Petilla 2008 Convention. [<http://www.columbia.edu/cu/biologfaculty/yuste/petilla/petilla-webpages/Nomenclature/PetillaNomenclaturefinal.pdf>].
2. Silberberg G, Markram H: Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells. *Neuron* 2007, 53:735-746.
3. Lundqvist M, Rehn M, Djurfeldt M, Lansner A: Attractor dynamics in a modular network of neocortex. *Network: Computation in Neural Systems* 2006, 17(3):253-276.
4. Pospisich M, Toledo-Rodriguez M, Monier C, Piwkowska Z, Bal T: Minimal Hodgkin-Huxley type models for different classes of cortical and thalamic neurons. *Biol Cybern* 2008, 99:427-441.
5. The Neuron simulation environment. [<ftp://ftp.neuron.yale.edu/neuron/papers/nsimenv.ps>].
6. Tsodyks M, Pawelzik K, Markram H: Neural networks with dynamics synapses. *Neural Comput* 1998, 10:821-835.

doi:10.1186/1471-2202-12-S1-P187

Cite this article as: Krishnamurthy et al.: A cortical attractor network with dynamic synapses. *BMC Neuroscience* 2011 12(Suppl 1):P187.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

