



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Short communication

## Kinetics of SARS-CoV-2 anti-S IgG after BNT162b2 vaccination

Daniel Grupel<sup>a,\*</sup>, Sivan Gazit<sup>b</sup>, Licita Schreiber<sup>c</sup>, Varda Nadler<sup>c</sup>, Tamar Wolf<sup>c</sup>, Rachel Lazar<sup>c</sup>, Lia Supino-Rosin<sup>c</sup>, Galit Perez<sup>b</sup>, Asaf Peretz<sup>d</sup>, Amir Ben Tov<sup>b,e</sup>, Miri Mizrahi-Reuveni<sup>f</sup>, Gabriel Chodick<sup>e,g</sup>, Tal Patalon<sup>b</sup>

<sup>a</sup> Ben-Gurion University, Beer Sheva, Israel

<sup>b</sup> Kahn Sagol Maccabi (KSM) Research & Innovation Center, Maccabi Healthcare Services, Israel

<sup>c</sup> Maccabi Healthcare Services, Central Laboratory, Rehovot, Israel

<sup>d</sup> Internal Medicine COVID-19 Ward, Samson Assuta Ashdod University Hospital, Ashdod, Israel

<sup>e</sup> Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

<sup>f</sup> Health Division, Maccabi Healthcare Services, Tel Aviv, Israel

<sup>g</sup> Maccabitech Institute for Research and Innovation, Maccabi Healthcare Services, Tel Aviv, Israel



## ARTICLE INFO

## Article history:

Received 8 March 2021

Received in revised form 4 August 2021

Accepted 7 August 2021

Available online 11 August 2021

## Keywords:

SARS-CoV-2

BNT162b2

IgG kinetics

Serology

## ABSTRACT

Deployment of the BNT162b2 mRNA Covid-19 Vaccine in Israel began in December 2020.

This is a retrospective analysis of serological data, showing SARS-CoV-2 anti-S IgG kinetics in 116 Israeli health care workers receiving BNT162b2.

Sero-conversion occurred in 14 days in all study participants, with IgG levels peaking approximately 30 days after initiation of the vaccination series.

A statistically significant difference was observed in IgG levels between subjects younger than 50 years and older participants, although in all cases, IgG levels were well above the level considered reactive by the test's manufacturer.

The importance of this difference needs to be studied further, but a potential difference in vaccine efficacy and vaccine effect length could possibly be present between these two groups.

© 2021 Elsevier Ltd. All rights reserved.

## 1. Introduction

Deployment of the BNT162b2 mRNA Covid-19 Vaccine in Israel began on December 19th 2020, following high efficacy shown in a randomized placebo-controlled trial (RCT) [1].

Recently, analyses based on real world data have been published [2,3], as different approaches attempt to overcome the difficulty in demonstrating vaccine efficacy in real world non-controlled settings.

The ability to infer immunological protection by serologic tests, as assessed in other infectious conditions, is yet unknown, as data pertaining to antibody levels in relation to clinical outcomes remains limited [4]. This possible correlation (or lack thereof) has significant implications on healthcare policy making.

This descriptive real-world study aims to assess antibody response of vaccinated individuals, by analyzing data collected from healthcare workers in Israel, among the first worldwide to

be vaccinated with BNT162b2 in two-dose course 21 days apart. Though analyses of antibody kinetics in response to COVID-19 have been examined rather extensively [5–7], data regarding antibody kinetics after vaccine administration is emerging but not yet complete: While data presented by Danese et al. [8] shows early kinetics, it is based on 3 subjects. A Larger cohort of 155 subjects is described by Favresse [9], with some sero-positive individuals, but without stating the effects of age on the response. A more comprehensive antibody panel is reported by Jalkanen [10], including IgA and IgM, but the data is less granular, and is tested at weeks 3 and 6 post vaccination. We present data highlighting the initial time period after vaccination in healthcare workers.

## 2. Methods

## 2.1. Setting and data collection

Maccabi Healthcare Services (MHS) is the second largest state-mandated, not-for-profit, healthcare provider in Israel with over 2.5 million members. Vaccine rollout in MHS began in December 2020.

\* Corresponding author at: Ben Gurion University of the Negev, MAGICAL Lab, Beer Sheva, Israel.

E-mail address: [daniel@grupel.com](mailto:daniel@grupel.com) (D. Grupel).

As part of the ongoing follow-up and quality control efforts by MHS' Health Division, serologic tests were made available to its employees in MHS' central laboratory (MegaLab). Healthcare workers over the age of 18 were serially tested for SARS-CoV-2 specific anti-s antibodies between January 14, 2020 and February 24, 2021 (six weeks).

At the time of vaccine roll-out in Israel, previously COVID-19 infected individuals were not eligible to be vaccinated, thus only employees without documented history of COVID-19 infection participated in serology collection.

The serology data was retrospectively analyzed, examining vaccine exposure (calculated as days from the first dose of vaccine administration), additional information collected was age and sex. The study was approved by the MHS's institutional review board.

### 3. Laboratory methods

A commercial assay to detect IgG antibodies against SARS-CoV-2 RBD portion of the spike protein was used - Quant II IgG anti-Spike 2-CoV-SARS by Abbott (Illinois, U.S.A.) and reported as AU/

**Table 1**  
Baseline characteristics.

	All (n = 116)	Age ≥ 50 (n = 68)	Age < 50 (n = 48)
Age y (mean [±SD])	50.3 (11.0)	57.9 (5.0)	39.6 (7.6)
Sex, male (%)	9 (7.8%)	5 (7.4%)	4 (8.3%)
Serology tests, n (mean [±SD])	3.3 (1.7)	3.3 (1.7)	3.3 (1.7)
Days between vaccine doses (mean [±SD])	21.5 (1.0)	21.3 (0.8)	21.7 (1.1)

3 individuals had not yet been eligible to receive the second dose of BNT162b2 by the time of data analysis. Two of these are older than 50.

mL (Arbitrary units). The cutoff for serology reactivity is 50 AU/mL according to the manufacturer's instructions.

Values below 21 AU/mL were truncated to 21 AU/mL and similarly for values above 40,000 AU/mL.

### 4. Statistical analyses

Geometric mean concentrations, along with associated 95% confidence intervals, were computed for each week and age group. Mann–Whitney *U* test was used to ascertain the difference in serological response between age groups using SPSS version 23 (IBM inc.). Two-sided confidence interval for weighted mean of data was calculated using Python. Categorical variables were compared using Chi-squared test or Fisher's exact test.


### 5. Results

A total of 382 serology tests were performed to 116 MHS' personnel. For each individual, there was no more than 1 test per week. Baseline characteristics are presented in Table 1. The mean age of individuals was 50.3 with 48 younger than 50 years of age and 68 were older.


Seroconversion occurred in all individuals in the studied cohort. All samples taken after day 14 were reactive (i.e. - more than 50 AU/mL), no samples were reactive before day 13.

In 16 individuals, AU values higher than the upper limit of the assay were observed. The mean day for these occurrences was 30 (SD 9.81).

Differences were observed in the anti-S IgG levels in members under 50 and over 50. Median peak IgG levels in the under 50 group were 23,801 AU/ml (IQR 14773–38471) and 13,031 (IQR 7427–21856) in the over 50 group (*P* = 0.00045). These peak values occurred in the 5th week for both groups – day 31.0 (SD 8.3) and 32.4 (SD 8.3) respectively.



**Fig. 1.** Serology results by days from first vaccine (age ≥ 50 in blue, age < 50 in orange). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 2.** Mean (95% CI) serology result per range of days from first vaccine (age ≥ 50 in blue, age < 50 in orange). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Individual IgG levels for each test are presented in Fig. 1, divided by age group.

Analysis on a per-week basis shows a lasting and statistically significant difference starting at day 15, with the population under 50 having higher IgG levels. It is important to note that while differences do exist, the values are well beyond the cutoff for reactivity. At the last week of follow-up geometric mean IgG titers for the under 50 group was 12336 (95% CI 5907–25765) and 6726 (95% CI 4015–11268) for the over 50 group.

Fig. 2 shows serology results per week (with day 0 being the day of the first BNT162b2 dose).

### 6. Discussion

In our cohort of Israeli healthcare workers, a robust antibody response is demonstrated, starting roughly two weeks after the first dose of BNT162b2 and peaking around day 30 (approximately 10 days after the second dose).

Decline is slow after peaking, though data was limited to six weeks.

A statistically significant difference in IgG levels was observed between employees over the age of 50 and younger individuals, nevertheless, levels remained well above the threshold for serological reactivity. Although our results are not in line with a previous work evaluating a single time point after natural infection, and showed higher IgG titers with advancing age [11], a non linear correlation to age was shown by Yang et al and very low levels in young adults – alluding to a possible confounder in these results. Our data are in accordance with not-yet peer reviewed data from a study evaluating BNT162b2-induced antibody responses [12].

The importance of this difference needs to be studied further, but a potential difference in vaccine efficacy and vaccine effect length could possibly be present between these two groups.

This study has several limitations. The available cohort is small and not representative of the general population (especially male to female ratio).

Since baseline serology was available for only a single individual, previous infection in other subjects can not be completely ruled out. Had such an infection occurred, a rapid increase in IgG [13,14,15] would have been observed, skewing our results. However, we believe previous infection to be unreasonable since vaccine eligibility at the time of the study solely included people with no documented past infection and while our data demonstrated marked heterogeneity in the 7–14 days period, none of the subjects reached levels similar to the peak levels at this point (as would have been expected with a previous exposure [9]).

### 7. Conclusion

This study demonstrated the kinetics of the antibody response post vaccination with BNT162b2. The robustness of seroconversion was observed, alongside a significant age dependent difference in antibody response. Further research is required in order to examine whether the difference persists over time.

More importantly, further investigations should be conducted to evaluate the correlation between seroconversion and clinical course of COVID-19, as well as examining other factors that might affect magnitude of seroconversion as larger datasets become available.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### References

[1] Polack FP et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med* 2020;383:2603–15.  
 [2] Dagan N et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med* 1–12 (2021). doi: 10.1056/NEJMoa2101765.

- [3] Amit S, Regev-yochay G, Afek A, Kreiss Y, Leshem E. Early rate reductions of and COVID-19 in BNT162b2 vaccine recipients. *Lancet* 2021;6736:18–9.
- [4] SF Lumley et al. Antibody status and incidence of SARS-CoV-2 infection in health care workers. *N Engl J Med*. 2020;1–8. doi:10.1056/nejmoa2034545.
- [5] Guo L et al. Profiling early humoral response to diagnose novel coronavirus disease (COVID-19). *Clin. Infect. Dis.* 2020;71:778–85.
- [6] Okba NMA et al. Severe acute respiratory syndrome coronavirus 2-specific antibody responses in coronavirus disease patients. *Emerg Infect Dis* 2020;26:1478–88.
- [7] Long QX et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nat Med* 2020;26:845–8.
- [8] Danese E et al. Comprehensive assessment of humoral response after Pfizer BNT162b2 mRNA Covid-19 vaccination: a three-case series. *Clin Chem Lab Med* 2021. doi:10.1515/CCLM-2021-0339.
- [9] Favresse J et al. Antibody titres decline 3-month post-vaccination with BNT162b2. <https://doi.org/10.1080/22221751.2021.1953403> 10, 1495–1498 (2021).
- [10] Jalkanen P et al. COVID-19 mRNA vaccine induced antibody responses against three SARS-CoV-2 variants. *Nat Commun* 2021;121(12):1–11.
- [11] Yang HS et al. Association of age With SARS-CoV-2 antibody response. *JAMA Netw Open* 4, e214302 (2021).
- [12] Müller L et al. Age-dependent immune response to the Biontech/Pfizer BNT162b2 COVID-19 vaccination. medRxiv 2021.03.03.21251066 (2021). doi: 10.1101/2021.03.03.21251066.
- [13] Ebinger JE et al. Antibody responses to the BNT162b2 mRNA vaccine in individuals previously infected with SARS-CoV-2. *Nat Med* 1–4 (2021). doi: 10.1038/s41591-021-01325-6.
- [14] Lombardi A et al. SARS-CoV-2 anti-spike antibody titres after vaccination with BNT162b2 in naive and previously infected individuals. *J Infect Public Health* 2021;14:1120–2.
- [15] Fraley E et al. Humoral immune responses during SARS-CoV-2 mRNA vaccine administration in seropositive and seronegative individuals. *BMC Med* 2021;19(19):1–12.