

RESEARCH

Open Access



Molecular docking and identification of G-protein-coupled receptor 120 (GPR120) agonists as SARS COVID-19 MPro inhibitors

Sellappan Mohan^{1*}, Jayagopal Dharani², Ramanathan Natarajan³ and Arumugam Nagarajan¹

Abstract

COVID-19 has become a pandemic, and any new drug for treating the disease could save millions of lives. Several drugs already in use for other diseases and medical conditions are repurposed for treating COVID-19 in an attempt to find treatment for the disease without spending research time on ADME TOX and other studies on side effects. In this exercise, the drugs repurposed are from antiviral, antibiotics, antiviral for HIV and HCV, anti-cancer, natural medicines, etc. Possible repurposing anti-diabetic GPR-120 agonists used as for SAR-CoV-2 is attempted in the study by carrying out docking of 68 GPR-120 agonists. Ten of these compounds were found to have docking scores -8.3 to -8.0 , and the best docking score was observed for an arylsulfonamide and a biarylpropanoic acid belonging to GPR120 agonists previously evaluated for the treatment of type II diabetes. These GPR120 agonists could serve as start point for novel inhibitors for the discovery of drugs to treat COVID-19.

Background

After suffering from a devastating spell of COVID-19, the world is slowly limping back to normalcy. This is one of the pandemics that has a better public awareness owing to the Internet and social media. As per the data reported to WHO Globally, as of 14 October 2021, there have been 239,007,759 confirmed cases of COVID-19, including 4,871,841 deaths. As of 13 October 2021, a total of 6,471,051,151 vaccine doses have been administered (<https://covid19.who.int/> accessed on October 15, 2021). The world is undergoing the largest vaccination program to guard the people from any further spells of the deadly virus.

In addition to the vaccines, the pharmaceutical companies and the scientists in various organization are trying to develop drugs to combat the SARS CoV-2. There are several targets that could be explored to develop new drugs for COVID-19 [1]. The therapeutic targets include

both structural and non-structural proteins [2]. Some of the targets considered to develop inhibitors are as follows:

- Spike protein (S-protein) [3–5]
- Angiotensin-converting enzyme-2 (ACE-2) [6, 7]
- Human proteases: Transmembrane protease, serine 2 (TMPRSS2) [7, 8], Furin [9], Papain like protease-2 (PLpro) [10–12] 3-chymotrypsin like protease (3-CLpro) or the main protease M^{Pro} [13]
- Viral proteases (RNA-dependent RNA-polymerase (RdRp) [14]

One of the steps taken by the scientific community to combat the pandemic was to repurpose drugs already known and in use. This provides a shortcut and reduces the considerable amount of time spent on ADME Tox studies and the burden on assessing the new drug molecule's therapeutic efficacy, side effects, and risks. Several small molecules were considered [15]. The repurposed drugs are usually broad-spectrum antivirals that fall under the two therapeutic classes namely,

*Correspondence: selvaamohan@gmail.com

¹ Karpagam College of Pharmacy, Coimbatore, Tamil Nadu 641032, India
Full list of author information is available at the end of the article

protease inhibitor and nucleosides. Among the repurposed drugs favipiravir, remdesivir, molnupiravir, galidesivir, sobosbivir, and azivudine are examples of nucleosides while boceprevir, nardaprevir, simeprevir, and calpain inhibitors belong to protease inhibitors.

The race in finding an antiviral for COVID-19 was given momentum by computer-aided drug design approach, especially with the aid of docking software such as Auto Dock and Schrödinger. Of the several targets mentioned above, the main protease (M^{Pro}) has been the most explored for the development of inhibitors. One of the main reasons for exploring the M^{Pro} inhibitors is its important role played in the replication and transcription of SARS CoV-2 [16]. The main protease (M^{Pro}) is one of the proteins encoded in SARS-CoV-2 genome and is a dimer of cystine protease. This is called the 3-chymotrypsin-like protease (3-CL Pro). M^{Pro} presents a highly conserved active site in several coronaviruses, such as SARS-CoV and MERS-CoV. M^{Pro} plays an important role in the cleavage of precursor polyproteins translated from viral RNA, and no other human protease does have a similar cleavage specificity. This makes M^{Pro} an attractive target for developing inhibitors, and the inhibitor may thus be non-toxic.

The development of inhibitors targeting the main protease appears to have not left any stone unturned and these repurposed molecules may be grouped into (1) inhibitors of other CoV, (2) antiviral therapeutics of human immunodeficiency virus (HIV), (3) anti-viral that are being used in hepatitis C virus (HCV), (4) antimalarial and other antivirals for influenza, (5) anti-bacterial, (6) anti-cancer drugs, (7) Traditional Chinese medicines, and (8) chemicals in traditional spices and other natural compounds from marine origin.

Based on the action of the main protease, Yang et al. has designed several inhibitors in 2005. The authors found molecule N3 (number assigned by Yang et al. [17]) (see Fig. 1) as the most potent inhibitor of CoV. This molecule was studied by Jin et al. [13] for SARS CoV-2 and from the results of docking study the authors proposed that N3 binds in 3CL Pro binding pockets in an irreversible manner, and they thus exhibited good inhibitory potency. The 3CL Pro complex with N3 molecule was used to identify new inhibitors and one such molecule is ebselen [13, 18]. Ebselen, a drug used for the treatment of stroke containing a selenium atom is repurposed for SARS-CoV-2. Crystal structure of M^{Pro} without any ligand bound to the protein was reported by Zhang et al. [19] and Zhang et al. [20] studied the binding affinity of alpha-ketamide to 3CL Pro and identified three binding pockets in the protein. By varying the four substructures (marked as A, B,




C, & D in Fig. 1), they obtained the best fitting into the protein pockets from the inhibitory potency (Fig 1).

Calligari et al. [21] investigated thirteen proteinase inhibitors that are used as antiviral for human immunodeficiency virus (HIV) and hepatitis C virus (HCV). The ten anti-HIV drugs are saquinavir, indinavir, tipranavir, ritonavir, lopinavir, atazanavir, nelfinavir, amprenavir, darunavir, and fosamprenavir while the three anti-HCV are simeprevir, faldaprevir, and asunaprevir. Among these simeprevir was found to have the highest docking score. Lopinavir/ritonavir, coformulation is sold under the brand name Kaletra as an antiretroviral medication for the treatment and prevention of HIV/AIDS. Repurposing of Kaletra for SARS CoV-2 was found to be effective. Nutho et al. [22] could explain the inhibitory efficacy of Kaletra based on the docking studies of lopinavir and ritonavir with 3CL Pro . Chang et al. [23] showed that indinavir binds with 3CL Pro stronger than lopinavir and ritonavir and Calligari et al. [21] had also inferred this in their study. Nelfinavir was identified to be a potential inhibitor for CoV-2 from a docking that used 1903 candidates [24]. These authors went on to determine the inhibitory potency of nelfinavir [25]. Atazanavir the HIV antiviral was found to be a potential inhibitor of 3CL Pro [26], and its ability to inhibit SAR CoV-2 Vero cells was studied by Fintelman_Rodrigues et al. [27].

In addition to the three HCV drugs mentioned above, ledipasvir and velpatasvir were reported by Chen in 2020 [28]. Li et al. [29] ended up with four molecules namely, prulifloxacin, bictegravir, nelfinavir, and tegobuvir by high throughput screening of 8000 clinical drug libraries based on the binding affinity with M^{Pro} . Khan et al. [30] screened 123 antiviral drugs to identify inhibitors of 3CL Pro as well as 2'-O-MTase (2'-O-ribose methyltransferase). Paritaprevir and Raltegravir were found to have high binding affinity for 3CL Pro .

Talluri [31] carried out virtual screening of several clinically approved antiviral and the crystal structure of M^{Pro} (PDB if 6LU7) and found saquinavir and beclabuvir as the best protease inhibitor candidate SARS CoV-2 among the compounds studied by them. Other anti-viral drugs that had been tested for repurposing by molecular docking and virtual screening include oseltamivir [32] and zanamivir [33].

Some of the antibiotics that have been identified to be effective based on computer-aided virtual screening are the quinoline antibiotic prulifloxacin [29], tetracycline antibiotics eravacycline [34], and the polypeptide antibiotic colistin [35]. Non-steroidal anti-inflammatory drugs (NSAID) were also repurposed as potential inhibitors of M^{Pro} [36] by docking studies. In a similar study on NSAIDs, Gimeno et al. [37] identified Perampanel, Carprofen, Celecoxib, Alprazolam, Trovafloxacin,

N3 (C₃₅H₄₈N₅O₈) α -ketamide (the four substructures subjected to change) α -ketamide with best inhibitory potency (see substructures A, B, C and D for the best fitting molecule in the M^{Pro} pockets)**Fig. 1** Structure of N3, α -ketamide with sub-structures marked

Sarafloxacin, and ethyl biscoumacetate as possible inhibitors of M^{Pro} by docking studies. The two compounds namely, Carprofen, a NSAID no longer use in human medicine but used for veterinary purpose, and Celecoxib another NSAID and a COX-2 inhibitor, were subjected to in vitro testing at 50 μ M, and they showed 3.97% and 11.90% M^{Pro} inhibition, respectively.

Dipyridamole (brand name Persantine) is a platelet inhibitor and is used to prevent blood clots after heart surgery was repurposed for CoV-2 by Liu et al. [38] and the inhibitory potency (IC₅₀) was studied targeting

3CL^{Pro}. Odhar et al. [39] studied the drug molecules from US-FDA-approved drugs library from ZINC 5 database, and from their docking on to the M^{Pro} (PDB id 6LU7), they identified ten hits that included drug that are used for cancer, epilepsy, and insomnia. The top ten hits based on the docking score are Perampanel (epilepsy), conivaptan (hyponatremia), sonidegib (basal-cell carcinoma), azelastine (allergy), idelalisib (leukemia and lymphoma), suvorexant (insomnia) olaparib (ovarian, breast, and pancreatic cancers), ponatinib (leukemia), loxapine (schizophrenia), and tolvaptan (hyponatremia). Wang et al. [34]

in the computational drug repurposing study identified carfilzomib (antineoplastic agent), valrubicin (chemotherapy drug), and elbasvir (antiviral for HCV) as inhibitors based on the docking with M^{Pro}, in addition to the antibiotic eravacycline.

Traditional Chinese medicine (TCM) and traditional Indian medicines that fall under Ayurveda and Sidha were used as immune boosters to fight against COVID-19. Zhang et al. [19, 20] carried out docking studies of about 100 constituents of the Lung-toxin Dispelling Formula No. 1 (LDFN1) of TCM and found 22 of these chemicals are inhibitors of 3CL^{Pro}. Of the several chemicals, baicalin and baicalein were found to have antiviral activities against 3CL^{Pro} [40] with EC₅₀ values of 10.27 μM and 1.69 μM, respectively. Liu et al. [41] studied the inhibitory activity of the ethanol extract of the herbal plant *Scutellaria baicalensis* and its major component, baicalein. They found that the plant extract and the constituent baicalein inhibited SARS-CoV-2 3CL^{Pro} activity in vitro with IC₅₀ of 8.52 mg/mL and 0.39 mM, respectively. The replication of SARS-CoV-2 in Vero cells were inhibited with EC₅₀s of 0.74 mg/ml and 2.9 mM, respectively. In their study on screening several natural compounds that are constituents of TCM, Zhang et al. [19] and Zhang et al. [20] identified betulinic acid, coumaroyltyramine, cryptotanshinone, desmethoxyreserpine, dihomog-γ-linolenic acid, kaempferol, lignan, *N*-cis-feruloyltyramine, quercetin, sugiol, and tanshinoneia to inhibit 3CL^{Pro}. Cherrak et al. [42] studied several glycosylated flavonoids by docking them on the M^{Pro} (6LU7) and identified quercetin-3-O-rhamnoside to have the highest binding affinity. Myricetin 3-rutinoside and rutin were also identified as potential inhibitors of 3CL^{Pro}, and the binding affinities of these three compounds were greater than that of N3 with 3CL^{Pro}. Shivanika et al. [43] carried out docking studies of several natural products that have been used as antiviral on to 6LU7 the 3CL^{Pro} protein structure and found theaflavin-3-3'-digallate, rutin, hypericin, robustaflavone, and (-)-solenolide as the compounds with highest binding energy. It might be noted that identification of rutin as a potential inhibitor is independently confirmed by two groups. Bhaliya and Shah [44] carried out docking studies of mono-carbonyl analogs of curcumin with 3CL^{Pro} and found one of the curcumin analogs was found to have potential to be used as an inhibitor. Joshi et al. [45] screened a library of ~7100 molecules that comprises of flavonoids, glucosinolates, anti-tussive, anti-influenza, anti-viral, terpenes, terpenoids, alkaloids, and other compounds predicted as potential therapeutic candidates against M^{Pro}. Molecules such as δ-viniferin, myricitrin, taiwanhomo-flavone A, lactucopicrin 15-oxalate, nympholide A,

afzelin, bio Robin, hesperidin, and phyllaemblicin B were found to bind strongly with M^{Pro} and hence suggested as potential inhibitors. Andrographolide a natural compound from *Andrographis paniculata* was studied [46] via docking on to M^{Pro}, and the in silico studies on ADME and toxicity prediction were also carried out. The molecule was predicted to have good solubility. Ramaiah et al. [47] studied the binding of natural molecules that are present in Indian spices and curry against M^{Pro} (6LU7). A similar study identified [48] carnosol a natural molecule as an inhibitor by docking studies using the protein structure PDBID: 6Y84, M^{Pro}. Bioactive compounds in medicinal plants were screened as potential M^{Pro} inhibitors [49] and natural compounds such as kaempferol, quercetin, luteolin-7-glucoside, demethoxycurcumin, naringenin, apigenin-7-glucoside, oleuropein, curcumin, catechin, and epicatechin-gallate as potential molecules for further exploration.

According to the latest report of the pharmaceutical company Merck, molnupiravir pills are able to reduce the hospitalization and deaths of people affected by COVID-19 [50]. They reported the results of Phase 2a trial ([ClinicalTrials.govNCT04405570](https://clinicaltrials.gov/NCT04405570)) in which safety, tolerability, and antiviral efficacy of molnupiravir in the treatment of COVID-19. Merck applied on October 11, 2021, for US-FDA emergency use authorization for the molnupiravir based-oral antiviral pill for COVID-19. This will not stop the hunt for new inhibitors, and the search for new molecules will continue.

In one of the studies of repurposing drugs [51], virtually screened 1615 FDA approved drugs by docking each of them on to M^{Pro} and then refined the selection by employing molecular dynamics to identify nine compounds. The nine drugs selected as potential inhibitors vary from vasoconstrictor to microscopy dye. The potential inhibitors identified and their original use are:

1. Dihydroergotamine: vasoconstrictor
2. Midostaurin: treatment of acute myeloid leukemia
3. Ziprasidone: antipsychotic
4. Etoposide: antineoplastic
5. Apixaban: used to reduce the risk of stroke and blood clots
6. Fluorescein: a dye used in microscopy
7. Tadalafil: used to treat erectile dysfunction (ED), benign prostatic hyperplasia (BPH), and pulmonary arterial hypertension)
8. Rolapitant: used along with an antiemetic (anti-vomiting) agent in adults for the prevention of delayed nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy
9. Palbociclib: used to treat HR-positive and HER2-negative breast cancer

The above discussion indicates that repurposing of drugs belonging to different classes have been evaluated for CoV-2. Human GPR120 is a transmembrane protein, characterized by the interactions with the endogenous ligand linoleic acid and docosahexaenoic acid. Apart from the key role played by GPR 120 in diabetes, it is also involved in many other disease conditions,

including cancer, inflammation, and central nervous system (CNS) disorders. GPR 120 presents itself in many metabolic pathways, and its pivotal role in controlling obesity and diabetes is worth mentioning. Using gene knockdown studies, GPR120 has been shown to induce chemoresistance in breast cancer treatment with epirubicin and cisplatin-highlighting the relevance of GPR 120




Fig. 2 **a** Flowchart for docking procedure to obtain binding score for docking a ligand. **b** Flowchart for to view and save the protein-ligand complex. **c** Structural similarity and drug like properties of docked GPR 120 agonists versus Rolapitant and Fluvastatin. **d** Toxicity prediction for compound 34 (ProTox-II - Prediction of TOXicity of chemicals (charite.de))

antagonists for chemotherapy [52]. Further, Toelzer et al. recently identified a linoleic acid binding pocket in the SARS-CoV-2 spike protein which prompted us to look for alternate drugs for binding with *COVID-19 MPro*. The present study reports the results of docking studies carried out using G-protein-coupled receptor (GPR) agonists against the MPro to identify any potential inhibitor of SARS CoV-2.

Methods

Auto Dock 4.2.6 was used to perform docking study. Chemical structures were drawn using Chemoffice 2002. Three-dimensional structures of proteins were downloaded from protein data bank (PDB id: 6LU7) (<https://www.rcsb.org/>).

Protein preparation

The protein was prepared for docking process according to the standard protein preparation procedure integrated in Accelry's Discovery Studio 4 which is shown in the flow chart (Fig. 2a).

Ligand 2D structures were drawn using ChemDraw Ultra 8.0 (ChemOffice 2002) and converted into 3D structure using chem3D Ultra 8.0. The 68 molecules were used as ligands, and each one of them was docked on to the crystal structure of M^{Pro} with PDB id 6LU7. The procedure for ligand preparation and docking is given as a flow chart in Fig. 2a, b. Docking scores were obtained to understand any inhibitory potential of the 68 GPR120 agonists.

Zhang and Macielag [53] discussed the patented GPR 120 agonists for the treatment of diabetes. They reviewed the therapeutic patents of ten different classes of compounds that amounted to 68 therapeutic molecules. The 68 GPR-120 agonists collected by the authors from different patents and journals are grouped into ten classes. The ten classes and the number assigned in this paper along with abbreviation are given below:

1. Natural GPR 120 agonists and early synthetic GPR 120 agonists (1–9)
2. Bi-aryl-based phenyl propionic acid derivatives as GPR 120 agonists (BiAr-PPA 10–14)
3. Cycloalkenyl and heterocycloalkenyl-based phenyl propionic acid derivatives (CycA_Hcyc--PPA 15–23)
4. Dihydrobenzofuran derivatives (Metabolex) and benzo-fused heterocyclic derivatives (Metabolex 24 and Jansesen 25–27)
5. Chemical scaffolds claimed by Merck (Merck 28–35)
6. Various carboxylic acid scaffolds claimed by Bristol-Meyers Squibb (BMS 36–45)

7. Patented structures by Piramal Enterprises Limited (PEL 46–52)
8. Other carboxylic acid-based GPR 120 agonists (Calden (53–54; LG 55–59; Ajinomoto 60; DOMPE 61–62)
9. Non-acid-containing structures claimed by AXXAM (AXXAM 63–65)
10. GPR120 agonists/antagonists in the peer-reviewed journals (GSK 65; U of B 66; GSK 67; Ch Pharm U 68).

Different classes of these 68 molecules are patented as GPR120 agonists to treat type-2 diabetes by various pharmaceutical companies namely, Janssen Pharmaceutica NV, Merck Sharp & Dohme Corp., Bristol Myers Squibb, Piramal Enterprises Limited, Caldan Therapeutics Limited, LG Life Sciences Ltd., Ajinomoto Company, Accepted Manuscript Inc., Dompe' Farmaceutici S.P.A., and AXXAM S. P. A. Structures of these compounds with their abbreviated id number used in this paper are given in Table 1.

Results and discussions

Binding scores for each of the 68 ligands are listed in Table 1 along with their molecular structures. Docking images each of the ligands in MPro (PDB ID 6LU7) are given in the [Supplementary material](#) while the ligands with best scores (≥ -8.0) are given in Fig. 3. The type of docking interaction for these ligands are presented in Table 2. Some of the ligands are having a docking score of ≥ -8.0 . The lowest value is -8.3 for the molecule with id GSK (65). This is a sulfonamide patented by GlaxoSmithKline as a selective antagonist against free fatty acid 4 (FFA4/GPR120) [54, 55] and to be used with the agonist GSK 137647A which is also a sulfonamide (id in this paper GSK137647A(8)). The compound with binding score -8.2 is a biaryl-based phenylpropanoic acid (13) [56] patented by Janssen Pharmaceutica [57]. The compound 17 with binding score -8.0 is also phenylpropanoic acid derivative namely, cyclohexenyl-based phenyl propionic acid [58]. Three of the other compounds 47, 48, and 52, with binding score -8.0 are patented by Piramal Enterprises Limited as GPR120 agonists [47, 59–61] and phenylbutanoic acid with biarylsubstituent wherein one of the aryl groups is a heterocyclic or fused heterocyclic system. Cyclopropane carboxylic acid derivative with a phenoxybiphenyl substituent **40** is also found to have a binding score 8.0. This molecule is patented by Bristol-Meyers Squibb Company [62] as GPR120 modulators useful for treatment of diabetes and related diseases.

Based on the binding score of the top 10 compounds investigated here, their role in blocking the binding site through Glu 166 and Cys 145 could be considered

relevant for their potential role as novel ligands for Sars-COVID-19 virus protein. The observation that Remdesivir, Nelfinavir, and other antiviral compounds show similar interaction support our inference [63–65]. Additional support for such a claim has been found in the paper describing docking study of metocurine with

M-Pro 6LUZ that indicate the drug occupies the binding site [66]. The important residues observed in the docking study of our GPR120 agonists as well as the above molecules studied by others including that of chlorquine [61] highlight the role of NH, COOH groups in manifesting pi bond formation with Glu 166 and aromatic pi

Table 1 GPR120 agonists, ligand ids (used in this paper), docking scores, and molecular structures








Ligand id	Docking Score	Structure
DHA (1) (3E,6E,9E,12E,15E,18E)-heneicosanoic acid	−5.6	
ALA (2) Alpha linoleic acid	−4.6	
EPA (3) Eicosapentaenoic acid	−5.7	
Grifolic acid (4)	−6.4	
Grifolic acid methyl ether (5)	−5.6	
GW9508 (6) 4-(3-Phenoxybenzylamino)phenylpropionic acid	−7.3	
NCG21 (7) 4-(4-(2-(phenyl(pyridin-2-yl)amino)ethoxy)phenyl)butanoic acid	−6.9	

Table 1 (continued)







GSK137647A (8) N-mesityl-4-methoxybenzenesulfonamide	-6.4	
TUG891 (9) 3-(4-((4-fluoro-4'-methyl-[1,1'-biphenyl]-2-yl)methoxy)phenyl)propanoic acid	-6.8	
BiAr-PPA (10) 3-(4-((3-(4-chlorophenyl)-5-(trifluoromethyl)isoxazol-4-yl)methoxy)phenyl)propanoic acid	-7.9	
BiAr-PPA (11) 3-(4-((3-(4-ethylphenyl)-5-(trifluoromethyl)isoxazol-4-yl)methoxy)phenyl)propanoic acid	-7.4	
BiAr-PPA (12) 3-(4-((3-(5-chloro-2-fluorophenyl)-5-(trifluoromethyl)isothiazol-4-yl)methoxy)-3,5-difluorophenyl)propanoic acid	-7.5	
BiAr-PPA (13) 3-(4-((4-(4-chlorophenyl)-2-(trifluoromethyl)thiophen-3-yl)methoxy)-3,5-difluorophenyl)propanoic acid	-8.2	

Table 1 (continued)







<p>BiAr-PPA (14)</p> <p>3-(4-((1-(4-chlorophenyl)-3-(trifluoromethyl)-1H-imidazol-2-yl)methoxy)-2,3-difluorophenyl)propanoic acid</p>	-7.1	
<p>CycA_Hcyc-PPA (15)</p> <p>3-(2,3-dimethyl-4-((2-(p-tolyl)cyclopent-1-en-1-yl)methoxy)phenyl)propanoic acid</p>	-5.6	
<p>CycA_Hcyc--PPA (16)</p> <p>3-(4-((2-(4-chlorophenyl)-4,4-difluorocyclopent-1-en-1-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-7.9	
<p>CycA_Hcyc--PPA (17)</p> <p>3-(4-((4'-chloro-4,4-difluoro-3,4,5,6-tetrahydro-[1,1'-biphenyl]-2-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-8.0	
<p>CycA_Hcyc--PPA (18)</p> <p>3-(4-((2-(4-chlorophenyl)cyclohept-1-en-1-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-7.5	
<p>CycA_Hcyc--PPA (19)</p> <p>3-(4-([1,1'-bi(cyclopentane)]-1,2'-dien-2-ylmethoxy)-2-(trifluoromethyl)phenyl)propanoic acid</p>	-7.3	

Table 1 (continued)







CycA_Hcyc--PPA (20) 3-(4-((2-isopropylcyclohex-1-en-1-yl)methoxy)-2-(trifluoromethyl)phenyl)propanoic acid	-7.2	
CycA_Hcyc--PPA (21) 3-(4-((4-(4-chlorophenyl)-5,6-dihydro-2H-pyran-3-yl)methoxy)-2,3-dimethylphenyl)propanoic acid	-7.8	
CycA_Hcyc--PPA (22) 3-(4-((4-(1-phenylvinyl)-5,6-dihydro-2H-pyran-3-yl)methoxy)-2-(trifluoromethyl)phenyl)propanoic acid	-6.8	
CycA_Hcyc--PPA (23) 3-(4-((4-(cyclopent-2-en-1-yl)-5,6-dihydro-2H-pyran-3-yl)methoxy)-2-(trifluoromethyl)phenyl)propanoic acid	-7.7	
Metabolex (24) 3-(4-((5-fluoro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)phenyl)propanoic acid	-7.6	
Janssen (25) 3-(4-((5-chloro-2,6-dimethylbenzofuran-7-yl)methoxy)-2,3-dimethylphenyl)propanoic acid	-7.3	

Table 1 (continued)







<p>Janssen (26)</p> <p>3-(4-((5-chloro-2-ethylbenzofuran-7-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-7.4	
<p>Janssen (27)</p> <p>3-(4-((6-chloro-2-methylbenzo[d]thiazol-4-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-7.0	
<p>Merck (28)</p> <p>2-(3-(2-chloro-5-(trifluoromethoxy)phenyl)-3-azaspiro[5.5]undecan-9-yl)acetic acid</p>	-7.7	
<p>Merck (29)</p> <p>3-(8-(5-cyclobutoxy-2-fluoro-4-methoxyphenyl)-8-azaspiro[4.5]decan-2-yl)propanoic acid</p>	-7.1	
<p>Merck (30)</p> <p>3-(2-(3-chloro-2-fluoro-5-((5-methylthiazol-2-yl)oxy)phenyl)-5-fluoro-1,2,3,4-tetrahydroisoquinolin-6-yl)propanoic acid</p>	-7.3	
<p>Merck (31)</p> <p>3-(2-(6-cyclobutoxy-3-fluoro-4-methoxypyridin-2-yl)-5-fluoro-1,2,3,4-tetrahydroisoquinolin-6-yl)propanoic acid</p>	-7.5	

Table 1 (continued)







Merck (32) 3-(5-(2-fluoro-5-(trifluoromethoxy)phenyl)benzofuran-2-yl)propanoic acid	-7.3	
Merck (33) 3-((R)-6-(2,3-difluoro-5-((1r,3R)-3-methoxycyclobutoxy)phenyl)chroman-2-yl)propanoic acid	-7.3	
Merck (34) 5-(5-((1s,3s)-3-(4-(difluoromethyl)-3-fluorophenyl)cyclobutoxy)pyrazin-2-yl)isoxazol-3-ol	-7.9	
Merck (35) 5-(6-((1s,3s)-3-(phenoxy)methyl)cyclobutoxy)pyridin-3-yl)isoxazol-3-ol	-7.6	
BMS (36) 2-(4-((1R,3S)-3-(2-fluoro-3-methylphenoxy)cyclohexyl)phenethoxy)acetic acid	-7.3	
BMS (37) 2-(4-(3-phenoxy)cycloheptyl)phenethoxy)acetic acid	-7.1	

Table 1 (continued)






<p>BMS (38)</p> <p>(S)-5-(4-(3-phenoxy-pyrrolidin-1-yl)phenyl)pentanoic acid</p>	-6.7	
<p>BMS (39)</p> <p>(1S,2R)-2-(4-((R)-3-phenoxy-piperidin-1-yl)phenethyl)cyclopropane-1-carboxylic acid</p>	-7.3	
<p>BMS (40)</p> <p>(1R,2R)-2-(((2',3,3'-trifluoro-5'-(3-fluorophenoxy)-[1,1'-biphenyl]-4-yl)oxy)methyl)cyclopropane-1-carboxylic acid</p>	-8.0	
<p>BMS (41)</p> <p>2-(((1s,3s)-3-(2'-fluoro-5'-phenoxy-[1,1'-biphenyl]-4-yl)cyclobutoxy)acetic acid</p>	-7.5	
<p>BMS (42)</p> <p>2-(((1r,4r)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid</p>	-7.3	

Table 1 (continued)






<p>BMS (43)</p> <p>(1S,2R)-2-(((1s,4R)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropane-1-carboxylic acid</p>	-7.8	
<p>BMS (44)</p> <p>3-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.2]octan-1-yl)propanoic acid</p>	-7.9	
<p>BMS (45)</p> <p>3-(4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.2]octan-1-yl)propanoic acid</p>	-7.3	
<p>PEL (46)</p> <p>4-(4-(5-fluoro-2-(4,5,6,7-tetrahydrobenzo[d]thiazol-2-yl)phenethyl)phenyl)butanoic acid</p>	-7.6	
<p>PEL (47)</p> <p>4-(4-(1,1-difluoro-2-(5-fluoro-2-(5-methoxypyridin-2-yl)phenyl)ethyl)phenyl)butanoic acid</p>	-8.0	

Table 1 (continued)







<p>PEL (48)</p> <p>4-(4-(5-fluoro-2-(5-(1-methylcyclopropyl)thiophen-2-yl)styryl)phenyl)butanoic acid</p>	-8.0	
<p>PEL (49)</p> <p>4-(4-((2-(5-cyclopropylthiophen-2-yl)-5-fluorophenyl)ethynyl)phenyl)butanoic acid</p>	-7.4	
<p>PEL (50)</p> <p>4-(4-((2-(5,6,7,8-tetrahydronaphthalen-1-yl)cyclohex-1-en-1-yl)methoxy)phenyl)butanoic acid</p>	-7.5	
<p>PEL (51)</p> <p>4-(4-((2-(o-tolyl)cyclopent-1-en-1-yl)methoxy)phenyl)butanoic acid</p>	-6.0	
<p>PEL (52)</p> <p>4-(2-(2-(5-cyclopropylthiophen-2-yl)-4-fluorophenyl)benzofuran-5-yl)butanoic acid</p>	-8.0	
<p>Calden (53)</p> <p>3-(2-(5-cyclobutoxy-2,3-difluorophenyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)propanoic acid</p>	-7.6	

Table 1 (continued)








<p>Calden(54)</p> <p>3-(3-(2-fluoro-5-(trifluoromethoxy)phenyl)-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)propanoic acid</p>	-7.0	
<p>LG (55)</p> <p>4-((2'-(cyclobutylmethoxy)-3,5-difluoro-[1,1'-biphenyl]-4-yl)oxy)butanoic acid</p>	-7.1	
<p>LG (56)</p> <p>4-((2'-cyclobutoxy-[1,1'-biphenyl]-4-yl)thio)butanoic acid</p>	-6.6	
<p>LG (57)</p> <p>3-(6-(2-(isopropylthio)pyridin-3-yl)quinolin-2-yl)propanoic acid</p>	-6.8	
<p>LG (58)</p> <p>3-(6-(2-(cyclopentylloxy)pyridin-3-yl)chroman-2-yl)propanoic acid</p>	-7.5	
<p>LG (59)</p> <p>3-(4-((2-(isopropylthio)pyridin-3-yl)methoxy)-2,3-dimethylphenyl)propanoic acid</p>	-6.8	
<p>Ajinamoto (60)</p> <p>(Z)-3-(3-chloro-4-((3,7-dimethylocta-2,6-dien-1-yl)oxy)phenyl)propanoic acid</p>	-6.8	

Table 1 (continued)









<p>DOMPE (61)</p> <p>7-(3-(N-(2,6-dimethyl-4-(trifluoromethyl)phenyl)sulfamoyl)phenyl)heptanoic acid</p>	-6.8	
<p>DOMPE (62)</p> <p>7-(3-(N-(2,6-dimethyl-4-(2-(methylsulfonyl)ethoxy)phenyl)sulfamoyl)phenyl)heptanoic acid</p>	-6.4	
<p>AXXAM (63)</p> <p>1-cyclopentyl-5-((2,6-difluoro-4-(3-hydroxypropyl)phenoxy)methyl)-3-methyl-1H-pyrazole-4-carbonitrile</p>	-7.2	
<p>AXXAM (64)</p> <p>5-((2,6-difluoro-4-(3-hydroxypropyl)phenoxy)methyl)-3-methyl-1-(5-methylpyridin-2-yl)-1H-pyrazole-4-carbonitrile</p>	-7.4	
<p>GSK (65)</p> <p>4-methyl-N-(9H-xanthen-9-yl)benzenesulfonamide</p>	-8.3	

Table 1 (continued)

Uof B (66) 2-(3-fluoro-5-(pyridin-2-yloxy)phenyl)- 2,3-dihydrobenzo[d]isothiazole 1,1- dioxide	-7.7	
GSK (67) 3-(4-((2-methoxy-5- (trifluoromethyl)benzyl)oxy)-3- methylphenyl)propanoic acid	-7.1	
ChPharmU (68) 4-((2',3,3',5-tetrafluoro-5'-((1r,3r)-3- methoxycyclobutoxy)-[1,1'-biphenyl]-4- yl)oxy)butanoic acid	-7.3	

interaction with Cys 145, respectively. Dock score for the reference compounds (Fig. 3i–l) evaluated along with the GPR120 agonists ALPHA KETOMIDE (-7.4), LOPINAVIR (-7.9), SHIKONIN (-7.0), and TIDEGLUSIB (-8.0) indicate the interactions with Glu 166 and Cys 145 are present in these drugs also.

Compounds 13, 16, 17, 44, and 50 identified with high dock score are hydrophobic compounds having thiophenyl, cyclopentenyl, cyclohexenyl, norbornyl, and cyclopentenyl groups along with a phenylpropanoic acid function. Compound 40 is a phenoxyphenyl ether having a cyclopropane carboxylic acid group while 34 has an isoxazolyl and pyrimidine compound with difluoromethane function. Compound 65 is a tricyclic compound having arylsulfonamide function. Compounds 13 and 52 are highly lipophilic having a log *P* value of 7.1 and 6.25, respectively, that might require vigorous optimization to make them orally available. On the other hand, linoleic acid has a log *P* value of 5.65 while that of LOPINAVIR is 4.56 and the value is 4.86 for TIDEGLUSIB. Hosseini et al. screened several classes of drugs and identified inhibitors

for SARS-CoV-2 MPro and highlighted H bond interactions with Thr 26, Phe 140, Gly 143, Glu 166, and Gln 189 in addition to pi stacking interaction through His 140 as key contributors for receptor binding [67]. Our GPR 120 agonists, 40, 47, and 65 revealed H bonding interaction with Glu 189 in the docking against MPro while hydrophobic interactions with His 41, Met 165, and Glu 166 were shown by compounds 40, 48, 52, and 65. Fluvastatin on the other hand was found to interact with Thr 26 and Gly 143 by Maryam et al., and a similar interaction was observed in our compound 40, which also interacted with Glu 166 and Cys 145. Compounds 65 and 34 have a log *P* value of 4.45 and 3.26, respectively, indicating that they might have a good oral bioavailability, although they require further optimization. The comparison of topological polar surface area (TPSA) of the compounds evaluated show that only compound 34 has a TPSA value above 75, whereas the reference compounds have TPSA of 120 for lopinavir, a-ketoamide has a value of 113. This suggests that compounds 34 and 40 could be used as start points, and further optimizations could result in finding a drug

(See figure on next page.)

Fig. 3 **a** GSK (65); Docking score -8.3. **b** BiAr-PPA (13); Docking score -8.2. **c** CycA_Hcyc--PPA (17); Docking score -8.0. **d** Merck (34); Docking score -7.9. **e** BMS (40); Docking score -8.0. **f** PML (47); Docking score -8.0. **g** PML (48); Docking score -8.0. **h** PML (52); Docking score -8.0. Docking of the ligands onto MPro (6LU7). Docking images of ligands with docking score ≥ -7.9 are given (for other ligands please see [Supplementary Information](#)); **i** ALPHA KETOMIDE = -7.4; **j** LOPINAVIR = -7.9; **k** SHIKONIN = -7.0; **l** TIDEGLUSIB = -8.0




Fig. 3 (See legend on previous page.)

Table 2 Docking interactions for the ligands with docking score ≥ -8.0

Ligand id	Docking score	Docking details		
		Conventional H-bond	Alkyl and pi-alkyl	others
GSK (65)	-8.3	GLN:189, GLU:166	MET:49	CYS:145 (Pi-donor hydrogen bond)
BiAr-PPA (13)	-8.2	THR:24, SER:46, GLY:143	MET:49, CYS:145	MET:49 (Pi-sigma) THR:26, LEU:141, ASN:142 (Halogen)
CycA_Hcyc-PPA (17)	-8.0	TYR:54	CYS:145, MET:165	PHE:140, LEU:141, ASN:142, GLU:166 (Halogen)
Merck (34)	-7.9	Asp 187, Tyr 54, His 164, His 41	Cys 145, Leu 27	
BMS (40)	-8.0	THR:25,26, GLY:143, SER:144, CYS:145, HIS:163 GLN:189, GLU:166	CYS:145, MET:165	GLN:189, CYS:145 (carbon hydrogen bond); GLY:143 (Pi-donor hydrogen bond); LEU:141, ARG:188 (Halogen)
PEL (47)	-8.0	SER:46, CYS:145, HIS:164	MET:49	THR:24 (unfavorable acceptor-acceptor); MET:49 (Pi-sigma) LEU:141, PHE:140 (carbon hydrogen bond); GLN:189, ASP:187 (Halogen)
PEL (48)	-8.0	THR:45	HIS:41, MET:49, CYS:145, MET:165	HIS:163 (Pi-cation); SER:46 (unfavorable donor-donor); PHE:140, LEU:141 (Halogen)
PEL (52)	-8.0	HIS:41, CYS:145	MET:49, MET:165	MET:49 (Pi-sigma); CYS:145 (Pi-sulfur); ARG:188 (Halogen)

for Sars-COVID-19. Compound 34 possesses a hydroxyl isoxazole group that would mimic COOH function and also the presence of polar heterocyclic ring providing it an ideal choice to improve its physicochemical characteristics. The compound 34 is in fact well anchored through H-bonded interactions, Pi interactions as shown in Fig. 3d. Similar to compound 40, having a cyclopropane carboxylic acid function could be optimized further to refine its log P to make this eligible as a lead compound. Obviously the GPR 120 agonists, designed as agonists for free fatty acid receptors, have functional groups and lipophilic characters designed for their receptor need to be tweaked to suit the binding interactions with MPro. These compounds identified through the present study are functionally similar to linoleic acid, a free fatty acid that has been found to occupy the binding pocket of spike protein in SARS-CoV-2 [68]. Comparing the free fatty acid, linoleic acid, the GPR 120 agonists identified herein possess several beneficial physicochemical properties in terms of favourable log P values and topological polar surface area, making them suitable for oral administration (Fig. 2c).

Toxicity prediction for compound 34, using online tool "protox_II," indicates that the molecule is safe for all the targets except showing carcinogenicity and hepatotoxicity of 0.51 and 0.64, respectively, further requiring structural modification. This compound also has an LD 50 value of 300 mg/kg and falls under predicted toxicity class 3, indicating it is only slightly toxic and slightly irritating.

Conclusions

The present data supports the possibility of repurposing free fatty acid GPR 120 receptor agonists as potential inhibitors of Sars-COVID-19 M-Protein. Based

on docking score and key interactions with the amino acid residues in the target protein, compound 34 could be used as a lead compound. The presence of COOH mimicking hydroxyl isoxazole group could provide necessary drug-like property in addition to maintaining H-bond and pi interactions with the receptor. Favorable logP and available physicochemical and toxicity data of compound 34 could shorten the drug development time to position the compound as an early lead candidate to overcome the hurdles in identifying therapeutic drugs in coronavirus infection.

Abbreviations

COVID-19 MPro: Severe acute respiratory syndrome coronavirus 2 main protease; GPR-120 agonists: G-protein-coupled receptor or free fatty acid receptor-4 agonists; Docking: Phenyl propanoic acid.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s43141-022-00375-8>.

Additional file 1. Docking images of the inhibitors (ligands) on M^{Pro}.

Acknowledgements

The authors wish to thank Karpagam Management Trust for their support for the facilities.

Authors' contributions

Dr. Mohan, Sellappan conceived the idea and gave directions. Dr. Nagarajan Arumugam gave information about GPR 120 agonists and their bioactivities and relation to COVID-19 MPro. Dr. Natarajan Ramanathan collected GPR agonists and guided the workflow for docking study. Dr. Tharani Jayagopal did the docking study and collected the images of docking and scoring details. Dr. Nagarajan Arumugam and Dr. Natarajan R wrote the manuscript. Dr. Mohan S revised the manuscript. The author(s) read and approved the final manuscript.

Authors' information

Not applicable.

Funding

There is no funding received from any source.

Availability of data and materials

Yes. Attached as Supporting information

Declarations**Ethics approval and consent to participate**

Not applicable.

Consent for publication

All authors consent for publication in the *Journal of Genetic Engineering and Biotechnology*.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Karpagam College of Pharmacy, Coimbatore, Tamil Nadu 641032, India. ²Sri Sarada Niketan College for Women, Karur, Tamilnadu, India. ³Saranathan College of Engineering, Panjappur, Tiruchirappalli, India.

Received: 28 February 2022 Accepted: 2 June 2022

Published online: 18 July 2022

References

- Prajapat M, Sarma P, Shekhar N, Avti P, Sinha S, Kaur H, Kumar S, Bhat-tacharyya A, Kumar H, Bansal S, Medhi B (2020) Drug targets for corona virus: a systematic review. *Indian J Pharm* 52:56–65. https://doi.org/10.4103/ijp.IJP_115_20
- Yadav R, Chaudhary JK, Jain N, Chaudhary PK, Khanra S, Dhamija P, Sharma A, Kumar A, Handu S (2021) Role of structural and non-structural proteins and therapeutic targets of SARS-CoV-2 for COVID-19. *Cells* 10:821. <https://doi.org/10.3390/cells10040821>
- Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D (2020) Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell* 181:281–292. <https://doi.org/10.1016/j.cell.2020.02.058>
- Xia S, Liu M, Wang C, Xu W, Lan Q, Feng S, Qi F, Bao L, Du L, Liu S, Qin C, Sun F, Shi Z, Zhu Y, Jiang S, Lu L (2020a) Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a highly potent pan-coronavirus fusion inhibitor targeting its spike protein that harbors a high capacity to mediate membrane fusion. *Cell Res* 30:343–355. <https://doi.org/10.1038/s41422-020-0305-x>
- Xia S, Zhu Y, Liu M, Lan Q, Xu W, Wu Y, Lu L (2020b) Fusion mechanism of 2019-nCoV and fusion inhibitors targeting HR1 domain in spike protein. *Cell Mol Immunol* 17:765–767. <https://doi.org/10.1038/s41423-020-s0374-2>
- Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF (2020) The proximal origin of SARS-CoV-2. *Nat Med* 26:450–452. <https://doi.org/10.1038/s41591-020-0820-9>
- Hoffmann M, KleineWeber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A, Müller MA, Drosten C, Pöhlmann S (2020) SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cells* 181:210–271. <https://doi.org/10.1016/j.cell.2020.02.052>
- Bestle D, Heindl MR, Limburg H, Van TVL, Pilgram O, Moulton H, Böttcher-Friebertshäuser E (2020) TMPRSS2 and furin are both essential for proteolytic activation and spread of SARS-CoV-2 in human airway epithelial cells and provide promising drug targets. *BioRxiv*. <https://doi.org/10.1101/2020.04.15.042085>
- Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E (2020) The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. *Antiviral Res* 176:104742. <https://doi.org/10.1016/j.antiviral.2020.104742>
- Amin SA, Banerjee S, Ghosh K, Gayen S, Jha T (2021) Protease targeted COVID-19 drug discovery and its challenges: insight into viral main protease (Mpro) and papain-like protease (PLpro) inhibitors. *Bioorg Med Chem* 29:115860. <https://doi.org/10.1016/j.bmc.2020.115860>
- McClain CB, Vabret N (2020) SARS-CoV-2: the many pros of targeting PLpro. *Signal Transduct Target Ther* 5:223. <https://doi.org/10.1038/s41392-020-00335-z>
- Shen Z, Ratia K, Cooper L, Lee H, Kwon Y, Li Y, Algarni S, Huang F, Dubrovskiy O, Thatcher GRJ, Xiong R (2021) Design of SARS-CoV-PLpro inhibitors for COVID-19 antiviral therapy leveraging binding cooperativity. *J Med Chem*. <https://doi.org/10.1021/acs.jmedchem.1c01307>
- Jin Z, Du X, Xu Y, Deng Y, Liu M, Zhao Y, Zhang B, Li X, Zhang L, Peng C, Duan Y, Yu J, Wang L, Yang K, Liu F, Jiang R, Yang X, You T, Liu X, Yang X, Bai F, Liu H, Liu X, Guddat LW, Xu W, Xiao G, Qin C, Shi Z, Jiang H, Rao Z, Yang H (2020) Structure of M^{pro} from SARS-CoV-2 and discovery of its inhibitors. *Nature* 582:89–293. <https://doi.org/10.1038/s41586-020-2223-y>
- Zhu W, Chen CZ, Gorshkov K, Xu M, Lo DC, Zheng W (2020) RNA-dependent RNA polymerase as a target for COVID-19 drug discovery. *SLAS Discov: Adv Sci Drug Discov* 25:1141–1151. <https://doi.org/10.1177/2472555220942123>
- Namchuk MN (2021) Early returns on small molecule therapeutics for SARS-CoV-2. *ACS Infect Dis* 7:1298–1302. <https://doi.org/10.1021/acscinfecdis.0c00874>
- Keretsu S, Bhujbal SP, Cho SJ (2020) Rational approach toward COVID-19 main protease inhibitors via molecular docking, molecular dynamics simulation and free energy calculation. *Sci Rep* 10:17716. <https://doi.org/10.1038/s41598-020-74468-0>
- Yang H, Xie W, Xue X, Yang K, Ma J, Liang W, Zhao Q, Zhou Z, Pei D, Ziebuhr J, Hilgenfeld R, Yuen KY, Wong L, Gao G, Chen S, Chen Z, Ma D, Bartlam M, Rao Z (2005) Design of wide-spectrum inhibitors targeting coronavirus main proteases. *PLoS Biol* 3:e324. <https://doi.org/10.1371/journal.pbio.0030324>
- Ampornnanai K, Meng X, Shang W, Jin Z, Rogers M, Zhao Y, Rao Z, Liu ZJ, Yang H, Zhang L, Neill PM, Samar Hasnain S (2012) Inhibition mechanism of SARS-CoV-2 main protease by ebsele and its derivatives. *Nat Commun* 12:306. <https://doi.org/10.1038/s41467-021-23313-7>
- Zhang L, Lin D, Sun X, Curth U, Drosten C, Sauerhering L, Becker S, Rox K, Hilgenfeld R (2020a) Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved α -ketoamide inhibitors. *Science (New York, NY)* 368:409–412. <https://doi.org/10.1126/science.abb3405>
- Zhang ZJ, Wu WY, Hou JJ, Zhang LL, Li FF, Gao L, Wu XD, Shi JY, Zhang R, Long HL, Lei M, Wu WY, Guo DA, Chen KX, Hofmann LA, Ci ZH (2020b) Active constituents and mechanisms of Respiratory Detox Shot, a traditional Chinese medicine prescription, for COVID-19 control and prevention: Network-molecular docking-LC-MS analysis. *J Integrat Med* 18:229–241. <https://doi.org/10.1016/j.joim.2020.03.004>
- Calligari P, Bobone S, Ricci G, Bocedi A (2020) Molecular investigation of SARS-CoV-2 proteins and their interactions with antiviral drugs. *Viruses* 12:445. <https://doi.org/10.3390/v12040445>
- Nutho B, Mahalapbutr P, Hengphasatporn K, Pattarangoon NC, Simanon N, Shigeta Y, Hannongbua S, Rungrotmongkol T (2020) Why are lopinavir and ritonavir effective against the newly emerged coronavirus 2019? Atomistic Insights into the Inhibitory Mechanisms. *Biochemistry*. 59:1769–1779. <https://doi.org/10.1021/acs.biochem.0c00160>
- Chang Y, Tung Y, Lee K, Chen T, Hsiao Y, Chang H, Hsieh T, Su C, Wang S, Yu J, Shih S, Lin Y, Lin Y, Tu YE, Hsu C, Juan H, Tung C, Chen C (2020) Potential therapeutic agents for COVID-19 based on the analysis of protease and RNA polymerase docking. *Preprints*. <https://doi.org/10.20944/preprints202002.0242.v2>
- Xu Z, Peng C, Shi Y, Zhu Z, Mu K, Wang X, Zhu W (2020a) Nelfinavir was predicted to be a potential inhibitor of 2019-nCoV main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. *bioRxiv*. <https://doi.org/10.1101/2020.01.27.921627>
- Xu Z, Shen J, Wu N, Xu Y, Lu X, Zhu W, Li LJ (2020b) Nelfinavir is active against SARS-CoV-2 in Vero E6 cells. *ChemRxiv*. <https://doi.org/10.26434/chemrxiv.12039888.v1>
- Beck BR, Shin B, Choi Y, Park S, Kang K (2020) Predicting commercially available antiviral drugs that may act on the novel coronavirus (2019-nCoV), Wuhan, China through a drug-target interaction deep learning model. *Comput Struct Biotechnol J* 18:784–790. <https://doi.org/10.1016/j.csbj.2020.03.025>
- Fintelman-Rodrigues N, Sacramento CQ, Ribeiro Lima C, Souza da Silva F, Ferreira AC, Mattos M, de Freitas CS, Cardoso Soares V, da Silva Gomes

- Dias S, Temerozo JR, Miranda MD, Matos AR, Bozza FA, Carels N, Alves CR, Siqueira MM, Bozza PT, Souza TML (2020) Atazanavir, alone or in combination with ritonavir, inhibits SARS-CoV-2 replication and proinflammatory cytokine production. *Antimicrob Agents Chemother* 64:e00825–e00820. <https://doi.org/10.1128/AAC.00825-20>
28. Chen YW, Yiu CB, Wong KY (2020) Prediction of the SARS-CoV-2 (2019-nCoV) 3C-like protease (3CL^{PRO}) structure: virtual screening reveals velpatasvir, ledipasvir, and other drug repurposing candidates. *F1000Research* 9:129. <https://doi.org/10.12688/f1000research.22457.2>
29. Li Y, Zhang J, Wang N, Li H, Shi Y, Guo G, Liu K, Zeng H, Zou Q (2020) Therapeutic drugs targeting 2019-nCoV main protease by high-throughput screening. *bioRxiv*. <https://doi.org/10.1101/2020.01.28.922922>
30. Khan RJ, Jha RK, Amara GM, Jain M, Singh E, Pathak A, Singh RP, Muthukumar J, Singh AK (2021) Targeting SARS-CoV-2: a systematic drug repurposing approach to identify promising inhibitors against 3C-like proteinase and 2'-O-ribose methyltransferase. *J Biomol Struct Dyn* 39:2679–2692. <https://doi.org/10.1080/07391102.2020.1753577>
31. Talluri S (2021) Molecular docking and virtual screening based prediction of drugs for COVID-19. *Comb Chem High Throughput Screen* 24:716–728. <https://doi.org/10.2174/1386207323666200814132149>
32. Muralidharan N, Sakthivel R, Velmurugan D, Gromiha MM (2021) Computational studies of drug repurposing and synergism of lopinavir, oseltamivir and ritonavir binding with SARS-CoV-2 protease against COVID-19. *J Biomol Struct Dyn* 39:2673–2678. <https://doi.org/10.1080/07391102.2020.1752802>
33. Hall DC, Ji HF (2020) A search for medications to treat COVID-19 via in silico molecular docking models of the SARS-CoV-2 spike glycoprotein and 3CL protease. *Travel Med Infect Dis* 35:101646. <https://doi.org/10.1016/j.tmaid.2020.101646>
34. Wang C, Horby PW, Hayden FG, Gao GF (2020) A novel coronavirus outbreak of global health concern. *Lancet* 395:470–473. [https://doi.org/10.1016/S0140-6736\(20\)30185-9](https://doi.org/10.1016/S0140-6736(20)30185-9)
35. Liu X, Wang XJ (2020) Potential inhibitors for 2019-nCoV coronavirus M protease from clinically approved medicines. *J Genet Genomics* 47:119–121. <https://doi.org/10.1016/j.jgg.2020.02.001>
36. Sisakht M, Solhjo A, Mahmoodzadeh A, Fathalipour M, Kabiri M, Sakhteman A (2021) Potential inhibitors of the main protease of SARS-CoV-2 and modulators of arachidonic acid pathway: non-steroidal anti-inflammatory drugs against COVID19. *Comput Biol Med* 136:104686
37. Gimeno A, Mestres-Trujol J, Ojeda-Montes MJ, Macip G, Saldívar-Espinoza B, Cereto-Massagué A, Pujadas G, García-Vallvé S (2020) Prediction of novel inhibitors of the main protease (M-pro) of SARS-CoV-2 through consensus docking and drug reposition. *Int J Mol Sci* 21:3793. <https://doi.org/10.3390/ijms21113793>
38. Liu X, Li Z, Liu S, Sun J, Chen Z, Jiang M, Zhang Q, Wei Y, Wang X, Huang YY, Shi Y, Xu Y, Xian H, Bai F, Ou C, Xiong B, Lew AM, Cui J, Fang R, Huang H, Zhao J, Hong X, Zhang Y, Zhou F, Luo HB (2020) Potential therapeutic effects of dipyridamole in the severely ill patients with COVID-19. *Acta Pharmaceut Sinica B* 10:1205–1215. <https://doi.org/10.1016/j.apsb.2020.04.008>
39. Odhar HA, Ahjel SW, Albeer A, Hashim AF, Rayshan AM, Humadi SS (2020) Molecular docking and dynamics simulation of FDA approved drugs with the main protease from 2019 novel coronavirus. *Bioinformation* 16:236–244. <https://doi.org/10.6026/97320630016236>
40. Su H, Yao S, Zhao W, Li M, Liu J, Shang W, Xie H, Ke C, Hu H, Gao M, Yu K, Liu H, Shen J, Tang W, Zhang L, Xiao G, Ni L, Wang D, Zuo J, Jiang H, Bai F, Wu Y, Ye Y, Xu Y (2020) Anti-SARS-CoV-2 activities in vitro of Shuanghuanglian preparations and bioactive ingredients. *Acta Pharmacol Sin* 41:1167–1177. <https://doi.org/10.1038/s41401-020-0483-6>
41. Liu H, Ye F, Sun Q, Liang H, Li C, Lu R, Huang B, Tan W, Lai L (2021) *Scutellaria baicalensis* extract and baicalin inhibit replication of SARS-CoV-2 and its 3C-like protease in vitro. *J Enzyme Inhib Med Chem* 36:497–503. <https://doi.org/10.1080/14756366.2021.187397>
42. Cherrak SA, Merzouk H, Mokhtari-Soulimane N (2020) Potential bioactive glycosylated flavonoids as SARS-CoV-2 main protease inhibitors: a molecular docking and simulation studies. *PLoS One* 15. <https://doi.org/10.1371/journal.pone.0240653>
43. Shivanika C, Deepak Kumar S, Ragunathan V, Tiwari P, Sumitha A, Brindha Devi P (2020) Molecular docking, validation, dynamics simulations, and pharmacokinetic prediction of natural compounds against the SARS-CoV-2 main-protease. *J Biomol Struct Dyn* 8:1–27. <https://doi.org/10.1080/07391102.2020.1815584>
44. Bhaliya J, Shah V (2020) Identification of potent COVID-19 main protease (MPro) inhibitor from curcumin analogues by molecular docking analysis. *Int J Adv Res, Ideas Innov Technol* 6:664–672
45. Joshi RS, Jagdale SS, Bansode SB, Shankar SS, Tellis MB, Pandya VK, Chugh A, Giri AP, Kulkarni MJ (2021) Discovery of potential multi-target-directed ligands by targeting host-specific SARS-CoV-2 structurally conserved main protease. *J Biomol Struct Dyn* 39:3099–3114. <https://doi.org/10.1080/07391102.2020.1760137>
46. Enmozhi SK, Raja K, Sebastine I, Joseph J (2021) Andrographolide as a potential inhibitor of SARS-CoV-2 main protease: an in silico approach. *J Biomol Struct Dyn* 39:3092–3098. <https://doi.org/10.1080/07391102.2020.1760136>
47. Ramaiah MHS, Vishwakarma R, Umashaanker R (2020) Molecular docking analysis of selected natural products from plants for inhibition of SARS-CoV-2 main protease. *Curr Sci* 118:1087–1092
48. Umesh KD, Selvaraj C, Singh SK, Dubey VK (2021) Identification of new anti-CoV drug chemical compounds from Indian spices exploiting SARS-CoV-2 main protease as target. *J Biomol Struct Dyn* 39:3428–3434. <https://doi.org/10.1080/07391102.2020.1763202>
49. Khaerunnisa S, Kurniawan H, Awaluddin R, Suhartati S, Soetjipto S (2020) Potential inhibitor of COVID-19 main protease (Mpro) from several medicinal plant compounds by molecular docking study. *Preprints* 2020030226. <https://doi.org/10.20944/preprints202003.0226.v1>
50. Fischer W, Eron JJ, Holman W, Cohen MS, Fang L, Szwedczyk LJ, Sheahan TP, Baric R, Mollan KR, Wolfe CR, Duke ER, Azizad MM, Borroto-Esoda K, Wohl DA, Loftis AJ, Alabanza P, Lipansky F, Painter WP, Molnupiravir, an Oral Antiviral Treatment for COVID-19. *medRxiv* [Preprint]. 2021. <https://doi.org/10.1101/2021.06.17.21258639>
51. Gupta A, Zhou HX (2020) Profiling SARS-CoV-2 Main Protease (M^{PRO}) Binding to repurposed drugs using molecular dynamics simulations in classical and neural network-trained force fields. *ACS Comb Sci* 22:826–832. <https://doi.org/10.1021/acscombsci.0c00140>
52. Pal A, Curtin JF, Kinsella GK (2021) Structure based prediction of a novel GPR120 antagonist based on pharmacophore screening and molecular dynamics simulations. *Comput Struct Biotechnol J* 19:6050–6063
53. Zhang X, Macielag MJ (2020) GPR120 agonists for the treatment of diabetes: a patent review (2014-present). *Expert Opin Ther Pat* 30:729–742. <https://doi.org/10.1080/13543776.2020.181185>
54. Sparks SM, Chen G, Collins JL, Danger D, Dock ST, Jayawickreme C, Jenkinson S, Laudeman C, Leesnitzer MA, Liang X, Maloney P, McCoy DC, Moncol D, Rash V, Rimele T, Vulimiri P, Way JM, Ross S (2014) Identification of diarylsulfonamides as agonists of the free fatty acid receptor 4 (FFA4/GPR120). *Bioorg Med Chem Lett* 24:3100–3103
55. Sparks SM, Aquino C, Banker P, Collins JL, Cowan D, Diaz C, Dock ST, Hertzog DL, Liang X, Swiger ED, Yuen J, Chen G, Jayawickreme C, Moncol D, Nyström C, Rash V, Rimele T, Roller S, Ross S (2017) Exploration of phenylpropanoic acids as agonists of the free fatty acid receptor 4 (FFA4): identification of an orally efficacious FFA4 agonist. *Bioorg Med Chem Lett* 27:1278–1283
56. Zhang X, Cai C, Winters M, Wells M, Wall M, Lanter J, Sui Z, Ma J, Novack A, Nashashibi I, Wang Y, Yan W, Suckow A, Hua H, Bell A, Haug P, Clapper W, Jenkinson C, Gunnet J, Leonard J, Murray WV (2017) Design, synthesis and SAR of a novel series of heterocyclic phenylpropanoic acids as GPR120 agonists. *Bioorg Med Chem Lett* 27:3272–3278
57. Janssen Pharmaceutica NV (2014) Preparation of GPR120 agonists for the treatment of type II diabetes. *US* 20140275179 A1
58. Janssen Pharmaceutica NV (2019) Cycloalkenyl derivatives useful as agonists of the GPR120 and/or GPR40 receptors and their preparation. *WO* 2019171277 A1
59. Piramal Enterprises Limited. Preparation of fused heterocyclic compounds as GPR120 agonists. *WO* 2016038540 A1; 2016b.
60. Piramal Enterprises Limited (2016c) Preparation of thiazole and thiophene derivatives containing carbon-carbon linker as GPR120 agonists for disease therapy. *WO* 2016125182 A1, p 74
61. da Silva Arouche T, Reis AF, Martins AY, Costa JFS, Junior RNC, Antonio Maia JC (2020) Neto interactions between Remdesivir, Ribavirin, Favipiravir, Galidesivir, Hydroxychloroquine and Chloroquine with fragment molecular of the COVID-19 main protease with inhibitor N3

- complex (PDB ID:6LU7) using molecular docking. *J Nanosci Nanotechnol* 20(12):7311–23. <https://doi.org/10.1166/jnn.2020.18955>
62. Bristol-Meyers Squibb Company. Preparation of novel cyclopropane carboxylic acid compounds as GPR120 modulators useful for treatment of diabetes and related diseases. WO 2016040222 A1; 2016.
 63. Selami E, Çınar E (2021) A molecular docking study of potential inhibitors and repurposed drugs against SARS-CoV-2 main protease enzyme. *J Indian Chem Soc* 98:100041
 64. Smarajit M, Banerjee A, Nazmeen A, Kanwar M, Das S (2020) Active-site molecular docking of Nigellidine with nucleocapsid- NSP2-MPro of COVID-19 and to human IL1R-IL6R and strong antioxidant role of Nigella-sativa in experimental rats. *J Drug Target*. <https://doi.org/10.1080/1061186X.2020.1817040>
 65. Rai H, Barik A, Singh YP, Suresh A, Singh L, Singh G, Nayak UY, Dubey VK, Modi G (2021) Molecular docking, binding mode analysis, molecular dynamics, and prediction of ADMET/toxicity properties of selective potential antiviral agents against SARS-CoV-2 main protease: an effort toward drug repurposing to combat COVID-19. *Mol Divers* 25(3):1905–1927. <https://doi.org/10.1007/s11030-021-10188-5>. Epub 2021 Feb 13. PMID: 33582935; PMCID: PMC7882058
 66. Rashi J, Somdutt M (2020) Repurposing metocurine as main protease inhibitor to develop novel antiviral therapy for COVID-19. *Struct Chem* 31:2487–2499
 67. Hosseini M, Chen W, Xiao D, Wang C (2021) Computational molecular docking and virtual screening revealed promising SARS-CoV-2 drugs. *Precis Clin Med* 4(1):1–16. <https://doi.org/10.1093/pcmedi/pbab001> PMID: 33842834; PMCID: PMC7928605
 68. Toelzer C, Gupta K, Yadav SKN, Borucu U, Davidson AD, Kavanagh Williamson M, Shoemark DK, Garzoni F, Staufer O, Milligan R, Capin J, Mulholland AJ, Spatz J, Fitzgerald D, Berger I, Schaffitzel C (2020) Free fatty acid binding pocket in the locked structure of SARS-CoV-2 spike protein. *Science* 370(6517):725–730. <https://doi.org/10.1126/science.abd3255> Epub 2020 Sep 21. PMID: 32958580; PMCID: PMC8050947
 69. <https://doi.org/10.1016/j.combiomed.2021.104686>
 70. <https://doi.org/10.1101/2021.06.17.21258639>
 71. Piramal Enterprises Limited (2015) Preparation of carboxylic acid derivatives for use as GPR120 agonists. WO 2015125085 A1, p 73
 72. Piramal Enterprises Limited (2016a) Preparation of phenyl alkanoid acid derivatives for use as GPR120 agonists. WO 2016012965 A2, p 75
 73. Usha YN, Dubey VK, Modi G Molecular docking, binding mode analysis, molecular dynamics, and prediction of ADMET/toxicity properties of selective potential antiviral agents against SARS-CoV-2 main protease: an effort toward drug repurposing to combat COVID-19. *Mol Divers*. <https://doi.org/10.1007/s11030-021-10188-5>
 74. Wang J (2020) Fast identification of possible drug treatment of coronavirus disease-19 (COVID-19) through computational drug repurposing study. *J Chem Inf Model* 60:3277–3286. <https://doi.org/10.1021/acs.jcim.0c00179>
 75. Willyard C (2021) How antiviral pill molnupiravir shot ahead in the COVID drug hunt? *Nature*. <https://doi.org/10.1038/d41586-021-02783-1>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
