

# The protective effects of ischemic preconditioning on rats with renal ischemia-reperfusion injury and the effects on the expression of Bcl-2 and Bax

SHENG SHEN, JIEXUE ZHOU, SHANDONG MENG, JIAQING WU,  
JUAN MA, CHUNLI ZHU, GENGGUO DENG and DONG LIU

Department of Organ Transplantation, Guangdong Second Provincial General Hospital,  
Guangzhou, Guangdong 510317, P.R. China

Received June 29, 2017; Accepted August 18, 2017

DOI: 10.3892/etm.2017.5047

**Abstract.** The aim of the present study was to investigate the protective effects of ischemic preconditioning on rats with renal ischemia-reperfusion injury and the effects on the expression of Bcl-2 and Bax. Thirty-six SD rats were randomly divided into three groups (n=12) including sham operation (S) group, ischemia-reperfusion group (I/R) group and ischemic preconditioning (IP) group. After anesthesia with intraperitoneal injection of chloral hydrate, bilateral renal pedicles were clipped for 45 min, followed by perfusion for 6 h to establish the I/R model. Both kidneys in rats of S group were separated and exposed for 45 min, but renal pedicles were not clipped. In IP group, bilateral renal pedicles were clipped for 5 min, followed by perfusion for 5 min, this procedure was repeated 3 times. Then bilateral renal pedicles were clipped for 45 min, followed by perfusion for 6 h. Blood samples were collected and rats were sacrificed to collect renal tissue. Levels of serum creatinine (Cr) and blood urea nitrogen (BUN) were measured. Activity of superoxide dismutase (SOD) was measured by xanthine oxidase assay. Degree of renal injury was evaluated by H&E staining. TUNEL kit was used to detect the number of apoptotic cells in renal tissue. Expression levels of Bcl-2 and Bax were detected by semi-quantitative PCR and western blot analysis at mRNA and protein levels, respectively. Results showed that levels of Cr and BUN in I/R and IP groups were significantly higher than those in S group, and levels of Cr and BUN in I/R group were significantly higher than that in IP group (P<0.05). Activity of SOD in I/R group and IP group were significantly lower than those in S group, and activity of SOD in I/R group were significantly lower than those in IP group (P<0.05). H&E staining showed that, compared

with S group, renal injury in the I/R and IP groups was more serious than that in the S group, and I/R group was more serious than the IP group (P<0.05). TUNEL apoptosis assay showed that number of apoptotic cells in IP and I/R groups were significantly higher than that in the S group (P<0.01). Semi-quantitative PCR and western blot analysis showed that, compared with the S group, expression levels of Bcl-2 mRNA and protein were significantly decreased, expression levels of Bax mRNA and protein were significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in the IP and I/R groups (P<0.01). Compared with the I/R group, expression level of Bcl-2 was significantly increased, the level of Bax was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in the IP group (P<0.01). As a result, ischemic preconditioning can protect rats with renal ischemia-reperfusion injury possibly by increasing the expression level of Bcl-2 and decreasing the expression level of Bax.

## Introduction

As a common renal disease in clinical practice, acute renal injury is caused by renal ischemia-reperfusion. Ischemia/reperfusion is the determinant factor for the early functional recovery of renal transplantation (1,2). As a highly perfused organ, kidney is vulnerable to ischemic injury. Studies have shown that renal ischemia for more than 20 min will cause damage, ischemia for 20-40 min is reversible, but reperfusion after ischemia for more than 40 min can cause permanent damage (3). Studies have shown that the development of tolerance of organs to ischemia can inhibit or reduced the ischemia-reperfusion injury (4). Takahashi-Sato *et al* (5) carried out a study to explore the effects of different preconditioning and postconditioning methods on ischemia/reperfusion in the heart. They found that postconditioning had a protective effect on reperfusion after cardiac ischemia (6-8). However, there is no research on the effect of preconditioning and postconditioning on renal ischemia-reperfusion. It was found (9) that Bcl-2 and Bax were involved in the process of ischemia-reperfusion in the heart, and Bcl-2 and Bax may be related to the protective effect of postconditioning on ischemia-reperfusion injury in the heart. In this study, rat ischemia-reperfusion model was established

---

*Correspondence to:* Dr Dong Liu, Department of Organ Transplantation, Guangdong Second Provincial General Hospital, 466 Xingang Middle Road, Haizhu, Guangzhou, Guangdong 510317, P.R. China  
E-mail: kr62ho@163.com

*Key words:* renal ischemia-reperfusion, ischemic preconditioning, protective effect, Bcl-2, Bax

to investigate the protective effects of preconditioning on rats with ischemia-reperfusion, and to test whether Bcl-2 and Bax were involved in that process. This study aimed to provide new insights into the pathogenesis and clinical treatment of renal ischemia-reperfusion injury.

## Materials and methods

**Instruments and materials.** Creatinine (Cr) assay kit (R&D Systems, Minneapolis, MN, USA); blood urea nitrogen (BUN) assay kit (R&D Systems); xanthine oxidase assay kit used to determine the activity of superoxide dismutase (SOD; R&D Systems); TUNEL kit (R&D Systems); dimethyl sulfoxide (DMSO; Sigma-Aldrich, St. Louis, MO, USA), TRIzol kit (Invitrogen, Carlsbad, CA, USA), reverse transcription kit (Invitrogen); rabbit anti-Bax-beta, rabbit anti-Bax and rabbit anti-GAPDH primary antibodies, and horseradish peroxidase labeled anti-rabbit secondary antibody (Cell Signaling Technology, Danvers, MA, USA); ECL luminescent substrate (Invitrogen); color developing powder (Invitrogen); pipettes (Eppendorf, Hamburg, Germany); PCR instrument (Applied Biosystems, Inc., Foster City, CA, USA); UV imaging system (Biometra GmbH, Göttingen, Germany); electronic balance (Sartorius BP121S; Sartorius AG, Goettingen, Germany); -80°C refrigerator (Thermo Fisher Scientific, Schwerte, Germany); low temperature centrifuge (Thermo Fisher Scientific) and frozen slicers (Leica Microsystems, Wetzlar, Germany). The sources of other related instruments and reagents are described in the relevant section.

**Experimental animals and grouping.** Male Sprague-Dawley (SD) rats (220-250 g) were purchased from the Shanghai Medical Laboratory Animal Center. Laboratory animal certificate number: SCXK (Shanghai) 2013-0015. Rats were raised in quiet environment with free access to food and water following the circadian rhythm for 1 week to adapt to the environment. Thirty-six SD rats were randomly divided into three groups (n=12) including sham operation (S) group, ischemia-reperfusion group (I/R) group and ischemic preconditioning (IP) group.

**Establishment of renal ischemia-reperfusion model.** Rats were fasted for 12 h before surgery. After anesthesia with intraperitoneal injection of 4% chloral hydrate at a dose of 10 ml/kg, abdominal hair was removed and an incision was made on abdomen along the midventral line to open abdominal cavity. Renal capsule was bluntly separated to separate the kidneys. Arterial clip was then used to clip bilateral kidney pedicles. The dark red color of kidneys indicated the successfully established ischemia model. In I/R group, bilateral renal pedicles were clipped for 45 min, followed by perfusion for 6 h to establish I/R model. Both kidneys in rats of S group were separated and exposed for 45 min, but renal pedicles were not clipped. In IP group, bilateral renal pedicles were clipped for 5 min, followed by perfusion for 5 min, this procedure was repeated 3 times, after that rats were subjected to the treatment described in the I/R group.

**Sample collection and preparation.** Peripheral blood was collected after reperfusion for 6 h. Blood samples were centri-

Table I. Primers used in PCR.

Sequences	
Bax	Forward: 5'-ATCCAGAGACAAGACATGTAC-3' Reverse: 5'-TTCAGATGTTCTAAGCCTACGG-3'
Bcl-2	Forward: 5'-TGGCGGTTTTCGGTGGAC-3' Reverse: 5'-CCAGTGCAGGGTCCGAGGT-3'
GAPDH	Forward: 5'-GATGATTGGCATGGCTTT-3' Reverse: 5'-CACCTTCCGTTCCAGTTT-3'

fuged 2,650 x g for 15 min to collect supernatant. Supernatant was transferred to tubes and stored at -80°C. After blood sample collection, rats in each group were sacrificed, and both kidneys were separated and fixed in 10% formalin for subsequent pathologic examination and other studies.

**Renal tissue injury scoring.** After H&E staining, pathological changes were observed under optical microscope (BX-42; Olympus, Tokyo, Japan) and scores of renal pathological damage were evaluated. Scoring was performed according to the standard of renal injury: normal renal tissue, 0 point; area of renal tissue damage in renal tubular area <25%, 1 point; area of renal tissue damage in renal tubular between 25 and 50%, 2 points; area of renal tissue damage in renal tubular between 50 and 75%, 3 points; area of renal tissue damage in renal tubular between >75%, 4 points.

**Determination of levels of Cr and BUN and activity of SOD in serum.** Contents of Cr and BUN in serum of rats in each group were determined according to the instructions of the Cr assay kit and BUN assay kit. Cr content was expressed as number +  $\mu$ mg/ml and BUN content expressed as number + mmol/ml. SOD activity in serum of rats in each group was determined according to the instructions of SOD activity assay kit and expressed as number + U/mg.

**TUNEL to detect the apoptosis of rat renal cells.** TUNEL assay was performed according to the instructions of kit and apoptotic cells were observed under an optical microscope. Cells with brown nucleus were positive cells. Five visual fields were randomly selected (magnification, x400) to count the number of apoptotic cells. The proportion of the number apoptotic cells to the number of total cells was used as apoptotic index.

**Semi-quantitative PCR to detect the expression of related mRNA.** Total RNA was extracted from renal tissue using the TRIzol kit. The quality of total RNA was checked by agarose gel electrophoresis. Only RNA samples with satisfactory qualities were used in subsequent experiments. Reverse transcription was performed using a kit to synthesize cDNA and expression levels of Bax and Bcl-2 were detected by semi-quantitative PCR with GAPDH as endogenous control. Reaction conditions were: 95°C for 4 min, followed by 35 cycles of 95°C for 30 sec, 64°C for 25 sec and 72°C for 30 sec. Primers were synthesized by Tiangen Biotech Co., Ltd. (Beijing, China). All primers are listed in Table I. PCR products were subjected to agarose gel




Figure 1. Levels of Cr in peripheral blood of rats in each group. Compared with group S, level of Cr was significantly increased in I/R and IP groups (\*\* $P<0.01$ ). Compare with IP group, level of Cr was significantly increased in I/R group (# $P<0.01$ ).




Figure 2. Levels of BUN in peripheral blood of rats in each group. Compared with group S, level of BUN was significantly increased in I/R and IP groups (\*\* $P<0.01$ ). Compared with IP group, level of BUN was significantly increased in I/R group (# $P<0.01$ ).

electrophoresis and results were observed using a UV imaging system. The relative expression levels of Bax and Bcl-2 were represented by the ratio of Bax/GAPDH and Bcl-2/GAPDH.

**Western blot analysis to detect the expression of related proteins.** Renal tissue was mixed with RIPA lysate with a ratio of 100 mg:1 ml. The mixture was centrifuged (10,500  $\times$  g) at 4°C for 10 min to collect the supernatant. Protein concentration was measured using the DAB Protein Quantitative kit (Invitrogen). After that, protein samples were subjected to SDS-PAGE gel electrophoresis, followed by transmembrane to PVDF membrane. After blocking with 5% skim milk, membranes were incubated with primary rabbit polyclonal Bax antibody (dilution, 1:500; cat. no. ab53154) and rabbit monoclonal Bcl-2 antibody (dilution, 1:500; cat. no. ab32124) overnight at 4°C. After washing with TBST 3 times, 5 min for each time, membranes were incubated with secondary goat anti-rabbit (HRP) IgG antibody (dilution, 1:2000; cat. no. ab6721) for 2 h. All antibodies were all purchased from Abcam (Cambridge, MA, USA). After washing with TBST 3 times, proper amount of ECL luminescent substrate (ratio of A and B was 1:1) was added and incubated with the membranes in the dark. The results were scanned and processed by ImageJ to calculate the expression level of each protein.

**Statistical analysis.** Data were expressed as mean  $\pm$  standard deviation, and processed using the SPSS 19.0 software (SPSS, Inc., Chicago, IL, USA). Comparisons between 2 groups were performed using t-test, and comparisons among multiple groups were performed using variance analysis. If the variance was homogeneous, the comparisons between two groups were performed using Bonferroni method. If the variance was not homogeneous, Welch method was used. Multiple comparisons were performed using Dunnett's T3 method.  $P<0.05$  was considered to be statistically significant.

## Results

**Changes in renal function in rats.** Levels of Cr and BUN in peripheral blood of rats in each group were detected by Cr and BUN detection kit. As shown in Figs. 1 and 2, compared with group S, levels of Cr and BUN were significantly increased in I/R group and IP group ( $P<0.01$ ). Compared with IP group,




Figure 3. Activity of SOD in rats of each group. Activity of SOD in I/R group and IP group was significantly lower than those in S group (\*\* $P<0.01$ ), and activity of SOD in I/R group was significantly lower than those in IP group (# $P<0.05$ ).

levels of Cr and BUN were significantly increased in the I/R group ( $P<0.01$ ).

**Changes in SOD activity.** Activity of SOD was detected by SOD activity test kit. As shown in Fig. 3, activity of SOD in the I/R and IP groups was significantly lower than in the S group ( $P<0.01$ ), and activity of SOD in the I/R group was significantly lower than in the IP group ( $P<0.05$ ).

**Pathological changes of the kidneys in rats.** Pathological changes of renal tissue in each group after H&E staining were observed under an optical microscope. Structure of renal tissue in group S was clear, renal tubular and glomerular were normal, degeneration and atrophy were not observed in renal tubular epithelial cells, and the cavity was not expanded. In the I/R and IP groups, renal tissue glomerular telangiectasia and congestion, renal tubular epithelial cell edema, and granular degeneration or vacuolar degeneration were observed and the cavity was narrowed. After scoring (Table II), renal injury was found to be more serious in the I/R and IP groups than in the S group ( $P<0.01$ ), and renal injury was more serious in the I/R group than in the IP group ( $P<0.05$ ).

**Renal cell apoptosis.** Apoptotic cells were detected using TUNEL kit. Number of positive cells was counted and the

Table II. Pathological changes of kidney in rats of each group.

Groups	Renal injury score
S group	0.68±0.19
I/R group	2.39±0.29 <sup>a</sup>
IP group	1.89±0.33 <sup>a,b</sup>

<sup>a</sup>P<0.01 compared with the S group; <sup>b</sup>P<0.05 compared with the I/R group.




Figure 4. (A and B) Cell apoptosis detected by TUNEL. Compared with S group, number of apoptotic cells in I/R group and IP group was significantly increased (\*\*P<0.01). Compared with IP group, number of apoptotic cells in I/R group was significantly increased (##P<0.01).

apoptotic index was calculated. As shown in Fig. 4, compared with S group, number of apoptotic cells in I/R group and IP group was significantly increased (P<0.01). Compared with IP group, the number of apoptotic cells in I/R group was significantly increased (P<0.01).

**mRNA expression detected by semi-quantitative PCR.** Expression of Bcl-2 and Bax mRNA was detected by semi-quantitative PCR. As shown in Fig. 5, compared with S group, expression level of Bcl-2 mRNA was significantly decreased, expression level of Bax mRNA was significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in IP group and I/R group (P<0.01). Compared with I/R group, expression level of Bcl-2 mRNA was significantly increased, expression level of Bax mRNA was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in IP group (P<0.05).

**Protein expression detected by western blot analysis.** Expression of Bcl-2 and Bax protein was detected by western blot analysis. As shown in Fig. 6, compared with S group, expression level of Bcl-2 protein was significantly decreased, expression level of Bax protein was significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in IP group




Figure 5. (A and B) Expression of Bcl-2 and Bax mRNA detected by semi-quantitative PCR. Compared with S group, expression level of Bcl-2 mRNA were significantly decreased, expression level of Bax mRNA was significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in IP group and I/R group (\*\*P<0.01). Compared with I/R group, expression level of Bcl-2 mRNA was significantly increased, expression level of Bax mRNA was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in IP group (#P<0.05).




Figure 6. (A and B) Expression of Bcl-2 and Bax protein detected by western blot analysis. Compared with S group, expression level of Bcl-2 protein were significantly decreased, expression level of Bax protein was significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in IP group and I/R group (\*\*P<0.01). Compared with I/R group, expression level of Bcl-2 protein was significantly increased, expression level of Bax protein was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in IP group (##P<0.01).

and I/R group (P<0.01). Compared with I/R group, expression level of Bcl-2 protein was significantly increased, expression level of Bax protein was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in IP group (P<0.01).

**Discussion**

Renal blood flow needs to be blocked in renal transplantation, nephrolithotomy and surgical resection of renal tumors and other complex surgeries, and renal reperfusion after blood

flow blocking can cause renal damage (10). Numerous studies have shown that ischemia-reperfusion injury may be related to the production of oxygen free radicals, adenine nucleotide metabolic disorders and cell apoptosis (11). Drugs (ulinastatin and propofol), low temperature, preconditioning and postconditioning can play a protective role in ischemia-reperfusion injury, and all these methods can be used to reduce ischemia-reperfusion injury to the kidneys (12-14).

In this study, rat ischemia-reperfusion model was established and levels of Cr and BUN in serum were detected to evaluate the degree of renal injury. Results showed that levels of Cr and BUN in the serum of rats in I/R group were significantly higher than those of S group, and the difference was statistically significant ( $P < 0.01$ ), indicating the successful establishment of rat ischemia-reperfusion model. Ischemia-reperfusion model was established by clamping bilateral renal pedicle. Reperfusion time and clamping sites can seriously affect the stability of the model. Preliminary experiment showed that reperfusion could significantly affect renal function and the mode was stable. This study showed that preconditioning performed by clamping bilateral renal pedicle for 5 min and reperfusion for 5 min could improve the renal ischemia-reperfusion injury in rats. Levels of Cr and BUN in IP group were higher than those in the I/R group ( $P < 0.01$ ). Ischemic preconditioning can induce the production of endogenous protective substances in the body to resist ischemia and reperfusion injury to organs and tissues, and this protective effect in multiple organs have been proved by various studies (15,16). Li and Liu (17) found that ischemic preconditioning can effectively reduce the effects of ischemia-reperfusion on myocardial function. Lipid peroxidation caused by oxygen free radical is one of the factors that cause ischemia-reperfusion injury. SOD activity can be used to indirectly assess the ability of the body to scavenge oxygen free radicals (18). In this study, compared with S group and IP group, SOD activity in I/R group was significantly reduced, indicating that ischemia-reperfusion could cause renal injury and ischemic preconditioning can protect the body. As the currently most commonly used method to study cell apoptosis, TUNEL can be used to directly observe apoptotic cells, and tissue or cell apoptosis can be evaluated by calculating apoptosis index. Bcl gene family, which contains anti-apoptotic gene Bcl-2 and pro-apoptotic gene Bax, is closely related to cell apoptosis. Bcl-2 can inhibit cell apoptosis by inhibiting the production of free radicals, intracellular calcium overload, reducing the permeability of mitochondrial membranes and maintaining the oxidative function of mitochondria, while Bax has opposite functions (19,20). This study found that, compared with S group, cell apoptosis was significantly increased in I/R group, and results of semi-quantitative PCR and western blot analysis showed that, compared with S group, expression levels of Bcl-2 mRNA and protein were significantly decreased, expression levels of Bax mRNA and protein were significantly increased, and the ratio of Bcl-2/Bax was significantly decreased in IP group and I/R group. Compared with I/R group, expression level of Bcl-2 was significantly increased, expression level of Bax was significantly decreased, and the ratio of Bcl-2/Bax was significantly increased in IP group. Those results suggest that ischemia-reperfusion can lead to increased cell apoptosis, while ischemic preconditioning can increase the membrane

stability and protect the kidneys by upregulating the expression of Bcl-2 and downregulating the expression of Bax in renal tissue.

In conclusion, ischemic preconditioning can protect rats with renal ischemia-reperfusion injury possibly by increasing the expression level of Bcl-2 and decreasing the expression level of Bax. The repeated short-term ischemia-reperfusion treatment can be applied in clinical practice to induce the injury to kidneys caused by ischemia-reperfusion.

## References

- Kim Y, Ge Y, Tao C, Zhu J, Chapman AB, Torres VE, Yu AS, Mrug M, Bennett WM, Flessner MF, *et al*: Consortium for Radiologic Imaging Studies of Polycystic Kidney Disease (CRISP): Automated segmentation of kidneys from MR images in patients with autosomal dominant polycystic kidney disease. *Clin J Am Soc Nephrol* 11: 576-584, 2016.
- Fox CS, Matsushita K, Woodward M, Bilo HJ, Chalmers J, Heerspink HJ, Lee BJ, Perkins RM, Rossing P, Sairenchi T, *et al*: Chronic Kidney Disease Prognosis Consortium: Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: A meta-analysis. *Lancet* 380: 1662-1673, 2012.
- Haynes R, Staplin N, Emberson J, Herrington WG, Tomson C, Agodoa L, Tesar V, Levin A, Lewis D, Reith C, *et al*: SHARP Collaborative Group: Evaluating the contribution of the cause of kidney disease to prognosis in CKD: Results from the Study of Heart and Renal Protection (SHARP). *Am J Kidney Dis* 64: 40-48, 2014.
- Ta MH, Schwensen KG, Foster S, Korgaonkar M, Ozimek-Kulik JE, Phillips JK, Peduto A and Rangan GK: Effects of TORC1 inhibition during the early and established phases of polycystic kidney disease. *PLoS One* 11: e0164193, 2016.
- Takahashi-Sato K, Murakawa M, Kimura J, Ito MA and Matsuoka I: Loss of ectonucleotidases from the coronary vascular bed after ischemia-reperfusion in isolated rat heart. *BMC Cardiovasc Disord* 68: 2577-2584, 2016.
- Chua S, Lee FY, Tsai TH, Sheu JJ, Leu S, Sun CK, Chen YL, Chang HW, Chai HT, Liu C, *et al*: Inhibition of dipeptidyl peptidase-IV enzyme activity protects against myocardial ischemia-reperfusion injury in rats. *J Transl Med* 19: 285-297, 2016.
- Wang K, Zhang J, Liu J, Tian J, Wu Y, Wang X, Quan L, Xu H, Wang W and Liu H: Variations in the protein level of Omi/HtrA2 in the heart of aged rats may contribute to the increased susceptibility of cardiomyocytes to ischemia/reperfusion injury and cell death: Omi/HtrA2 and aged heart injury. *Age (Dordr)* 30: 266-271, 2016.
- Huang P, Sun Y, Yang J, Chen S, Liu AD, Holmberg L, Huang X, Tang C, Du J and Jin H: The ERK1/2 signaling pathway is involved in sulfur dioxide preconditioning-induced protection against cardiac dysfunction in isolated perfused rat heart subjected to myocardial ischemia/reperfusion. *Int J Mol Sci* 206: 26-32, 2016.
- O'Neill KL, Huang K, Zhang J, Chen Y and Luo X: Inactivation of prosurvival Bcl-2 proteins activates Bax/Bak through the outer mitochondrial membrane. *Genes Dev* 30: 973-988, 2016.
- Lindqvist LM, Heinlein M, Huang DC and Vaux DL: Prosurvival Bcl-2 family members affect autophagy only indirectly, by inhibiting Bax and Bak. *Invest New Drugs* 34: 713-725, 2016.
- Zhang WZ, Li R, Liu S, Zhang JD, Ning XF and Cai SL: Effects of renal ischemic postconditioning on myocardial ultrastructural organization and myocardial expression of Bcl-2/Bax in rabbits. *BioMed Res Int* 2016: 9349437, 2016.
- Shen S, Wang JF, Wu JQ, Zhou JX, Meng SD, Ma J, Zhu CL, Deng GG and Liu D: GC/MS-based metabolomic analysis of alleviated renal ischemia-reperfusion injury induced by remote ischemic preconditioning. *Eur Rev Med Pharmacol Sci* 21: 765-774, 2017.
- Tawfik MK: Renoprotective activity of telmisartan versus pioglitazone on ischemia/reperfusion induced renal damage in diabetic rats. *Eur Rev Med Pharmacol Sci* 16: 600-609, 2012.
- Park SY, Shrestha S, Youn YJ, Kim JK and Kim SY: Role of Toll-like receptor-4 in renal graft ischemia-reperfusion injury. *Asian Pac J Trop Med* 9: 2-9, 2016.

15. Kierulf-Lassen C, Kristensen ML, Birn H, Jespersen B and Nørregaard R: No effect of remote ischemic conditioning strategies on recovery from renal ischemia-reperfusion injury and protective molecular mediators. *BMC Cancer* 16: 686-691, 2016.
16. Chen YT, Tsai TH, Yang CC, Sun CK, Chang LT, Chen HH, Chang CL, Sung PH, Zhen YY, Leu S, *et al*: Exendin-4 and sitagliptin protect kidney from ischemia-reperfusion injury through suppressing oxidative stress and inflammatory reaction. *J Transl Med* 11: 270, 2013.
17. Li Y and Liu S: The effect of dexmedetomidine on oxidative stress response following cerebral ischemia-reperfusion in rats and the expression of intracellular adhesion molecule-1 (ICAM-1) and S100B. *Med Sci Monit* 23: 867-873, 2017.
18. Batinic-Haberle I, Tovmasyan A, Roberts ER, Vujaskovic Z, Leong KW and Spasojevic I: SOD therapeutics: Latest insights into their structure-activity relationships and impact on the cellular redox-based signaling pathways. *Antioxid Redox Signal* 20: 2372-2415, 2014.
19. Zhang L, Gan W and An G: Influence of Tanshinone IIa on heat shock protein 70, Bcl-2 and Bax expression in rats with spinal ischemia/reperfusion injury. *Neural Regen Res* 7: 2882-2888, 2012.
20. Ma Z, Wei Q, Dong G, Huo Y and Dong Z: DNA damage response in renal ischemia-reperfusion and ATP-depletion injury of renal tubular cells. *Biochim Biophys Acta* 1842: 1088-1096, 2014.