



Corrigendum: A Multicenter Phase II RCT to Compare the Effectiveness of EMDR Versus TAU in Patients With a First-Episode Psychosis and Psychological Trauma: A Protocol Design

Alicia Valiente-Gómez^{1,2,3,4*}, Nuria Pujol^{2,3,5}, Ana Moreno-Alcázar^{1,2,3,4}, Joaquim Radua^{6,7,8}, Eila Monteagudo-Gimeno^{4,5}, Itxaso Gardoki-Souto^{1,2,4}, Bridget Hogg^{1,2,4}, Maria José Álvarez⁹, Gemma Safont^{3,4,10}, Walter Lupo¹, Victor Pérez^{2,3,4,5}, Benedikt L. Amann^{1,2,3,4} and the FEP-EMDR Research Group

¹ Centre Forum Research Unit, Institut de Neuropsiquiatria i Addiccions (INAD), Parc de Salut Mar, Barcelona, Spain, ² IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain, ³ CIBERSAM, Madrid, Spain, ⁴ Department of Psychiatry and Forensic Medicine, School of Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain, ⁵ Institut de Neuropsiquiatria i Addiccions (INAD), Hospital del Mar, Barcelona, Spain, ⁶ Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain, ⁷ Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom, ⁸ Department of Clinical Neuroscience, Centre for Psychiatry Research, Karolinska Institutet, Stockholm, Sweden, ⁹ Mental Health Department, Vic Hospital Consortium, Barcelona, Spain, ¹⁰ University Hospital Mutua Terrassa, Barcelona, Spain

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA,
Switzerland

*Correspondence:

Alicia Valiente-Gómez
62648@parcdesalutmar.cat

Specialty section:

This article was submitted to
Schizophrenia,
a section of the journal
Frontiers in Psychiatry

Received: 05 February 2020

Accepted: 23 March 2020

Published: 14 April 2020

Citation:

Valiente-Gómez A, Pujol N, Moreno-Alcázar A, Radua J, Monteagudo-Gimeno E, Gardoki-Souto I, Hogg B, Álvarez MJ, Safont G, Lupo W, Pérez V, Amann BL and the FEP-EMDR Research Group (2020) Corrigendum: A Multicenter Phase II RCT to Compare the Effectiveness of EMDR Versus TAU in Patients With a First-Episode Psychosis and Psychological Trauma: A Protocol Design. *Front. Psychiatry* 11:283. doi: 10.3389/fpsy.2020.00283

Keywords: first episode psychosis, psychological trauma, post-psychotic posttraumatic stress, comorbidity, EMDR therapy, treatment as usual

A Corrigendum on

A Multicenter Phase II RCT to Compare the Effectiveness of EMDR Versus TAU in Patients With a First-Episode Psychosis and Psychological Trauma: A Protocol Design

by Valiente-Gómez A, Pujol N, Moreno-Alcázar A, Radua J, Monteagudo-Gimeno E, Gardoki-Souto I, Hogg B, Álvarez MJ, Safont G, Lupo W, Pérez V., Amann BL and the FEP-EMDR Research Group (2020). *Front. Psychiatry* 10:1023. doi: 10.3389/fpsy.2019.01023

An author's name was incorrectly spelled as "Daniel Vergé" in the FEP-EMDR Research Group section. The correct spelling is "Daniel Bergé." The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way.

The original article has been updated.

Copyright © 2020 Valiente-Gómez, Pujol, Moreno-Alcázar, Radua, Monteagudo-Gimeno, Gardoki-Souto, Hogg, Álvarez, Safont, Lupo, Pérez, Amann and the FEP-EMDR Research Group. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.