

Association with HLA-DR β 1 position 37 distinguishes juvenile dermatomyositis from adult-onset myositis

Claire T. Deakin^{1,2,3,*}, John Bowes^{4,5}, Lisa G. Rider⁶, Frederick W. Miller⁶, Lauren M. Pachman⁷, Helga Sanner^{8,9}, Kelly Rouster-Stevens¹⁰, Gulnara Mamyrova¹¹, Rodolfo Curiel¹¹, Brian M. Feldman¹², Adam M. Huber¹³, Ann M. Reed¹⁴, Heinrike Schmeling¹⁵, Charlotte G. Cook¹, Lucy R. Marshall^{1,2,3}, Meredyth G. Ll. Wilkinson^{1,2,3}, Stephen Eyre^{4,5}, Soumya Raychaudhuri^{4,5,16,17}, Lucy R. Wedderburn^{1,2,3} and on behalf of the Juvenile Dermatomyositis Cohort and Biomarker Study, the Childhood Myositis Heterogeneity Study Group, and the Myositis Genetics Consortium (MYOGEN)

¹Infection, Immunity and Inflammation Research and Teaching Department, UCL Great Ormond Street Institute of Child Health, London, UK

²Centre for Adolescent Rheumatology Versus Arthritis at UCL, UCL Hospital and Great Ormond Street Hospital, London, UK

³NIHR Biomedical Research Centre at Great Ormond Street Hospital, London, UK

⁴Centre for Genetics and Genomics Versus Arthritis, Centre for Musculoskeletal Research, Manchester Academic Health Science Centre, The University of Manchester, Manchester, UK

⁵National Institute of Health Research Manchester Biomedical Research Centre, Manchester Academic Health Science Centre, Manchester University NHS Foundation Trust, Manchester, UK

⁶Environmental Autoimmunity Group, Clinical Research Branch, National Institute of Environmental Health Sciences, National Institutes of Health, Bethesda, MD, USA

⁷Ann & Robert H. Lurie Children's Hospital of Chicago, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

⁸Department of Rheumatology, University of Oslo, Oslo, Norway

⁹Oslo New University College, Oslo, Norway

¹⁰Emory University School of Medicine, Atlanta, GA, USA

¹¹Division of Rheumatology, George Washington University School of Medicine and Health Sciences, Washington, DC, USA

¹²Division of Rheumatology, Department of Pediatrics, The Hospital for Sick Children, Toronto, ON, Canada

¹³IWK Health Centre and Dalhousie University, Halifax, NS, Canada

¹⁴Pediatrics, Duke University, Durham, NC, USA

¹⁵Department of Pediatrics, Alberta Children's Hospital, University of Calgary, Calgary, AB, Canada

¹⁶Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

¹⁷Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

*To whom correspondence should be addressed at: Infection, Immunity and Inflammation Research and Teaching Department, UCL Great Ormond Street Institute of Child Health, 30 Guilford Street, London WC1N 1EH, UK. Tel: +442079052671; Email: c.deakin@ucl.ac.uk

†The authors wish it to be known that, in their opinion, the final two authors should be regarded as joint Senior Authors.

Abstract

Juvenile dermatomyositis (JDM) is a rare, severe autoimmune disease and the most common idiopathic inflammatory myopathy of children. JDM and adult-onset dermatomyositis (DM) have similar clinical, biological and serological features, although these features differ in prevalence between childhood-onset and adult-onset disease, suggesting that age of disease onset may influence pathogenesis. Therefore, a JDM-focused genetic analysis was performed using the largest collection of JDM samples to date. Caucasian JDM samples ($n = 952$) obtained via international collaboration were genotyped using the Illumina HumanCoreExome chip. Additional non-assayed human leukocyte antigen (HLA) loci and genome-wide single-nucleotide polymorphisms (SNPs) were imputed. HLA-DRB1*03:01 was confirmed as the classical HLA allele most strongly associated with JDM [odds ratio (OR) 1.66; 95% confidence interval (CI) 1.46, 1.89; $P = 1.4 \times 10^{-14}$], with an independent association at HLA-C*02:02 (OR = 1.74; 95% CI 1.42, 2.13, $P = 7.13 \times 10^{-8}$). Analyses of amino acid positions within HLA-DRB1 indicated that the strongest association was at position 37 (omnibus $P = 3.3 \times 10^{-19}$), with suggestive evidence this association was independent of position 74 (omnibus $P = 5.1 \times 10^{-5}$), the position most strongly associated with adult-onset DM. Conditional analyses also suggested that the association at position 37 of HLA-DRB1 was independent of some alleles of the Caucasian HLA 8.1 ancestral haplotype (AH8.1) such as HLA-DQB1*02:01 (OR = 1.62; 95% CI 1.36, 1.93; $P = 8.70 \times 10^{-8}$), but not HLA-DRB1*03:01 (OR = 1.49; 95% CR 1.24, 1.80; $P = 2.24 \times 10^{-5}$). No associations outside the HLA region were identified. Our findings confirm previous associations with AH8.1 and HLA-DRB1*03:01, HLA-C*02:02 and identify a novel association with amino acid position 37 within HLA-DRB1, which may distinguish JDM from adult DM.

Received: September 10, 2021. Revised: January 14, 2022. Accepted: January 17, 2022

© The Author(s) 2022. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Juvenile dermatomyositis (JDM) is a rare, severe autoimmune disease and the most prevalent idiopathic inflammatory myopathy with proximal muscle weakness and skin rash as typical features. Clinical features are heterogeneous and can include serious complications such as calcinosis, ulceration, treatment-resistant rash and involvement of major organs, including gut, lungs and brain. Although some patients achieve remission following standard disease management, which consists of long-term immunosuppression with glucocorticoids, methotrexate and other medications, others respond poorly.

While JDM and adult-onset dermatomyositis (DM) share similar clinical and biological features, there are differences in prevalence of clinical features (1). The incidence of JDM is approximately one-tenth of the incidence of DM (2). DM can be associated with cancer, but this has not been reported in JDM. Conversely, calcinosis is a major cause of morbidity in JDM but has a lower prevalence in DM. The prevalence of myositis-specific autoantibodies (MSAs), which are linked to different clinical features of disease, also differs between the adult and juvenile forms of the disease. Anti-nuclear matrix protein-2 is one of the more abundant MSAs in JDM (reported in 20–25% of patients (3–5)) but has a lower prevalence in DM (reported in 1.6–17% of different patient populations) (6–8). The most prevalent MSA in DM, anti-histidyl tRNA synthetase (anti-Jo-1), is rare in JDM. These differences in the distribution of MSA and clinical features suggest an influential role for age of disease onset in the pathogenesis of disease. However, little is known at the mechanistic level about the influence of age on JDM phenotypes and pathogenesis. Knowledge about how disease mechanisms differ between patient subgroups and interact with patient age to result in different complications may enable targeting of novel molecular pathways, more accurate modelling of life-long risk and more stratified therapeutic approaches to address this risk.

Candidate gene and genome-wide studies of myositis have established the strongest genetic association within the Caucasian 8.1 ancestral haplotype (AH8.1; HLA A1-B8-DR3-DQ2) of the major histocompatibility complex (MHC), also associated with many other immune-mediated diseases (9–13). Distinct human leukocyte antigen (HLA) alleles have been identified as associated with serological subphenotypes of myositis in different ethnic populations. Most notably associations between the development of anti-Jo-1 autoantibodies and HLA-DRB1*03:01, HLA-DQB1*02:01 and HLA-B*08, consistent with AH8.1, have been identified in Caucasian and African-American patients (11,12,14). In adult myositis, gene–environment interactions have been found between HLA-DRB1*03, smoking and the presence of anti-Jo-1 autoantibodies, and between HLA-DRB1*11:01 and anti-3-hydroxy-3-methylglutaryl-CoA reductase (anti-HMGCR)-positive statin-induced immune-mediated

necrotising myopathy (15,16). To date, the rarity of JDM has meant that candidate gene studies in JDM have been small and subgroup analyses of JDM in genome-wide studies have had limited statistical power relative to other myositis phenotypes. The aim of this research was to identify novel genetic loci associated with JDM using a larger cohort of patients.

Results

Samples and genotyping quality control

Samples of Caucasian ancestry ($n=952$) were obtained via international collaboration including samples from the UK Juvenile Dermatomyositis Cohort & Biomarker Study, the Childhood Myositis Heterogeneity Study Group and the Myositis Genetics Consortium (Table 1). Many of these cases have contributed to previous analyses (10,17). Demographic features are described in Supplementary Material, Table S1. After genotyping and stringent quality control (QC), $n=178\,164$ single-nucleotide polymorphisms (SNPs) remained (Supplementary Material, Table S2) on $n=851$ JDM samples (Supplementary Material, Tables S3 and S4). The proportion of phenotypic variance explained by these markers was estimated as 0.18 (0.02).

HLA-DRB1*03:01 confirmed as allele most strongly associated with JDM

Case–control analysis of assayed SNPs confirmed that the strongest association with JDM was within the HLA region (Fig. 1; Supplementary Material, Fig. S1). Analysis of imputed markers within the HLA region indicated that the strongest association was with SNP rs3117103 [odds ratio (OR)=1.87; 95% confidence interval (CI) 1.64, 2.13; $P=1.79 \times 10^{-20}$] and the classical allele HLA-DRB1*03:01, consistent with previous reports (9,10). The OR for HLA-DRB1*03:01 itself was 1.66 (95% CI 1.46, 1.89; $P=1.4 \times 10^{-14}$; Supplementary Material, Table S5).

Conditioning on this allele, an independent association with HLA-C*02:02 was identified (OR=1.74; 95% CI 1.42, 2.13, $P=7.13 \times 10^{-8}$) as reported previously (10). The next most associated allele was HLA-B*44:02, but the P -value for this association was above the threshold for statistical significance (OR=1.31, 95% CI 1.15, 1.50; $P=6.04 \times 10^{-5}$). No alleles within HLA-A were associated with JDM (all P -values above 0.08).

Analysis of amino acid positions within HLA-DRB1 identifies position 37 as the most significantly associated with JDM

Since the strongest association with JDM was identified within HLA-DRB1, further analysis sought to resolve this association at the level of amino acid positions. Position 37 was the most strongly associated with disease (omnibus $P=3.3 \times 10^{-19}$; Fig. 2 and Table 2). Residues at position 37 are located within the P9 pocket of the antigen-binding groove (Supplementary Material, Fig. S2) (18,19). Relative to Tyr-37, Ser-37 was the most significantly associated residue at this position (OR=0.65; 95% CI 0.57, 0.75; $P=7.34 \times 10^{-10}$), followed

Table 1. Sources of samples from Caucasian patients with JDM

Country	Source	Genotyped samples (n = 952)	Genotyped samples after quality control (n = 851)
UK	UK Juvenile Dermatomyositis Cohort & Biomarker Study	365	326
USA	National Institute of Environmental Health Sciences	262	224
	Northwestern University	140	116
	Emory University	41	31
	George Washington University	32	30
	Mayo Clinic	13	12
Canada	The Hospital for Sick Children, Toronto	24	24
	IWK Health Centre	16	16
	Alberta University	10	9
Norway	Oslo University Hospital	49	46




Figure 1. Manhattan plot of the association of assayed SNPs with JDM. SNPs ($n = 178\,164$) were available for $n = 851$ JDM samples and $n = 12\,232$ controls of Caucasian origin. Association was tested using logistic regression, with the first 10 principal components included as covariates to account for population stratification. The red dotted line indicates genome-wide level of significance (5×10^{-8}).

by Phe-37 (OR=0.63; 95% CI 0.54, 0.75; $P = 1.14 \times 10^{-7}$) (Table 3). Asn-37, which is found on HLA-DRB1*03:01, was not associated relative to Tyr-37 (OR=1.13; 95% CI 1.06, 1.19; $P = 0.05$). Conditioning on position 37, there was suggestive evidence of possible independent associations at positions 74 ($P = 5.1 \times 10^{-5}$) and 26 ($P = 5.9 \times 10^{-5}$), although the P -values for these associations were not significant (Table 2).

To evaluate whether position 37 explained the association within HLA-DRB1 more convincingly than the classical HLA-DRB1*03:01 allele and AH8.1, conditional analyses were performed. Conditioning on HLA-DRB1*03:01, the association of position 37 was above the threshold for significance (omnibus $P = 4.42 \times 10^{-5}$), although there may be weak evidence of an independent association. After conditioning on all residues at position 37, there was evidence of independent effects for

HLA-DQB1*02:01 within AH8.1 (OR=1.62; 95% CI 1.36, 1.93; $P = 8.70 \times 10^{-8}$) and HLA-C*02:02 (OR=1.72; 95% CI 1.40, 2.10; $P = 1.58 \times 10^{-7}$). However, the effect for HLA-DRB1*03:01 did not meet the threshold for significance (OR=1.49; 95% CR 1.24, 1.80; $P = 2.24 \times 10^{-5}$). Taken together, these results indicate an effect of position 37 that is independent of some of the previously reported effects within AH8.1 and at HLA-C*02:02, but that is not independent of HLA-DRB1*03:01.

Genome-wide imputation identifies possible loci associated at a suggestive level of significance

Following genome-wide imputation, there were no additional loci identified at the genome-wide level of statistical significance ($P = 5 \times 10^{-8}$). Two loci with $P < 1 \times 10^{-6}$ and minor allele frequency (MAF) over 0.05 were identified (Table 4; Fig. 3). rs6501160 is an intronic




Figure 2. Representation of the P-values for omnibus tests of individual amino acid positions within *HLA-DRB1* for association with JDM. Blue circles represent P-values for amino acid positions tested alone. Green circles represent P-values for amino acid positions after conditioning on position 37. Red circles represent P-values for 4-digit classical alleles. The grey dashed line represents the threshold for statistical significance ($P = 6.8 \times 10^{-6}$).

Table 2. Analysis of amino acid positions within *HLA-DRB1* associated with JDM^a

Position	χ^2 ^b	P-value	Effects independent of position 37	
			χ^2	P-value
37	89.2	3.3×10^{-19}	–	–
74	76.7	1.6×10^{-16}	22.5	5.1×10^{-05}
26	59.5	1.2×10^{-13}	19.5	5.9×10^{-05}
10	47.5	5.4×10^{-12}		
67	46.0	1.2×10^{-11}		
–17	41.0	1.5×10^{-10}		
–16	37.5	9.1×10^{-10}		
30	39.5	2.6×10^{-9}		
–25	34.4	4.5×10^{-9}		
11	40.2	9.9×10^{-9}		
13	40.0	1.1×10^{-08}		
233	32.3	1.3×10^{-8}		
71	27.8	1.4×10^{-7}		
–1	26.1	3.2×10^{-7}		
181	28.9	5.3×10^{-7}		
57	24.2	8.8×10^{-7}		
60	25.4	3.1×10^{-6}		

^aAlpha threshold for statistical significance in the MHC region taken as 6.8×10^{-6} (47). ^bTest statistics compared using a chi-squared distribution.

variant within *transmembrane protein 114* (TMEM114), a transmembrane protein. rs6892006 is an intronic variant within *myocyte enhancer factor 2C antisense RNA 1* (MEF2C-AS1), an anti-sense RNA gene (Supplementary Material, Fig. S3). Associations with clinical phenotypes have not been reported for either variant. Although additional loci

were identified at $P < 5 \times 10^{-8}$ (Supplementary Material, Table S6 and Supplementary Material, Fig. S4), these had MAF between 0.01 and 0.05 and may represent imputation artefacts.

Discussion

This analysis represents the largest international genetic study of JDM to date. We found that the strongest association was in the HLA region within *HLA-DRB1*03:01*, consistent with previous studies (9,10,17). We confirmed an independent association with *HLA-C*02:02* and found some evidence of a possible additional independent association at *HLA-B*44:02*, which would need to be confirmed in a future study. Interestingly, we did not find evidence of an association with *HLA-A*, even though associations within this gene are well known for multiple other autoimmune diseases including rheumatoid arthritis (RA), juvenile idiopathic arthritis (JIA), psoriatic arthritis and type I diabetes (20–23).

In this analysis focused on juvenile-onset disease, we found amino acid position 37 within *HLA-DRB1* had the strongest association with JDM, with Ser-37 and Phe-37 having protective effects relative to Tyr-37. We also found evidence that the association with position 37 was independent of AH8.1, a well-established association with myositis, but there was only weak evidence of independence from *HLA-DRB1*03:01*, and so this needs to be followed up in future studies with greater numbers of both juvenile-onset and adult-onset patients using the

Table 3. Association of residues within position 37 with JDM, with effects of classical HLA alleles linked to these residues^{a,b}

Residue	Classical HLA allele linked to residue	Allele frequency in JDM	Allele frequency in controls	OR	95% CI	P-value
Tyr-37	–	0.37	0.31	–	–	–
Ser-37	–	0.21	0.26	0.65	0.57, 0.75	7.3×10^{-10}
–	HLA-DRB1*01:01	0.10	0.12	0.76	0.65, 1.46	1.7×10^{-3}
–	HLA-DRB1*15:01	0.11	0.14	0.66	0.57, 0.78	6.5×10^{-7}
Phe-37	–	0.11	0.15	0.63	0.54, 0.75	1.1×10^{-7}
–	HLA-DRB1*07:01	0.09	0.13	0.67	0.57, 0.79	2.8×10^{-6}
–	HLA-DRB1*14:01	0.02	0.02	0.86	0.60, 1.24	0.42
Asn-37	–	0.31	0.24	1.13	1.06, 1.19	0.05
–	HLA-DRB1*03:01	0.19	0.12	1.66	1.46, 1.89	1.4×10^{-14}

^aSummary statistics for individual residues derived from the omnibus model for position 37, with Tyr-37 used as the reference residue. Summary statistics for classical four-digit HLA alleles derived from case-control analysis of imputed HLA data. Alleles linked to these residues at position 37 but not represented were filtered out during QC. ^bAlpha threshold for statistical significance in the MHC region taken as 6.8×10^{-6} (47).

Table 4. Genome-wide imputed loci displaying potential association with JDM, with P-value cut-offs of 1×10^{-6} and MAF cut-off of 0.01^a

Rsid ^b	Chromosome	Position	Allele A	Allele B	MAF	OR	95% CI	P-value	Imputation R ² ^c	Information measure ^d	Association information measure ^e
rs6501160	16	8 607 139	G	A	0.27	0.72	0.65, 0.81	5.8×10^{-8}	0.91	0.91	0.91
rs6892006	5	88 450 541	T	G	0.13	0.68	0.58, 0.79	8.7×10^{-7}	0.87	0.87	0.89

^aLoci with P-values below a suggestive level of significance (1×10^{-6}) are displayed for MAF cut-offs of 0.05 and 0.01; only additional loci captured by the lower MAF cut-off are shown for that cut-off. ^bRsid, Reference SNP cluster ID. There were no assayed markers within 100 000 kb of either of these loci. ^cSquared correlation of imputation of genotypes with true unmeasured genotypes, as estimated by Minimac3 during imputation via the Michigan Imputation Server: https://genome.sph.umich.edu/wiki/Minimac3_Info_File#Rsq. ^dImpute INFO measure calculated by SNPTEST: https://mathgen.stats.ox.ac.uk/genetics_software/snpctest/snpctest.html#info_measures. ^eRelative information measure about parameters of the model fitted during association testing: https://mathgen.stats.ox.ac.uk/genetics_software/snpctest/snpctest.v2.p.




Figure 3. Manhattan plot of the association of imputed SNPs with JDM. Loci were filtered according to MAF of 0.05. Data were imputed for $n = 851$ JDM samples and $n = 12\,232$ controls of Caucasian origin. The first 10 principal components were adjusted for during analysis to account for population stratification. SNPs were weighted by the information score to account for imputation uncertainty. The red dotted line indicates genome-wide level of significance (5×10^{-8}). The degree of transparency of each data-point represents the R² value for imputation accuracy, with more solid colours representing higher certainty.

same genotyping chip. Amino acid position 37 is within the P9 pocket of the antigen-binding groove (18,19). Substitution of Ser-37 to Tyr-37 has been shown to be sufficient to alter alloantigenicity and stimulate a T-cell response (24). Tyr-37 promotes a stronger response to streptococcal protein (25).

The previous ImmunoChip analysis of adult and juvenile myositis combined identified the strongest association within HLA-DRB1 at amino acid position 74 (10). Subsequent analyses of that dataset identified position 74 as having an association with anti-Jo1, anti-PM/Scl and Anti-cytosolic 5'-nucleotidase 1A (anti-cN1A)

autoantibodies (26), which are less prevalent in juvenile-onset disease. We found weak evidence of independent association at position 74 after conditioning on position 37, above the threshold for significance and so this finding needs to be followed up with greater number of patients. Nonetheless, it raises an intriguing possibility that children and adults may have different dominant autoantigenic peptides being presented in the antigen-binding groove, although we recognize that multiple antigens with different allelic associations are likely to be involved in disease. While the Immunochip analyses benefitted from a relatively large cohort ($n=2544$), the number of JDM patients was approximately half that of our analysis ($n=493$). These patients are represented in our analysis, along with additional cases. It may be that the effects of juvenile-onset disease were obscured by larger number of adult-onset patients and other phenotypes present in the combined cohort. Future work and greater numbers are required to better dissect the genetic influence on age effects in juvenile and adult-onset disease, as well as on clinical and serological heterogeneity.

Associations between other positions within *HLA-DRB1* and autoimmune disease have been reported. For example, position 11 has the strongest effect in RA, with independent effects at positions 71 and 74 (27). In JIA, position 13 has the strongest effect, although in systemic JIA the strongest effect is at position 58 (21,28). Positions 71/74 have also been implicated in anti-fibrillarin-positive systemic sclerosis, Crohn's disease, multiple sclerosis, type I diabetes and Grave's disease (29–33). In systemic lupus erythematosus (SLE), Ser-1, Phe47 and Ala71 are associated with disease (34). Position 37 has been found to be associated with primary sclerosing cholangitis, ulcerative colitis in Asians, ACPA⁺ RA in Han Chinese, SLE in Asians and psoriasis vulgaris in Taiwanese (18,19,35–37). Interestingly, a recent report of Japanese patients with RA identified an association between position 37 and younger age of disease onset (defined as 16–30 years of age), but not with older age of onset (defined as over 60 years of age) (38). At present, it is not well understood how these differing positional effects relate to the spectrum of phenotypes represented in autoimmune disease, and how these effects interact with ethnicity and age of disease onset.

The major limitation of our study is limited statistical power for identifying associations outside the HLA region, which is a practical challenge for rare disease research. Although our combined cohort represents the largest-ever assembled international cohort of Caucasian patients with JDM, it is a relatively small sample size for a genetic study. As such, the findings of suggestive associations will need to be confirmed in future studies with greater number of patients. Nonetheless, we were able to confirm the findings of previous studies in a larger cohort dedicated to JDM. Future studies with greater numbers may better define the relationship between the associations at *HLA-DRB1*03:01* and amino

acid positions within *HLA-DRB1* such as position 37. Loci outside the HLA region identified at genome-wide and suggestive levels of significance in previous analyses (including *PTPN22*, *UBE2L3*, *CD28*, *TRAF6*, *STAT4*) were not replicated in this study, although signals at these loci were not specifically tested for. This may reflect the smaller sample size of this study and the dominance of adult patients in those analyses whose characteristics may differ from paediatric disease (10). JDM is a heterogeneous phenotype; however, the small sample size restricted our ability to analyse more clinically homogeneous subgroups, such as autoantibody subgroups. It may be that future developments in methodologies for genetic analysis will enable further insights to be derived from this dataset. Many of the patients in this study overlapped with a previous analysis, which identified associations with major MSA subtypes, including anti-Jo-1, anti-PM/Scl, anti-cN1A, anti-Mi-2 and anti-TIF1 γ (26). As more patients become available for inclusion in genetic studies, analyses of further MSA subtypes may become possible. It will also be critical to study genetic associations with MSA subtypes in different ethnic populations.

Analysis following genome-wide imputation identified 2 possible suggestive loci, an intronic variant within *TMEM114* and an intronic variant within *MEF2C-AS1*. *TMEM114* is a glycosylated transmembrane protein, and knowledge of its cellular function is limited, although missense mutations in this gene and a chromosomal translocation in its promoter are associated with congenital and juvenile cataract disorders, respectively (39). *MEF2C-AS1* is a non-coding anti-sense RNA gene with no known function. It is unclear how these loci relate to JDM, but it may be these alleles function as epigenetic marks.

In summary, we have confirmed the association between JDM and *HLA-DRB1*03:01* and shown that within *HLA-DRB1*, position 37 is most strongly associated with disease in a population of patients with juvenile-onset myositis.

Materials and Methods

Genotyping and genotype calling

Genotyping of Caucasian JDM samples ($n=952$) was performed using the Illumina (Cambridge, UK) HumanCore-Exome chip across three batches at a single centre (University College London). This cohort included the majority of JDM cases included in previous analyses, as well as additional cases recruited subsequent to those analyses including cases from more centres (10,17). GenomeStudio 2.0 (version 2.0.4 of the Genotyping Module and the GenTrain 3.0 Cluster Algorithm) was used for genotype clustering and calling for each separate batch. Samples with less than 90% call rate were excluded, and genotype clustering and calling were repeated, before data were exported in PLINK format for QC.

Data on healthy individuals ($n=12\,474$) of European ancestry who had also been genotyped using the

HumanCoreExome chip as part of the International Age-Related Macular Degeneration Genetic Consortium were obtained from dbGaP (dbGAP Study Accession phs001039.v1.p1).

Quality control

QC of markers and samples was performed as described previously (22). The following steps were done separately for each of the three batches of JDM data and the control data using PLINK 1.07 (40). Mitochondrial and Y chromosome SNPs were excluded. SNPs were also excluded if they had elevated missing rates (over 2%), had a MAF below 1% and deviated significantly from Hardy–Weinberg equilibrium (HWE; threshold of $P < 0.0001$). Samples with elevated missing genotypes (over 5%) and outlying heterozygosity rates (above or below 5 standard deviations from the mean rate). Data were aligned to the Haplotype Reference Consortium (HRC) before merging for further QC using the HRC checking tool (41).

Duplicated or related individuals in the merged dataset were identified using identity-by-descent, performed on a linkage disequilibrium (LD)-pruned dataset of 65 862 SNPs with MAF over 5%. A PI_HAT threshold of 0.2 was used, with the individual with the most missing data excluded. Principal component analysis was performed using PLINK 1.9 (42), to evaluate population stratification in the LD-pruned data merged with the International HapMap 3 data (43). Cases and controls were retained if they were within 10 standard deviations of the mean value for the first two principal components (PCs) for the HapMap CEU population (Supplementary Material, Fig. S5).

Following QC, there were 178 164 SNPs for $n = 851$ JDM cases and $n = 12 232$ controls (Supplementary Material, Tables S1–S4).

Analysis of assayed markers

Autosomal markers were analysed for association with JDM using logistic regression in PLINK with adjustment for the first 10 PCs to control for population stratification. Manhattan plots were generated using R version 6.3 and the 'ggplot2' package (version 3.3.2). The proportion of phenotypic variance explained by the SNPs and a standard error for that estimated proportion were estimated using the genome-based restricted maximum likelihood (GREML) method using GTAC version 1.93.3, assuming an approximate prevalence of 0.00004 for JDM (2,44,45).

Imputation of HLA loci

Classical HLA alleles, amino acids and SNPs within the HLA region were imputed using SNP2HLA (version 1.0.3) and the Type 1 Diabetes Genetics Consortium reference panel ($n = 5225$) (46). QC of imputed markers used the following criteria: imputation information score (R^2) over 0.9, MAF over 0.01 and significant departures from HWE in controls ($P < 0.001$). Imputed markers were coded as present or absent.

Analysis of HLA and amino acid positions

Case–control analysis of all imputed HLA was performed assuming an additive model using logistic regression in PLINK, also with adjustment for the first 10 PCs, to identify the most strongly associated locus. Clinical covariates were not adjusted for. For HLA analyses, P -values below 6.8×10^{-6} were considered statistically significant, using a Bonferroni correction for the number of imputed markers as reported previously (47). The two- and four-digit classical alleles identified by this locus were subsequently conditioned on in further logistic regression analyses to identify any independent associations.

Amino acid positions were interrogated using likelihood ratio tests (LRTs) in R as follows. At each position, logistic regression models were fitted with all residues in the model, except the residue that was most prevalent in the controls and served as the reference. The first 10 PCs were adjusted for in each of these models. To evaluate whether amino acid positions were associated with disease, LRTs were performed to compare the model that was fitted for the residues at each position against a null model, which comprised the first 10 PCs only. The effect of each amino acid position evaluated using a LRT is represented by the LRT P -value, but there is no estimated effect size for the amino acid position generated by this test. Allele frequencies, odds ratios and P -values for amino acid residues at key positions are also reported. Possible independent effects were identified by conditional analysis as above.

Genome-wide imputation and analysis

Genome-wide imputation of SNPs was performed using the Michigan Imputation Server (version 1.2.4) and the HRC reference panel (version r1.1; Supplementary Material, Fig. S6) (41,48). Imputed SNPs with MAF below 0.01 or imputation information score below 0.5 were filtered out.

Imputed SNPs were analysed using SNPTEST version 2.5.4-beta3 (49), with adjustment for the first 10 PCs and weighting for the imputation information score to account for imputation uncertainty. Since imputation artefacts are enriched in rare variants, a stringent MAF threshold of 0.05 was used during analysis, although loci with MAF 0.01–0.05 are also reported in Supplementary Material.

Supplementary Material

Supplementary Material is available at HMG online.

Acknowledgements

C.T.D. would like to thank the study administrators and research assistants at each international centre who assisted with extraction of clinical data and preparation and shipment of DNA samples, including Dr Terry O'Hanlon and Payam Noroozi Farhadi (NIH), Siri Tennebø Flåm (University of Oslo); Gabrielle Morgan (Northwestern University); Dwaraka Veeramreddy

(Alberta Children's Hospital); Hayyah Clairman (Hospital for Sick Children, Toronto). C.T.D. would also like to thank Dr Socrates Varakliotis (Centre for Adolescent Rheumatology Versus Arthritis at UCL, UCLH and GOSH) for setting up remote access to a Linux desktop computer which enabled analyses to continue during the COVID-19 pandemic. C.T.D. thanks Drs Michael Ombrello and Elaine Remmers for useful comments on the manuscript.

The Juvenile Dermatomyositis Research Group would like to thank all of the patients and their families who contributed to the Juvenile Dermatomyositis Cohort & Biomarker Study & Repository. We thank all local research coordinators and principal investigators who have made this research possible. The JDRG members were as follows:

Dr Kate Armon, Mr Joe Ellis-Gage, Ms Holly Roper, Ms Vanja Briggs and Ms Joanna Watts (Norfolk and Norwich University Hospitals); Dr Liza McCann, Mr Ian Roberts, Dr Eileen Baildam, Ms Louise Hanna, Ms Olivia Lloyd, Susan Wadeson and Michelle Andrews (The Royal Liverpool Children's Hospital, Alder Hey, Liverpool); Dr Phil Riley, Ms Ann McGovern, Verna Cuthbert (Royal Manchester Children's Hospital, Manchester); Dr Clive Ryder, Mrs Janis Scott, Mrs Beverley Thomas, Professor Taunton Southwood, Dr Eslam Al-Abadi and Ruth Howman (Birmingham Children's Hospital, Birmingham); Dr Sue Wyatt, Mrs Gillian Jackson, Dr Mark Wood, Dr Tania Amin, Dr Vanessa VanRooyen and Ms Deborah Burton, Louise Turner & Sarah Hanson (Leeds General Infirmary, Leeds); Dr Joyce Davidson, Dr Janet Gardner-Medwin, Dr Neil Martin, Ms Sue Ferguson, Ms Liz Waxman and Mr Michael Browne, Ms Roisin Boyle, Ms Emily Blyth (The Royal Hospital for Sick Children, Yorkhill, Glasgow); Dr Mark Friswell, Professor Helen Foster, Mrs Alison Swift, Dr Sharmila Jandial, Ms Vicky Stevenson, Ms Debbie Wade, Dr Ethan Sen, Dr Eve Smith, Ms Lisa Qiao, Mr Stuart Watson and Ms Claire Duong (Great North Children's Hospital, Newcastle); Dr Helen Venning, Dr Rangaraj Satyapal, Mrs Elizabeth Stretton, Ms Mary Jordan, Dr Ellen Mosley, Ms Anna Frost, Ms Lindsay Crate, Dr Kishore Warriar, Stefanie Stafford, Kelly Sandhu & Tracey Dandy (Queens Medical Centre, Nottingham); Professor Lucy Wedderburn, Dr Clarissa Pilkington, Dr Nathan Hasson, Dr Muthana Al-Obadi, Dr Giulia Varnier, Dr Sandrine Lacassagne, Mrs Sue Maillard, Mrs Lauren Stone, Ms Elizabeth Halkon, Ms Virginia Brown, Ms Audrey Juggins, Dr Sally Smith, Mrs Sian Lunt, Ms Elli Enayat, Mrs Hemlata Varsani, Miss Laura Kassoumeri, Miss Laura Beard, Miss Katie Arnold, Mrs Yvonne Glackin, Ms Stephanie Simou, Dr Beverley Almeida, Dr Kiran Nistala, Dr Raquel Marques, Dr Claire Deakin, Dr Parichat Khaosut, Ms Stefanie Dowle, Dr Charalampia Papadopoulou, Dr Shireena Yasin, Dr Christina Boros, Dr Meredyth Wilkinson, Dr Chris Piper, Mrs Cerise Johnson-Moore, Ms Lucy Marshall, Ms Kathryn O'Brien, Ms Emily Robinson (Great Ormond Street Hospital, London); Dr Kevin Murray (Princess Margaret Hospital, Perth, Western Australia); Dr Coziana Ciurtin, Dr John Ioannou, Mrs

Caitlin Clifford and Ms Linda Suffield (University College London Hospital, London); Ms Helen Lee, Ms Sam Leach, Ms Helen Smith, Dr Anne-Marie McMahon, Ms Heather Chisem, Jeanette Hall, Ruth Kingshott and Maxine Mutten (Sheffield's Children's Hospital, Sheffield); Dr Nick Wilkinson, Ms Emma Inness, Ms Eunice Kendall, Mr David Mayers, Ruth Etherton, Danielle Miller and Dr Kathryn Bailey (Oxford University Hospitals, Oxford); Dr Jacqui Clinch, Ms Natalie Fineman, Ms Helen Pluess-Hall and Suzanne Sketchley (Bristol Royal Hospital for Children, Bristol); Dr Joyce Davidson, Margaret Connon and Ms Lindsay Vallance (Royal Aberdeen Children's Hospital); Dr Kirsty Haslam, Charlene Bass-Woodcock, Trudy Booth and Ms Louise Akeroyd (Bradford Teaching Hospitals); Dr Alice Leahy, Amy Collier, Rebecca Cutts, Emma Macleod, Dr Hans De Graaf, Dr Brian Davidson, Sarah Hartfree, Danny Pratt, Elizabeth Fofana and Lorena Caruana (University Hospital Southampton) and all the Children, Young people and their families who have contributed to this research.

The Childhood Myositis Heterogeneity Study Group is supported by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences. The Childhood Myositis Heterogeneity Study Group thanks the following contributors to this study:

Daniel Albert (Dartmouth-Hitchcock Medical Center), Kathy Amoroso (Carilion Roanoke Memorial Hospital), Bitu Arabshahi (Fairfax Inova Medical Center), Edsel Arce-Hernandez (Valley Children's Healthcare), Alan Baer (Johns Hopkins University School of Medicine), Susan Ballinger (Riley Children's Hospital, Indiana University), Lilliana Barillas (Albany Medical Center), Mara Becker (Children's Mercy, Kansas City), Catherine April Bingham (Penn State Children's Hospital), John Bohnsack (University of Utah), Jon M Burnham (Children's Hospital of Philadelphia), Ruy Carrasco (Child Neurology Consultants of Austin), Victoria Cartwright (Legacy Emanuel Medical Center), Gail Cawkwell (Johns Hopkins All Children's Hospital), Wendy De La Pena (Loma Linda University), Fatma Dedeoglu (Boston Children's Hospital), Barbara Anne Eberhard (Northwell Health), Kaleo Ede (Phoenix Children's Hospital), Barbara Edelheit (University of Connecticut), Andrew Eichenfield (New York-Presbyterian Hospital), Terri Finkel (University of Colorado), Robert Fuhlbrigge (Boston Children's Hospital), Abraham Gedalia (Children's Hospital New Orleans), Stephen George (University of Maryland), Harry Gewanter (Virginia Commonwealth University), Ellen Goldmuntz (Children's National Medical Center), Donald Goldsmith (St. Christopher's Hospital for Children, Drexel University), Beth Gottlieb (Northwell Health), Brent Graham (Vanderbilt University), Thomas Griffin (Carolinas Medical Center), Hilary Haftel (University of Michigan), Melissa Hawkins-Holt (University of Maryland), Theresa Hennon (John R. Oishei Children's Hospital, University of Buffalo), Michael Henrickson (Cincinnati Children's Hospital), Aimee Hersh (University of Utah), Gloria Higgins (Nationwide Children's Hospital),

Lisa Imundo (Columbia University), Christi Inman (University of Utah), Anna Jansen (NIEHS, NIH), Rita Jerath (Medical College of Georgia), Olcay Y Jones (Walter Reed National Military Medical Center), Lawrence Jung (Children's National Medical Center), Ildy Katona (Uniformed University of the Health Sciences), Philip Kahn (New York University), Hanna Kim (NIAMS, NIH), Susan Kim (University of California, San Francisco), Yukiko Kimura (Hackensack Meridian School of Medicine), Daniel Kingsbury (Legacy Emanuel Medical Center), W Patrick Knibbe (St Luke's Clinic), Bianca Lang (Dalhousie University), Melissa Lerman (Children's Hospital of Philadelphia), Carol Lindsley (University of Kansas), Daniel Lovell (Cincinnati Children's Hospital), Azadeh Majlessi (Loma Linda University), Gulnara Mamyrova (George Washington University), Paul McCarthy (Yale University), Diana Milojevic (Johns Hopkins All Children's Hospital), Stephen R. Mitchell (Georgetown University), Renee Modica (Arnold Palmer Hospital for Children, University of Florida), Chihiro Morishima (University of Washington), Simona Nativ (Atlantic Health System), Edward Oberle (Nationwide Children's Hospital), Kiem Oen (University of Manitoba), Judyann Olson (Children's Wisconsin, Medical College of Wisconsin), Elif Oral (University of Michigan), Barbara Ostrov (Penn State Children's Hospital), Lauren M Pachman (Lurie Children's Hospital, Northwestern University), Murray Passo (The Medical University of South Carolina), Maria Perez (Cook Children's Medical Center), Marilynn Punaro (University of Texas at Dallas), C. Eglia Rabinovich (Duke University), Linda Ray (University of Mississippi), Robert Rennebohm (Cleveland Clinic), Rafael Rivas-Chacon (Nicklaus Children's Hospital, University of Miami), Tova Ronis (Children's National Hospital), Margalit Rosenkranz (University of Pittsburgh Medical Center), Deborah Rothman (Albany Medical College), Sara Sabbagh (NIAMS, NIH), Adam Schiffenbauer (NIEHS), Bracha Shaham (Children's Hospital of Los Angeles, University of Southern California), Robert Sheets (Rady Children's Hospital, University of California at San Diego), Susan Shenoi (Seattle Children's, University of Washington), David Sherry (Seattle Children's, University of Washington), Abigail Smukler (The Toledo Clinic), Jennifer Soep (University of Colorado), Sangeeta Sule (Children's National Medical Center), Robert Sundel (Boston Children's Hospital), Stacey Tarvin (Riley Hospital for Children, University of Indiana), Mohamad Tolaymat (University of Florida), Cagri Toruner (Texas Children's Hospital), Rita Volochayev (NIEHS, NIH), Jennifer Wargula (Seattle Children's), Peter Weiser (University of Alabama), Andrew White (Washington University), Patience White (George Washington University), Andrew Zeft (Cleveland Clinic), Lawrence Zemel (Connecticut Children's Medical Center).

Conflict of Interest statement. The authors have declared that no conflict of interest exists.

Funding

TMA Fellowship and a Cure JM grant (to C.T.D.). UK JDM Cohort and Biomarker study has been provided by generous grants from the Wellcome Trust UK (085860); Action Medical Research UK (SP4252); the Myositis Support Group UK, Arthritis Research UK now Versus Arthritis (14518, 20164, 21593); the Henry Smith Charity and Great Ormond Street Children's Charity (V1268); Tiny Hearts Society, the Myositis Association, Cure JM, the Medical Research Council (MR/N003322/1) and the National Institute for Health Research (NIHR) via the Translational Research Collaboration (TRC) Rare Diseases and the NIHR-Biomedical Research Centre at GOSH. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

References

1. Tansley, S. and Wedderburn, L.R. (2015) Comparing and contrasting clinical and serological features of juvenile and adult-onset myositis. *Curr. Opin. Rheumatol.*, **27**, 601–607.
2. Meyer, A., Meyer, N., Schaeffer, M., Gottenberg, J.-E., Geny, B. and Sibilia, J. (2015) Incidence and prevalence of inflammatory myopathies: a systematic review. *Rheumatology*, **54**, 50–63.
3. Tansley, S.L., Betteridge, Z.E., Shaddick, G., Gunawardena, H., Arnold, K., Wedderburn, L.R. and McHugh, N.J. (2014) Calcinosis in juvenile dermatomyositis is influenced by both anti-NXP2 autoantibody status and age at disease onset. *Rheumatology (Oxford)*, **53**, 2204–2208.
4. Rider, L.G., Shah, M., Mamyrova, G., Huber, A.M., Rice, M.M., Targoff, I.N. and Miller, F.W. (2013) The myositis autoantibody phenotypes of the juvenile idiopathic inflammatory myopathies. *Medicine*, **92**, 223–243.
5. Espada, G., Maldonado Cocco, J.A., Fertig, N. and Oddis, C.V. (2009) Clinical and serologic characterization of an Argentine pediatric myositis cohort: identification of a novel autoantibody (anti-mj) to a 142-kda protein. *J. Rheumatol.*, **36**, 2547–2551.
6. Ichimura, Y., Matsushita, T., Hamaguchi, Y., Kaji, K., Hasegawa, M., Tanino, Y., Inokoshi, Y., Kawai, K., Kanekura, T., Habuchi, M. et al. (2012) Anti-NXP2 autoantibodies in adult patients with idiopathic inflammatory myopathies: possible association with malignancy. *Ann. Rheum. Dis.*, **71**, 710–713.
7. Fiorentino, D.F., Chung, L.S., Christopher-Stine, L., Zaba, L., Li, S., Mammen, A.L., Rosen, A. and Casciola-Rosen, L. (2013) Most patients with cancer-associated dermatomyositis have antibodies to nuclear matrix protein NXP-2 or transcription intermediary factor 1 γ . *Arthritis Rheum.*, **65**, 2954–2962.
8. Ceribelli, A., Fredi, M., Taraborelli, M., Cavazzana, I., Franceschini, F., Quinzanini, M., Tincani, A., Ross, S.J., Chan, J.Y.F., Pauley, B.A. et al. (2012) Anti-MJ/NXP-2 autoantibody specificity in a cohort of adult Italian patients with polymyositis/dermatomyositis. *Arthritis Res. Ther.*, **14**, R97.
9. Miller, F.W., Chen, W., O'Hanlon, T.P., Cooper, R.G., Vencovsky, J., Rider, L.G., Danko, K., Wedderburn, L.R., Lundberg, I.E., Pachman, L.M. et al. (2015) Genome-wide association study identifies HLA 8.1 ancestral haplotype alleles as major genetic risk factors for myositis phenotypes. *Genes Immun.*, **16**, 470–480.
10. Rothwell, S., Cooper, R.G., Lundberg, I.E., Miller, F.W., Gregersen, P.K., Bowes, J., Vencovsky, J., Danko, K., Limaye, V., Selva-O'Callaghan, A. et al. (2016) Dense genotyping of immune-related

- loci in idiopathic inflammatory myopathies confirms HLA alleles as the strongest genetic risk factor and suggests different genetic background for major clinical subgroups. *Ann. Rheum. Dis.*, **75**, 1558–1566.
11. Arnett, F.C., Targoff, I.N., Mimori, T., Goldstein, R., Warner, N.B. and Reveille, J.D. (1996) Interrelationship of major histocompatibility complex class II alleles and autoantibodies in four ethnic groups with various forms of myositis. *Arthritis Rheum.*, **39**, 1507–1518.
 12. O'Hanlon, T.P., Carrick, D.M., Targoff, I.N., Arnett, F.C., Reveille, J.D., Carrington, M., Gao, X., Oddis, C.V., Morel, P.A., Malley, J.D. et al. (2006) Immunogenetic risk and protective factors for the idiopathic inflammatory myopathies: distinct HLA-A, -B, -Cw, -DRB1, and -DQA1 allelic profiles distinguish European American patients with different myositis autoantibodies. *Medicine*, **85**, 111–127.
 13. Chinoy, H., Salway, F., Fertig, N., Shephard, N., Tait, B.D., Thomson, W., Isenberg, D.A., Oddis, C.V., Silman, A.J., Ollier, W.E.R.R. et al. (2006) In adult onset myositis, the presence of interstitial lung disease and myositis specific/associated antibodies are governed by HLA class II haplotype, rather than by myositis subtype. *Arthritis Res. Ther.*, **8**, R13.
 14. O'Hanlon, T.P., Rider, L.G., Mamuyrova, G., Targoff, I.N., Arnett, F.C., Reveille, J.D., Carrington, M., Gao, X., Oddis, C.V., Morel, P.A. et al. (2006) HLA polymorphisms in African Americans with idiopathic inflammatory myopathy: allelic profiles distinguish patients with different clinical phenotypes and myositis autoantibodies. *Arthritis Rheum.*, **54**, 3670–3681.
 15. Chinoy, H., Adimulam, S., Marriage, F., New, P., Vincze, M., Zilahi, E., Kapitány, A., Gyetvai, A., Ekholm, L., Novota, P. et al. (2012) Interaction of HLA-DRB1*03 and smoking for the development of anti-Jo-1 antibodies in adult idiopathic inflammatory myopathies: a European-wide case study. *Ann. Rheum. Dis.*, **71**, 961–965.
 16. Mammen, A.L., Gaudet, D., Brisson, D., Christopher-Stine, L., Lloyd, T.E., Leffell, M.S. and Zachary, A.A. (2012) Increased frequency of DRB1*11:01 in anti-HMG-CoA reductase-associated autoimmune myopathy. *Arthritis Care Res.*, **64**, 1233–1237.
 17. Miller, F.W., Cooper, R.G., Vencovský, J., Rider, L.G., Danko, K., Wedderburn, L.R., Lundberg, I.E., Pachman, L.M., Reed, A.M., Ytterberg, S.R. et al. (2013) Genome-wide association study of dermatomyositis reveals genetic overlap with other autoimmune disorders. *Arthritis Rheum.*, **65**, 3239–3247.
 18. Guo, J., Zhang, T., Cao, H., Li, X., Liang, H., Liu, M., Zou, Y., Zhang, Y., Wang, Y., Sun, X. et al. (2019) Sequencing of the MHC region defines HLA-DQA1 as the major genetic risk for seropositive rheumatoid arthritis in Han Chinese population. *Ann. Rheum. Dis.*, **78**, 773–780.
 19. Hov, J.R., Kosmoliaptsis, V., Traherne, J.A., Olsson, M., Boberg, K.M., Bergquist, A., Schrupf, E., Bradley, J.A., Taylor, C.J., Lie, B.A. et al. (2011) Electrostatic modifications of the human leukocyte antigen-DR P9 peptide-binding pocket and susceptibility to primary sclerosing cholangitis. *Hepatology*, **53**, 1967–1976.
 20. Han, B., Diogo, D., Eyre, S., Kallberg, H., Zhernakova, A., Bowes, J., Padyukov, L., Okada, Y., González-Gay, M.A., Rantapää-Dahlqvist, S. et al. (2014) Fine mapping seronegative and seropositive rheumatoid arthritis to shared and distinct HLA alleles by adjusting for the effects of heterogeneity. *Am. J. Hum. Genet.*, **94**, 522–532.
 21. Hinks, A., Bowes, J., Cobb, J., Ainsworth, H.C., Marion, M.C., Comeau, M.E., Sudman, M., Han, B., Becker, M.L., Bohnsack, J.F. et al. (2017) Fine-mapping the MHC locus in juvenile idiopathic arthritis (JIA) reveals genetic heterogeneity corresponding to distinct adult inflammatory arthritic diseases. *Ann. Rheum. Dis.*, **76**, 765–772.
 22. Bowes, J., Budu-Aggrey, A., Huffmeier, U., Uebe, S., Steel, K., Hebert, H.L., Wallace, C., Massey, J., Bruce, I.N., Bluett, J. et al. (2015) Dense genotyping of immune-related susceptibility loci reveals new insights into the genetics of psoriatic arthritis. *Nat. Commun.*, **6**, 1–11.
 23. Nejentsev, S., Howson, J.M.M., Walker, N.M., Szeszeko, J., Field, S.F., Stevens, H.E., Reynolds, P., Hardy, M., King, E., Masters, J. et al. (2007) Localization of type 1 diabetes susceptibility to the MHC class I genes HLA-B and HLA-A. *Nature*, **450**, 887–892.
 24. Kaneko, T. and Obata, F. (1996) Allogeneic recognition of HLA-DRB1(*0406) by T cells with HLA-DRB1(*0403): role of amino acid residue 37 on the β sheet in T cell recognition. *Immunobiology*, **195**, 261–270.
 25. Chen, Y.Z., Matsushita, S. and Nishimura, Y. (1997) A single residue polymorphism at DR β 37 affects recognition of peptides by T cells. *Hum. Immunol.*, **54**, 30–39.
 26. Rothwell, S., Chinoy, H., Lamb, J.A., Miller, F.W., Rider, L.G., Wedderburn, L.R., McHugh, N.J., Mammen, A.L., Betteridge, Z.E., Tansley, S.L. et al. (2019) Focused HLA analysis in Caucasians with myositis identifies significant associations with autoantibody subgroups. *Ann. Rheum. Dis.*, **78**, 996–1002.
 27. Raychaudhuri, S., Sandor, C., Stahl, E.A., Freudenberg, J., Lee, H.-S., Jia, X., Alfredsson, L., Padyukov, L., Klareskog, L., Worthington, J. et al. (2012) Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nat. Genet.*, **44**, 291–296.
 28. Ombrello, M.J., Remmers, E.F., Tachmazidou, I., Grom, A., Foell, D., Haas, J.P., Martini, A., Gattorno, M., Özen, S., Pahalad, S. et al. (2015) HLA-DRB1*11 and variants of the MHC class II locus are strong risk factors for systemic juvenile idiopathic arthritis. *Proc. Natl. Acad. Sci. U. S. A.*, **112**, 15970–15975.
 29. Goyette, P., Boucher, G., Mallon, D., Ellinghaus, E., Jostins, L., Huang, H., Ripke, S., Gusareva, E.S., Annesse, V., Hauser, S.L. et al. (2015) High-density mapping of the MHC identifies a shared role for HLA-DRB1*01:03 in inflammatory bowel diseases and heterozygous advantage in ulcerative colitis. *Nat. Genet.*, **47**, 172–179.
 30. Patsopoulos, N.A., Barcellos, L.F., Hintzen, R.Q., Schaefer, C., van Duijn, C.M., Noble, J.A., Raj, T., Gourraud, P.-A., Stranger, B.E., Oksenberg, J. et al. (2013) Fine-mapping the genetic association of the major histocompatibility complex in multiple sclerosis: HLA and non-HLA effects. *PLoS Genet.*, **9**, e1003926.
 31. Menconi, F., Osman, R., Monti, M.C., Greenberg, D.A., Concepcion, E.S. and Tomer, Y. (2010) Shared molecular amino acid signature in the HLA-DR peptide binding pocket predisposes to both autoimmune diabetes and thyroiditis. *Proc. Natl. Acad. Sci. U. S. A.*, **107**, 16899–16903.
 32. Ban, Y., Davies, T.F., Greenberg, D.A., Concepcion, E.S., Osman, R., Oashi, T. and Tomer, Y. (2004) Arginine at position 74 of the HLA-DR β 1 chain is associated with Graves' disease. *Genes Immun.*, **5**, 203–208.
 33. Gourh, P., Safran, S.A., Alexander, T., Boyden, S.E., Morgan, N.D., Shah, A.A., Mayes, M.D., Doumatey, A., Bentley, A.R., Shriner, D. et al. (2020) HLA and autoantibodies define scleroderma subtypes and risk in African and European Americans and suggest a role for molecular mimicry. *Proc. Natl. Acad. Sci. U. S. A.*, **117**, 552–562.
 34. Langefeld, C.D., Ainsworth, H.C., Graham, D.S.C., Kelly, J.A., Comeau, M.E., Marion, M.C., Howard, T.D., Ramos, P.S., Croker, J.A., Morris, D.L. et al. (2017) Transancestral mapping and genetic loci in systemic lupus erythematosus. *Nat. Commun.*, **8**, 1–18.

35. Han, B., Akiyama, M., Kim, K.K., Oh, H., Choi, H., Lee, C.H., Jung, S., Lee, H.S., Kim, E.E., Cook, S. *et al.* (2018) Amino acid position 37 of HLA-DR β 1 affects susceptibility to Crohn's disease in Asians. *Hum. Mol. Genet.*, **27**, 3901–3910.
36. Molineros, J.E., Looger, L.L., Kim, K., Okada, Y., Terao, C., Sun, C., Zhou, X., Raj, P., Kochi, Y., Suzuki, A. *et al.* (2019) Amino acid signatures of HLA class-I and II molecules are strongly associated with SLE susceptibility and autoantibody production in Eastern Asians. *PLoS Genet.*, **15**, e1008092.
37. Jee, S.-H., Tsai, T.-F., Tsai, W.-L., Liaw, S.-H., Chang, C.-H. and HU, C.-Y. (1998) HLA-DRB1*0701 and DRB1*1401 are associated with genetic susceptibility to psoriasis vulgaris in a Taiwanese population. *Br. J. Dermatol.*, **139**, 978–983.
38. Oka, S., Furukawa, H., Shimada, K., Hashimoto, A., Komiya, A., Tsunoda, S., Saisho, K., Tsuchiya, N., Katayama, M., Shinohara, S. *et al.* (2019) Association of HLA-DRB1 genotype with younger age onset and elder age onset rheumatoid arthritis in Japanese populations. *Medicine*, **98**, e18218.
39. Jamieson, R.V., Farrar, N., Stewart, K., Perveen, R., Mihelec, M., Carette, M., Grigg, J.R., McAvoy, J.W., Lovicu, F.J., Tam, P.P.L. *et al.* (2007) Characterization of a familial t(16;22) balanced translocation associated with congenital cataract leads to identification of a novel gene, TMEM114, expressed in the lens and disrupted by the translocation. *Hum. Mutat.*, **28**, 968–977.
40. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., De Bakker, P.I.W., Daly, M.J. *et al.* (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.*, **81**, 559–575.
41. McCarthy, S., Das, S., Kretschmar, W., Delaneau, O., Wood, A.R., Teumer, A., Kang, H.M., Fuchsberger, C., Danecek, P., Sharp, K. *et al.* (2016) A reference panel of 64,976 haplotypes for genotype imputation. *Nat. Genet.*, **48**, 1279–1283.
42. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M. and Lee, J.J. (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, **4**, 7.
43. Altshuler, D.M., Gibbs, R.A., Peltonen, L., Schaffner, S.F., Yu, F., Dermitzakis, E., Bonnen, P.E., De Bakker, P.I.W., Deloukas, P., Gabriel, S.B. *et al.* (2010) Integrating common and rare genetic variation in diverse human populations. *Nature*, **467**, 52–58.
44. Yang, J., Benyamin, B., McEvoy, B.P., Gordon, S., Henders, A.K., Nyholt, D.R., Madden, P.A., Heath, A.C., Martin, N.G., Montgomery, G.W. *et al.* (2010) Common SNPs explain a large proportion of the heritability for human height. *Nat. Genet.*, **42**, 565–569.
45. Lee, S.H., Wray, N.R., Goddard, M.E. and Visscher, P.M. (2011) Estimating missing heritability for disease from genome-wide association studies. *Am. J. Hum. Genet.*, **88**, 294–305.
46. Jia, X., Han, B., Onengut-Gumuscu, S., Chen, W.-M., Concannon, P.J., Rich, S.S., Raychaudhuri, S. and de Bakker, P.I.W. (2013) Imputing amino acid polymorphisms in human leukocyte antigens. *PLoS One*, **8**, e64683.
47. Okada, Y., Han, B., Tsoi, L.C., Stuart, P.E., Ellinghaus, E., Tejasvi, T., Chandran, V., Pellett, F., Pollock, R., Bowcock, A.M. *et al.* (2014) Fine mapping major histocompatibility complex associations in psoriasis and its clinical subtypes. *Am. J. Hum. Genet.*, **95**, 162–172.
48. Das, S., Forer, L., Schönherr, S., Sidore, C., Locke, A.E., Kwong, A., Vrieze, S.I., Chew, E.Y., Levy, S., McGue, M. *et al.* (2016) Next-generation genotype imputation service and methods. *Nat. Genet.*, **48**, 1284–1287.
49. Marchini, J., Howie, B., Myers, S., McVean, G. and Donnelly, P. (2007) A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat. Genet.*, **39**, 906–913.