

# The association of smoking status with SARS-CoV-2 infection, hospitalization and mortality from COVID-19: a living rapid evidence review with Bayesian meta-analyses (version 7)

David Simons<sup>1</sup> , Lion Shahab<sup>2</sup> , Jamie Brown<sup>2</sup>  & Olga Perski<sup>2</sup> 

Centre for Emerging, Endemic and Exotic Diseases, Royal Veterinary College, London, UK<sup>1</sup> and Department of Behavioural Science and Health, University College London, London, UK<sup>2</sup>

## ABSTRACT

**Aims** To estimate the association of smoking status with rates of (i) infection, (ii) hospitalization, (iii) disease severity and (iv) mortality from SARS-CoV-2/COVID-19 disease. **Design** Living rapid review of observational and experimental studies with random-effects hierarchical Bayesian meta-analyses. Published articles and pre-prints were identified via MEDLINE and medRxiv. **Setting** Community or hospital, no restrictions on location. **Participants** Adults who received a SARS-CoV-2 test or a COVID-19 diagnosis. **Measurements** Outcomes were SARS-CoV-2 infection, hospitalization, disease severity and mortality stratified by smoking status. Study quality was assessed (i.e. 'good', 'fair' and 'poor'). **Findings** Version 7 (searches up to 25 August 2020) included 233 studies with 32 'good' and 'fair' quality studies included in meta-analyses. Fifty-seven studies (24.5%) reported current, former and never smoking status. Recorded smoking prevalence among people with COVID-19 was generally lower than national prevalence. Current compared with never smokers were at reduced risk of SARS-CoV-2 infection [relative risk (RR) = 0.74, 95% credible interval (CrI) = 0.58–0.93,  $\tau = 0.41$ ]. Data for former smokers were inconclusive (RR = 1.05, 95% CrI = 0.95–1.17,  $\tau = 0.17$ ), but favoured there being no important association (21% probability of  $RR \geq 1.1$ ). Former compared with never smokers were at somewhat increased risk of hospitalization (RR = 1.20, CrI = 1.03–1.44,  $\tau = 0.17$ ), greater disease severity (RR = 1.52, CrI = 1.13–2.07,  $\tau = 0.29$ ) and mortality (RR = 1.39, 95% CrI = 1.09–1.87,  $\tau = 0.27$ ). Data for current smokers were inconclusive (RR = 1.06, CrI = 0.82–1.35,  $\tau = 0.27$ ; RR = 1.25, CrI = 0.85–1.93,  $\tau = 0.34$ ; RR = 1.22, 95% CrI = 0.78–1.94,  $\tau = 0.49$ , respectively), but favoured there being no important associations with hospitalization and mortality (35% and 70% probability of  $RR \geq 1.1$ , respectively) and a small but important association with disease severity (79% probability of  $RR \geq 1.1$ ). **Conclusions** Compared with never smokers, current smokers appear to be at reduced risk of SARS-CoV-2 infection, while former smokers appear to be at increased risk of hospitalization, increased disease severity and mortality from COVID-19. However, it is uncertain whether these associations are causal.

**Keywords** COVID-19, e-cigarettes, hospitalization, infection, living review, mortality, nicotine replacement therapy, SARS-CoV-2, smoking, tobacco.

*Correspondence to:* Olga Perski, Department of Behavioural Science and Health, University College London, 1–19 Torrington Place, London WC1E 7HB, UK.  
E-mail: olga.perski.14@ucl.ac.uk

Submitted 24 April 2020; initial review completed 8 July 2020; final version accepted 25 September 2020

## INTRODUCTION

COVID-19 is a respiratory disease caused by the SARS-CoV-2 virus. Large age and gender differences in case severity and mortality have been observed in the ongoing COVID-19 pandemic [1]; however, these differences are currently unexplained. SARS-CoV-2 enters epithelial cells

through the angiotensin-converting enzyme 2 (ACE-2) receptor [2]. Some evidence suggests that gene expression and subsequent receptor levels are elevated in the airway and oral epithelium of current smokers [3,4], thus putting smokers at higher risk of contracting SARS-CoV-2. Other studies, however, suggest that nicotine down-regulates the ACE-2 receptor [5]. These uncertainties

notwithstanding, both former and current smoking is known to increase the risk of respiratory viral [6,7] and bacterial [8,9] infections and is associated with worse outcomes once infected. Cigarette smoke reduces the respiratory immune defence through peri-bronchiolar inflammation and fibrosis, impaired mucociliary clearance and disruption of the respiratory epithelium [10]. There is also reason to believe that behavioural factors (e.g. regular hand-to-mouth movements) involved in smoking may increase SARS-CoV-2 infection and transmission in current smokers. However, early data from the COVID-19 pandemic have not provided clear evidence for a negative impact of current or former smoking on SARS-CoV-2 infection or COVID-19 disease outcomes, such as hospitalization or mortality [11]. It has also been hypothesized that nicotine might protect against a hyperinflammatory response to SARS-CoV-2 infection, which may lead to adverse outcomes in patients with COVID-19 disease [12].

There are several reviews that fall within the scope of smoking and COVID-19 [11,13–18]. We aimed to produce a rapid synthesis of available evidence pertaining to the rates of infection, hospitalization, disease severity and mortality from SARS-CoV-2/COVID-19 stratified by smoking status. Given the increasing availability of data on this topic, this is a living review with regular updates. As evidence accumulates, the review will be expanded to include studies reporting COVID-19 outcomes by alternative nicotine use (e.g. nicotine replacement therapy or e-cigarettes).

## METHODS

### Study design

This is a living evidence review, which is updated as new evidence becomes available [19]. We adopted recommended best practice for rapid evidence reviews, which involved limiting the search to main databases and having one reviewer extract the data and another verify [20]. This study was not pre-registered, but evolved from a report written for a UK medical society [21]. The most recent (and all future) version(s) of this living review is <https://www.qeios.com/read/latest-UJR2AW>. A completed Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist is included in Supporting information, Fig. S1.

### Eligibility criteria

Studies were included if they:

1. Were primary research studies using experimental (e.g. randomized controlled trial), quasi-experimental (e.g. pre- and post-test;) or observational (e.g. case-control, retrospective cohort, prospective cohort) study designs;
2. Included adults aged 16 + years;

3. Recorded as outcome (i) results of a SARS-CoV-2 diagnostic test (including antibody assays), (ii) clinical diagnosis of COVID-19, (iii) hospitalization with COVID-19, (iv) severity of COVID-19 disease in those hospitalized or (v) mortality from COVID-19;
4. Reported any of the outcomes of interest by self-reported or biochemically verified smoking status (e.g. current smoker, former smoker, never smoker) or current vaping or nicotine replacement therapy (NRT) use;
5. Were available in English; and
6. Were published in a peer-reviewed journal, as a pre-print or a public health report by reputable agents (e.g. governments, scientific societies).

### Search strategy

The following terms were searched for in Ovid MEDLINE (2019-search date) as free text or Medical Subject Headings:

1. Tobacco Smoking/ or Smoking Cessation/ or Water Pipe Smoking/ or Smoking/ or Smoking Pipes/ or Cigar Smoking/ or Smoking Prevention/ or Cigarette Smoking/ or smoking.mp. or Pipe Smoking/ or Smoking, Non-Tobacco Products/ or Smoking Water Pipes/
2. Nicotine/or nicotine.mp. or Electronic Nicotine Delivery Systems/ or Nicotine Chewing Gum/
3. vaping.mp. or Vaping/
4. 1 or 2 or 3
5. Coronavirus/ or Severe Acute Respiratory Syndrome/ or Coronavirus Infections/ or covid.mp.
6. 4 and 5

The following terms were searched for in titles, abstracts and full texts in medRxiv no time limitations):

1. covid (this term captures both covid and SARS-CoV-2)
2. covid AND smoking
3. covid AND nicotine
4. covid AND vaping

Additional articles/reports of interest were identified through mailing lists, Twitter, the International Severe Acute Respiratory and Emerging Infection Consortium (ISARIC) and the US Centers for Disease Control and Prevention (CDC). Where updated versions of pre-prints or public health reports were available, old versions were superseded.

### Selection of studies

One reviewer screened titles, abstracts and full texts against the inclusion criteria.

### Data extraction

Data were extracted by one reviewer and verified (i.e. independently checked against pre-prints and published reports) by another on (i) author (year); (ii) date published;

(iii) country; (iv) study design; (v) study setting; (vi) sample size; (vii) sex; (viii) age; (ix) smoking status (e.g. current, former, never, not stated, missing); (x) use of alternative nicotine products; (xi) SARS-CoV-2 testing; (xii) SARS-CoV-2 infection; (xiii) diagnosis of COVID-19; (xiv) hospitalization with COVID-19; (xv) disease severity in those hospitalized with COVID-19; and (xvi) mortality.

### Quality appraisal

The quality of included studies was assessed to determine suitability for inclusion in meta-analyses. Studies were judged as 'good' quality if they: (i) had < 20% missing data on smoking status and used a reliable self-report measure that distinguished between current, former and never smoking status; AND (ii) used biochemical verification of smoking status and reported results from adjusted analyses; OR reported data from a representative/random sample. Studies were rated as 'fair' if they fulfilled only criterion (i) and were otherwise rated as 'poor'. The quality appraisal was conducted by one reviewer and verified by a second.

### Evidence synthesis

A narrative synthesis was conducted. Data from 'good' and 'fair' quality studies were pooled in R version 3.6.3 [22]. In a living review where new data are regularly added to the analyses, it may be more appropriate to use a Bayesian (as opposed to frequentist) approach where prior knowledge is used in combination with new data to estimate a posterior risk distribution. A Bayesian approach mitigates against the issue of performing multiple statistical tests, which can inflate family-wise error. A series of random-effects hierarchical Bayesian meta-analyses were performed with the *brms* [23] package to estimate the relative risk for each comparison with accompanying 95% credible intervals (CrIs). We first defined prior distributions for the true pooled effect size ( $\mu$ ) and the between-study heterogeneity ( $\tau$ ), with  $\mu$  specified as a normal distribution with a mean equal to the derived point estimate from each comparison of interest in the immediately preceding version of this living review [24], and  $\tau$  specified as a half-Cauchy distribution with a mean of 0 and standard deviation of 1. The half-Cauchy distribution was selected to reflect prior knowledge that high levels of between-study heterogeneity are more likely than lower levels. Markov chain Monte Carlo methods (20000 burn-ins followed by 80000 iterations) were then used to generate a risk distribution for each study, in addition to a pooled effect for the posterior risk distribution. We report forest plots with the pooled effect for the posterior risk distribution displayed as the median relative risk (RR) with an accompanying 95% CrIs. We used the empirical cumulative distribution function

(ECDF) to estimate the probability of there being a 10% reduction or 10% increase in the RR (i.e.  $RR \geq 1.1$  or  $RR \leq 0.9$ ). Due to a lack of indication as to what constitutes a clinically or epidemiologically meaningful effect (e.g. with regard to onward disease transmission or requirements for intensive care beds), we deemed a 10% change in risk as small, but important. Where data were inconclusive (as indicated by CrIs crossing  $RR = 1.0$ ), to disambiguate whether data favoured no effect or there being a small but important association, we estimated whether there was  $\geq 75\%$  probability of  $RR \geq 1.1$  or  $RR \leq 0.9$ .

Two sensitivity analyses were performed. First, a minimally informative prior for  $\mu$  was specified as a normal distribution with a mean of 0 and standard deviation of 1 and  $\tau$  as described above. Second, an informative prior as described above for  $\mu$  was used with  $\tau$  specified as a half-Cauchy distribution with a mean of 0.3 and standard deviation of 1 to reflect greater between-study heterogeneity.

To aid in the visualization of smoking prevalence in the included studies, 95% bootstrap percentile confidence intervals (CIs) were calculated for each study. We performed 1000 bootstrap replications, with the 2.5th and 97.5th percentiles of the empirical distribution forming the 95% bootstrap percentile CIs [25]. It should be noted that prevalence estimates in the included studies were not adjusted for age, sex, socio-economic position or region within countries.

## DATA AVAILABILITY


All data contributing to the current and future review versions are <https://doi.org/10.6084/m9.figshare.12756020>. All code required to reproduce the current and future analyses are <https://doi.org/10.5281/zenodo.4002046>.

## RESULTS

In the current review (version 7) with searches up to 25 August 2020, a total of 347 new records were identified, with 233 studies included in a narrative synthesis and 32 studies included in meta-analyses (see Fig. 1).

### Study characteristics

Characteristics of included studies are presented in Table 1. Studies were conducted across 33 countries. Sixty-two studies were conducted in the United States, 53 in China, 26 in the United Kingdom, 13 in Spain, 12 in Mexico, 11 in France, seven in Italy, six across multiple international sites, four in Brazil and Iran, three in Israel and Turkey, two in Bangladesh, Chile, Denmark, Finland, India, Japan and Qatar and one from 15 further countries (see



**FIGURE I** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of included studies.

Supporting information, Fig. S1). The majority of studies used observational designs (see Supporting information, Table S1). One hundred and fifty-five studies were conducted in hospital settings, 62 studies included a community component in addition to hospitalized patients, 14 studies were conducted exclusively in the community, one study was conducted in a quarantine centre and one did not state the study setting. Studies had a median of 404 (interquartile range = 115–1631) participants. The majority of studies (93.5%) used reverse transcriptase–polymerase chain reaction (RT–PCR) for confirmation of SARS-CoV-2 infection, 2.6% used an antibody test to confirm prior infection and 3.9% further studies relied on a combination of RT–PCR and clinical diagnosis (see Supporting information, Table S1).

#### Smoking status

Categorization of smoking status was heterogeneous (see Table 1). One hundred and forty-five studies collected data on smoking status through routine electronic health records (EHRs), 59 studies used a bespoke case report form for COVID-19 and 29 studies did not state the source for information on smoking status. None of the studies verified smoking status biochemically. Notably, only 57 (24.4%) studies reported current, former and never smoking status (see Supporting information, Table S2a), with a further 17

studies reporting ever and never smoking status (see Supporting information, Table S2b). The remaining 159 studies reported current, current/former or current and former smoking status but did not explicitly state whether remaining participants were never smokers or if data were missing on smoking status (see Supporting information, Table S2c). Seventy-eight studies explicitly reported the proportion with missing data on smoking status, which ranged from 0.08 to 96.4%.

#### Use of alternative nicotine products

Five studies recorded the use of alternative nicotine products in current and/or former smokers but did not report COVID-19 outcomes stratified by nicotine use [26–30].

#### Quality appraisal

One study was performed in a random, representative population sample and was rated as 'good' quality. Forty-six studies were rated as 'fair' quality. The remaining 186 studies were rated as 'poor' quality (see Table 1).

#### Smoking prevalence by country

Unadjusted smoking prevalence compared with overall estimates for national adult smoking prevalence split by country and study setting is presented in Fig. 2a,b. Lower

TABLE 1 Characteristics of included studies.

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality	
[1]	Guan, Ni	2020–02–28	China	1099	Hospital	47 (35–58)	41.9	12.5	1.9	—	84.3	—	1.27	Fair	
[50]	Guan, Liang	2020–03–26	China	1590	Hospital	49 (33–64)	42.7	—	—	7.0	93.0	—	0.00	Poor	
[51]	Lian	2020–03–25	China	788	Hospital	NA	38.5	6.9	—	—	—	—	93.15	Poor	
[52]	Jin	2020–03–24	China	651	Hospital	46 (32–60)	49.2	6.3	—	—	—	—	93.70	Poor	
[53]	Chen	2020–03–26	China	548	Hospital	62 (44–70)	37.6	4.4	2.6	—	—	—	93.07	Poor	
[54]	Zhou, Yu	2020–03–11	China	191	Hospital	56 (46–67)	38.0	5.8	—	—	—	—	94.24	Poor	
[55]	Mo	2020–03–16	China	155	Hospital	54 (53–66)	44.5	3.9	—	—	—	—	96.13	Poor	
[56]	Zhang, Dong	2020–02–19	China	140	Hospital	57 <sup>†</sup> (25–87)	46.3	1.4	5.0	—	—	—	93.57	Poor	
[57]	Wan	2020–03–21	China	135	Hospital	47 (36–55)	46.7	6.7	—	—	—	—	93.33	Poor	
[58]	Liu, Tao	2020–02–28	China	78	Hospital	38 (33–57)	50.0	—	—	6.4	—	—	—	93.59	Poor
[59]	Huang, Wang	2020–01–24	China	41	Hospital	49 (41–58)	27.0	7.3	—	—	—	—	92.68	Poor	
[60]	Zhang, Cai	2020–03–20	China	645	Hospital	NA	49.1	6.4	—	—	—	—	93.64	Poor	
[61]	Guo	2020–03–27	China	187	Hospital	59 (45–73)	51.3	9.6	—	—	—	—	90.37	Poor	
[62]	Liu, Ming	2020–03–12	China	41	Hospital	39 (30–48)	58.5	9.8	—	—	—	—	90.24	Poor	
[63]	Huang, Yang	2020–03–05	China	36	Hospital	69 (60–78)	30.6	—	—	11.1	—	—	88.89	Poor	
[64]	Xu	2020–03–08	China	53	Hospital	NA	47.2	11.3	—	—	—	—	88.68	Poor	
[65]	Li	2020–02–12	China	17	Hospital	45 (33–57)	47.1	17.6	—	—	—	—	82.35	Poor	

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[31]	Rentsch	2020–04–14	USA	3528	Community and Hospital	66 (60–70)	4.6	27.2	30.6	—	36.9	—	5.30	Fair
[66]	Hu	2020–03–25	China	323	Hospital	61 <sup>†</sup> (23–91)	48.6	—	—	11.8	—	—	88.24	Poor
[67]	Wang, Pan	2020–03–24	China	125	Hospital	41 (26–66)	43.2	—	—	12.8	—	—	87.20	Poor
[68]	Chow (US CDC)	2020–03–31	USA	7162	Community and Hospital	NA	—	1.3	2.3	—	—	—	96.36	Poor
[69]	Dong, Cao	2020–03–20	China	9	Hospital	44 (30–46)	66.7	11.1	—	—	—	—	88.89	Poor
[70]	Kim	2020–04–01	South Korea	28	Hospital	43 (30–56)	46.4	17.9	—	—	—	—	82.14	Poor
[71]	Shi, Yu	2020–03–18	China	487	Hospital	46 (27–65)	46.8	—	—	8.2	—	—	91.79	Poor
[72]	Yang, Yu	2020–02–24	China	52	Hospital	60 (47–73)	37.0	3.8	—	—	—	—	96.15	Poor
[73]	Argenziano	2020–05–29	USA	1000	Hospital	63 (50–75)	40.4	4.9	17.9	—	77.2	—	0.00	Fair
[74]	Solis	2020–04–25	Mexico	650	Hospital	46 (NA)	42.1	9.4	—	—	—	—	90.62	Poor
[75]	Richardson	2020–04–22	USA	5700	Hospital	63 (52–75)	39.7	—	—	9.8	52.8	—	37.42	Poor
[76]	Fontanet	2020–04–23	France	661	Community and Hospital	37 (16–47)	62.0	10.4	—	—	—	89.6	0.00	Poor
[77]	Zheng, Gao	2020–04–19	China	66	Hospital	47 <sup>†</sup> (NA)	25.8	12.1	—	—	—	—	87.88	Poor
[78]	Liao, Feng	2020–04–24	China	1848	Hospital	55 (48–61)	54.7	—	—	0.4	—	—	99.57	Poor
[79]	Gil-Agudo	2020–04–24	Spain	7	Hospital	68 (34–75)	28.6	—	—	42.9	57.1	—	0.00	Poor
[80]	Shi, Ren	2020–04–23	China	134	Hospital	46 (34–58)	51.5	—	—	10.4	—	—	89.55	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[81]	Hadjadj	2020–04–23	France	50	Hospital	55 (50–63)	22.0	2.0	18.0	—	80.0	—	0.00	Fair
[82]	Gold (US CDC)	2020–04–20	USA	305	Hospital	NA	50.5	5.2	—	—	—	—	94.75	Poor
[83]	Yu, Cai	2020–04–27	China	95	Hospital	NA	44.2	8.4	—	—	—	—	91.58	Poor
[84]	Zheng, Xiong	2020–04–30	China	73	Hospital	43 <sup>†</sup> (NA)	45.2	—	—	11.0	89.0	—	0.00	Poor
[85]	de la Rica	2020–05–11	Spain	48	Hospital	66 <sup>†</sup> (33–88)	33.0	—	—	20.8	—	—	79.17	Poor
[86]	Yin, Yang	2020–05–10	China	106	Hospital	73 (61–85)	39.6	—	—	17.0	—	—	83.02	Poor
[87]	Shi, Zuo	2020–05–17	USA	172	Hospital	63 <sup>†</sup> (44–82)	44.0	—	—	26.2	—	—	73.84	Poor
[88]	Cho	2020–05–11	UK	322	Community and Hospital	NA	49.2	14.2	21.4	—	64.4	—	0.00	Fair
[89]	Allenbach	2020–05–08	France	152	Hospital	77 (60–83)	31.1	—	—	6.6	—	—	93.42	Poor
[90]	Robilotti	2020–05–08	USA	423	Hospital	NA	50.0	2.1	37.6	—	58.6	—	1.65	Fair
[91]	The OpenSAFELY Collaborative	2020–07–01	UK	17278	Community and Hospital	NA	50.1	17.0	32.9	—	45.9	—	4.17	Fair
[92]	Borobia	2020–05–06	Spain	392	Community and Hospital	61 (46–78)	52.0	7.1	—	—	—	—	92.95	Poor
[93]	Giacomelli	2020–05–06	Italy	233	Hospital	61 (50–72)	31.9	—	—	30.0	70.0	—	0.00	Poor
[94]	Shah	2020–05–06	USA	316	Hospital	63 (43–72)	48.1	16.5	17.7	—	42.1	—	23.73	Poor
[95]	Kolin	2020–05–05	UK	502	Community and Hospital	56.5 (48–64)	54.4	10.5	34.4	—	54.4	—	0.59	Fair
[96]	Lubetzy	2020–05–08	USA	54	Hospital	57 (29–83)	62.0	—	—	22.2	—	—	77.78	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[97]	Goyal	2020–04–17	USA	393	Hospital	62.2 (49–74)	39.3	5.1	—	—	—	—	94.91	Poor
[98]	Feng	2020–04–10	China	476	Hospital	53 (40–64)	43.1	9.2	—	—	—	—	90.76	Poor
[99]	Yao	2020–04–24	China	108	Hospital	52 (37–58)	60.2	3.7	—	—	—	—	96.30	Poor
[100]	Sami	2020–05–19	Iran	490	Hospital	56.6 (41–71)	39.0	14.1	—	—	—	85.9	0.00	Poor
[101]	Almazedi	2020–05–15	Kuwait	1096	Hospital	41 (25–57)	19.0	4.0	—	—	—	96.0	0.00	Poor
[102]	Carillo-Vega	2020–05–14	Mexico	10544	Community and Hospital	46.5 <sup>†</sup> (30–62)	42.3	8.9	—	—	—	—	91.12	Poor
[103]	Yanover	2020–05–13	Israel	4353	Community and Hospital	35 (22–54)	44.5	11.8	3.0	—	85.2	—	0.00	Fair
[104]	Hamer	2020–05–13	UK	387	Hospital	56.2 (48–64)	55.1	9.7	34.8	—	55.5	—	0.00	Fair
[105]	Regina	2020–05–14	Switzerland	109	Hospital	70 (55–81)	40.0	4.5	—	—	—	—	95.50	Poor
[39]	de Lusignan	2020–05–15	UK	3802	Community and Hospital	58 (34–73)	57.6	10.9	46.1	—	29.6	—	13.44	Fair
[106]	Targher	2020–05–13	China	339	Hospital	48.4 <sup>†</sup> (NA)	52.8	8.3	—	—	—	—	91.74	Poor
[107]	Valenti	2020–05–18	Italy	789	Community	40.7 <sup>†</sup> (NA)	35.0	25.9	—	—	—	—	74.14	Poor
[108]	Feuth	2020–05–18	Finland	28	Hospital	56 (47–72)	46.0	10.7	28.6	—	60.7	—	0.00	Fair
[109]	Ge	2020–05–18	China	51	Hospital	70 (58–79)	27.5	13.7	—	—	—	—	86.27	Poor
[110]	Parrotta	2020–05–18	USA	76	Community and Hospital	44.9 (13–71)	61.8	2.6	26.3	—	68.4	—	2.63	Fair
[111]	Shekhar	2020–05–18	USA	50	Hospital	55.5 (20–85)	54.0	48.0	—	—	—	—	52.00	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[112]	Mejia-Vilet	2020–05–16	Mexico	329	Hospital	49 (41–60)	36.0	–	–	7.0	–	–	93.01	Poor
[113]	Chen, Jiang	2020–05–16	China	135	Hospital	NA	42.2	–	–	9.6	–	–	90.37	Poor
[114]	Li, Chen	2020–05–16	China	1008	Hospital	55 (44–65)	43.6	5.7	–	–	–	–	94.35	Poor
[27]	Rimland	2020–05–19	USA	11	Hospital	59 (48–65)	18.2	9.1	–	–	–	–	81.82	Poor
[115]	Palaiodimos	2020–05–15	USA	200	Hospital	64 (50–73.5)	51.0	–	–	32.5	67.5	–	0.00	Poor
[116]	Ip	2020–05–25	USA	2512	Hospital	64 (52–76)	37.6	3.1	17.8	–	64.5	–	14.61	Fair
[117]	Heili-Frades	2020–05–25	Spain	4712	Hospital	62 (47–77)	50.5	4.9	17.4	–	–	66.5	11.16	Poor
[118]	Vaquero-Roncero	2020–05–24	Spain	146	Hospital	66 <sup>†</sup> (59–72)	32.2	–	–	6.8	–	–	93.15	Poor
[119]	Kim, Garg	2020–05–22	USA	2491	Hospital	62 (50–75)	46.8	6.0	25.8	–	–	68.1	0.08	Poor
[120]	Wu	2020–05–21	Italy	174	Hospital	61.2 <sup>†</sup> (50–71)	30.5	–	–	33.3	–	–	66.67	Poor
[121]	Shi, Zhao	2020–05–20	China	101	Hospital	71 (59–80)	40.6	–	–	5.0	–	–	95.05	Poor
[122]	Al-Hindawi	2020–05–20	UK	31	Hospital	61 (NA)	12.9	3.2	71.0	–	25.8	–	0.00	Fair
[123]	Basse	2020–05–19	France	141	Hospital	62 (52–72)	72.0	17.7	–	–	–	–	82.27	Poor
[124]	Freites	2020–05–19	Spain	123	Hospital	59.88 <sup>†</sup> (44–74)	69.9	3.3	–	–	–	–	96.75	Poor
[125]	Alshami	2020–05–19	Saudi Arabia	128	Quarantine Centre	39.6 <sup>†</sup> (24–55)	53.9	15.6	2.3	–	–	–	82.03	Poor
[126]	Berumen	2020–05–26	Mexico	102	Hospital	NA	49.1	–	–	9.6	–	90.4	0.00	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[127]	Gianfrancesco	2020–05–29	Multiple	600	Community and Hospital Not Stated	56 (45–67) 49 <sup>†</sup> (13–80)	71.0 61.0	— —	— 5.5	21.5 —	64.8 —	— —	13.67 94.48	Poor Poor
[128]	Li, Long	2020–05–28	China	145	Hospital	57.27 <sup>†</sup> (48–66)	44.3	11.2	—	— 70.5	— —	— —	88.77 0.00	Poor Fair
[129]	Batty	2020–06–17	UK	908	Community and Hospital	40 (27–59)	48.7	16.8	12.7	— —	— 70.5	— —	— 71.16	Poor
[130]	Israel	2020–06–01	Israel	24906	Community and Hospital	62 (52–72)	40.6	5.5	23.3	— —	— —	— —	— 85.00	Poor
[131]	del Valle	2020–05–30	USA	1484	Hospital	52 (45.5–61)	60.0	— —	— 15.0	— 45.5	— —	— —	— 54.55	Poor
[132]	Chaudhry	2020–05–29	USA	40	Community and Hospital	66.5 <sup>†</sup> (55–77)	36.4	— —	— —	— —	— —	— —	— 88.00	Poor
[133]	Louis	2020–05–28	USA	22	Hospital	NA	30.0	— —	— 12.0	— —	— —	— —	— 71.88	Poor
[134]	Soto-Mota	2020–06–05	Mexico	400	Hospital	63 (49–75)	47.0	5.5	22.6	— —	— —	— —	— 29.55	Poor
[135]	Garibaldi	2020–05–26	USA	832	Hospital	72.9 (58–82)	40.0	4.2	21.7	— —	— —	— —	— 96.71	Poor
[136]	Docherty	2020–05–22	Multiple	20133	Hospital	40 (33–50)	51.6	3.3	— —	— —	— —	— —	— 9.70	Poor
[137]	Boulware	2020–06–03	Multiple	821	Community	66 (57–76)	50.0	4.6	35.1	— —	— 26.5	— —	— 73.53	Poor
[138]	Kuderer	2020–05–28	Portugal	928	Community and Hospital	41 <sup>†</sup> (26–66)	67.7	— —	— —	— —	— —	— —	— 90.9	Fair
[139]	Romao	2020–06–08	Mexico	34	Community	42.5 <sup>†</sup> (25–59)	49.1	9.1	— —	— —	— —	— —	0.00 0.00	Poor Poor
[140]	Giannouchos	2020–06–07	Mexico	236	Community and Hospital	52 (34.7–69.5)	55.2	— —	— 26.8	— 73.2	— —	— —	— 63.79	Poor
[141]	Ramlall	2020–06–06	USA	11116	Community and Hospital	67 (NA)	48.0	— —	— 36.2	— —	— —	— —	— 63.79	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[143]	Perrone	2020–06–05	Italy	1189	Hospital	NA	21.2	–	–	21.9	–	–	78.13	Poor
[144]	Sharma	2020–06–05	India	501	Hospital	35.1 <sup>†</sup> (18–51)	36.0	–	4.2	–	–	–	95.81	Poor
[145]	Eugen-Olsen	2020–06–02	Denmark	407	Hospital	64 (47–77)	57.7	20.6	36.9	–	39.6	–	2.95	Fair
[146]	Martinez-Portilla	2020–06–02	Mexico	224	Community and Hospital	29 (26–33)	100.0	–	–	3.1	–	–	96.88	Poor
[147]	Raisi-Estabragh	2020–06–02	UK	4510	Hospital	NA	48.8	–	–	51.8	–	–	48.20	Poor
[148]	Luo	2020–06–02	China	625	Hospital	46 (NA)	47.7	3.0	–	–	–	–	96.96	Poor
[149]	Houlihan	2020–06–09	UK	200	Community	34 (29–44)	61.0	11.0	16.5	–	66.5	–	6.00	Fair
[150]	Cen	2020–06–08	China	1007	Hospital	61 (49–68)	51.0	–	–	8.7	–	–	91.26	Poor
[151]	Klang	2020–05–23	USA	3406	Hospital	NA	61.8	–	–	23.3	–	–	76.72	Poor
[152]	Maraschini	2020–06–12	Italy	146	Hospital	32.5 <sup>†</sup> (27–38)	100.0	–	9.6	–	80.8	–	9.59	Poor
[153]	Wang, Zhong	2020–06–12	USA	7592	Community and Hospital	NA	45.1	3.6	17.1	–	51.9	–	27.42	Poor
[154]	McQueenie	2020–06–12	UK	428	Community and Hospital	NA	54.9	–	–	44.4	55.0	–	0.59	Poor
[26]	Miyara	2020–06–12	France	479	Community and Hospital	NA	44.7	6.7	31.6	–	59.5	–	1.87	Fair
[155]	Apea	2020–06–12	UK	1737	Hospital	63.4 <sup>†</sup> (NA)	30.4	–	–	10.0	–	–	90.04	Poor
[156]	Woolford	2020–06–11	UK	4510	Community and Hospital	70.5 (NA)	51.2	13.0	38.1	–	48.1	–	0.80	Fair
[157]	Hultcrantz	2020–06–11	USA	127	Community and Hospital	68 (41–91)	46.0	–	–	26.8	72.4	–	0.79	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[158]	Rajter	2020–06–10	USA	280	Hospital	59.6 <sup>†</sup> (41–77)	45.5	5.7	10.7	—	74.6	—	8.93	Fair
[159]	Lan	2020–06–09	USA	1044	Community	49 <sup>†</sup> (34–63)	47.1	—	—	24.0	—	—	75.96	Poor
[160]	Zeng	2020–06–16	China	1031	Hospital	60.3 <sup>†</sup> (46–74)	47.8	—	—	10.2	—	—	89.82	Poor
[161]	Suleyman	2020–06–16	USA	463	Hospital	57.5 <sup>†</sup> (40–74)	55.9	—	—	34.6	—	—	65.44	Poor
[162]	Chen, Yu	2020–06–16	China	1859	Hospital	59 (45–68)	50.0	2.4	3.6	—	94.0	—	0.00	Fair
[163]	Garasino	2020–06–12	Multiple	200	Community and Hospital	68 (61.8–75)	30.0	24.0	55.5	—	18.5	—	2.00	Fair
[164]	Hernandez-Gardano	2020–06–11	Mexico	32583	Community and Hospital	45 (34–56)	48.7	—	—	11.0	—	88.8	0.15	Poor
[165]	Govind	2020–06–20	UK	6309	Community and Hospital	46.5 <sup>†</sup> (31–61)	38.3	66.3	26.8	—	5.5	—	1.49	Fair
[166]	Siso-Almirall	2020–06–20	Spain	322	Community and Hospital	56.7 <sup>†</sup> (38–74)	50.0	—	—	25.2	—	—	74.84	Poor
[167]	Gu	2020–06–18	USA	5698	Community and Hospital	47 <sup>†</sup> (26–67)	62.0	7.0	24.7	—	50.8	—	17.53	Fair
[168]	Kibler	2020–06–16	France	702	Community and Hospital	82 <sup>†</sup> (75–88)	56.0	3.7	—	—	—	—	96.30	Poor
[169]	Ikitimur	2020–06–03	Turkey	81	Hospital	55 <sup>†</sup> (38–72)	44.0	—	—	28.4	—	—	71.60	Poor
[170]	Sierpinska	2020–06–03	Poland	1942	Community	50 (NA)	60.0	6.3	—	—	—	49.7	44.03	Poor
[171]	Zhou, He	2020–06–10	China	238	Hospital	55.5 (35–67)	57.0	2.9	—	—	—	—	97.06	Poor
[172]	Crovetto	2020–06–19	Spain	874	Community and Hospital	33.7 <sup>†</sup> (28–38)	100.0	1.1	—	—	—	—	85.70	Poor
[173]	Veras	2020–06–09	Brazil	32	Hospital	58.9 <sup>†</sup> (40–77)	47.0	—	—	25.0	—	—	75.00	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[174]	Sterlin	2020–06–11	France	135	Hospital	61 (50–72)	41.0	3.7	38.5	—	57.8	—	0.00	Fair
[175]	Rossi	2020–06–09	France	246	Hospital	68 <sup>†</sup> (53–83)	39.0	—	—	25.2	—	—	74.80	Poor
[176]	Duan	2020–06–22	China	616	Hospital	64 (53–70)	57.5	3.7	—	—	—	—	96.27	Poor
[177]	Martin-Jimenez	2020–06–09	Spain	339	Hospital	81.6 (72–87)	39.5	—	—	30.7	—	—	69.32	Poor
[178]	Elezkurtaj	2020–06–17	Germany	26	Hospital	70 (61.8–78.3)	34.6	—	—	19.2	—	—	80.77	Poor
[179]	Lenka	2020–06–22	USA	32	Hospital	62.2 <sup>†</sup> (51–73)	37.5	—	—	50.0	—	—	50.00	Poor
[180]	Olivares	2020–06–16	Chile	21	Hospital	61 <sup>†</sup> (26–85)	76.2	—	—	9.5	—	—	90.48	Poor
[181]	Salton	2020–06–20	Italy	173	Hospital	64.4 <sup>†</sup> (NA)	34.9	—	—	29.5	—	—	70.52	Poor
[182]	Wei	2020–06–18	USA	147	Hospital	52 <sup>†</sup> (34–70)	41.0	14.3	—	—	—	—	85.71	Poor
[183]	Zuo, Estes	2020–06–17	China	172	Hospital	61 <sup>†</sup> (25–95)	44.0	—	—	26.2	—	—	73.84	Poor
[184]	Killerby	2020–06–17	USA	531	Community and Hospital	51.6 (38–62)	57.1	—	—	17.1	71.4	—	11.49	Poor
[185]	Petrilli	2020–05–22	USA	5279	Community and Hospital	54 (38–66)	51.5	5.5	17.1	—	61.9	—	15.55	Fair
[186]	Magagnoli	2020–06–05	USA	807	Hospital	70 (60–75)	4.3	—	—	15.9	—	—	84.14	Poor
[33]	Niedzwiedz	2020–05–29	UK	392	Community and Hospital	NA	54.9	9.8	34.8	—	55.4	—	0.00	Fair
[187]	Bello-Chavolla	2020–05–31	Mexico	177	Community and Hospital	42.6 (26–59)	48.9	—	—	9.3	—	—	90.72	Poor
[188]	Zuo, Yalavarthi	2020–04–24	USA	50	Hospital	61 (46–76)	34.0	—	—	36.0	—	—	64.00	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[189]	Sigel	2020–06–28	USA	493	Hospital	60 (55–67)	24.1	—	—	28.6	—	—	71.40	Poor
[190]	Nguyen	2020–06–29	USA	689	Community and Hospital	55 (40–68)	57.0	—	—	24.8	—	—	75.18	Poor
[191]	de Melo	2020–06–29	Brazil	181	Hospital	55.3 <sup>†</sup> (34–76)	60.8	9.9	12.2	—	38.1	—	39.78	Poor
[192]	Auvinen	2020–06–29	Finland	61	Hospital	53 (41–67)	36.0	18.0	27.9	—	54.1	—	0.00	Fair
[193]	Souza	2020–06–28	Brazil	8443	Hospital	NA	53.0	—	—	1.7	—	96.3	2.01	Poor
[194]	Mendy	2020–06–27	USA	689	Community and Hospital	49.5 (35.2–67.5)	47.0	—	—	24.7	—	—	75.33	Poor
[195]	Pongpirul	2020–06–26	Thailand	193	Hospital	37 (29–53)	41.5	—	—	15.0	66.3	—	18.65	Poor
[196]	Jin, Gu	2020–06–25	China	6	Hospital	60.5 <sup>†</sup> (51–75)	33.3	33.3	—	—	—	—	66.67	Poor
[197]	Favara	2020–05–23	UK	70	Community and Hospital	41 (23–64)	87.1	10.0	—	—	—	—	90.00	Poor
[198]	Fisman	2020–06–23	Canada	21922	Community and Hospital	NA	57.0	—	—	2.3	—	—	97.65	Poor
[199]	Madariaga	2020–06–23	USA	103	Community and Hospital	41.8 <sup>†</sup> (27–55)	48.5	—	—	25.2	74.8	—	0.00	Poor
[200]	Senkal	2020–07–07	Turkey	611	Hospital	57 <sup>†</sup> (18–98)	40.6	11.3	—	—	—	—	88.71	Poor
[201]	Mohamud	2020–07–02	USA	6	Hospital	65.8 <sup>†</sup> (55–78)	16.7	—	—	16.7	—	—	83.33	Poor
[202]	Magleby	2020–06–30	USA	678	Hospital	68 (50–81)	38.9	—	—	28.6	—	—	71.39	Poor
[203]	Kimmig	2020–07–06	USA	111	Hospital	63 <sup>†</sup> (48–78)	44.1	7.2	36.0	—	56.8	—	0.00	Fair
[204]	Bello-Chavolla, Antonio-Villa	2020–07–04	Mexico	60121	Community and Hospital	45.5 <sup>†</sup> (29–61)	47.0	—	—	10.5	—	—	89.52	Poor
[205]	Zacharioudakis	2020–07–04	USA	314	Hospital	64 (54–72)	34.7	—	—	22.8	—	—	77.22	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[206]	Antonio-Villa	2020–07–04	Mexico	34263	Community and Hospital	40 <sup>†</sup> (29–50)	62.9	9.7	—	—	—	—	90.32	Poor
[207]	Patel	2020–07–03	USA	129	Hospital	60.8 <sup>†</sup> (47–74)	45.0	37.2	—	—	—	55.8	6.98	Poor
[208]	Merzon	2020–07–03	Israel	7807	Community and Hospital	46.2 <sup>†</sup> (NA)	58.6	—	—	16.2	—	—	83.82	Poor
[34]	Trubiano	2020–07–02	Australia	2935	Community and Hospital	39 (29–53)	63.5	—	—	8.8	—	—	91.18	Poor
[209]	Fan	2020–07–11	UK	1425	Community and Hospital	NA	46.7	12.2	40.1	—	46.9	—	0.84	Fair
[210]	Shi, Resurreccion	2020–07–11	UK	1521	Community and Hospital	61.5 <sup>†</sup> (57–66.8)	45.9	—	—	54.9	—	—	45.10	Poor
[211]	Maucourant	2020–07–10	Sweden	27	Hospital	57 (18–78)	22.2	11.1	25.9	—	40.7	—	22.22	Poor
[212]	Elmunzer	2020–07–09	Multiple	1992	Hospital	60 <sup>†</sup> (43–76)	43.0	6.3	28.6	—	59.0	—	6.12	Fair
[213]	Alizadehsani	2020–07–09	Iran	3119	Hospital	45.48 <sup>†</sup> (26–63)	55.5	—	—	0.3	—	—	99.69	Poor
[214]	Xie	2020–07–07	China	6119	Hospital	NA	52.0	—	—	8.2	—	—	91.76	Poor
[36]	Merkely	2020–07–17	Hungary	10474	Community	48.7 <sup>†</sup> (30–66)	53.6	28.0	20.5	—	51.4	—	0.16	good
[215]	Fox	2020–07–17	UK	55	Community and Hospital	63 (23–88)	31.0	1.8	10.9	—	56.4	—	30.91	Poor
[56]	Zhang, Cao	2020–07–14	China	289	Hospital	57 (22–88)	46.6	3.5	6.2	—	—	—	90.31	Poor
[216]	Martinez-Resendez	2020–07–20	Mexico	8	Hospital	57 (48–69)	25.0	—	—	12.5	—	—	87.50	Poor
[217]	Hoertel	2020–07–20	France	12612	Hospital	58.7 <sup>†</sup> (39–77)	49.6	—	—	9.3	—	—	90.72	Poor
[218]	McGrail	2020–07–19	USA	209	Hospital	62.5 (NA)	38.8	—	—	18.7	—	—	81.34	Poor
[219]	Pandolfi	2020–07–17	Italy	33	Hospital	62 (52–65)	21.1	3.0	24.2	—	72.7	—	0.00	Fair

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[28]	Girardeau	2020–07–17	France	10	Community	30 (29–33)	50.0	40.0	10.0	—	—	—	40.00	Poor
[220]	Kurashima	2020–07–17	Japan	53	Hospital	62.9 <sup>†</sup> (49–76)	35.8	—	—	50.9	—	—	49.06	Poor
[221]	Zhan	2020–07–16	China	75	Hospital	57 (25–75)	48.0	—	—	12.0	—	—	88.00	Poor
[222]	Omraní	2020–07–16	Qatar	1409	Community and Hospital	39 (30–50)	17.2	—	—	9.2	—	—	90.77	Poor
[223]	Gupta	2020–07–16	USA	496	Hospital	70 (60–78)	46.0	—	—	7.3	—	31.7	61.09	Poor
[87]	Shi, Zuo	2020–07–15	USA	172	Hospital	61.48 <sup>†</sup> (25–96)	44.0	—	—	26.2	—	—	73.84	Poor
[224]	Hussein	2020–07–15	USA	502	Hospital	60.9 <sup>†</sup> (45–76)	52.0	9.0	22.1	—	—	68.9	0.00	Poor
[225]	Bian	2020–07–15	China	28	Hospital	56 <sup>†</sup> (42–67)	42.9	7.1	—	—	—	—	92.86	Poor
[226]	Eiros	2020–07–14	Spain	139	Community and Hospital	52 (41–57)	72.0	4.3	50.4	—	—	—	45.32	Poor
[227]	Marcos	2020–07–14	Spain	918	Hospital	72.8 <sup>†</sup> (58–87)	42.2	6.1	—	15.3	—	—	78.65	Poor
[228]	Hoertel, Sanchez-Rico	2020–07–14	France	7345	Hospital	NA	49.3	8.5	—	—	—	—	91.52	Poor
[229]	Soares	2020–07–16	Brazil	10713	Community and Hospital	NA	55.0	2.0	—	—	—	98.0	0.00	Poor
[230]	Zobairy	2020–07–28	Iran	203	Community and Hospital	49.2 <sup>†</sup> (32–65)	44.8	5.9	—	—	—	94.1	0.00	Poor
[231]	Altamimi	2020–07–27	Qatar	68	Hospital	49 <sup>†</sup> (40–58)	2.0	16.4	—	—	—	83.6	0.00	Poor
[232]	Thompson	2020–07–27	UK	470	Hospital	71 (57–82)	46.0	14.0	27.2	—	58.7	—	0.00	Fair
[233]	Reiter	2020–07–26	Austria	235	Community	44.2 <sup>†</sup> (32–55)	70.0	22.6	22.6	—	54.7	—	0.00	Fair

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[234]	Motta	2020–07–26	USA	374	Hospital	64.7 <sup>†</sup> (46–82)	41.4	—	—	33.2	66.8	—	0.00	Poor
[235]	Santos	2020–07–25	USA	43	Community and Hospital	50 (34–73)	63.0	—	—	4.7	—	—	95.35	Poor
[236]	Schneeweiss	2020–07–22	USA	24313	Community and Hospital	67 <sup>†</sup> (53–80)	53.0	—	—	2.9	—	—	97.12	Poor
[237]	Concha-Mejia	2020–07–24	Colombia	72	Community and Hospital	46 (28–64)	47.0	8.3	11.1	—	—	—	80.56	Poor
[238]	Izquierdo	2020–07–24	Spain	71192	Community and Hospital	42 <sup>†</sup> (18–66)	59.0	10.0	—	—	—	90.0	0.00	Poor
[239]	Bernaola	2020–07–21	Spain	1645	Hospital	NA	38.5	2.5	10.9	—	86.6	—	0.00	Fair
[30]	Islam	2020–08–18	Bangladesh	1016	Community and Hospital	37 (28–49)	35.9	18.2	—	—	—	—	77.85	Poor
[240]	Qi	2020–03–03	China	267	Hospital	48 (35–65)	45.2	19.9	—	—	—	—	80.1	Poor
[241]	Peters	2020–08–15	Netherlands	1893	Hospital	66.8 <sup>†</sup> (52–81)	39.4	4.9	—	—	—	—	95.14	Poor
[242]	Ouyang	2020–08–14	China	217	Hospital	46.5 <sup>†</sup> (30–62)	53.5	16.6	—	—	—	—	83.41	Poor
[47]	Ward	2020–08–21	UK	99908	Community	NA	56.1	10.6	—	—	—	88.4	0.98	Poor <sup>*</sup>
[243]	Valenzuela	2020–08–14	Chile	29	Hospital	56.9 <sup>†</sup> (43–70)	6.9	17.2	—	—	—	82.8	0.00	Poor
[244]	Monteiro	2020–08–14	USA	112	Hospital	61 (45–74)	34.0	6.2	17.9	—	68.8	—	7.14	Fair
[245]	Philipose	2020–08–14	UK	466	Hospital	67 (6–97)	41.8	6.0	73.2	—	16.5	—	4.29	Fair
[246]	Weerahandi	2020–08–14	USA	394	Community	63 (55–70)	37.0	5.3	25.9	—	55.8	—	12.94	Fair
[29]	Ebinger	2020–08–04	USA	6062	Community	41.5 <sup>†</sup> (29–53)	67.8	1.7	—	—	—	—	96.88	Poor

(Continues)

Table 1. (Continued)


Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[247]	Altibi	2020–08–11	USA	706	Hospital	66.7 <sup>†</sup> (51–81)	43.0	4.0	37.3	—	58.8	—	0.00	Fair
[248]	Izzi-Engbeaya	2020–08–11	UK	889	Hospital	65.8 <sup>†</sup> (48–83)	40.0	—	—	21.3	33.2	—	45.6	Poor
[249]	Rizzo	2020–08–11	USA	76819	Hospital	54 (38–67)	55.2	6.7	20.8	—	50.4	—	22.05	Poor
[250]	Dashü	2020–08–11	USA	4140	Community and Hospital	52 (36–65)	55.0	—	—	28.4	51.6	—	19.95	Poor
[251]	Morshed	2020–08–04	Bangladesh	103	Community	37 (31–53)	28.2	31.1	—	—	68.9	—	0.00	Poor
[252]	Jun	2020–08–01	USA	3086	Hospital	66 (56–77)	40.9	3.7	21.3	—	52.8	—	22.23	Poor
[253]	Higuchi	2020–07–30	Japan	57	Hospital	52 (35–70)	43.9	12.3	29.8	—	57.9	—	0.00	Fair
[254]	Zhou, Sun	2020–07–29	China	144	Hospital	47 (38–56)	46.5	9.0	—	—	—	91.0	0.00	Poor
[255]	Salerno	2020–08–22	USA	15920	Hospital	49 (30–65)	57.0	—	—	36.8	55.9	—	7.29	Poor
[256]	Kumar	2020–07–29	India	91	Hospital	47 <sup>†</sup> (41–52)	21.0	44.0	—	—	—	—	56.04	Poor
[257]	Hao	2020–06–01	China	788	Hospital	46 (35–56)	48.4	6.9	—	—	—	—	93.15	Poor
[258]	Iversen	2020–08–03	Denmark	28792	Community and Hospital	44.4 <sup>†</sup> (31–57)	78.9	16.0	6.5	—	76.8	—	0.67	Fair
[259]	Hippisley-Cox	2020–07–13	UK	8275	Community and Hospital	48.5 <sup>†</sup> (30–66)	50.3	17.2	21.4	—	57.3	—	4.04	Fair
[260]	Fillmore	2020–08–24	USA	22914	Community and Hospital	NA	—	37.5	40.7	—	15.5	—	6.38	Fair
[261]	Rashid	2020–08–22	UK	517	Hospital	72.8 <sup>†</sup> (59–86)	31.9	9.9	29.0	—	29.4	—	31.72	Poor
[262]	Pan	2020–08–22	USA	12084	Community and Hospital	45.5 <sup>†</sup> (27–63)	54.3	—	—	17.5	—	—	82.49	Poor

(Continues)

Table 1. (Continued)

Ref.	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %	Never/unknown smokers %	Missing %	Study quality
[263]	Alkurt	2020–08–20	Turkey	932	Community and Hospital	34.8 <sup>†</sup> (25–44)	64.4	24.5	—	—	—	—	75.54	Poor
[264]	Zhao, Chen	2020–07–30	USA	641	Hospital	60 (NA)	40.1	21.7	—	—	—	—	78.32	Poor
[265]	Holman	2020–08–13	UK	10989	Community and Hospital	NA	38.8	5.5	42.6	—	49.0	—	2.82	Fair
[266]	Qu	2020–07–29	China	246	Hospital	53.6 <sup>†</sup> (38–68)	53.3	42.3	—	—	—	—	57.72	Poor
[267]	Chand	2020–08–19	USA	300	Hospital	58.2 <sup>†</sup> (45–70)	39.3	22.3	—	—	—	—	77.67	Poor

NA Age not provided for total sample. - Not reported for total sample.<sup>†</sup>Denotes mean ± standard deviation. This study was rated as 'poor' quality as the manuscript only presents data for current (but not former) smokers despite having obtained complete smoking status, thus resulting in > 20% missing data on smoking status.



**FIGURE 2** (a) Weighted mean prevalence of current smoking in included studies with 95% bootstrap confidence intervals (CIs) compared with national current smoking prevalence (solid red lines), split by country. Shape corresponds to study setting (community, community and hospital, hospital) and shape size corresponds to relative study sample size. (b) Weighted mean prevalence of former smoking in included studies (where this was reported) with 95% bootstrap CIs compared with national former smoking prevalence (solid red lines), split by country. Shape corresponds to study setting (community, community and hospital, hospital) and shape size corresponds to relative study sample size. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

than expected current smoking prevalence was generally observed. Former smoking prevalence was more similar to expected prevalence when reported. National smoking prevalence estimates used for comparison are presented in Supporting information, Table S3.


#### SARS-CoV-2 testing by smoking status

Three studies provided data on access to SARS-CoV-2 diagnostic testing for those meeting local testing criteria by smoking status. In a cohort study of US military veterans aged 54–75 years [31], current smokers were more likely to receive a test: 42.3% (1603 of 3789) of the sample were current smokers compared with 23.8% of all veterans aged 50+ years using any tobacco product between 2010 and 2015 [32]. In the UK Biobank cohort [33], former (RR = 1.29, 95% CI = 1.14–1.45,  $P < 0.001$ ) and current (RR = 1.44, 95% CI = 1.20–1.71,  $P < 0.001$ ) compared

with never smokers were more likely to receive a test in a multivariable analysis. In an Australian rapid assessment screening clinic for COVID-19 [34], 9.4% (397 of 4226) of the self-referred sample (subsequently assessed by a health-care professional to decide on testing) were current smokers. Current compared with former or never smokers were less likely to require a test (RR = 0.93, 95% CI = 0.86–1.0,  $P = 0.045$ ).

#### SARS-CoV-2 infection by smoking status

Forty-five studies provided data on SARS-CoV-2 infection for people meeting local testing criteria by smoking status (see Table 2). Meta-analyses were performed for one ‘good’ and 16 ‘fair’ quality studies (see Figs 3 and 4). Current smokers were at reduced risk of testing positive for SARS-CoV-2 compared with never smokers (RR = 0.74, 95% CrI = 0.58–0.93,  $\tau = 0.41$ , 95% CI = 0.24–0.64). The

**FIGURE 2** Continued.

probability of current smokers being at reduced risk of infection compared with never smokers ( $RR \leq 0.9$ ) was 95%. Former compared with never smokers were at increased risk of testing positive, but data were inconclusive ( $RR = 1.05$ , 95% CrI = 0.95–1.17,  $\tau = 0.17$ , 95% CI = 0.10–0.26) and favoured there being no important association. The probability of former smokers being at increased risk of infection ( $RR \geq 1.1$ ) compared with never smokers was 21%. Results were materially unchanged in the two sensitivity analyses (see Supporting information, Fig. S2).

#### Hospitalization for COVID-19 by smoking status

Twenty-nine studies examined hospitalization for COVID-19 disease stratified by smoking status (see Table 3). Meta-analyses were performed for eight ‘fair’ quality studies (see Figs 5 and 6). Current ( $RR = 1.06$ , CrI = 0.82–1.35,  $\tau = 0.27$ , 95% CI = 0.08–0.55) and former ( $RR = 1.20$ , CrI = 1.03–1.44,  $\tau = 0.17$ , 95% CI = 0.06–0.37) compared with never smokers were at increased risk of hospitalization with COVID-19, but data for current smokers were inconclusive, and favoured there being no important association. The probability of current

and former smokers being at increased risk of hospitalization compared with never smokers was 35 and 89%, respectively. Results were materially unchanged in two sensitivity analyses (see Supporting information, Fig. S3).

#### Disease severity by smoking status

Sixty studies reported disease severity in hospitalized patients stratified by smoking status (see Table 4). Severe (as opposed to non-severe) disease was broadly defined as requiring intensive treatment unit (ITU) admission, requiring oxygen as a hospital inpatient or in-hospital death. Meta-analyses were performed for eight ‘fair’ quality studies (see Figs 7 and 8). Current ( $RR = 1.25$ , CrI = 0.85–1.93,  $\tau = 0.34$ , 95% CI = 0.01–0.86) and former ( $RR = 1.52$ , CrI = 1.13–2.07,  $\tau = 0.29$ , 95% CI = 0.47–0.66) compared with never smokers were at increased risk of greater disease severity; data for current smokers were inconclusive, but favoured there being a small but important association. The probability of current and former smokers having increased risk of greater disease severity compared with never smokers was 79 and 98%, respectively. Results were materially unchanged in two sensitivity analyses (see Supporting information, Fig. S4).

TABLE 2 SARS-CoV-2 infection by smoking status.

Author	SARS-CoV-2-negative						SARS-CoV-2-positive					
	Total population tested	n (%)	Current smoker (%)	Former smoker (%)	Never smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)	Never smoker (%)	Not stated (%)
Rentsch	3528	2974 (84.30%)	1444 (48.55%)	704 (23.67%)	—	826 (27.77%)	—	554 (15.70%)	179 (28.70%)	—	216 (38.99%)	—
Fontanet	661	490 (74.13%)	64 (13.06%)	—	—	426 (86.94%)	—	171 (25.87%)	5 (2.92%)	—	166 (97.08%)	—
Cho	1331	793 (59.58%)	142 (17.91%)	214 (26.99%)	—	437 (55.11%)	—	538 (40.42%)	111 (20.63%)	145 (26.95%)	282 (52.42%)	—
Shah	243	212 (87.24%)	52 (24.53%)	47 (22.17%)	—	113 (53.30%)	—	29 (0.00%)	0 (0.00%)	9 (31.03%)	20 (68.97%)	—
Kölin	1474	805 (54.61%)	141 (17.52%)	307 (38.14%)	—	354 (43.98%)	3 (0.37%)	669 (45.39%)	72 (10.76%)	285 (42.60%)	303 (45.29%)	9 (1.35%)
de Lusignan	3291	2740 (83.26%)	366 (13.36%)	1450 (52.92%)	—	924 (33.72%)	—	551 (16.74%)	47 (8.53%)	303 (54.99%)	201 (36.48%)	—
Valenti	789	689 (87.33%)	197 (28.59%)	—	—	—	492 (71.41%)	40 (5.07%)	7 (17.50%)	—	—	33 (82.50%)
Parrotta	76	39 (51.32%)	10 (2.56%)	27 (25.64%)	—	27 (69.23%)	1 (2.56%)	37 (48.68%)	10 (27.03%)	—	25 (25.75%)	1 (2.70%)
Berumen	102875	71353 (69.36%)	—	7173 (10.05%)	64180 (89.95%)	—	31522 (30.64%)	—	—	2748 (165.23%)	28774 (0.99%)	—
Israel	24906	20755 (83.33%)	3783 (18.23%)	2671 (12.87%)	—	14301 (68.90%)	—	41151 (30.64%)	406 (165.23%)	—	3262 (7.93%)	—
del Valle	1108	143 (12.91%)	27 (18.88%)	53 (37.06%)	—	63 (44.06%)	—	965 (87.09%)	55 (5.70%)	293 (30.36%)	617 (1.17%)	—
Romao	34	20 (58.82%)	—	5 (25.00%)	—	15 (75.00%)	—	14 (75.00%)	—	—	4 (41.18%)	—
Ramlall	111116	4723 (42.49%)	—	—	—	—	—	6393 (57.51%)	—	—	1643.001 (25.70%)	—
Sharma	501	267 (53.29%)	—	—	1 (0.37%)	—	266 (99.63%)	234 (46.71%)	—	20 (8.55%)	4749.999 (74.30%)	214 (91.45%)
Eugen-Olsen	407	290 (71.25%)	76 (26.21%)	104 (35.86%)	—	102 (35.17%)	—	117 (28.75%)	8 (6.84%)	46 (39.32%)	59 (50.43%)	—

(Continues)

Table 2. (Continued)


Author	Total population tested	SARS-CoV-2-negative						SARS-CoV-2-positive					
		n (%)	Current smoker (%)	Former smoker (%)	Never smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Never smoker (%)	Current/former smoker (%)	Former/former smoker (%)	Never smoker (%)
Raisi-Estabragh	4510	3184 (70.60%)	—	—	1653 (51.92%)	—	1531 (48.08%)	1326 (29.40%)	—	—	683 (51.51%)	—	643 (48.49%)
Houlihan	177	97 (54.80%)	14 (14.43%)	14 (14.43%)	—	69 (71.13%)	—	80 (45.20%)	7 (8.75%)	19 (23.75%)	—	54 (67.50%)	—
McQueenie	428199	424355 (99.10%)	—	—	189299 (44.61%)	235056 (55.39%)	—	1311 (0.31%)	—	—	669 (51.03%)	642 (48.97%)	—
Woolford	4474	3161 (70.65%)	441 (13.95%)	1194 (37.77%)	—	1526 (48.28%)	—	1313 (29.35%)	145 (11.04%)	525 (39.98%)	—	643 (48.97%)	—
Lan	104	83 (79.81%)	—	—	24 (28.92%)	—	59 (71.08%)	21 (20.19%)	—	—	1 (4.76%)	—	20 (95.24%)
Hernandez-Garduno	32583	20279 (62.24%)	—	—	2399 (11.83%)	17861 (88.08%)	—	12304 (37.76%)	—	—	1191 (9.68%)	11083 (90.08%)	—
Govind	6215	6207 (99.87%)	4104 (66.12%)	1669 (26.89%)	—	342 (5.51%)	—	102 (1.64%)	78 (7.647%)	20 (19.61%)	—	2 (1.96%)	—
Gu	4699	3815 (81.19%)	360 (9.44%)	1142 (29.93%)	—	2313 (60.65%)	—	884 (18.81%)	40 (4.52%)	264 (29.86%)	—	580 (65.61%)	—
Kibler	702	680 (96.87%)	25 (3.68%)	—	—	655 (96.32%)	22 (3.13%)	1 (4.55%)	—	—	—	—	21 (95.45%)
Petrilli	10620	5341 (50.29%)	3454 (64.67%)	816 (15.28%)	—	541 (10.13%)	(9.92%)	5279 (49.71%)	3268 (61.91%)	902 (17.09%)	—	288 (54.6%)	821 (15.55%)
Bello-Chavolla	150200	98567 (65.62%)	—	—	9624 (9.76%)	—	88943 (90.24%)	51633 (34.38%)	—	4366 (17.09%)	—	47267 (91.54%)	—
Auvinen	61	33 (54.10%)	10 (30.30%)	8 (24.24%)	—	15 (45.45%)	—	28 (45.90%)	1 (3.57%)	9 (32.14%)	—	18 (64.29%)	—
Favara	70	55 (78.57%)	5 (9.09%)	—	—	—	—	50 (90.91%)	15 (21.43%)	2 (13.33%)	—	—	13 (86.67%)
Antonio-Villa	34263	23338 (68.11%)	2293 (9.83%)	—	—	—	—	21045 (90.17%)	10925 (31.89%)	1023 (9.36%)	—	—	9902 (90.64%)
Merzon	7807	7025 (89.98%)	—	—	1136 (16.17%)	—	5889 (83.83%)	782 (10.02%)	—	127 (16.24%)	—	655 (83.76%)	—

(Continues)


Table 2. (Continued)

Author	Total population tested	SARS-CoV-2-negative						SARS-CoV-2-positive					
		n (%)	Current smoker (%)	Former smoker (%)	Never smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Never smoker (%)	Current/former smoker (%)	Former smoker (%)	Never smoker (%)
Trubiano	2935	2827 (96.66%)	—	—	256 (9.06%)	—	2586 (91.48%)	108 (3.68%)	—	—	3 (2.78%)	—	105 (97.22%)
Shi,	1521	1265	—	—	681 (53.83%)	—	584 (46.17%)	256 (16.83%)	—	—	154 (60.16%)	—	102 (39.84%)
Resurreccion		(83.17%)					19914 (16.55%)	97953 (81.32%)	159 (0.13%)	3 (1.89%)	—	—	17 (10.69%)
Riley	120620	120461 (99.87%)	2594 (2.15%)	—	—	—	196	123	—	—	1 (0.81%)	—	139 (87.42%)
Alizadehsani	319	196	—	—	—	—	—	—	—	—	—	—	122 (99.19%)
Merkely	10474	10336 (61.44%)	2904 (28.10%)	2107 (20.39%)	—	5310 (51.37%)	15 (0.15%)	70 (0.67%)	16 (22.86%)	15 (21.43%)	—	—	38 (54.29%)
McGrail	209	118 (56.46%)	—	31 (26.27%)	—	87	91 (73.73%)	—	—	—	8 (8.79%)	—	83 (91.21%)
Izquierdo	71192	NA	—	—	—	—	—	1006 (43.54%)	111 (11.03%)	111 (11.03%)	—	—	895 (88.97%)
Ward	99908	94416 (94.50%)	10202 (10.81%)	—	—	—	84214 (89.19%)	5492 (5.50%)	433 (7.88%)	—	—	—	5059 (92.12%)
Ebinger	60622	5850 (96.50%)	99 (1.69%)	—	—	—	5668 (96.89%)	212 (3.50%)	3 (1.42%)	—	—	—	205 (96.70%)
Salerno	15920	14753 (92.67%)	—	5517 (37.40%)	8278 (56.11%)	958 (6.49%)	1167 (7.33%)	—	—	—	339 (29.05%)	626 (53.64%)	202 (17.31%)
Iversen	28792	27629 (95.96%)	4430 (16.03%)	1799 (6.51%)	—	21217 (76.79%)	246 (0.89%)	1163 (4.04%)	177 (15.22%)	78 (6.71%)	—	898 (77.21%)	10 (0.86%)
Hippisley-Cox	8275949	NA	—	—	—	—	—	19486 (0.24%)	1354 (6.95%)	5715 (29.33%)	—	12036 (61.77%)	381 (1.96%)
Fillmore	22914	21120 (92.17%)	8137 (38.53%)	8416 (39.85%)	—	3227 (15.28%)	1340 (6.34%)	1794 (7.83%)	452 (25.20%)	899 (50.11%)	—	322 (17.95%)	121 (6.74%)
Alkurt	119	NA	—	—	—	—	—	119 (100.00%)	14 (11.76%)	—	—	—	105 (88.24%)

Niedzwiedz et al. reported on SARS-CoV-2 infection by smoking status in multivariable analyses but did not present raw data. NA = not available.



**FIGURE 3** Forest plot for risk of testing positive for SARS-CoV-2 in current versus never smokers. \*This was a 'good' quality study. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 4** Forest plot for risk of testing positive for SARS-CoV-2 in former versus never smokers. \*This was a 'good' quality study. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

TABLE 3 Hospitalization with COVID-19 by smoking status.


Author	Population with outcome	Community						Hospitalized					
		Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)
Rentsch	554	269 (48%)	69 (25.65%)	90 (33.46%)	—	110 (40.89%)	—	—	285 (51%)	90 (31.58%)	89 (31.23%)	—	106 (37.19%)
Chow (US CDC)	6637 (77%)	5143 (1.19%)	61 (1.56%)	80 (1.19%)	—	—	—	5002 (97.24%)	1494 (22%)	27 (1.81%)	78 (5.22%)	—	— (92.97%)
Argenziano	1000	151 (15%)	14 (9.27%)	18 (11.92%)	—	119 (78.81%)	—	—	849 (84%)	35 (4.12%)	161 (18.96%)	—	653 (76.91%)
Lubetzy	54	15 (27%)	—	4 (26.67%)	—	—	—	11 (73.33%)	39 (72%)	—	8 (20.51%)	—	— (79.49%)
Carillo-Vega	9946	3922 (39%)	408 (10.40%)	—	—	—	—	3514 (89.60%)	6024 (60%)	486 (8.07%)	—	—	— (91.93%)
Yanover	4353	4180 (96%)	484 (11.58%)	118 (2.82%)	—	3578 (85.60%)	—	—	173 (3%)	30 (17.34%)	11 (6.36%)	—	132 (76.30%)
Hamer	387109	386 (9.66%)	37333 (9.66%)	134542 (34.82%)	—	214474 (55.51%)	—	—	760 (0%)	93 (12.24%)	313 (41.18%)	—	354 (46.58%)
Heli-Frades	4712	1973 (41%)	121 (6.13%)	222 (11.25%)	—	—	1630 (82.62%)	1630 (82.62%)	2739 (58%)	112 (4.09%)	598 (21.83%)	—	2029 (74.08%)
Freites	123	69 (56%)	1 (1.45%)	—	—	—	—	68 (98.55%)	54 (43%)	3 (5.56%)	—	—	— (94.44%)
Berumen	102875	18832 (18%)	—	1546 (8.21%)	—	17286 (91.79%)	—	12 (690)	—	—	1202 (12%)	—	11488 (90.53%)
Gianfrancesco	600	323 (53%)	—	—	61 (18.89%)	—	—	262 (8.11%)	277 (46%)	—	68 (24.55%)	—	209 (75.45%)
Chaudhry	40	19 (47%)	—	—	0 (0.00%)	—	—	19 (100.00%)	21 (52%)	—	6 (28.57%)	—	15 (71.43%)
Giannouchos	89756	58485 (65%)	4679 (8.00%)	—	—	53806 (92.00%)	—	31 (34%)	2721 (8.70%)	—	—	28550 (91.30%)	—
Wang, Oekelen	57	22 (38%)	—	—	6 (27.27%)	—	—	16 (72.73%)	36 (63%)	—	15 (41.67%)	—	20 (55.56%)

(Continues)


Table 3. (Continued)

Author	Population with outcome	Community						Hospitalized					
		n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)
Miyata	470	132 (28%)	14 (10.61%)	41 (31.06%)	—	77 (58.33%)	—	—	338 (71%)	18 (53.33%)	111 (32.84%)	—	209 (61.83%)
Suleyman	463	108 (23%)	—	—	23 (21.30%)	—	—	85 (78.70%)	355 (76%)	—	137 (38.59%)	—	— (61.41%)
Garassino	196	48 (24%)	10 (20.83%)	27 (56.25%)	—	11 (22.92%)	—	—	152 (77%)	38 (25.00%)	84 (55.26%)	—	26 (17.11%)
Siso-Almirall	260	119 (45%)	—	—	31 (26.05%)	—	—	88 (73.95%)	141 (54%)	—	50 (35.46%)	—	— (64.54%)
Gu	884	511 (57%)	30 (5.87%)	126 (24.66%)	—	355 (69.47%)	—	—	373 (42%)	10 (2.68%)	138 (37.00%)	—	225 (60.32%)
Killerby	531	311 (58%)	—	—	37 (11.90%)	222 (71.38%)	—	52 (16.72%)	220 (41%)	—	54 (24.55%)	157 (71.36%)	— (4.09%)
Petrilli	5279	2538 (48%)	147 (5.79%)	337 (13.28%)	—	1678 (66.12%)	—	376 (14.81%)	2741 (51%)	141 (5.14%)	565 (20.61%)	—	1590 (58.01%)
Nguyen	689	333 (48%)	—	—	57 (17.12%)	—	—	276 (82.88%)	356 (51%)	—	114 (32.02%)	—	— (16.23%)
Mendy	689	473 (68%)	—	—	84 (17.76%)	—	—	389 (82.24%)	216 (31%)	—	86 (39.81%)	—	— (44.5%)
Soares	10713	9561 (89%)	132 (1.38%)	—	—	—	9429 (98.62%)	—	1152 (10%)	77 (6.68%)	—	—	1075 (93.32%)
Zobairy	203	65 (32%)	1 (1.54%)	—	—	—	64 (98.46%)	—	138 (6.7%)	11 (7.97%)	—	—	127 (92.03%)
Izquierdo	1006	743 (73%)	52 (7.00%)	—	—	—	691 (93.00%)	—	263 (26%)	16 (6.08%)	—	—	247 (93.92%)
Rizzo	76819	60039 (78%)	3931 (6.55%)	11379 (18.95%)	—	30042 (50.04%)	—	14687 (24.46%)	16 (2.1%)	1254 (7.47%)	4585 (27.32%)	—	8693 (51.81%)
Dashfi	4140	2759 (66%)	—	—	600 (21.75%)	1541 (55.85%)	—	618 (22.40%)	1381 (33%)	—	577 (41.78%)	—	596 (43.16%)
Pan	12084	8548 (70%)	—	—	1263 (14.78%)	—	—	7285 (85.22%)	3536 (29%)	—	874 (24.72%)	—	— (75.28%)

NA = not available; CDC = Centers for Disease Control



**FIGURE 5** Forest plot for risk of hospitalization in current versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 6** Forest plot for risk of hospitalization in former versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

TABLE 4 Disease severity by smoking status.

Author	Population with severity	Non-severe disease						Severe disease					
		n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Not stated (%)
Guan, Ni	1085	913 (84%)	108 (11.83%)	12 (1.31%)	—	793 (86.86%)	—	—	172 (15%)	29 (16.86%)	9 (5.23%)	—	134 (77.91%)
Zhang, Dong	9	3 (0.00%)	0 (0.00%)	3 (100.00%)	—	0 (0.00%)	—	—	6 (66%)	2 (33.33%)	4 (66.67%)	—	0 (0.00%)
Wan	9	8 (88%)	8 (100.00%)	0 (0.00%)	—	0 (0.00%)	—	—	1 (11%)	1 (100.00%)	0 (0.00%)	—	0 (0.00%)
Huang, Wang	3	3 (100%)	0 (100.00%)	0 (0.00%)	—	0 (0.00%)	—	—	0 (0%)	0 (0%)	0 (0%)	—	0 (0%)
Rentsch	285	168 (58%)	47 (27.98%)	53 (31.55%)	—	68 (40.48%)	—	—	117 (41%)	43 (36.75%)	36 (30.77%)	—	38 (32.48%)
Hu	323	151 (46%)	—	—	12 (7.95%)	—	139 (92.05%)	—	172 (53%)	—	26 (15.12%)	—	146 (84.88%)
Wang, Pan	125	100 (80%)	—	—	9 (9.00%)	—	91 (91.00%)	—	25 (20%)	—	7 (28.00%)	—	18 (72.00%)
Kim	27	21 (77%)	3 (14.29%)	—	—	—	18 (85.71%)	—	6 (22%)	2 (33.33%)	0 (0.00%)	—	4 (66.67%)
Shi, Yu	474	425 (89%)	—	—	34 (8.00%)	—	391 (92.00%)	—	49 (10%)	—	6 (12.24%)	—	43 (87.76%)
Liao, Feng	148	92 (62%)	—	—	5 (5.43%)	—	—	87 (94.57%)	56 (37%)	3 (5.36%)	—	—	—
Shi, Ren	134	88 (65%)	—	—	8 (9.09%)	—	—	80 (90.91%)	46 (34%)	—	6 (13.04%)	—	40 (86.96%)
Hadjadj	50	15 (30%)	1 (6.67%)	2 (13.33%)	—	12 (80.00%)	—	—	35 (70%)	0 (0.00%)	7 (20.00%)	—	—
Zheng, Xiong	73	43 (58%)	—	6 (13.95%)	37 (86.05%)	—	—	30 (41%)	—	—	2 (6.67%)	28 (93.33%)	—
de la Rica	48	26 (54%)	—	—	6 (23.08%)	—	—	20 (41%)	—	—	4 (20.00%)	—	16 (80.00%)
Yin, Yang	106	47 (44%)	—	—	6 (12.77%)	—	—	41 (59%)	—	—	12 (20.34%)	—	47 (79.66%)
Allenbach	147	100 (68%)	—	—	9 (9.00%)	—	—	91 (47%)	—	—	0 (0.00%)	—	47 (100.00%)

(Continues)

Table 4. (Continued)

(Continues)


Table 4. (Continued)

Author	Population with severity	Non-severe disease						Severe disease					
		n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)
Giannouchos	89 756	78 050	6322 (8.10%)	—	—	71 728 (91.90%)	—	11 706 (9.30%)	—	—	—	10 617 (90.70%)	—
Cen	1007	720 (71%)	—	—	70 (9.72%)	—	650 (90.28%)	287 (28%)	—	18 (6.27%)	—	—	269 (93.73%)
Maraschini	132	89 (67%)	—	11 (12.35%)	—	78 (87.64%)	—	43 (32%)	—	3 (6.98%)	—	40 (93.02%)	—
Siso-Almirall	260	212 (81%)	—	—	60 (28.30%)	—	152 (71.70%)	48 (18%)	—	—	21 (43.75%)	—	—
Gu	884	511 (57%)	30 (5.87%)	126 (24.66%)	—	355 (69.47%)	—	134 (15%)	3 (2.24%)	61 (45.52%)	—	70 (52.24%)	—
Petrilli	2729	1739 (63%)	97 (5.58%)	325 (18.69%)	—	1067 (61.36%)	—	250 (14.38%)	990 (36%)	44 (4.44%)	—	236 (23.84%)	—
Mendy	689	598 (86%)	—	—	133 (22.24%)	—	—	465 (77.76%)	91 (13%)	—	37 (40.66%)	—	517 (52.22%)
Pongpirul	193	161 (83%)	—	—	25 (15.53%)	106 (65.84%)	—	30 (18.63%)	32 (16%)	—	4 (41.25%)	21 (65.62%)	—
Jin, Gu	6	2 (33%)	—	—	0 (0.00%)	—	—	4 (200.00%)	4 (66%)	—	2 (50.00%)	—	—
Senkal	611	446 (73%)	48 (10.76%)	—	—	—	—	398 (89.24%)	165 (27%)	21 (12.73%)	—	—	—
Patel	129	89 (68%)	26 (29.21%)	—	—	58 (65.17%)	—	5 (5.62%)	40 (40%)	22 (11.76%)	—	—	—
Maucourtant	27	10 (37%)	1 (10.00%)	2 (20.00%)	—	2 (20.00%)	—	5 (50.00%)	17 (31%)	2 (55.00%)	—	—	—
Xie	619	469 (75%)	—	—	32 (6.82%)	—	—	5 (62%)	437 (42%)	—	19 (12.67%)	—	—
Fox	55	30 (54%)	1 (3.33%)	4 (13.33%)	—	17 (56.67%)	—	8 (26.67%)	25 (93.18%)	0 (24%)	—	14 (52.94%)	—
Zhang, Cao	240	162 (67%)	2 (1.23%)	6 (3.70%)	—	—	—	154 (45%)	78 (45.13%)	4 (8.00%)	—	14 (56.00%)	—
								(95.06%)	(32%)	(5.13%)	—	—	—


(Continues)

Table 4. (Continued)

Author	Population with severity	Non-severe disease						Severe disease						
		n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Not stated	n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Not stated	
Kurashima	53	10 (18%)	—	—	3 (30.00%)	—	—	7 (70.00%)	43 (81%)	—	—	24 (55.81%)	—	
Zhan	75	NA	—	—	—	—	—	75 (100%)	52 (6%)	—	—	9 (12.00%)	—	
Omraní	858	806 (93%)	—	—	121 (15.01%)	—	—	685 (84.99%)	18 (6%)	—	9 (17.31%)	—	—	
Marcos	918	555 (60%)	38 (6.85%)	—	69 (12.43%)	—	—	448 (80.72%)	363 (39%)	18 (4.96%)	—	71 (19.56%)	—	—
Hoertel, Sanchez- Rico	7345	6014 (81%)	433 (7.29%)	—	—	—	—	5581 (92.80%)	1331 (18%)	190 (14.27%)	—	—	—	—
Qi	267	217 (81%)	22 (10.14%)	—	—	195 (89.86%)	—	50 (18%)	31 (62.00%)	—	—	19 (38.00%)	—	—
Monteiro	112	84 (75%)	3 (3.57%)	14 (16.67%)	—	63 (75.00%)	—	4 (4.76%)	28 (25%)	4 (14.29%)	6 (21.43%)	14 (50.00%)	—	4 (14.29%)
Dashti	1381	619 (44%)	—	—	239 (38.61%)	292 (47.17%)	—	88 (14.22%)	762 (55%)	—	338 (44.36%)	304 (39.90%)	—	120 (15.75%)
Morshed	103	87 (84%)	28 (32.18%)	—	—	59 (67.82%)	—	16 (15%)	4 (25.00%)	—	—	12 (75.00%)	—	—
Zhou, Sun	144	108 (75%)	11 (10.19%)	—	—	—	—	97 (89.81%)	36 (25%)	2 (5.56%)	—	—	—	34 (94.44%)
Hippisley- Cox	—	NA	—	—	—	—	—	—	1286 (4.35%)	56 (33.20%)	427 (61.51%)	791 (0.93%)	—	12 (0.93%)
Zhao, Chen	641	398 (62%)	87 (21.86%)	—	—	—	—	311 (78.14%)	195 (30%)	52 (26.67%)	—	—	—	143 (73.33%)
Qu	246	226 (91%)	90 (39.82%)	—	—	—	—	136 (60.18%)	20 (8%)	14 (70.00%)	—	—	—	6 (30.00%)



**FIGURE 7** Forest plot for the risk of severe disease in current versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 8** Forest plot for the risk of severe disease in former versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

### Mortality by smoking status

Fifty studies reported mortality from COVID-19 by smoking status (see Table 5), with nine 'fair' quality studies included in meta-analyses (see Figs 9 and 10). Current (RR = 1.22, 95% CrI = 0.78–1.94,  $\tau$  = 0.49, 95% CI = 0.16–0.99) and former (RR = 1.39, 95% CrI = 1.09–1.87,  $\tau$  = 0.27, 95% CI = 0.05–0.58) compared with never smokers were at increased risk of in-hospital mortality from COVID-19. Data for current smokers were inconclusive, but favoured there being no important association. The probability of current and former smokers being at greater risk of in-hospital mortality compared with never smokers was 70 and 97%, respectively. Results were materially unchanged in two sensitivity analyses (see Supporting information, Fig. S5).

## DISCUSSION

This living rapid review found uncertainty in the majority of 233 studies arising from the recording of smoking status. Notwithstanding these uncertainties, compared with overall adult national prevalence estimates, recorded current smoking rates in most countries were lower than expected. In a subset of better quality studies ( $n = 17$ ), current smokers had a reduced risk of testing positive for SARS-CoV-2 but appeared more likely to present for testing and/or receive a test. Data for current smokers on the risk of hospitalization, disease severity and mortality were inconclusive, but favoured there being no important associations with hospitalization and mortality and a small but important increase in the risk of severe disease. Former smokers were at increased risk of hospitalization, disease severity and mortality compared with never smokers.

### Issues complicating interpretation

Interpretation of results from studies conducted during the first phase of the SARS-CoV-2 pandemic is complicated by several factors (see Fig. 11):

1. Exposure to SARS-CoV-2 is heterogeneous, with different subgroups at heightened risk of infection at different stages of the pandemic. This will likely introduce bias in studies assessing the rate of infection by smoking status conducted early on.
2. Current and former smokers may be more likely to meet local criteria for community testing due to increased prevalence of symptoms consistent with SARS-CoV-2 infection, such as cough, increased sputum production or altered sense of smell or taste [35]. Evidence from a small number of studies indicates that current smokers may be more likely to present for testing, hence increasing the denominator in comparisons with never smokers and potentially inflating the rate of negative tests in current smokers. Infection positivity rates estimated among random samples will be more informative than currently available data. We identified one population study conducted in Hungary reporting on seroprevalence and smoking status [36]; however, the response rate was fairly low, at 58.8%, and the current smoking rate was 10 percentage points below national prevalence estimates, thus questioning the representativeness of the final sample. Smoking status is being collected in at least two large representative infection and antibody surveys in the United Kingdom [37,38].
3. Testing for acute infection requires swabbing of the mucosal epithelium, which may be disrupted in current smokers, potentially altering the sensitivity of assays [39].
4. Diagnostic criteria for SARS-CoV-2 infection and COVID-19 have changed during the course of the pandemic [40]. It was not possible to extract details on the specific RT-PCR technique or platforms used across the included studies due to reporting gaps. Different platforms have varying sensitivity and specificity to detect SARS-CoV-2 infection.
5. Most included studies relied on EHRs as the source of information on smoking status. Research shows large discrepancies between EHRs and actual behaviour [41]. Known failings of EHRs include implausible longitudinal changes, such as former smokers being recorded as never smokers at subsequent hospital visits [41]. Misreporting on the part of the patient (perhaps due to perceived stigmatization) has also been observed, with biochemical measures showing higher rates of smoking compared with self-report in hospitalized patients in the United States [42]. It is hence possible that under-reporting of current and former smoking status in hospitals occurred across the included studies.
6. Individuals with severe COVID-19 symptoms may have stopped smoking immediately before admission to hospital and may therefore not have been recorded as current smokers (i.e. reverse causality).
7. Smokers with COVID-19 may be less likely to receive a SARS-CoV-2 test or present to hospital due to lack of access to healthcare, and may be more likely to die in the community from sudden complications (i.e. self-selection bias) and thus not be recorded.
8. If there is a protective effect of nicotine on COVID-19 disease outcomes, abrupt nicotine withdrawal upon hospitalization may lead to worse outcomes [12].
9. During periods of heightened demand of limited healthcare resources, current and former smokers with extensive comorbidities may have reduced priority for intensive care admission, thus leading to higher in-hospital mortality.

TABLE 5 Mortality by smoking status.

Author	Population with mortality	Recovered						Died					
		Current smoker n (%)	Former smoker n (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker n (%)	Not stated n (%)	Current/ former smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker n (%)	Not stated n (%)
Chen	274	161 (58%)	5 (3.11%)	5 (3.11%)	—	—	—	151 (93.79%)	113 (41%)	7 (6.19%)	2 (1.77%)	—	—
Zhou, Yu	191	137 (71%)	6 (4.38%)	—	—	—	—	131 (95.62%)	54 (28%)	5 (9.26%)	—	—	49 (90.74%)
Yang, Yu	52	20 (79%)	2 (38%)	—	—	—	18 (90.00%)	—	32 (61%)	—	—	—	—
Borobia	2226	1766 (79%)	113 (6.40%)	—	—	—	—	1653 (93.60%)	460 (20%)	44 (9.57%)	—	—	416 (90.43%)
Giacomelli	233	185 (79%)	—	53 (28.65%)	132 (71.35%)	—	—	—	48 (20%)	—	17 (35.42%)	31 (64.58%)	—
Yao	108	96 (88%)	1 (1.04%)	—	—	—	—	95 (98.96%)	12 (11%)	3 (25.00%)	—	—	9 (0.00%)
Carillo-Vega	9946	8983 (90%)	795 (8.85%)	—	—	—	—	8188 (91.15%)	963 (9%)	99 (10.28%)	—	—	864 (89.72%)
Heng	51	39 (76%)	6 (15.38%)	—	—	—	—	33 (84.62%)	12 (23%)	1 (8.33%)	—	—	11 (91.67%)
Chen, Jiang	135	NA	—	—	—	—	—	—	31 (22%)	—	4 (12.90%)	—	—
Heili-Frades	4712	4086 (86%)	210 (5.14%)	659 (16.13%)	—	—	3217 (78.73%)	—	626 (13%)	23 (3.67%)	161 (25.72%)	—	442 (70.61%)
Kim, Garg	2490	2070 (83%)	128 (6.18%)	481 (23.24%)	—	—	1461 (70.58%)	—	420 (16%)	22 (5.24%)	161 (38.33%)	—	236 (56.19%)
Al-Hindawi	31	15 (48%)	0 (50%)	10 (0.00%)	—	5 (66.67%)	—	—	16 (51%)	1 (6.25%)	12 (75.00%)	—	—
Louis	22	16 (72%)	—	7 (43.75%)	—	—	9 (33.33%)	—	6 (51%)	—	—	3 (18.75%)	—
Soto-Mota	400	200 (50%)	—	23 (11.50%)	—	—	—	—	—	—	3 (50.00%)	—	3 (50.00%)
Garibaldi	747	634 (84%)	36 (5.68%)	129 (20.35%)	—	—	—	469 (88.50%)	113 (50%)	6 (46.90%)	36 (87.50%)	—	71 (17.50%)
								(73.97%)	(15%)	(5.31%)	(31.86%)		(62.83%)

(Continues)

Table 5. (Continued)

Author	Population with mortality	Recovered						Died						
		Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	Not stated (%)	Current smoker (%)	Former smoker (%)	Former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	Not stated (%)	
Docherty	13 364	8199 (61%)	370 (4.51%)	1832 (22.34%)	— (50.97%)	4179 (52.06%)	— (22.17%)	1818 (3.84%)	5165 (4.13%)	214 (4.14%)	1350 (26.14%)	— (40.76%)	2105 (40.76%)	1496 (28.96%)
Kuderer	928	807 (86%)	38 (4.71%)	262 (32.47%)	— (26.40%)	425 (73.60%)	— (73.60%)	31 —	121 —	5 64	— —	44 —	— —	2 (1.65%)
Ramlal	11116	10498 (94%)	— —	2771 —	— (14.32.56%)	7727 —	— —	— 29	618 14	— —	208 7 (50.00%)	— —	410 7 (50.00%)	— —
Wang, Okele Martinez- Portilla Cen	57 (75%) 224 (96%) 1007 (95%)	— — — —	— — — —	7 (3.23%) — 87 (9.02%) —	— — — —	— — — —	— — — —	210 877	7 (3%) 43	— —	— —	0 (0.00%) 1 (2.33%)	— —	7 (50.00%) 7 (100.00%) 42 (97.67%)
Klang	3406 (66%)	2270 —	— —	492 (21.67%)	— —	— —	— —	1778 (78.33%)	1136 (33%)	— —	— —	301 (26.50%)	— —	835 (73.50%)
Wang, Zhong Miyara	5510 (88%) 338 (62%)	4874 (5.07%) 211 (6.16%)	247 13 —	1083 (22.22%) 58 (27.49%)	— — 28 (13.40%)	3544 (72.71%) 141 (66.82%)	— — —	— 636	28 (11%)	214 (4.40%)	— (33.65%)	— —	394 (61.95%)	— —
Rajier	255 (81%)	209 —	— —	— 69 (7.97%)	— —	— 181	— —	— 53	— —	— —	— 18 (33.96%)	— 28 (52.83%)	— —	— —
Zeng	1031 (84%)	866 —	— —	— 69 (7.97%)	— —	— 797	— 165	— —	— —	— —	36 (21.82%)	— —	129 (78.18%)	— —
Chen, Yu	1859	1651 (88%)	32 (1.94%)	54 (3.27%)	— —	1565 (94.79%)	— —	— 66	208 —	13 61	12 —	— —	183 (87.98%)	— —
Garassino	190	124 (65%)	— —	— —	92 (74.19%) —	32 (25.81%)	— —	— (34%)	— —	— —	— —	5 —	— —	— —
Gu	884	864 (97%)	40 (4.63%)	250 (28.94%)	— —	219 (25.35%)	— —	— —	20 —	0 —	14 —	— —	6 —	— —
Sigel	88	70 (79%)	— —	— 37 (52.86%)	— —	— —	— —	33 —	18 —	— —	11 (61.11%)	— —	7 (38.89%)	— —

(Continues)

Table 5. (Continued)

Author	Population with mortality	Recovered						Died					
		n (%)	Current smoker (%)	Former smoker (%)	Current/ former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)	Not stated (%)	n (%)	Current smoker (%)	Former smoker (%)	Never/ unknown smoker (%)	Never/ unknown smoker (%)
Nguyen	356	308 (86%)	—	—	91 (29.55%)	—	—	217 (70.45%)	45 (12%)	—	—	23 (51.11%)	—
de Souza	8443	7826 (92%)	—	—	95 (1.21%)	—	7571 (96.74%)	160 (2.04%)	617 (7%)	—	47 (7.62%)	—	560 (90.75%)
Mendy	532	663 (124%)	—	—	160 (24.13%)	—	—	502 (75.72%)	26 (4%)	—	—	10 (38.46%)	—
Shi,	256	210 (82%)	—	—	128 (60.95%)	—	—	82 (39.05%)	46 (17%)	—	—	26 (56.52%)	—
Resurreccion	619	591 (95%)	—	—	43 (7.28%)	—	—	548 (92.72%)	28 (4%)	—	—	8 (28.57%)	—
Fox	54	35 (64%)	1 (2.86%)	4 (11.43%)	—	18 (51.43%)	—	12 (34.29%)	19 (35%)	0 (0.00%)	2 (10.53%)	—	12 (63.16%)
Zhang, Cao	289	240 (83%)	10 (4.17%)	6 (2.50%)	—	—	—	224 (93.33%)	49 (16%)	4 (8.16%)	8 (16.33%)	—	—
Gupta	496	255 (51%)	—	—	15 (5.88%)	—	80 (31.37%)	160 (62.75%)	241 (48%)	—	—	21 (8.71%)	77 (31.95%)
Soares	1075	696 (64%)	38 (5.46%)	—	—	—	658 (94.54%)	—	456 (42%)	39 (8.55%)	—	—	—
Thompson	470	301 (64%)	39 (12.96%)	79 (26.25%)	—	183 (60.80%)	—	169 (35%)	27 (15.98%)	49 (28.99%)	—	93 (55.03%)	—
Bernaola	1645	1382 (84%)	35 (2.53%)	146 (10.56%)	—	1201 (86.90%)	—	—	263 (15%)	6 (2.28%)	33 (12.55%)	—	218 (82.89%)
Islam	654	631 (96%)	103 (16.32%)	—	—	—	—	507 (80.35%)	23 (3%)	3 (13.04%)	—	—	—
Philipose	466	267 (57%)	19 (7.12%)	204 (76.40%)	—	44 (16.48%)	—	—	199 (42%)	9 (4.52%)	137 (68.84%)	—	33 (16.58%)
Dashti	4140	3953 (95%)	—	—	1068 (27.02%)	2078 (52.57%)	—	804 (20.34%)	187 (4%)	—	—	109 (58.29%)	56 (29.95%)
Filmore	1794	1566 (87%)	408 (26.05%)	758 (48.40%)	—	279 (17.82%)	—	98 (6.26%)	228 (12%)	44 (19.30%)	141 (61.84%)	—	43 (18.86%)

(Continues)

Table 5. (Continued)

Author	Population with mortality	Recovered				Died			
		Current smoker n (%)	Former smoker n (%)	Current/ former smoker n (%)	Never/ unknown smoker n (%)	Not stated n (%)	Current smoker n (%)	Former smoker n (%)	Current/ former smoker n (%)
Pan	3536	3302 (93%)	—	862 (26.11%)	—	—	2440 (73.89%)	234 (6%)	—
Zhao, Chen	474	398 (83%)	87 (21.86%)	—	—	311 (78.14%)	82 (17%)	36 (43.90%)	—
Holman	10989	NA	—	—	—	—	10989 (100%)	609 (5.54%)	4684 (42.62%)
Chand	300	143 (47%)	23 (16.08%)	—	—	—	120 (83.92%)	44 (52%)	—
							(28.03%)	(28.03%)	(71.97%)

Solis *et al.* and the OpenSAFELY Collaborative reported on mortality by smoking status in a multivariable analysis but did not present raw data for both the exposure and outcome variables.

10. Given the lack of knowledge of the disease progression and long-term outcomes of COVID-19, it is unclear whether studies conducted thus far in the pandemic have monitored patients for a sufficient time-period to report complete survival outcomes or whether they are subject to early censoring.

11. Reasons for hospitalization vary by country and time in the pandemic. For example, early cases may have been hospitalized for isolation and quarantine reasons and not due to medical necessity. It is plausible that this may have skewed early data towards less severe cases. In addition, the observed association between former smoking and greater disease severity may be explained by collider bias [43], where conditioning on a collider (e.g. testing or hospitalization) by design or analysis may introduce a spurious association between current or former smoking (a potential cause of testing or hospitalization) and SARS-CoV-2 infection/adverse outcomes from COVID-19 (potentially exacerbated by smoking) [44].

### Limitations


This living rapid evidence review was limited by having a single reviewer extracting data with a second independently verifying the data extracted to minimize errors, restricting the search to one electronic database and one pre-print server and by not including at least three large population surveys due to their reliance upon self-reported suspected or confirmed SARS-CoV-2 infection (which means they do not meet our eligibility criteria) [35,45,46]. We also did not include a large, UK-based, representative seroprevalence study [47] in our meta-analyses, as the odds of testing positive in former smokers was not reported. However, the odds of infection for current smokers (odds ratio = 0.64, 95% CI = 0.58–0.71) was in concordance with the pooled estimate in our meta-analysis. Population surveys—particularly with linked data on confirmed infection or antibodies—will be included in future review versions to help mitigate some of the limitations of healthcare based observational studies. The comparisons of current and former smoking prevalence in the included studies with national prevalence estimates did not adjust observed prevalence for the demographic profile of those tested/admitted to hospital. Other reviews focused on this comparison have applied adjustments for sex and age, and continue to find lower than expected prevalence—notwithstanding the issues complicating interpretation described above [17].

### Implications for research, policy and practice


Further scientific research is needed to resolve the mixed findings summarized in our review. First, clinical trials of



**FIGURE 9** Forest plot for the risk of mortality in current versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 10** Forest plot for the risk of mortality in former versus never smokers. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



**FIGURE 11** A schematic of some of the interpretation issues for the association of smoking and SARS-CoV-2/COVID-19. \*Indicates potential confounding with smoking status.

the posited therapeutic effect of nicotine could have important implications both for smokers and for improved understanding of how the SARS-CoV-2 virus causes disease in humans. Such trials should focus upon medicinal nicotine (as smoked tobacco is a dirty delivery mechanism that could mask beneficial effects) and potentially differentiate between different modes of delivery (i.e. inhaled versus ingested), as this can affect pharmacokinetics [48] and potential therapeutic effects. A second research priority would be a large, representative (randomly sampled) population survey with a validated assessment of smoking status which distinguishes between recent and long-term ex-smokers—ideally biochemically verified—and assesses seroprevalence and links to health records.

In the meantime, public-facing messages about the possible protective effect of smoking or nicotine are premature. In our view, until there is further research, the quality of the evidence does not justify the huge risk associated with a message likely to reach millions of people that a lethal activity, such as smoking, may protect against COVID-19. It continues to be appropriate to recommend smoking cessation and emphasize the role of alternative nicotine products to support smokers to stop as part of public health efforts during COVID-19. At the very least, smoking cessation reduces acute risks from cardiovascular disease and could reduce demands on the health-care system [49]. GPs and other health-care providers can play a crucial role—brief, high-quality and free on-line training is available at National Centre for Smoking Cessation and Training.

## CONCLUSION

Across 233 studies, recorded smoking prevalence was generally lower than national prevalence estimates. Current smokers were at reduced risk of testing positive for SARS-CoV-2 and former smokers were at increased risk of hospitalization, disease severity and mortality compared with never smokers.

## Declaration of interests

D.S. and O.P. have no conflicts of interest to declare. L.S. has received a research grant and honoraria for a talk and travel expenses from manufacturers of smoking cessation medications (Pfizer and Johnson & Johnson). J.B. has received unrestricted research funding to study smoking cessation from companies who manufacture smoking cessation medications. All authors declare no financial links with tobacco companies or e-cigarette manufacturers or their representatives.

## Acknowledgements

An original short review for the Royal College of Physicians was converted to an extended living review after a request by Martin Dockrell, Tobacco Control Lead, Public Health England. All scientific decisions were made by the authors independently of funders and external organizations. The authors would like to thank Rosemary Koper for her assistance in running the electronic searches and data extraction. D.S. is supported by a PhD studentship from the UK Biotechnology and Biological Sciences Research Council (BB/M009513/1). O.P. receives salary support from Cancer Research UK (C1417/A22962). J.B., L.S. and O.P. are members of SPECTRUM, a UK Prevention Research Partnership Consortium (MR/S037519/1). UKPRP is an initiative funded by the UK Research and Innovation Councils, the Department of Health and Social Care (England) and the UK devolved administrations, and leading health research charities.

## Author contributions

**David Simons:** Conceptualization; data curation; formal analysis; methodology; writing-original draft; writing-review & editing. **Lion Shahab:** Conceptualization; data curation; formal analysis; methodology; writing-original draft; writing-review & editing. **Jamie Brown:** Conceptualization; data curation; formal analysis; methodology; writing-original draft; writing-review & editing. **Olga Perski:** Conceptualization; data curation; formal analysis; methodology; writing-original draft; writing-review & editing.

## Future review versions

<https://www.qeios.com/read/latest-UJR2AW>

## Previous review versions

- Version 1:** <https://doi.org/10.32388/UJR2AW>
- Version 2:** <https://doi.org/10.32388/UJR2AW.3>
- Version 3:** <https://doi.org/10.32388/UJR2AW.4>
- Version 4:** <https://doi.org/10.32388/UJR2AW.5>
- Version 5:** <https://doi.org/10.32388/UJR2AW.6>
- Version 6:** <https://doi.org/10.32388/UJR2AW.7>

## References

- and mortality associated with COPD and smoking in patients with COVID-19: a rapid systematic review and meta-analysis. *medRxiv* 2020; **15**: e0233147.
15. Patanavanich R, Glantz SA. Smoking is associated with COVID-19 progression: a meta-analysis. *medRxiv* 2020. <https://doi.org/10.14171/2014.02.004>.
  16. Berlin I., Thomas D., Le Faou A.-L., Cornuz J. COVID-19 and smoking. *Nicotine Tob Res* 2020; **22**: 1650–2.
  17. Farsalinos K., Barbouni A., Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med* 2020; **15**: 845–52.
  18. Grundy E. J., Suddek T., Filippidis F. T., Majeed A., Coronini-Cronberg S. Smoking, SARS-CoV-2 and COVID-19: a review of reviews considering implications for public health policy and practice. *Tob Induc Dis* 2020; **18**. <https://doi.org/10.18332/tid/124788>.
  19. Elliott J. H., Turner T., Clavisi O., Thomas J., Higgins J. P. T., Maverganes C. *et al.* Living systematic reviews: an emerging opportunity to narrow the evidence-practice gap. *PLOS Med* 2014; **11**: e1001603.
  20. Tricco A. C., Antony J., Zarin W., Strifler L., Ghassemi M., Ivory J. *et al.* A scoping review of rapid review methods. *BMC Med* 2015; **13**: 224.
  21. Simons D., Brown J., Shahab L., Perski O. Smoking and COVID-19: Rapid evidence review for the Royal College of Physicians, London (UK). *Qeios* 2020; [10.32388/VGJCN](https://doi.org/10.32388/VGJCN).
  22. R Core Team *The R Project for Statistical Computing*. Vienna, Austria: R core Team; 2013, pp. 1–12.
  23. Bürkner P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *ArXiv170511123 Stat* 2017. <http://arxiv.org/abs/1705.11123> (accessed 26 July 2020).
  24. Simons D., Shahab L., Brown J., Perski O. The association of smoking status with SARS-CoV-2 infection, hospitalization and mortality from COVID-19: a living rapid evidence review (version 5). *Qeios* 2020; <https://doi.org/10.32388/UJR2AW.6>
  25. Efron B. Better bootstrap confidence intervals. *J Am Stat Assoc* 1987; **82**: 171–85.
  26. Miyara M., Tubach F., Martinez V., Morelot-Panzini C., Pernet J., Haroche J. *et al.* Low rate of daily smokers in patients with symptomatic COVID-19. *medRxiv* 2020; <https://doi.org/10.1101/2020.06.10.20127514>
  27. Rimland C. A., Morgan C. E., Bell G. J., Kim M. K., Hedrick T., Marx A. *et al.* Clinical characteristics and early outcomes in patients with COVID-19 treated with tocilizumab at a United States academic center. *medRxiv* 2020; <https://doi.org/10.1101/2020.05.13.20100404>
  28. Girardeau Y., Gallois Y., de Bonnecaze G., Escudé B., Lafont C., Chatellier G. *et al.* Confirmed central olfactory system lesions on brain MRI in COVID-19 patients with anosmia: a case-series. *medRxiv*. <https://doi.org/10.1101/2020.07.08.20148692> (accessed 25 August 2020).
  29. Ebinger J., Botwin G. J., Albert C. M., Alotaibi M., Ardití M., Berg A. H. *et al.* SARS-CoV-2 Seroprevalence across a diverse cohort of healthcare workers. *medRxiv* 2020; <https://doi.org/10.1101/2020.07.31.20163055>
  30. Islam M. Z., Riaz B. K., Islam A. S., Khanam F., Akhter J., Choudhury R. *et al.* Risk factors associated with morbidity and mortality outcomes of COVID-19 patients on the 14th and 28th day of the disease course: a retrospective cohort study in Bangladesh. *medRxiv* 2020; <https://doi.org/10.1101/10.1101/2020.08.17.20176586>

31. Rentsch C. T., Kidwai-Khan F., Tate J. P., Park L. S., King J. T., Skanderson M. *et al.* Covid-19 testing, hospital admission, and intensive care among 2,026,227 United States Veterans aged 54–75 years. *medRxiv* 2020; <https://doi.org/10.1101/2020.04.09.20059964>
32. Odani S. Tobacco product use among military Veterans—United States, 2010–2015. *Morb Mortal Wkly Rep* 2018; **67**: 7–12.
33. Niedzwiedz C. L., O'Donnell C. A., Jani B. D., Demou E., Ho F. K., Celis-Morales C. *et al.* Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK biobank. *BMC Med* 2020; **18**: 160.
34. Trubiano J. A., Vogrin S., Smibert O. C., Marhoon N., Alexander A. A., Chua K. Y. *et al.* COVID-MATCH65—a prospectively derived clinical decision rule for severe acute respiratory syndrome coronavirus 2. *medRxiv* 2020; <https://doi.org/10.1101/2020.06.30.20143818>
35. Hopkinson N. S., Rossi N. N., Moustafa J. E.-S. S. E., Laverty A. A., Quint J. K., Freydin M. B. *et al.* Current tobacco smoking and risk from COVID-19 results from a population symptom app in over 2.4 million people. *medrxiv* 2020; **44**; <https://doi.org/10.1101/2020.05.18.20105288>
36. Merkely B., Szabó A. J., Kosztin A., Berényi E., Sebestyén A., Lengyel C. *et al.* Novel coronavirus epidemic in the Hungarian population, a cross-sectional nationwide survey to support the exit policy in Hungary. *GeroScience* 2020; **42**: 1063–74.
37. Department of Health and Social Care. Major home testing programme for coronavirus will track levels of infection in the community—GOV.UK. <https://www.gov.uk/government/news/major-home-testing-programme-for-coronavirus-will-track-levels-of-infection-in-the-community> (accessed 22 May 2020).
38. Office for National Statistics. COVID-19 Infection Survey (CIS). <https://www.ons.gov.uk/surveys/informationforhouseholdsandindividuals/householdandindividualsurveys/covid19infectionssurveycis> (accessed 30 June 2020).
39. de Lusignan S., Dorward J., Correa A., Jones N., Akinyemi O., Amirthalingam G. *et al.* Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of general practitioners research and surveillance centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020; **20**: 1034–42.
40. World Health Organization. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases. <https://www.who.int/publications-detail-redirect/10665-331501> (accessed 29 July 2020).
41. Polubriaginof F., Salmasian H., Albert D. A., Vawdrey D. K. Challenges with collecting smoking status in electronic health records. *Annu Symp Proc AMIA Symp* 2017; **2017**: 1392–400.
42. Benowitz N. L., Schultz K. E., Haller C. A., Wu A. H. B., Dains K. M., Jacob P. Prevalence of smoking assessed biochemically in an urban public hospital: a rationale for routine cotinine screening. *Am J Epidemiol* 2009; **170**: 885–91.
43. Griffith G., Morris T. T., Tudball M., Herbert A., Mancano G., Pike L. *et al.* Collider bias undermines our understanding of COVID-19 disease risk and severity. *medRxiv* 2020; <https://doi.org/10.1101/2020.05.04.20090506>
44. Murray E. Causation in smoking and COVID-19. Twitter 2020. <https://twitter.com/EpiEllie/status/1258607277357006849?s=20> (accessed 8 May 2020).
45. Bowyer R. C. E., Varsavsky T., Carole H. Geo-social gradients in predicted COVID-19 prevalence and severity in Great Britain: results from affiliations: corresponding authors: understanding the geographical distribution of COVID-19 through the general population is key to the provision of ade. 2020. <https://doi.org/10.1101/2020.04.23.20076521>
46. Jackson S. E., Brown J., Shahab L., Steptoe A., Fancourt D. COVID-19, smoking, and inequalities: a cross-sectional survey of adults in the UK. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.30.20086074>
47. Ward H., Atchison C. J., Whitaker M., Ainslie K. E., Elliot J., Okell L. C. *et al.* Antibody prevalence for SARS-CoV-2 in England following first peak of the pandemic: REACT2 study in 100,000 adults. *medRxiv* 2020; <https://doi.org/10.1101/2020.08.12.20173690>
48. Shahab L., Brose L. S., West R. Novel delivery systems for nicotine replacement therapy as an aid to smoking cessation and for harm reduction: rationale, and evidence for advantages over existing systems. *CNS Drugs* 2013; **27**: 1007–19.
49. Stead L. E., Buitrago D., Preciado N., Sanchez G., Hartmann-Boyce J., Lancaster T. Physician advice for smoking cessation. *Cochrane Database Syst Rev* 2013; **5**: CD000165.
50. Guan W., Liang W., Zhao Y., Liang H. R., Chen Z. S., Li Y. M. *et al.* Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* 2020; **55**: 2000547.
51. Lian J., Jin X., Hao S., Cai H., Zhang S., Zheng L. *et al.* Analysis of epidemiological and clinical features in older patients with coronavirus disease 2019 (COVID-19) outside Wuhan. *Clin Infect Dis* 2020; **71**: 740–7.
52. Jin X., Lian J.-S., Hu J.-H., Gao J., Zheng L., Zhang Y. M. *et al.* Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut* 2020; **69**: 1002–9.
53. Chen T., Wu D., Chen H., Yan W., Yang D., Chen G. *et al.* Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020; **368**: m1091.
54. Zhou F., Yu T., Du R., Fan G., Liu Y., Liu Z. *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054–62.
55. Mo P., Xing Y., Xiao Y., Deng L., Zhao Q., Wang H. *et al.* Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis* 2020; <https://doi.org/10.1093/cid/ciaa270>
56. Zhang J., Dong X., Cao Y., Yuan Y. D., Yang Y. B., Yan Y. Q. *et al.* Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 2020; **75**: 1730–41.
57. Wan S., Xiang Y., Fang W., Zheng Y., Li B., Hu Y. *et al.* Clinical features and treatment of COVID-19 patients in Northeast Chongqing. *J Med Virol* 2020; **92**: 797–806.
58. Liu W., Tao Z.-W., Wang L., Yuan M. L., Liu K., Zhou L. *et al.* Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J* 2020; **133**: 1032–8.
59. Huang C., Wang Y., Li X., Ren L., Zhao J., Hu Y. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506.
60. Zhang X., Cai H., Hu J., Lian J., Gu J., Zhang S. *et al.* Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings. *Int J Infect Dis* 2020; **94**: 81–7.

61. Guo T., Fan Y., Chen M., Wu X., Zhang L., He T. *et al.* Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020; **5**: 811–8.
62. Liu R., Ming X., Xu O., Song L., Gao Z., Gao L. *et al.* Association of cardiovascular manifestations with in-hospital outcomes in patients with COVID-19: a hospital staff data. *medRxiv* 2020; <https://doi.org/10.1101/2020.02.29.20029348>
63. Huang Y., Yang R., Xu Y., Gong P. Clinical characteristics of 36 non-survivors with COVID-19 in Wuhan, China. *medRxiv* 2020. <https://doi.org/10.1101/2020.02.27.20029009>
64. Xu H., Hou K., Xu H., Li Z., Chen H., Zhang N. *et al.* Acute myocardial injury of patients with coronavirus disease 2019. *medRxiv* 2020; <https://doi.org/10.1101/2020.03.05.20031591>
65. Li J., Li S., Cai Y., Liu Q., Li X., Zeng Z. *et al.* Epidemiological and clinical characteristics of 17 hospitalized patients with 2019 novel coronavirus infections outside Wuhan, China. *medRxiv* 2020; <https://doi.org/10.1101/2020.02.11.20022053>
66. Hu L., Chen S., Fu Y., Gao Z., Long H., Ren H. W. *et al.* Risk factors associated with clinical outcomes in 323 COVID-19 patients in Wuhan, China. *medRxiv* 2020; <https://doi.org/10.1101/2020.03.25.20037721>
67. Wang R., Pan M., Zhang X., Han M., Fan X., Zhao F. *et al.* Epidemiological and clinical features of 125 hospitalized patients with COVID-19 in Fuyang, Anhui, China. *Int J Infect Dis* 2020; **95**: 421–8.
68. CDCMMWR Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019—United States, February 12–March 28, 2020. *Morb Mortal Wkly Rep* 2020; **69**: 382–6.
69. Dong X., Cao Y., Lu X., Zhang J. J., du H., Yan Y. Q. *et al.* Eleven faces of coronavirus disease 2019. *Allergy* 2020; **75**: 1699–709.
70. Kim E. S., Chin B. S., Kang C. K., Kim N. J., Kang Y. M., Choi J. P. *et al.* Clinical course and outcomes of patients with severe acute respiratory syndrome coronavirus 2 infection: a preliminary report of the first 28 patients from the Korean cohort study on COVID-19. *J Korean Med Sci* 2020; **35**: e142.
71. Shi Y., Yu X., Zhao H., Wang H., Zhao R., Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care* 2020; **24**: 108.
72. Yang X., Yu Y., Xu J., Shu H., Xia J., Liu H. *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; **8**: 475–81.
73. Argenziano M. G., Bruce S. L., Slater C. L., Tiao J. R., Baldwin M. R., Barr R. G. *et al.* Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020; **369**: m1996.
74. Solís P., Carreño H. COVID-19 fatality and comorbidity risk factors among confirmed patients in Mexico. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.21.20074591>
75. Richardson S., Hirsch J. S., Narasimhan M., Crawford J. M., McGinn T., Davidson K. W. *et al.* Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the new York City area. *JAMA* 2020; **323**: 2052–9.
76. Fontanet A., Tondeur L., Madec Y., Grant R., Besombes C., Jolly N. *et al.* Cluster of COVID-19 in northern France: a retrospective closed cohort study. *medRxiv* 2020; <https://doi.org/10.1101/2020.04.18.20071134>
77. Zheng K. I., Gao F., Wang X.-B., Sun Q. F., Pan K. H., Wang T. Y. *et al.* Letter to the editor: obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism* 2020; **108**: 154244.
78. Liao Y., Feng Y., Wang B., Wang H., Huang J., Wu Y. *et al.* Clinical characteristics and risk factors for developed COVID-19 patients transferring to designated hospital from Jianghan Fangcang Shelter Hospital: a retrospective, observational study. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.21.20074724>
79. Gil-Agudo A., Rodriguez-Cola M., Jimenez-Velasco I., Gutierrez-Henares E., Lopez-Dolado E., Gambarutta-Malfatti C. *et al.* Clinical features of coronavirus disease 2019 (COVID-19) in a cohort of patients with disability due to spinal cord injury. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.20.20072918>
80. Shi P., Ren G., Yang J., Li Z., Deng S., Li M. *et al.* Clinical characteristics of imported and second-generation COVID-19 cases outside Wuhan, China: a multicenter retrospective study. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.19.20071472>
81. Hadjadj J., Yatim N., Barnabei L., Corneau A., Boussier J., Pere H. *et al.* Impaired type I interferon activity and exacerbated inflammatory responses in severe Covid-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.19.20068015>
82. Gold J. A. W., Wong K. K., Szablewski C. M., Patel P. R., Rossow J., da Silva J. *et al.* Characteristics and clinical outcomes of adult patients hospitalized with COVID-19—Georgia, March 2020. *Morb Mortal Wkly Rep* 2020; **69**: 545–50.
83. Yu T., Cai S., Zheng Z., Cai X., Liu Y., Yin S. *et al.* Association between clinical manifestations and prognosis in patients with COVID-19. *Clin Ther* 2020; **42**: 964–72.
84. Zheng Y., Xiong C., Liu Y., Qian X., Tang Y., Liu L. *et al.* Epidemiological and clinical characteristics analysis of COVID-19 in the surrounding areas of Wuhan, Hubei Province in 2020. *Pharmacol Res* 2020; **157**: 104821.
85. de la Rica R., Borges M., Aranda M., del Castillo A., Socias A., Payeras A. *et al.* Low albumin levels are associated with poorer outcomes in a case series of COVID-19 patients in Spain: a retrospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.07.20094987>
86. Yin R., Yang Z., Wei Y., Li Y., Chen H., Ma D. *et al.* Clinical characteristics of 106 patients with neurological diseases and co-morbid coronavirus disease 2019: a retrospective study. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.29.20085415>
87. Shi H., Zuo Y., Yalavarthi S., Gockman K., Zuo M., Madison J. A. *et al.* Neutrophil calprotectin identifies severe pulmonary disease in COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.06.20093070>
88. Cho E. R., Slutsky A. S., Jha P. Smoking and the risk of COVID-19 infection in the UK Biobank Prospective Study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.05.20092445>
89. Allenbach Y., Saadoun D., Maalouf G., Vieira M., Hellio A., Boddaert J. *et al.* Multivariable prediction model of intensive

- care unit transfer and death: a French prospective cohort study of COVID-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.04.20090118>
90. Robilotti E. V., Babady N. E., Mead P. A., Rolling T., Perez-Johnston R., Bernardes M. et al. Determinants of severity in cancer patients with COVID-19 illness. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.04.20086322>
  91. Williamson E. J., Walker A. J., Bhaskaran K., Bacon S., Bates C., Morton C. E. et al. OpenSAFELY: factors associated with COVID-19 death in 17 million patients. *Nature* 2020; **584**: 430–6.
  92. Borobia A. M., Carcas A. J., Arnalich F., Álvarez-Sala R., Monserrat-Villatoro J., Quintana M. et al. A cohort of patients with COVID-19 in a major teaching hospital in Europe. *J Clin Med* 2020; **9**: 1733.
  93. Giacomelli A., Ridolfo A. L., Milazzo L., Oreni L., Bernacchia D., Siano M. et al. 30-day mortality in patients hospitalized with COVID-19 during the first wave of the Italian epidemic: a prospective cohort study. *Pharmacol Res* 2020; **158**: 104931.
  94. Shah S. J., Barish P. N., Prasad P. A., Kistler A., Neff N., Kamm J. et al. Clinical features, diagnostics, and outcomes of patients presenting with acute respiratory illness: a comparison of patients with and without COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.02.20082461>
  95. Kolin D. A., Kulm S., Elemento O. Clinical and genetic characteristics of Covid-19 patients from UK Biobank. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.05.20075507>
  96. Lubetzky M., Aull M., Craig-Shapiro R., Lee J., Lee J., Sultan S. et al. Kidney allograft recipients diagnosed with coronavirus disease-2019: a single center report. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.30.20086462>
  97. Goyal P., Choi J. J., Pinheiro L. C., Schenck E. J., Chen R., Jabri A. et al. Clinical characteristics of Covid-19 in new York City. *N Engl J Med* 2020; **382**: 2372–4.
  98. Feng Y., Ling Y., Bai T., Xie Y., Huang J., Li J. et al. COVID-19 with different severities: a multicenter study of clinical features. *Am J Respir Crit Care Med* 2020; **201**: 1380–8.
  99. Yao Q., Wang P., Wang X., Qie G., Chu Y. A retrospective study of risk factors for severe acute respiratory syndrome coronavirus 2 infections in hospitalized adult patients. *Polish Arch Intern Med* 2020; **130**: 390–9.
  100. Sami R., Soltaninejad E., Amra B., Naderi Z., Javanmard S. H., Iraj B. et al. A one-year hospital-based prospective COVID-19 open-cohort in the Eastern Mediterranean region: the Khorshid COVID cohort (KCC) study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.11.20096727>
  101. Almazeedi S., Youha S. A., Jamal M. H., Al-Haddad M., Al-Muhaini A., Al-Ghimplas F. et al. Clinical characteristics, risk factors and outcomes among the first consecutive 1,096 patients diagnosed with COVID-19: the Kuwait experience. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.09.20096495>
  102. Carrillo-Vega M. F., Salinas-Escudero G., Garcia-Peña C., Gutierrez-Robledo L. M., Parra-Rodriguez L. Early estimation of the risk factors for hospitalization and mortality by COVID-19 in Mexico. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.11.20098145>
  103. Yanover C., Mizrahi B., Kalkstein N., Marcus K., Akiva P., Barer Y. et al. What factors increase the risk of complications in SARS-CoV-2 positive patients? A cohort study in a nationwide Israeli health organization. *medRxiv* 2020; **6**: e20872.
  104. Hamer M., Kivimäki M., Gale C. R., Batty G. D. Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: a community-based cohort study of 387,109 adults in UK. *Brain Behav Immun* 2020; **87**: 184–7.
  105. Regina J., Papadimitriou-Olivgeris M., Burger R., Filippidis P., Tschopp J., Desgranges F. et al. Epidemiology, risk factors and clinical course of SARS-CoV-2 infected patients in a Swiss university hospital: an observational retrospective study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.11.20097741>
  106. Targher G., Mantovani A., Wang X.-B., Yan H. D., Sun Q. F., Pan K. H. et al. Patients with diabetes are at higher risk for severe illness from COVID-19. *Diabetes Metab* 2020; **46**: 335–7.
  107. Valenti L., Bergna A., Pelusi S., Facciotti F., Lai A., Tarkowski M. et al. SARS-CoV-2 seroprevalence trends in healthy blood donors during the COVID-19 Milan outbreak. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.11.20098442>
  108. Feuth T., Saarela T., Karlsson A., Valtonen M., Peltola V., Rintala E. et al. Is sleep apnoea a risk factor for Covid-19? Findings from a retrospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.14.20098319>
  109. Ge H., Zhu M., Du J., Zhou Y., Wang W., Zhang W. et al. Cardiac structural and functional characteristics in patients with coronavirus disease 2019: a serial echocardiographic study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.12.20095885>
  110. Parrotta E., Kister I., Charvet L., Sammarco C., Saha V., Charlson R. E. et al. COVID-19 outcomes in MS: observational study of early experience from NYU multiple sclerosis comprehensive care center. *Neuroimmunol Neuroinflammation* 2020; **7**: e835.
  111. Shekhar R., Sheikh A. B., Upadhyay S., Atencio J., Kapuria D. Early experience with COVID-19 patients at academic hospital in southwestern United States. *Infect Dis* 2020; **52**: 596–9.
  112. Mejia-Vilet J. M., Cordova-Sanchez B. M., Fernandez-Camargo D., Mendez-Perez R. A., Morales-Buenrostro L. E., Hernandez-Gilsoul T. A risk score to predict admission to intensive care unit in patients With COVID-19: the ABC-GOALS score. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.12.20099416>
  113. Chen C., Jiang J., Xu X., Hu Y., Hu Y., Zhao Y. Dynamic liver function indexes monitoring and clinical characteristics in three types of COVID-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.13.20099614>
  114. Li J., Chen Y., Chen S., Wang S., Zhang D., Wang J. et al. Derivation and validation of a prognostic model for predicting in-hospital mortality in patients admitted with COVID-19 in Wuhan, China: the PLANS (Platelet Lymphocyte Age Neutrophil Sex) model. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.13.20100370>
  115. Palaiodimos L., Kokkinidis D. G., Li W., Karamanis D., Ognibene J., Arora S. et al. Severe obesity, increasing age and male sex are independently associated with worse in-hospital outcomes, and higher in-hospital mortality, in a cohort of patients with COVID-19 in the Bronx, New York. *Metabolism* 2020; **108**: 154262.
  116. Ip A., Berry D. A., Hansen E., Goy A. H., Pecora A. L., Sinclair B. A. et al. Hydroxychloroquine and tocilizumab therapy in COVID-19 patients—an observational study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.21.20109207>
  117. Heili-Frades S. COVID-19 outcomes in 4712 consecutively confirmed SARS-CoV2 cases in the city of Madrid. *medRxiv*

2020. <https://doi.org/10.1101/2020.05.22.20109850> (accessed 27 July 2020).
118. Vaquero L. M., Barrado M. E. S., Escobar D., Arribas P., Gonzalez J. R., Bermejo J. F. *et al.* C-Reactive protein and SOFA score as early predictors of critical care requirement in patients with COVID-19 pneumonia in Spain. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.22.20110429>
119. Kim L., Garg S., O'Halloran A., Whitaker M., Pham H., Anderson E. J. *et al.* Interim analysis of risk factors for severe outcomes among a cohort of hospitalized adults identified through the U.S. coronavirus disease 2019 (COVID-19)-associated hospitalization surveillance network (COVID-NET). *medRxiv* 2020. <https://doi.org/10.1101/2020.05.18.20103390>
120. Wu M. A., Fossali T., Pandolfi L., Carsana L., Ottolina D., Frangipane V. *et al.* COVID-19: the key role of pulmonary capillary leakage. An observational cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.17.20104877>
121. Shi Q., Zhao K., Yu J., Jiang F., Feng J., Zhao K. *et al.* Clinical characteristics of 101 COVID-19 nonsurvivors in Wuhan, China: a retrospective study. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.04.20031039>
122. Al-Hindawi A., Sokhi J., Cuddihy J., Lockie C., Christie L., Davies R. *et al.* COVID-19 in London, a case series demonstrating late improvement in survivors. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.16.20103853>
123. Basse C., Diakite S., Servois V., Frelaut M., Noret A., Bellesoeur A. *et al.* Characteristics and outcome of SARS-CoV-2 infection in cancer patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.14.20101576>
124. Freites D., Leon L., Mucientes A., Rodriguez-Rodriguez L., Font J., Madrid A. *et al.* Risk factors for hospital admission related to COVID-19 in inflammatory rheumatic diseases. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.14.20101584>
125. Alshami A. A., Alattas R. A., Anan H. F., Al Qahtani H. S., Al Mulhim M. A., Alahilmi A. A. *et al.* Silent disease and loss of taste and smell are common manifestations of SARS-CoV-2 Infection in a quarantine facility: first report from Saudi Arabia. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.13.20100222>
126. Berumen J., Schmulson M., Alegre J., Guerrero G., Olaiz G., Wong-Chew R. M. *et al.* Risk of infection and hospitalization by Covid-19 in Mexico: a case-control study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.24.20104414>
127. Gianfrancesco M., Hyrich K. L., Al-Adely S., Carmona L., Danila M. I., Gossec L. *et al.* Characteristics associated with hospitalization for COVID-19 in people with rheumatic disease: data from the COVID-19 global rheumatology Alliance physician-reported registry. *Ann Rheum Dis* 2020; **79**: 859–66.
128. Li J., Long X., Zhu C., Wang H., Wang T., Lin Z. *et al.* Olfactory dysfunction in recovered coronavirus disease 2019 (COVID-19) patients. *Mov Disord* 2020; **35**: 1100–1.
129. Batty G. D., Deary I., Luciano M., Altschul D., Kivimaki M., Gale C. Psychosocial factors and hospitalizations for COVID-19: Prospective cohort study of the general population. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.29.20100735>
130. Israel A., Feldhamer I., Lahad A., Levin-Zamir D., Lavie G. Smoking and the risk of COVID-19 in a large observational population study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.01.20118877>
131. Valle D. M. D., Kim-schulze S., Hsin-hui H., Beckmann N. D., Nirenberg S., Wang B. *et al.* An inflammatory cytokine signature helps predict COVID-19 severity and death. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.28.20115758>
132. Chaudhry F., Bulkha H., Rathnam A. S., Said O. M., Lin J., Lorigan H. *et al.* COVID-19 in multiple sclerosis patients and risk factors for severe infection. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.27.20114827>
133. Louis S., Dhawan A., Newey C., Nair D., Jehi L., Hantus S. *et al.* Continuous electroencephalography (cEEG) characteristics and acute symptomatic seizures in COVID-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.26.20114033>
134. Soto-Mota A., Garza B. A. M., Rodriguez E. M., Rodriguez J. O., Romo A. E., Minutti P. A. *et al.* The low-harm score for predicting mortality in patients diagnosed with covid-19: a multicentric validation study. *medRxiv* 2020; <https://doi.org/10.1101/2020.05.26.20111120>
135. Garibaldi B. T., Fiksel J., Muschelli J., Robinson M. L., Rouhizadeh M., Nagy P. *et al.* Patient trajectories and risk factors for severe outcomes among persons hospitalized for COVID-19 in the Maryland/DC region. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.24.20111864>
136. Docherty A. B., Harrison E. M., Green C. A., Hardwick H. E., Pius R., Norman L. *et al.* Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO clinical characterization protocol: prospective observational cohort study. *BMJ* 2020; **369**: m1985.
137. Boulware D. R., Pullen M. F., Bangdiwala A. S., Pastick K. A., Lofgren S. M., Okafor E. C. *et al.* A randomized trial of Hydroxychloroquine as Postexposure prophylaxis for Covid-19. *N Engl J Med* 2020; **383**: 517–25.
138. Kuderer N. M., Choueiri T. K., Shah D. P., Shyr Y., Rubinstein S. M., Rivera D. R. *et al.* Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study. *Lancet* 2020; **395**: 1907–18.
139. Romão V. C., Oliveira-Ramos E., Cruz-Machado A. R., Martins P., Barreira S., Silva-Dinis J. *et al.* A COVID-19 outbreak in a rheumatology department upon the early days of the pandemic. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.05.20107011>
140. Giannouchos T., Sussman R., Mier J. M., Poulas K., Farsalinos K. Characteristics and risk factors for COVID-19 diagnosis and adverse outcomes in Mexico: an analysis of 89,756 laboratory-confirmed COVID-19 cases. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.04.20122481>
141. Ramlall V., Thangaraj P., Meydan C., Foox J., Butler D., May B. *et al.* Identification of Immune complement function as a determinant of adverse SARS-CoV-2 infection outcome. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.05.20092452>
142. Wang B., Oekelen O. V., Mouhieddine T., Del Valle D. M., Richter J., Cho H. J. *et al.* A tertiary center experience of multiple myeloma patients with COVID-19: lessons learned and the path forward. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.04.20122846>
143. Perrone F., Piccirillo M. C., Ascierto P. A., Salvarani C., Parrella R., Marata A. M. *et al.* Tocilizumab for patients with COVID-19 pneumonia. The TOCIVID-19 prospective phase 2 trial. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.01.20119149>
144. Sharma A. K., Ahmed A., Baig V. N., Dhakad P., Dalela G., Kacker S. *et al.* Characteristics and outcomes of hospitalized young adults with mild to moderate Covid-19 at a University

- Hospital in India. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.02.20106310>
145. Eugen-Olsen J., Altintas I., Tingleff J., Gamst-Jensen H., Lindstroem M. B., Rasmussen L. J. et al. Low levels of the prognostic biomarker suPAR are predictive of mild outcome in patients with symptoms of COVID-19—a prospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.27.20114678>
  146. Martinez-Portilla R. J., Sotiriadis A., Torres-Torres J., Christos C., Hawkins-Villarreal A., Villafan-Bernal J. R. et al. Risk factors for mortality in pregnant women with SARS-CoV-2 infection. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.31.20107276>
  147. Raisi-Estabragh Z., McCracken C., Bethell M. S., Cooper J., Cooper C., Caulfield M. J. et al. Greater risk of severe COVID-19 in black, Asian and minority ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: study of 1326 cases from the UK biobank. *J Public Health* 2002; **42**: 451–60.
  148. Luo H., Liu S., Wang Y., Phillips-Howard P. A., Yang Y., Ju S. et al. Age differences in clinical features and outcomes in patients with COVID-19, Jiangsu, China: a retrospective, multi-center cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.01.20086025>
  149. Houlihan C. F., Vora N., Byrne T., Lewer D., Kelly G., Heaney J. et al. Pandemic peak SARS-CoV-2 infection and seroconversion rates in London frontline health-care workers. *Lancet* 2020; **396**: e6–e7.
  150. Cen Y., Chen X., Shen Y., Zhang X. H., Lei Y., Xu C. et al. Risk factors for disease progression in patients with mild to moderate coronavirus disease 2019—a multi-Centre observational study. *Clin Microbiol Infect* 2020; **26**: 1242–7.
  151. Klang E., Kassim G., Soffer S., Freeman R., Levin M. A., Reich D. L. Morbid obesity as an independent risk Factor for COVID-19 mortality in hospitalized patients younger than 50. *Obesity* 2002; **28**: 1595–9.
  152. Marascini A., Corsi E., Salvatore M. A., Donati S. Coronavirus and birth in Italy: results of a national population-based cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.11.20128652>
  153. Wang A.-L., Zhong X., Hurd Y. Comorbidity and Sociodemographic determinants in COVID-19 Mortality in an US Urban Healthcare System. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.11.20128926>
  154. McQueenie R., Foster H., Jani B. D., Katikireddi S. V., Sattar N., Pell J. P. et al. Multimorbidity, polypharmacy, and COVID-19 infection within the UK Biobank cohort. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.10.20127563>
  155. Apea V. J., Wan Y. I., Dhairyawan R., Puthucheary Z. A., Pearse R. M., Orkin C. M. et al. Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: an observational cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.10.20127621>
  156. Woolford S. J., D'Angelo S., Curtis E. M., Parsons C. M., Ward K. A., Dennison E. M. et al. COVID-19 and associations with frailty and multimorbidity: a prospective analysis of UK Biobank participants. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.09.20126292>
  157. Hultcrantz M., Richter J., Rosenbaum C., Patel D., Smith E., Korde N. et al. COVID-19 infections and outcomes in patients with multiple myeloma in New York City: a cohort study from five academic centers. *medRxiv* 2020; <https://doi.org/10.1101/2020.06.09.20126516>
  158. Rajter J. C., Sherman M., Fatteh N., Vogel F., Sacks J., Rajter J.-J. ICON (Ivermectin in Covid nineteen) study: use of ivermectin is associated with lower mortality in hospitalized patients with COVID19. *medRxiv* 2020. <https://doi.org/10.1101/20.06.06.20124461>
  159. Lan F.-Y., Suharlim C., Kales S. N., Yang J. Association between SARS-CoV-2 infection, exposure risk and mental health among a cohort of essential retail workers in the United States. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.08.20125120>
  160. Zeng H., Zhang T., He X., Du Y., Tong Y., Wang X. et al. Impact of chronic comorbidities on progression and prognosis in patients with COVID-19: a retrospective cohort study in 1031 hospitalized cases in Wuhan, China. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.14.20125997>
  161. Suleyman G., Fadel R. A., Malette K. M., Hammond C., Abdulla H., Entz A. et al. Clinical characteristics and morbidity associated with coronavirus disease 2019 in a series of patients in metropolitan Detroit. *JAMA Netw Open* 2020; **3**: e2012270–e2012270.
  162. Chen L., Yu J., He W., Chen L., Yuan G., Dong F. et al. Risk factors for death in 1859 subjects with COVID-19. *Leukemia* 2020; **34**: 2173–83.
  163. Garassino M. C., Whisenant J. G., Huang L.-C., Trama A., Torri V., Agostoni F. et al. COVID-19 in patients with thoracic malignancies (TERAVOLT): first results of an international, registry-based, cohort study. *Lancet Oncol* 2020; **21**: 914–22.
  164. Hernández-Garduño E. Obesity is the comorbidity more strongly associated for Covid-19 in Mexico. A case-control study. *Obes Res Clin Pract* 2020; **14**: 375–9.
  165. Govind R., de Freitas D. F., Pritchard M. R., Hayes R. D., MacCabe J. H. Clozapine treatment and risk of COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.17.20133595>
  166. Sisó-Almirall A., Kostov B., Mas-Heredia M., Vilanova-Rotllan S., Sequeira-Aymar E., Sans-Corrales M. et al. Prognostic factors in Spanish Covid-19 patients: a case series from barcelona. *PLOS ONE* 2020; **15**: e0237960.
  167. Gu T., Mack J. A., Salvatore M., Sankar S. P., Valley T. S., Singh K. et al. COVID-19 outcomes, risk factors and associations by race: a comprehensive analysis using electronic health records data in Michigan Medicine. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.16.20133140>
  168. Kibler M., Carmona A., Marchandot B., Matsushita K., Trimaille A., Kanso M. et al. Risk and severity of COVID-19 and ABO blood group in transcatheter aortic valve patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.13.20130211>
  169. Ikitimur H., Borku Uysal B., Cengiz M., Ikitimur B., Uysal H., Ozcan E. et al. Determining host factors contributing to disease severity in a family cluster of 29 hospitalized SARS-CoV-2 patients: could genetic factors be relevant in the clinical course of COVID-19? *J Med Virol* 2020. <https://doi.org/10.1002/jmv.26106>
  170. Sierpiński R., Pinkas J., Jankowski M., Zgliczyński W. S., Wierzba W., Gujski M. et al. Sex differences in the frequency of gastrointestinal symptoms and olfactory or taste disorders among 1,942 non-hospitalized patients with COVID-19. *Pol Arch Intern Med* 2020; **13**: 15414.
  171. Zhou Y., He X., Zhang J., e Xue Y., Liang M., Yang B. et al. Prolonged SARS-CoV-2 viral shedding in patients with COVID-19 was associated with delayed initiation of arbidol

- treatment: a retrospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.09.20076646>
172. Crovetto F., Crispi F., Llurba E., Figueras F., Gomez-Roig M. D., Gratacos E. Seroprevalence and clinical spectrum of SARS-CoV-2 infection in the first versus third trimester of pregnancy. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.17.20134098>
  173. Veras F. P., Pontelli M., Silva C., Toller-Kawahisa J., de Lima M., Nascimento D. *et al.* SARS-CoV-2 triggered neutrophil extracellular traps (NETs) mediate COVID-19 pathology. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.08.20125823>
  174. Sterlin D., Mathian A., Miyara M., Mohr A., Anna F., Claer L. *et al.* IgA dominates the early neutralizing antibody response to SARS-CoV-2. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.10.20126532>
  175. Rossi B., Nguyen L. S., Zimmermann P., Boucenna F., Baucher L., Dubret L. *et al.* Effect of tocilizumab in hospitalized patients with severe pneumonia COVID-19: a cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.06.20122341>
  176. Duan L., Zhang S., Guo M., Zhou E., Fan J., Wang X. *et al.* Epidemiological and clinical characteristics in patients with SARS-CoV-2 antibody negative probable COVID-19 in Wuhan. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.18.20134619>
  177. Martin-Jimenez P., Munoz-Garcia M. I., Seoane D., Roca-Rodriguez L., Garcia-Reyne A., Lalueza A. *et al.* Cognitive impairment is a common comorbidity in COVID-19 deceased patients. A hospital-based retrospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.08.20125872>
  178. Elezkurtaj S., Greuel S., Ihlow J., Michaelis E., Bischoff P., Kunze C. A. *et al.* Causes of death and comorbidities in patients with COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.15.20131540>
  179. Lenka J., Chhabria M. S., Sharma N., Bryan E., Tan X., Boppana L. K. *et al.* Clinical characteristics and outcomes of critically ill patients with COVID-19 in a tertiary community hospital in upstate New York. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.18.20135046>
  180. Olivares E., Munoz D., Fica A., Delama I., Alvarez I., Navarrete M. *et al.* Covid-19 in Chile. The experience of a regional reference center. Preliminary report. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.14.20130898>
  181. Salton F., Confalonieri P., Santus P., Harari S., Scala R., Lanini S. *et al.* Prolonged low-dose methylprednisolone in patients with severe COVID-19 pneumonia. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.17.20134031>
  182. Wei W., Ortwine J. K., Mang N. S., Joseph C., Hall B. C., Prokesch B. C. Limited role for antibiotics in COVID-19: scarce evidence of bacterial coinfection. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.16.20133181>
  183. Zuo Y., Estes S. K., Gandhi A. A., Yalavarthi S., Ali R. A., Shi H. *et al.* Prothrombotic antiphospholipid antibodies in COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.15.20131607>
  184. Killerby M. E. Characteristics associated with hospitalization among patients with COVID-19—metropolitan Atlanta, Georgia, March–April 2020. *Morb Mortal Wkly Rep* 2020; **69**: 790–4.
  185. Petrilli C. M., Jones S. A., Yang J., Rajagopalan H., O'Donnell L., Chernyak Y. *et al.* Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020; **369**: m1966.
  186. Magagnoli J., Narendran S., Pereira F., Cummings T. H., Hardin J. W., Sutton S. S. *et al.* Outcomes of hydroxychloroquine usage in United States Veterans Hospitalized with COVID-19. *Med (NY)* 2020. <https://doi.org/10.1016/j.medj.2020.06.001>
  187. Bello-Chavolla O. Y., Bahena-López J. P., Antonio-Villa N. E., Vargas-Vázquez A., González-Díaz A., Márquez-Salinas A. *et al.* Predicting mortality due to SARS-CoV-2: a mechanistic score relating obesity and diabetes to COVID-19 outcomes in Mexico. *J Clin Endocrinol Metab* 2020; **105**: 2752–61.
  188. Zuo Y., Zuo M., Yalavarthi S. Neutrophil extracellular traps and thrombosis in COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.30.20086736>
  189. Sigel K., Swartz T., Golden E., Paranjepe I., Somani S., Richter F. *et al.* Covid-19 and people with HIV infection: outcomes for hospitalized patients in New York City. *Clin Infect Dis* 2020; <https://doi.org/10.1093/cid/ciaa880>
  190. Nguyen A. B., Upadhyay G. A., Chung B., Smith B., Besser S. A., Johnson J. A. *et al.* Outcomes and cardiovascular comorbidities in a predominantly African-American population with COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.28.20141929>
  191. de Melo A. C., Thuler L., da Silva J. L., de Albuquerque L. Z., Pecego A. C., de OR Rodrigues L. *et al.* Cancer inpatient with COVID-19: a report from the Brazilian National Cancer Institute. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.27.20141499>
  192. Auvinen R., Nohynek H., Syrjänen R., Ollgren J., Kerittula T., Mäntylä J. *et al.* Comparison of the clinical characteristics and outcomes of hospitalized adult COVID-19 and influenza patients: a prospective observational study. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.29.20140632>
  193. Souza F. S. H., Hojo-Souza N. S., Santos E. B., Silva C. M., Guidoni D. L. Predicting the disease outcome in COVID-19 positive patients through machine learning: a retrospective cohort study with Brazilian data. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.26.20140764>
  194. Mendy A., Apewokin S., Wells A. A., Morrow A. L. Factors associated with hospitalization and disease severity in a racially and ethnically diverse population of COVID-19 Patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.25.20137323>
  195. Pongpirul W. A., Wiboonchutikul S., Charoenpong L., Panitantum N., Vachiraphan A., Uttayamakul S. *et al.* Clinical course and potential predicting factors of pneumonia of adult patients with coronavirus disease 2019 (COVID-19): a retrospective observational analysis of 193 confirmed cases in Thailand. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.24.20139642>
  196. Jin C., Gu J., Yuan Y., Long Q., Zhang Q., Zhou H. *et al.* Treatment of six COVID-19 patients with convalescent plasma. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.21.20109512>
  197. Favara D. M., Cooke A., Doffinger R., Houghton S., Budriunaite I., Bossingham S. *et al.* First results from the UK COVID-19 Serology in Oncology Staff Study (CSOS). *medRxiv* 2020. <https://doi.org/10.1101/2020.06.22.20136838>
  198. Fisman D., Greer A. L., Tuite A. Derivation and validation of clinical prediction rule for COVID-19 mortality in Ontario,

- Canada. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.21.20136929>
199. Madariaga M. L. L., Guthmiller J., Schrantz S., Jansen M. O., Christensen C., Kumar M. *et al.* Clinical predictors of donor antibody titer and correlation with recipient antibody response in a COVID-19 convalescent plasma clinical trial. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.21.20132944>
  200. Senkal N., Chronic A. C. E. inhibitor use is associated with decreased odds of severe disease in patients with COVID-19. *Anatol J Cardiol* 2020; **24**: 21–9.
  201. Mohamud A. Y., Griffith B., Rehman M., Miller D., Chebl A., Patel S. C. *et al.* Intraluminal carotid artery thrombus in COVID-19: another danger of cytokine storm? *Am J Neuroradiol* 2020; **41**: 1677–82.
  202. Magleby R., Westblade L. F., Trzebucki A., Simon M. S., Rajan M., Park J. *et al.* Impact of SARS-CoV-2 viral load on risk of intubation and mortality among hospitalized patients with coronavirus disease 2019. *Clin Infect Dis* 2020; <https://doi.org/10.1093/cid/ciaa851>
  203. Kimmig L. M., Wu D., Gold M., Pettit N. N., Pittrak D., Mueller J. *et al.* IL6 inhibition in critically ill COVID-19 patients is associated with increased secondary infections. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.15.20103531>
  204. Bello-Chavolla O. Y., Antonio-Villa N. E., Vargas-Vázquez A., Fermín-Martínez C. A., Márquez-Salinas A., Bahena-López J. P. Profiling pre-symptomatic and asymptomatic cases with confirmed SARS-CoV-2 infection in Mexico City. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.02.20145516>
  205. Zacharioudakis I. M., Prasad P. J., Zervou F. N., Basu A., Inglima K., Weisenberg S. A. *et al.* Association of SARS-CoV-2 genomic load with COVID-19 patient outcomes. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.02.20145151>
  206. Antonio-Villa N. E., Bello-Chavolla O. Y., Vargas-Vazquez A., Fermín-Martínez C. A., Marquez-Salinas A., Bahena-Lopez J. P. Health-care workers with COVID-19 living in Mexico City: clinical characterization and related outcomes. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.02.20145169>
  207. Patel M., Chowdhury J., Mills N., Marron R., Gangemi A., Dorey-Stein Z. *et al.* ROX index predicts intubation in patients with COVID-19 pneumonia and moderate to severe hypoxic respiratory failure receiving high flow nasal therapy. *medRxiv* 2020. <https://doi.org/10.1101/2020.06.30.20143867>
  208. Merzon E., Tworowski D., Gorohovski A., Vinker S., Golan Cohen A., Green I. *et al.* Low plasma 25(OH) vitamin D3 level is associated with increased risk of COVID-19 infection: an Israeli population-based study. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.01.20144329>
  209. Fan X., Yin C., Wang J., Yang M., Ma H., Jin G. *et al.* Pre-diagnostic circulating concentrations of insulin-like growth factor-1 and risk of COVID-19 mortality: results from UK Biobank. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.09.20149369>
  210. Shi Z., Resurreccion W. K., Wang C.-H., Wei J., Na R., Zheng S. L. *et al.* Association of cancer with risk and mortality of COVID-19: results from the UK Biobank. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.10.20151076>
  211. Maucourant C., Filipovic I., Ponzetta A., Aleman S., Cornillet M., Hertwig L. *et al.* Natural killer cell activation related to clinical outcome of COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.07.20148478>
  212. Elmunzer B. J., Spitzer R. L., Foster L. D., Merchant A. A., Howard E. F., Patel V. A. *et al.* Digestive manifestations in patients hospitalized with COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.07.20143024>
  213. Alizadehsani R., Sani Z. A., Behjati M., Hussain S., Abedini N., Hasanzadeh F. *et al.* Risk factors prediction, clinical outcomes, and mortality of COVID-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.07.20148569>
  214. Xie Y., Chen S., Wang X., Li B., Zhang T., He X. *et al.* Early Diagnosis and Clinical Significance of acute cardiac injury —under the iceberg: a retrospective cohort study of 619 non-critically ill hospitalized COVID-19 pneumonia patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.06.20147256>
  215. Fox T. A., Troy-Barnes E., Kirkwood A. A., Chan W. Y., Day J. W., Chavda S. J. *et al.* Clinical outcomes and risk factors for severe COVID-19 infection in patients with haematological disorders receiving chemo- or immunotherapy. *Br J Haematol* 2020. doi: 10.1111/bjh.17027.
  216. Martinez-Resendez M. F., Castilleja-Leal F., Torres-Quintanilla A., Rojas-Martinez A., Garcia-Rivas G., Ortiz-Lopez R. *et al.* Initial experience in Mexico with convalescent plasma in COVID-19 patients with severe respiratory failure, a retrospective case series. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.14.20144469>
  217. Hoertel N., Rico M. S., Vernet R., Jannat A. S., Neuraz A., Blanco C. *et al.* observational study of haloperidol in hospitalized patients with Covid-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.15.20150490>
  218. McGrail D. E., Edwards D. COVID-19 case series at UnityPoint Health St. Luke's Hospital in Cedar Rapids, IA. *medRxiv* 2020; <https://doi.org/10.1101/2020.07.17.20156521>
  219. Pandolfi L., Fossali T., Frangipane V., Bozzini S., Morosini M., D'Amato M. *et al.* Broncho-alveolar inflammation in COVID-19 patients: a correlation with clinical outcome. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.17.20155978>
  220. Kurashima K., Kagiyama N., Ishiguro T., Takaku Y., Nakajima H., Shibata S. *et al.* IgG antibody seroconversion and the clinical progression of COVID-19 pneumonia: A retrospective, cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.16.20154088> (accessed 25 August 2020).
  221. Zhan Z., Yang X., Du H., Zhang C., Song Y., Ran X. *et al.* Early improvement of acute respiratory distress syndrome in patients with COVID-19: insights from the data of ICU patients in Chongqing, China. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.15.20154047>
  222. Omrani A. S., Almaslamani M. A., Daghfal J., Alattar R. A., Elgara M., Shaar S. H. *et al.* The first consecutive 5000 patients with Coronavirus disease 2019 from Qatar: a nation-wide cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.15.20154690>
  223. Gupta R., Agrawal R., Bukhari Z., Jabbar A., Wang D., Diks J. *et al.* Higher comorbidities and early death is characteristic of hospitalized African-American patients with COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.15.20154906>
  224. Hussein M. H., Toraih E. A., Attia A. S., Youssef M., Omar M., Burley N. *et al.* Asthma in COVID-19: an extra chain fitting around the neck? *medRxiv* 2020. <https://doi.org/10.1101/2020.07.13.20153130>

225. Bian H., Zheng Z.-H., Wei D., Zhang Z., Kang W.Z., Hao C.Q. et al. Meplazumab treats COVID-19 pneumonia: an open-labelled, concurrent controlled add-on clinical trial. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.21.20040691>
226. Eiros R., Barreiro-Perez M., Martin-Garcia A., Almeida J., Villacorta E., Perez-Pons A. et al. Pericarditis and myocarditis long after SARS-CoV-2 infection: a cross-sectional descriptive study in health-care workers. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.12.20151316>
227. Marcos M., Belhassen-Garcia M., Puente A. S., Sampedro-Gomez J., Azibeiro R., Dorado-Diaz P. I. et al. Development of a severity of disease score and classification model by machine learning for hospitalized COVID-19 patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.13.20150177>
228. Hoertel N., Rico M. S., Vernet R., Beeker N., Jannat A. S., Neuraz A. et al. Association between SSRI antidepressant use and reduced risk of intubation or death in hospitalized patients with Coronavirus disease 2019: a multicenter retrospective observational study. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.09.20143339>
229. Soares R. D., Mattos L. R., Raposo L. M. Risk factors for hospitalization and mortality due to COVID-19 in Espírito Santo State, Brazil. *Am J Trop Med* 2020; 103: 1184–90.
230. Zobairy H., Shamsoddin E., Rasouli M. A., Khodlan N. V., Moradi G., Zareie B. et al. Association of olfactory dysfunction with hospitalization for COVID-19: a multicenter study in Kurdistan. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.26.20158550>
231. Altamimi H., Alahmad Y., Khazal F., El Hassan M., Al Binali H., Arabi A. et al. The outcome of COVID-19 patients with acute myocardial infarction. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.21.20156349>
232. Thompson J. V., Meghani N., Powell B. M., Newell I., Craven R., Skilton G. et al. Patient characteristics and predictors of mortality in 470 adults admitted to a district general hospital in England with Covid-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.21.20153650>
233. Reiter T., Pajenda S., Wagner L., Gaggl M., Atamaniuk J., Holzer B. et al. Covid-19 serology in nephrology health care workers. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.21.20136218>
234. Motta J. K., Ogunnaike R. O., Shah R., Stroever S., Cedeno H. V., Thapa S. K. et al. Clinical outcomes with the use of prophylactic versus therapeutic anticoagulation in COVID-19. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.20.20147769>
235. Santos C., Rhee Y., Hollinger E., Olaitan O., Schadde E., Peev V. et al. Comparative incidence and outcomes of COVID-19 in kidney or kidney-pancreas transplant recipients versus kidney or kidney-pancreas waitlisted patients: a pilot study. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.20.20157990>
236. Schneeweiss M. C., Leonard S., Weckstein A., Schneeweiss S., Rassen J. Renin–angiotensin–aldosterone-system inhibitor use in patients with COVID-19 infection and prevention of serious events: a cohort study in commercially insured patients in the US. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.22.20159855>
237. Concha-Mejia A., Rincon-Sanchez R. A. CCOFEE-GI study: Colombian COVID19 first experience in gastroenterology. Characterization of digestive manifestations in patients diagnosed with COVID-19 at a highly complex institution in Bogota D.C., Colombia. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.24.20161604>
238. Izquierdo J. L., Almonacid C., Gonzalez Y., Del Rio-Bermudez C., Ancochea J., Cardenas R. et al. The impact of COVID-19 on patients with asthma. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.24.20161596>
239. Bernaola N., Mena R., Bernaola A., Lara A., Carballo C., Larranaga P. et al. Observational study of the efficiency of treatments in patients hospitalized with Covid-19 in Madrid. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.17.20155960>
240. Qi D., Yan X., Tang X., Peng J., Yu Q., Feng L. et al. Epidemiological and clinical features of 2019-nCoV acute respiratory disease cases in Chongqing municipality, China: a retrospective, descriptive, multiple-center study. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.01.20029397>
241. Peters E. J., Collard D., van Assen S., Beudel M., Bomers M. K., Buijs J. et al. Outcomes of persons with COVID-19 in hospitals with and without standard treatment with (hydroxy) chloroquine. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.14.20173369>
242. Ouyang J., Shan X., Wang X., Zhang X., Chen Y., Qi M. et al. Clinical characteristics of COVID-19 and the model for predicting the occurrence of critically ill patients: a retrospective cohort study. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.13.20173799>
243. Valenzuela O., Ibanez S. E., Poli M., Roessler P., Aylwin M., Roizen G. et al. First report of tocilizumab use in a cohort of Latin American patients hospitalized for severe COVID-19 pneumonia. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.12.20173104>
244. Monteiro A. C. C., Suri R., Emeruwa I. O., Stretch R. J., Lopez R. Y., Sherman A. et al. Obesity and smoking as risk factors for invasive mechanical ventilation in COVID-19: a retrospective, observational cohort study. *medRxiv* 2020; <https://doi.org/10.1101/2020.08.12.20173849>
245. Philipose Z., Smati N., Wong C. S. J., Aspey K., Mendall M. A. Obesity, old age and frailty are the true risk factors for COVID-19 mortality and not chronic disease or ethnicity in Croydon. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.12.20156257>
246. Weerahandi H., Hochman K. A., Simon E., Blaum C., Chodosh J., Duan E. et al. Post-discharge health status and symptoms in patients with severe COVID-19. *medRxiv* 2020; <https://doi.org/10.1101/2020.08.11.20172742>
247. Altibi A. M., Bhargava P., Liaqat H., Slota A. A., Sheth R., Al Jebbawi L. et al. Comparative clinical outcomes and mortality in prisoner and non-prisoner populations hospitalized with COVID-19: a cohort from Michigan. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.08.20170787>
248. Izzi-Engbeaya C., Distaso W., Amin A., Yang W., Idowu O., Kenkre J. S. et al. Severe COVID-19 and diabetes: a retrospective cohort study from three London teaching hospitals. *medRxiv* 2020; <https://doi.org/10.1101/2020.08.07.20160275>
249. Rizzo S., Chawla D., Zalocusky K., Keebler D., Chia J., Lindsay L. et al. Descriptive epidemiology of 16,780 hospitalized COVID-19 patients in the United States. *medRxiv* 2020; <https://doi.org/10.1101/2020.07.17.20156265>
250. Dashti H. T., Bates D., Fiskio J. M., Roche E. C., Mora S., Demler O. Clinical Characteristics and severity of COVID-19 disease in patients from Boston area hospitals. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.27.20163071>
251. Morshed M. S., Mosabbir A. A., Chowdhury P., Ashadullah S. M., Hossain M. S. Clinical manifestations of patients with

- Coronavirus disease 2019 (COVID-19) attending at hospitals in Bangladesh. *medRxiv* 2020. <https://doi.org/10.1101/2020.09.09.20191114>
252. Jun T., Nirenberg S., Kovatch P., Huang K. Sex-specificity of mortality risk factors among hospitalized COVID-19 patients in New York City: prospective cohort study. *medRxiv* 2020; <https://doi.org/10.1101/2020.07.29.20164640>
253. Higuchi T., Nishida T., Iwahashi H., Morimura O., Otani Y., Okauchi Y. *et al.* Early clinical factors predicting the development of critical disease in Japanese patients with COVID-19: a single-center retrospective, observational study. *medRxiv* 2020; <https://doi.org/10.1101/2020.07.29.20159442>
254. Zhou K., Sun Y., Li L., Zang Z., Wang J., Li J. *et al.* Eleven routine clinical features predict COVID-19 severity. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.28.20163022>
255. Salerno S., Zhao Z., Sankar S. P., Salvatore M., Gu T., Fritzsche L. G. *et al.* Understanding the patterns of repeated testing for COVID-19: Association with patient characteristics and outcomes. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.26.20162453>
256. Kumar A., Prasad G., Srivastav S., Gautam V. K., Sharma N. A retrospective study on efficacy and safety of Guduchi Ghan Vati for Covid-19 asymptomatic patients. *medRxiv* 2020. <https://doi.org/10.1101/2020.07.23.20160424>
257. Hao S.-R., Zhang S.-Y., Lian J.-S., Jin X., Ye C. Y., Cai H. *et al.* Liver enzyme elevation in coronavirus disease 2019: a multicenter, retrospective, cross-sectional study. *Am J Gastroenterol* 2020; <https://doi.org/10.14309/ajg.0000000000000717>
258. Iversen K., Bundgaard H., Hasselbalch R. B., Kristensen J. H., Nielsen P. B., Pries-Heje M. *et al.* Risk of COVID-19 in health-care workers in Denmark: an observational cohort study. *Lancet Infect Dis* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30589-2](https://doi.org/10.1016/S1473-3099(20)30589-2)
259. Hippisley-Cox J., Young D., Coupland C., Channon K. M., San Tan P., Harrison D. A. Risk of severe COVID-19 disease with ACE inhibitors and angiotensin receptor blockers: cohort study including 8.3 million people. *Heart* 2020; **106**: 1503–11.
260. Fillmore N. R., La J., Szalat R. E., Tuck D. P., Nguyen V., Yildirim C. *et al.* Prevalence and outcome of Covid-19 infection in cancer patients: a national VA study. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.21.20177923>
261. Rashid M., Wu J., Timmis A., Curzen N., Zaman A., Clarke S. *et al.* Clinical characteristics and outcomes of COVID-19 positive acute coronary syndrome patients: a multisource electronic healthcare records study from England. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.20.20175091>
262. Pan A., Khan O., Meeks J., Boom M., Masud F., Andrieni J. *et al.* Disparities in covid-19 hospitalizations and mortality among black and hispanic patients: cross-sectional analysis from the Greater Houston Metropolitan Area. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.19.20177956>
263. Alkurt G., Murt A., Aydin Z., Tatli O., Agaoglu N. B., Irven A. *et al.* seroprevalence of coronavirus disease 2019 (covid-19) among health care workers from three pandemic hospitals of Turkey. *medRxiv* 2020. <https://doi.org/10.1101/2020.08.19.20178095>
264. Zhao Z., Chen A., Hou W., Graham J. M., Li H., Richman P. S. *et al.* Prediction model and risk scores of ICU admission and mortality in COVID-19. *PLOS ONE* 2020; **15**: e0236618.
265. Holman N., Knighton P., Kar P., O'Keefe J., Curley M., Weaver A. *et al.* Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol* 2020; **8**: 823–33.
266. Qu J., Chang L. K., Tang X., Du Y., Yang X., Liu X. *et al.* Clinical characteristics of COVID-19 and its comparison with influenza pneumonia. *Acta Clin Belg* 2020; **75**: 348–56.
267. Chand S., Kapoor S., Orsi D., Fazzari M. J., Tanner T. G., Umeh G. C. *et al.* COVID-19-associated critical illness—report of the first 300 patients admitted to intensive care units at a new York City medical center. *J Intens Care Med* 2020; **35**: 963–70.

## Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**Figure S1** Map of countries where included studies were conducted. Six studies were performed in multiple countries and are not included here.

**Table S1** Study design, use of clinical diagnosis and stratification of smoking status by sex, age or socio-economic position.

**Table S2a** Studies reporting complete smoking status

**Table S2b** Studies reporting partially complete smoking status

**Table S2c** Studies reporting incomplete smoking status

**Table S3** Smoking prevalence in countries with included studies

**Figure S2** Supporting Information

**Figure S3** Supporting Information

**Figure S4** Supporting Information

**Figure S5** Supporting Information